
REFERENCES

- Abman, S.H.; Griebel, J.L.; Parker, D.K. (1994):**
Inhaled nitric oxide in children with severe hypoxaemic respiratory failure. *J. Pediatric.*, 124: 881-888.
- Adnots, D.; Reffestin, B.; Eddhibis (1995):**
NO in the lung. *Respir physiol.* 101-109.
- Allison, S.P.; Hinton, P.; Ghamberlain, M.J. (1998):**
Intravenous glucose tolerance insulin, and free fatty acid levels in burned patients. *Lancet*, 2: 1113.
- Alterra, M.; Bellet, J.; Scott, H. (1990):**
Preparedness for pediatric emergencies encountered in the practitioner's office. *Pediatrics*, 85: 710.
- Ambrus, J.L.; Ambrus, C.M. (1990):**
Changes in the fibrinolysin system in infantile and adult respiratory distress syndrome (ARDS), caused by trauma and/or septic shock in patients and in experimental animals. *J. Med.*, 21: 67.
- American Academy of Pediatrics (1998):**
141 Northwest Point Road, PO Box 927, Elk Grove Village, IL, 60007.
- Applebaum, D. (1985):**
Advanced prehospital care for pediatric emergencies. *Ann. Emerg. Med.*, 14: 656.
- Anaissie, E.J.; Darouiche, R.O.; Abi-Said, D. et al. (1996):**
Management of invasive candidal infections: Results of a prospective, randomized, multicenter study of fluconazole versus amphotericin B and review of the literature. *Clin. Infect. Dis.*, 23: 965-972.
- Anggard, E. (1994):**
Nitric oxide, mediator, murderer, medicine. *Lancet*, 343: 1199-1206.

- Aulick, I.H.; Wilmore, D.W. (1979):**
Increased peripheral amino acid release following burn injury. *Surgery*, 85: 560.
- Askanazi, J.; Carpentier, Y.A.; Michelson, C.B. et al. (1980):**
Muscle and plasma amino acids following injury: influence of intercurrent infection. *Ann. Surg.*, 792: 78.
- Astiz, M.E.; Rackow, E.C.; Flak, J.L. et al. (1987):**
Oxygen delivery and consumption in patients with hyperdynamic septic shock. *Crit Care Med.*, 15: 26-28.
- Astiz, M.E.; Rackow, E.C.; Kaufman, B. et al. (1988):**
Relationship of oxygen delivery and mixed venous oxygenation to lactic acidosis in patients with sepsis and acute myocardial infarction. *Crit Care Med.*, 16: 655-658.
- Astiz, M.E.; Rackow, E.C.; Schumer, W. et al. (1988):**
Early impairment of oxidative metabolism and energy production in severe sepsis. *Circ Shock*, 26: 311-320.
- Astiz, M.E.; Rackow, E.C.; Weil, M.H. (1986):**
Oxygen delivery and utilization during rapidly fatal septic shock in rats. *Circ Shock*, 20: 281-290.
- Astiz, M.E.; Tilly, E.; Rackow, E.C. et al. (1991):**
Peripheral vascular tone in sepsis. *Chest*, 99: 1072-1075.
- Avontuur, J.A.M.; Tutein, Nothenius, R.P.; van Bodegom, J.W. et al. (1998):**
Prolonged inhibition of nitric oxide synthesis in severe septic shock: A clinical study. *Crit Care Med.*, 26: 660-667.
- Bakker, J.; Coffernils, M.; Leon, M. et al. (1991):**
Blood lactate levels are superior to oxygen derived variables acting outcome in human septic shock. *Chest*, 99: 956-962.
- Banks, J.G.; Fouk, D.; Leaingham, I. et al. (1996):**
Liver function in septic shock. *J. Clin. Pathol.*, 26: 1082-1090.
- Barnes, P.J.; Belvis, M.E. (1993):**
Nitric oxide and lung disease. *Thorax.*, 48: 1034-1043.

- Barton, P.; Garcia, J.; Koutali, A. et al. (1996):**
Hemodynamic effects of IV milirnone lactate in pediatric patients with septic shock: A prospective, double-blinded, randomized, placebo-controlled interventional study. *Chest*, 109: 1302.
- Beca, J.; Butt, W. (1994):**
Extracorporeal membrane oxygenation for refractory septic shock in children. *Pediatrics*, 93: 726.
- Bedell, S.A.; Bush, B.T. (1986):**
Erythrocyte sedimentation rate from folklore to facts. *Am. J. Med.*, 78: 1001-1009.
- Bergstrom, J.; Fuerst, P.; Noree, L.O. et al. (1974):**
Intracellular free amino acid concentration in human muscle tissue. *J. appl. Physiol.*, 36: 693.
- Bessey, P.Q.; Brooks, D.C.; Black, P.R. et al. (1983):**
Epinephrine acutely mediales skeletal muscle insulin resistance. *Surgery*, 91: 172.
- Bessey, P.Q.; Watters, J.M.; Aoki, T.T. et al. (1985):**
Combined hormonal infusion simulates the metabolic response to injury. *Ann. Surg.*, 20: 264.
- Bihari, D.; Smithies, M.; Gimson, A. (1993):**
The effects of vasodilatation with prostacyclin on oxygen delivery and uptake in critically ill patients. *N. Engl. J. Med.*, 317: 397-402.
- Birke, G.; Carlson, L.A.; Von Euler, U.S. (1982):**
Lipid metabolism, catecholamine excretion, basal metabolic rate and water loss during treatment of burns with warm dry air. *Acta. Chir. Scand.*, 138: 321.
- Blomqvist, B.I.; Hammarqvist, F.; VonderDecken, A. et al. (1995):**
Glutamine and alpha-ketoglutarate prevent the decrease in muscle free glutamine concentration and influence protein synthesis eplacement. *Metabolism*, 11: 1215.

- Blumberg, R.M.; Cady, E.B.; Wigglesworth, J.S.; McKenzie, J.E.; Edwards, A.D. (1996):**
Relation between delayed impairment of cerebral energy metabolism and infarction following transient focal hypoxia ischaemia in the developing brain. Exp. Brain Res., In Press.
- Black, P.R.; Brook, D.C.; Bessy, P.Q. et al. (1982):**
Mechanism of insulin resistance following injury.
Ann. Surgery, 196-120.
- Bolotina, V.M.; Najibi, S.; Palacino, J.J.; Pagano, P.J.; Cohen, R.A. (1994):**
Nitric oxide directly activates calcium-dependent potassium channels in vascular smooth muscle.
Nature., 368: 850-853.
- Buntain, W.L.; Lynch, F.P.; Ramenofsky, M.L. (1987):**
Management of the actually injured child. Adv. Truma, 2: 42.
- Bonadio, W.; Wagner, V. (1988):**
Efficacy of TAC topical anesthetic for repair of pediatric lacerations. Am. J. Dis. Child., 142: 203.
- Bone, R. (1991):**
Sepsis, the sepsis syndrome, multi-organ failure: a plea for comparable definitions. Ann Intern. Med., 114:332-333.
- Bone, R.; Fischer, C.; Clemmer, T. et al. (1987):**
Methylprednisolone in severe sepsis study group: A controlled clinical trial of high dose methylprednisolone in the treatment of severe and septic shock. N. Engl. J. Med., 317: 653-658.
- Bone, R.; Fischer, C.; Clemmer, T. et al. (1987):**
Methylprednisolone in severe sepsis study group: Early methylprednisolone treatment for septic syndrome and the adult respiratory distress syndrome. Chest, 92: 1032-1036.
- Bone, R.; Fischer, C.; Clemmer, T.P. et al. (1989):**
Sepsis syndrome: A valid clinical entity. Crit Care Med., 17: 389-393.

- Brandtzaeg, P.; Hogasen, K.; Kierulf, P. et al. (1996):**
The excessive complement activation in fulminant meningococcal septicemia is predominantly caused by alternative pathway activation. *J. Infect. Dis.*, 173: 647.
- Bredius, R.G.; Derkx, B.H.; Fijen, C.A. et al. (1994):**
Fc gamma receptor IIa (CD32) polymorphism in fulminant meningococcal septic shock in children. *J. Infect. Dis.*, 170: 848.
- Bredt, D.S.; Snyder, S.H. (1989):**
Nitric oxide mediates glutamate-linked enhancement of cGMP levels in the cerebellum. *Proc. Natl. Acad. Sci. USA*, 86: 9030-9033.
- Brigham, K.L.; Meyrick, B. (1986):**
Endotoxin and lung injury. *Am. Rev. Respir. Dis.*
- Bryan, C.; Reynolds, K.; Brenner, E. (1983):**
Analysis of 1186 episodes of gram-negative bacteremia in non-university hospitals: the effects of antimicrobial therapy. *Rev. Infect. Dis.*, 5: 629-630.
- Caines, C.; Gill, M.V.; Cunha, B.A. (1997):**
Non-Clostridium difficile nosocomial diarrhea in the intensive care unit. *Heart Lung*, 26: 111.
- Camp, F.E.; Long, C.L.; Killin, P. (1984):**
Studies of glucose intolerance in septic injured patients. *J. Trauma*, 14: 375.
- Carpati, C.; Astiz, M.E.; Rackow, E.C. et al. (1992):**
Monophosphoryl lipid A attenuates the effects of endotoxic shock in pigs. *J. Lab. Clin. Med.*, 119: 346-353.
- Carroll, G.; Snyder, J. (1982):**
Hyperdynamic severe intravascular sepsis depends on fluid administration in cynomolgus monkey. *Am. J. Physiol.*, 243: R131-R141.
- Carroll, K.; Reimer, L. (1996):**
Microbiology and laboratory diagnosis of upper respiratory tract infections. *Clin. Infect. Dis.*, 23: 442-448.

- Carcillo, J.A.; Davis, A.I.; Zaritsky, A. (1991):**
Role of early fluid resuscitation in pediatric septic shock.
JAMA, 226: 1242.
- Carcillo, J.A.; Pollack, M.M.; Ruttiman, U.E. et al. (1989):**
Sequential physiologic interactions in pediatric cardiogenic and septic shock.
Crit. Care Med., 17: 12.
- Castillo, L.; Yu, Y.M.; Marchini, J.S. et al. (1994):**
Phenylalanine and tyrosine kinetics in critically ill children with sepsis. Pediatr. Res., 35: 580.
- Celemajor, D.S.; Cullen, S.; Ceanfield, J.E. (1993):**
Impairment of endothelium-dependent pulmonary artery relaxation in children with congenital heart disease and abnormal pulmonary haemodynamics. Circulation, 87: 440-446.
- Ceneviva, G.; Carcillo, J.A. (1996):**
Inotropes, vasopressors, and vasodilators in pediatric septic shock.
Pediatric Critical Care Colloquium.
- Chessare, J. (1986):**
Utilization of emergency services among patients of a pediatric group practice. Pediatr Emerg. Care., 2:27.
- Clancy, R.; Leszczynska, J.; Amin, A.; Levartovsky, D.; Abramson, S.B. (1995):**
Nitric oxide stimulates ADP ribosylation of actin in association with the inhibition of actin polymerization in human neutrophils. J. Leukoc. Biol., 38: 196-202.
- Corrigan, J.J. (1988):**
Disseminated intravascular coagulopathy. Pediatrics, 64: 37.
- Corrigan, J.J. (1974):**
Thrombocytopenia: A laboratory sign of septicemia in infants and children.
J. Pediatr., 85: 219.

- Cristine, C.; Corriveau, Partica, J. (1998):**
Effect of nitric oxide on chemotaxis and Endotoxin-induced interleukin-8 production in human neutrophils. JID, 177-116-126.
- Cronenwett, J.; Lindenauer, M. (1992):**
Hemodynamic effects of cecal ligation sepsis in dogs. J. Surg. Res., 33: 324-331.
- Cunha, R.E.; Schaer, G.L.; Parker, M.M. et al. (1993):**
The coronary circulation in human septic shock. Circulation, 73: 637-644.
- Dahn, M.S.; Lange, P.; Kubdel, K. et al. (1987):**
Splanchnic and total body oxygen consumption differences in septic and injured patients. Surgery, 101: 69-80.
- Dammas, P.; Reuter, A.; Gysen, P. et al. (1990):**
Tumor necrosis factor and interleukin-1 serum levels during severe sepsis in humans. Crit Care Med., 17: 975-978.
- Danner, R.L.; Elin, R.J.; Hosseini, J. et al. (1991):**
Endotoxemia in human septic shock. Chest, 99: 169-175.
- Danner, R.L.; Suffredini, A.F.; Watanason, C. et al. (1989):**
Microbial toxins: role in the pathogenesis of septic shock and multiple failure. In Bihari, D Cerra F (eds): New Horizons: Multiple Organ failure. Fullerton, CA, Society of Critical Care Medicine, PP. 151-191.
- Danner, R.L.; Cobb, J.P.; Van Dervort, A.L. (1995):**
Nitric oxide as a mediator of hypotension and inflammation in sepsis. In: Fink MP, Payen D, eds. Role of Nitric Oxide in Sepsis and ARDS. New York, NY: Springer-Verlag NY Inc.; 24: 155-180.
- Derkn, H.H.; Kuijper, E.J.; Fijen, C.A. et al. (1995):**
Inherited complement deficiency in children surviving fulminant meningococcal septic shock. Eur J. Pediatr., 154: 735.

- Dietch, E.; Specian, R.; Berg, R. (1991):**
Endotoxin-induced bacterial translocation and mucosal permeability: Role of xanthine oxidase complement activation and macrophage products. *Crit Care Med.*, 19: 758-791.
- Dinerman, J.L.; Lowenstein, C.J.; Snyder, S.H. (1993):**
Molecular mechanisms of nitric oxide regulation. Potential relevance to cardiovascular disease. *Circ. Res.*, 73: 217-222.
- Dorio, V.; Whalen, C.; Naldi, M. et al. (1989):**
Contribution of peripheral blood pooling to central hemodynamic disturbances during endotoxin insult in intact dogs. *Crit Care Med.*, 17: 1314.
- Doughty, L.A.; Carcillo, J.A.; Frye, R. (1997):**
Decreased P450 metabolism in pediatric sepsis and multiple organ failure. Society for Critical Care Medicine Symposium, San Diego, CA.
- Doughty, L.A.; Kaplan, S.S.; Carcillo, J.A. (1997):**
Inflammation and outcome in pediatric sepsis induced multiple organ failure. Society for Critical Care Medicine Symposium, San Diego, CA.
- Doughty, L.A.; Kaplan, S.S.; Carcillo, J.A. (1996):**
Inflammatory cytokine and nitric oxide responses in pediatric sepsis and organ failure. *Crit Care Med.*, 24: 1137.
- Duff, J.H.; Groves, A.C.; McLean, A.P. et al. (1969):**
Defective oxygen consumption in septic shock. *Surg. Gynecol. Obstet.*, 128: 1051-1060.
- Dupont, H.L.; Spink, W.W. (1989):**
Infections due to gram negative organisms: An analysis of 860 patients with bacteremia at the University of Minnesota Medical Center, 1958-1966. *Medicine*, 48: 507.
- Downey, R.S.; Monafio, W.W.; Karl, I.E. et al. (1996):**
Protein dynamics in skeletal muscle after trauma: local and systemic effects. *Surgery*, 99: 265.

- Edwards, A.D. (1995):**
The pharmacology of inhaled nitric oxide. Arch Dis. Child., 72: E127-E130.
- Edward, S.A.D.; Mupanemunda, R.H. (1990):**
Nitric oxide physiology, pathophysiology and potential clinical applications. Recent advance in pediatrics, 13: 119-136.
- Edwards, D.; Brown, c.; Nightengale, P. et al. (1989):**
Use of survivors cardiorespiratory varues as therapeutic goals in septic shock. Crit Care Med., 17: 1098-1102.
- Ellrodt, A.; Riedinger, M.; Kimichi, A. et al. (1985):**
Left ventricular performance in septic shock: reversible segmental and global abnormalities. Am. Heart J., 110: 402-409.
- El-Naggar, M. (1997):**
Life and death.
Ped. Crit. Care., 2-12.
- Falk, J.L.; Rackow, E.C.; Leavy, J. et al. (1985):**
Delayed lactate clearance in patients surviving circulatory shock. Acute Care, 11: 212-215.
- Feltes, T.F.; Pignatelli, R.; Kleinert, S. et al. (1994):**
Quantitated left entricular systolic mechanics in children with septic shock utilizing noninvasive wall-stress analysis. Crit Care Med., 22: 1647.
- Fiddian-Green, R. (1990):**
Studies in splanchnic ischemia and multiple organ failure. In Marston A, Bulkley G.; Fiddian-Geene RG, et al. (eds): Splanchnic Ischemia and Multiple Organ Failure. London, CV Mosby, PP. 350-363.
- Fifield, G.; Magnuson, C.; Carr, W.P. (1984):**
Pediatric emergency in a metropolitan area. J. Emerg. Med., 1:495.
- Fincher, R.M.E.; Page, M.I. (1986):**
Clinical significance of extreme values of the erythrocyte sedimentation rate. Arch Intern. Med., 146: 1581-1583.

- Fink, M.P. (1991):**
Gastrointestinal mucosal injury in experimental models of shock, trauma and sepsis. *Crit Care Med.*, 19: 627-641.
- Fink, M.P.; Cohn, S.M.; Lee, P.C. et al. (1989):**
Effect of lipopolysaccharide on intestinal intramucosal hydrogen in concentration in pigs: evidence of gut ischemia in a normodynamic model of septic shock. *Crit Care Med.*, 17: 641-646.
- Finkel, M.S.; Oddis, C.V.; Jacob, T.D.; Watkins, S.C.; Hattler, B.G.; Simmons, R.L. (1992):**
Negative inotropic effects of cytokines on the heart mediated by nitric oxide. *Science*. 257: 387-389.
- Fijen, C.A.; Derkx, B.H.; Kuijper, E.J. et al. (1995):**
Fulminant meningococcal septic shock in a boy with combined inherited properdin and protein C deficiency. *Clin. Exp. Immunol.*, 102: 290.
- Fijnvandraat, K.; Derkx, B.; Peters, M. et al. (1995):**
Coagulation activation and tissue necrosis in meningococcal septic shock: Severely reduced protein C levels predict a higher mortality. *Thromb Haemost.*, 73: 15.
- Fleck, A.; Raines, G.; Hawker, F. et al. (1985):**
Increased vascular permeability: a major cause of hypoalbuminemia in disease and injury. *Lancet*, 1:781-783.
- Forsterman, U.; Schmidt, H.H.H.W.; Pollock, H.S. (1991):**
Isoforms of nitric oxide synthase: Characterization and purification from different cell types. *Biochem. Pharmacol.*, 42: 1849-1856.
- Fourrier, F.; Chopin, C.; Huart, J.J. et al. (1993):**
Double blind, placebo-controlled trial of antithrombin III concentrates in septic shock with disseminated intravascular coagulation. *Chest*, 104: 882.
- Franson, T.; Hierholzer, W.; LaBrecque, D. (1985):**
Frequency and characteristics of hyperbilirubinemia associated with bacteremia. *Rev. Infect. Dis.*, 7: 1-9.

- Fry, D.E.; Kalein, C.R.; Giammara, B.L. et al. (1981):**
Alterations of oxygen metabolism in experimental bacteremia. *Adv. Shock Res.*, 6: 45.
- Fry, D.E.; Pearlsteen, L.; Fulton, R.L. et al. (1980):**
Multiple system organ failure: the role of uncontrolled infection. *Surgery*, 115: 136-140.
- Fry, D.E.; Silver, B.b.; Rink, R.D. et al. (1979):**
Hepatic cellular hypoxia in the murine peritonitis. *Surgery*, 85: 652-666.
- Furchgott, R.F.; Zawadzki, J.V. (1990):**
The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature*, 288: 373-376.
- Furchgott, R.F.; Vanhoutte, P.M. (1989):**
Endothelium derived relaxing and contractor factors. *FASEB, J.*, 3: 2007-2018.
- Garber, A.J.; Karl, I.E.; Kipnis, D.M. (1976):**
Alanine and glutamine synthesis and release from skeletal muscle. I. Glycolysis and amino acid release. *J. Biol. Chem.*, 251: 826.
- Garboursy, J.; Woodman, r.C.; Granger, D.N.; Reinhardt, P.; Kubes, P. (1993):**
Nitric oxide prevents leukocyte adherence: role of superoxide. *Am. J. Physiol.*, 265: H862-7.
- Gardlund, B.; Sjolín, J.; Nilsson, A. et al. (1993):**
Plasmapheresis in treatment of primary septic shock in humans. *Scand J. Infect. Dis.*, 25: 757.
- Gaston, B.; Drazen, J.M.; Loscalzo, J.; Stamler, J.S. (1994):**
The biology of nitrogen oxides in the airways. *Am. J. Respir. Crit Care Med.*, 149: 538-551.
- Gill, C.W.; Bush, W.S.; Burleigh, W.M. et al. (1981):**
An evaluation of C-reactive protein assay using a rate immunonephelometric procedure. *Am. J. Clin. Pathol.*, 75: 50-55.

Gray, G.; Schott, C.; Julou-Schaeffer, G. et al. (1991):

The effect of inhibitors of the L-arginine/nitric oxide pathway on endotoxin-induced loss of vascular responsiveness in anaesthetized rats. *Br. J. Pharmacol.*, 103: 1218-1244.

Greene, J.N. (1996):

The microbiology of colonization, including techniques for assessing and measuring colonization.

Infect. Control Hosp. Epidemiol., 17: 114-118.

Green, L.C.; Wagner, D.A.; Glogowski, J. et al. (1982):

Nitrate, nitrite, and ^{15}N nitrate in biologic. *Biochem.*, 126: 131-8.

Groeneveld, J.; Bronsveld, W.; Thijis, L. (1986):

Hemodynamic determinants of mortality in human septic shock, *Surgery*, 99: 140-159.

Guerrant, R.L.; Shields, D.S.; Thorson, S.M. et al. (1985):

Evaluation and diagnosis of acute diarrhea. *Am. J. Med.*, 78 (Suppl. 6b): 91-99.

Hagisawa, M.; Ogins, Goto, R. et al. (1983):

Endotoxin clearance by exchange blood transfusion in septic shock neonates. *Acta Paediatr. Scand*, 72: 87.

Haller, J.A.; Shorter, N.; Miller, D. et al. (1993):

Organization and function of a regional pediatric trauma center: Does a system of management improve outcome? *Trauma*, 23: 691.

Halperin, R.; Meyers, A.; Alpert, J. (1989):

Utilization of pediatric emergency services. *Pediatr. Clin. North, Am.* 26: 747/

Haque, K.M.; Remo, C.; Bahakim, H. et al. (1995):

Comparison of two types of intravenous immunoglobulins in the treatment of neonatal sepsis. *Clin. Exp. Immunol.*, 101: 328.

Hasselgren, P.O.; James, J.H.; Benson, D.W. et al. (1989):

Total and myofibrillar protein breakdown in different types of rat skeletal muscle: effects of sepsis and regulation by insulin. *Metabolism*, 38: 634.

- Hector, R.; Wong, Josphe, A.; Carcill, G.; Sandra, S. (1996):**
Nitric oxide production in critically ill patients. A review of disease in childhood, 74: 482-489.
- Hegesh, E.; Shiloah, J. (1993):**
Blood nitrates and infantile methemoglobinemia. Clin. Chin. Acta., 125: 107-121.
- Hermeck, A.; Thal, A. (1989):**
Mechanisms for high circulatory requirements in sepsis and septic shock. Ann. Surg., 179: 677-695.
- Heughan, C.; Niinikoski, J.; Hunt, T. (1982):**
Effect of excessive saline solution on tissue oxygen transport. Surg. Gynecol. Obstet., 135: 257.
- Hibbs, Jr. J.B. (1991):**
Synthesis of nitric oxide from L-arginine: a recently discovered pathway induced by cytokines with antitumor and antimicrobial activities. Res. Immunol., 142: 565-569.
- Hibbs, J.B.; Westenfelder, C.; Taintor, R. et al. (1992):**
Evidence for cytokine-induced nitric oxide synthesis from L-arginine in patients receiving interleukin-2 therapy. J. Clin. Invest., 89: 867-17.
- Hiker, T.L. (1988):**
Nonemergency visits to a pediatric emergency department. Journal of the American College of Emergency Physicians. 7:3.
- Hoffman, M.; Greenfield, L.; Sugerman, H. et al. (1992):**
Unsuspected right ventricular dysfunction in shock and sepsis. Ann. Surg., 198: 307-318.
- Holbrook, P.R.; Taylor, G.; Pollack, M.M. et al. (1983):**
Adult respiratory distress syndrome in children. Pediatr Clin. North Am., 27: 677.
- Hollenberg, S.M.; Cunnion, R.E.; Zimmerberg, J. (1993):**
Nitric oxide synthase inhibition reverses arteriolar hyporesponsiveness to catecholamines in septic rats. Am. J. Physiol., 263: H 660-H 663.

- Holmes, M.J.; Reyes, H.M. (1994):**
A critical review of urban pediatric trauma. *J. Trauma*, 24: 253.
- Horwitz, J.R.; Elerian, L.F.; Sparks, J.W. et al. (1995):**
Use of extracorporeal membrane oxygenation in the septic neonate. *J. Pediatr. Surg.*, 30: 813.
- Ignarro, L.J. (1989):**
Endothelium-derived nitric oxide: actions and properties. *FASEB, J.* 3: 31-36.
- Jacobs, R.F.; Sowell, M.K.; Moss, M.M. et al. (1990):**
Septic shock in children: Bacteria etiologies and temporal relationships. *Pediatr. Infect. Dis. J.*, 9: 196.
- Jenkins, R.D.; Fenn, J.P.; Matsen, J.M. (1986):**
Review of urine microscopy for bacteriuria. *JAMA*, 255: 397-3403.
- Jia, L.; Bonaventura, C.; Bonaventura, J.; Stamler, J. (1993):**
S-nitrosohaemoglobin: a dynamic activity. *Med*, 329: 2002-2012.
- Johnson, D.J.; Jiang, Z.M.; Colpoys, M. et al. (1986):**
Branched chain amino acid uptake and muscle free amino acid concentrations predict postoperative muscle nitrogen balance. *Am. Surg.*, 204: 513.
- Joseph, A.; Carillo, Robert, E. (1997):**
Septic shock.
Crit care Clinics, 13: 553-572.
- Kalter, E.S.; Carlson, R.W.; Thijs, L. et al. (1982):**
Effects of methylprednisolone on hemodynamics, arteriovenous oxygen differences. *Crit Care Med.*, 10: 662-666.
- Kaufman, B.S.; Rackow, E.C.; Falk, J.L. (1984):**
The relationship between oxygen delivery and consumption during fluid resuscitation of hypovolemic and septic shock. *Chest*, 8: 336-349.

- Kilbourn, R.G.; Jubran, A.; Gross, S.S. et al. (1990):**
Reversal of endotoxin-mediated shock by NG-methyl-L-arginine, an inhibitor of nitric oxide synthesis. *Biochem. Biophys. Res. Commun.* 172: 1132-1138.
- Kimichi, A.; Ellrodt, G. Berman, P. et al. (1993):**
Right ventricular performance in septic shock. A Combined radionucleotide hemodynamic study. *J. Am. Coll. Cardiol.*, 4: 945-951.
- Kinney, J.M. (1976):**
Surgical hyper-metabolism and nitrogen metabolism. In: Wilkinson, Cuthbertson D (eds). *Metabolism and the Response to Injury*. Tunbridge Wells, Pitman Medical, 237.
- Kinney, J.M. (1987):**
Injury. In: Richards JR, Kinney JM (eds). *Nutritional Aspects of Care in the Critically Ill*. Edinburgh, Churchill Livingstone, 95.
- Kinney, J.M.; Elwyn, D.H. (1993):**
Protein metabolism and injury. *Ann. Res. Nutr.*, 3: 133.
- Kinsella, J.P.; Neish, S.R.; Ivy, D.D.; Shaffer, E.; Abman, S.H. (1993):**
Clinical responses to prolonged treatment of persistent pulmonary hypertension of the newborn with low doses of inhaled nitric oxide. *J. Pediatr.*, 123: 103-108.
- Klein, B.S.; Perloff, W.H.; Maki, D.G. (1989):**
Reduction of nosocomial infection during pediatric intensive care by protective isolation. *N. Engl. J. Med.*, 320: 171.
- Klem, K.; Fuchs, G.J.; Kulapongs, P. et al. (1988):**
Endotoxemia in protein-energy malnutrition. *J. Pediatr. Gastroenterol. Nutr.*, 7: 225.
- Klemm, P.; Thiemermann, C.; Winklmaier, G.; Martorana, P.A.; Henning, R. (1995):**
Effects of nitric oxide synthase inhibition combined with nitric oxide inhalation in a porcine model of endotoxin shock. *Br. J. Pharmacol.*, 114: 363-368.

- Kooy, N.W.; Royall, J.A.; Ye, Y.Z. et al. (1995):**
Evidence from in vivo peroxynitrite production in human acute lung injury. *Am. J. Respir. Crit Care Med.*, 151: 1250.
- Kornelisse, R.F.; Hazelzet, J.A.; Savelkoul, H.F. et al. (1996):**
The relationship between plasminogen activator inhibitor-1 and proinflammatory and counterinflammatory mediators in children with meningococcal septic shock. *J. Infect. Dis.*, 173: 1148.
- Krafte-Jacobs, B.; Carver, J.; Wilkinson, J.D. (1995):**
Comparison of gastric intramuscular H and standard perfusional measurements in pediatric septic shock. *Chest*, 1008: 141.
- Kreger, B.E.; craven, D.E.; McCabe, W.R. (1980):**
Gram -negative bacteremia III: reassessment of etiology epidemiology in 612 patients. *Am. J. Med.*, 68: 332-343.
- Kupman, N.; Inkelis, S.H.; Saladino, R. (1994):**
The role of heparin in the protection of extremity and digit necrosis in meningococcal purpura fulminans. *Ped. Infect.*, 13: 867.
- Landmann, R.; Zimmerli, W.; Sansano, S. et al. (1995):**
Increased circulating solution D14 is associated with high mortality in gram negative septic shock. *J. Infect. Dis.* 155-169.
- Leclerc, F.; Hazelzet, J.; Jude, B. et al. (1992):**
Protein C and S deficiency in severe infectious purpura of children: A collaborative study of 40 cases. *Intensive Care Med.*, 18: 202.
- Lief, Nelin, eorge, M.; Hoffman (1998):**
The use of inhaled nitric oxide in a wide variety of clinical problems: *Ped. Clinic. Of North Americ.* Volume 45: 531-548.
- Liu, S.F.; Adcock, I.M.; Old, R.W.; Barnes, P.J.; Evans, T.W. (1996):**
Differential regulation of the constitutive and inducible nitric oxide synthase mRNA by lipopolysaccharide treatment in vivo in the rat. *Crit Care Med.*, 24: 1219-25.

- Long, C.L.; Kinney, J.M.; Geiger, J.W. (1986):**
Nonsuppressibility of gluconeogenesis in septic patients. *Metabolism*, 25: 193.
- Lorente, J.A.; Landin, L.; DePablo, R. et al. (1993):**
L-arginine pathway in the sepsis syndrome. *Crit Care Med.*, 21: 1287-1295.
- Losek, J.; Walsh-Ketty, C.; Glaeser, P. (1996):**
Pediatric emergency departments. *Pediatr. Emerg. Care*, 2: 215.
- Lucking, S.E.; Williams, T.E.; Chaden, F.C. et al. (1990):**
Dependence of oxygen consumption on oxygen delivery in children with hyperdynamic septic shock and low oxygen extraction. *Crit. Care Med.*, 18: 1316.
- Lundberg, J.O.N.; Hellstrom, P.M.; Lundberg, J.M.; Alving, K. (1994):**
Greatly increased luminal nitric oxide in ulcerative colitis. *Lancet*, 344: 1673-1674.
- Lunn, R.J. (1995):**
Inhaled nitric oxide therapy. *Mayo Clin Proc.*, 70: 247-255.
- McCarrhy, P.I.; Sharpe, M.R.; Spiesel, S.Z. et al. (1982):**
Observation scales to identify serious illness in children. *Pediatrics*, 70: 802.
- MacNaul, R.L.; Huthehinson, N.I. (1993):**
Differentiate in expression of iNOS mRNA in human aortic smooth muscle cells and endothelial cells under normal and inflammatory conditions. *Biochem. Biophys. Res. Commun.*, 196: 1330-1334.
- Marks, K.A.; Mallard, C.; Roberts, I.; Williams, C.; Gluckman, P.; Edwards, A.D. (1996):**
Nitric oxide synthase inhibition attenuates delayed vasoconstriction and increases injury following cerebral ischaemia in fetal sheep. *Pediatr. Res.*
- McCabe, R.E.; Remington, J.S. (1994):**
C-reactive protein in patients with bacteremia. *J. Clin. Microbiol.*, 20: 317-319.

- McCune, S.; Short, B.L.; Miller, M.K. et al. (1990):**
Extracorporeal membrane oxygenation therapy in neonates with septic shock. *J. Pediatr. Surg.*, 25: 479.
- McManus, M.L.; Churchwell, K.B. (1993):**
Coagulopathy as a predictor of outcome in meningococcal sepsis and the systemic inflammatory response syndrome with purpura. *Crit Care Med.*, 21: 706.
- McMillan, M.; Chernow, B.; Roth, B. (1983):**
Hepatic α -adrenergic receptor alteration in a rat model of chronic sepsis. *Circ. Shock*, 19: 185-193.
- Meadows, D.; Edwards, J.D.; Wilkins, R.G. et al. (1988):**
Reversal of intractable septic shock with norepinephrine therapy. *Crit Care Med.*, 16: 663.
- Meher, C.M.; Rackow, E.C.; Astiz, M.E. et al. (1991):**
Unaccounted anion in metabolic acidosis during septic shock in humans. *Crit Care Med.*, 19: 705-711.
- Mercier, J.C.; Beaufile, F.; Hartmann, J.F. et al. (1988):**
Hemodynamic patterns of meningococcal shock in children. *Crit Care Med.*, 16: 27.
- Meyer, D.M.; Jessen, M.E. (1995):**
Results of extracorporeal membrane oxygenation in neonates with sepsis: The extracorporeal life Support Organization experience. *J. Thorac. Cardiovasc. Surg.*, 109: 419.
- Meyer, D.M.; Jessen, M.E.; Eberhart, R.C. (1995):**
Neonatal extracorporeal membrane oxygenation complicated by sepsis. Extracorporeal life support Organization. *Ann. Thorac Surg.*, 59: 795.
- Midha, N.K.; Stratton, C.W. (1998):**
Microbiology and laboratory diagnosis of meningitis. *Clin. Infect. Disease*, 980-982.
- Minard, G.; Kudsk, K.A. (1994):**
Effect of feeding on the incidence of septic complications in critically ill patients. *Semin Respir. Infect.*, 9: 228.

- Mink, R.B.; Pollack, M.M. (1992):**
Effect of blood transfusion on oxygen consumption in pediatric.
- Moncada, S.; Palmer, R.M.J.; Higgs, E.A. (1991):**
Nitric oxide: physiology, pathology, and pharmacology.
Pharmacol. Rev., 43: 109-42.
- Moncada, S.; Higgs, A. (1993):**
The L-arginine-nitric oxide pathway. N. Engl. Med., 329:
2022-12.
- Moran, J.L.; O'Fathartaigh, M.S.; Peisach, A.R. et al. (1993):**
Epinephrine as an inotropic agent in septic shock: A dose
profile analysis. Crit Care Med., 21: 70.
- Motocarda, S.; Palmer, R.M.I.; Higgs, E.N. (1991):**
Nitric oxid physiology, Pathophysiology, and pharmacology.
Pharmacol. Rev.
- Mupanemunda, R.H.; Edwards, A.D. (1995):**
Treatment of newborn infants with inhaled nitric oxide. Arch.
Dis. Child., 72: F131-F134.
- Mupanmunda, R.H. (1995):**
Nitric oxide in the perinatal period. Contemp Rev. Obstet.
Gynecol., 7: 210-214.
- Myers, P.R.; Zhong, Q.; Jones, J.J.; Tanner, M.A.; Adams, H.R.;
Parker, J.L. (1995):**
Release of EDRF and NO in ex vivo perfused aorta: inhibition
by in vivo E. coli endotoxemia. Am. J. Physiol., 268: H955-61.
- Nasrawy, S.A.; Rackow, E.C.; Astiz, M.E. et al. (1989):**
Ionotrope response to digoxin and dopamine in patients with
severe sepsis. Cardiac Failure and systemic hypoperfusion.
Chest, 95: 612-615.
- Nathan, C.F.; Stuphr, D.J. (1990):**
Does endothelium derived nitric oxide have a role in cytokine-
induced hypotension?
J. Nat. Cancer Inst., 726-728.

- Nathan, C. (1992):**
Nitric oxide as a secretory product of mammalian cells. *FASEB, J.*, 6: 3051-64.
- Natanson, C.; Eichenholz, P.; Danner, R. et al. (1989):**
Endotoxin and tumor necrosis factor challenges in dogs simulate the cardiovascular profile of human septic shock. *J. Exp. Med.*, 169: 823-832.
- Nelson, D.; Samsel, R.; Wood, L.D.H. et al. (1988):**
Pathological supply dependence of systemic and intestinal O₂ uptake during endotoxemia. *J. Appl. Physiol.*, 64: 2410-2419.
- Nishijima, H.; Weil, M.H.; Shubin, H. et al. (1983):**
Hemodynamic and metabolic studies on shock associated with gram-negative bacteremia. *Medicine* 52: 287-294.
- Nguyen, P.; Reynaud, J.; Pouzol, P. et al. (1994):**
Varicella and thrombotic complications associated with transient protein C and protein S deficiencies in children. *Eur. J. Pediatr.*, 153: 646.
- Norrby-Teglund, A.; Pauksens, K.; Norgren, M. et al. (1995):**
Correlation between serum TNF alpha and IL-6 levels and severity of group A streptococcal infections. *Scand J. Infect. Dis.*, 27: 125.
- Norris, M.K.; Steinhorn, D.M. (1994):**
Nutritional management during critical illness in infants and children. *AACN Clin. Issues in Crit Care Nurs.*, 4: 485.
- Ochoa, J.; Udekwa, A.; Billiar, T. et al. (1991):**
Nitrogen oxide levels in patients after trauma and during sepsis. *Ann. Surg.*, 214: 621-626.
- Ochoa, J.B.; Curti, B.; Peitzman, A.B. et al. (1992):**
Increased circulating nitrogen oxides after human tumor immunotherapy: correlation with toxic hemodynamic changes. *J. Natl Cancer Inst.*, 84: 864-867.
- O'Krent, D.; Abraham, E.; Winston, D. (1987):**
Cardiorespiratory patterns in viral septicemia. *Am. J. Med.*, 83: 683-686.

- Palmer, B.M.J.; Ferrige, A.G.; Moncada, S. (1997):**
Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature.*, 327: 524-526.
- Palmer, B.M.J.; Ress, D.D.; Ashton, D.S.; Moncada, S. (1988):**
L-arginine is the physiological precursor for the formation of nitric oxide in endothelium-dependent relaxation. *Biochem. Biophys. Res. Commun.*, 153: 1251-1256.
- Pang, D.; Wilberger, J.E. (1992):**
Spinal cord injury without radiographic abnormalities in children. *J. Neurosurg.*, 57: 114.
- Parker, M.; McCarthy, K.; Ognibene, F. et al. (1990):**
Right ventricular dysfunction and dilation similar to left ventricular changes characterize the cardiac depression of septic shock in humans. *Chest*, 97: 126-131.
- Parker, M.M.; Ognibene, F.P.; Parriloo, J.E. (1995):**
Peak systolic pressure/ end-systolic volume ratio, a load-independent measure of ventricular function, is reversibly decreased in human septic shock. *Crit Care Med.*, 23: 1791.
- Parker, M.M.; Shelhamer, J.H.; Natanson, C. et al. (1987):**
Serial cardiovascular variables in survivors and nonsurvivors of human septic shock: Heart rate as an early predictor of prognosis. *Crit Care Med.*, 15: 923.
- Parker, M.M.; Shelhamer, J.; Natanson, C. et al. (1987):**
Serial cardiovascular variables in survivors and nonsurvivors of human septic shock: heart rate, an early predictor of prognosis. *Crit Care Med.*, 15: 923-929.
- Parrella, M.A.; Yoshizumi, M.; Fen, Z. et al. (1994):**
Transferring growth factor but not dexamethasone, down-regulates nitric oxide synthase after its induction by interleukin-1 β in rat smooth muscle cells. *Chem.*, 269: 14595-600.
- Peitzman, A.; Udekwa, A.; Ochoa, J. et al. (1991):**
Bacterial translocation in trauma patients. *J. Trauma*, 31: 1083-1087.

- Peltola, H.O. (1982):**
C-reactive protein for rapid monitoring of infections of the central nervous system. *Lancet*, 1: 980-982.
- Perren Cobb, Robert, L.; Danner, (1996):**
Nitric oxide and septic shock.
JAMA- 275: 43-48.
- Persico, M.; Barker, G.A.; Mitchel, D.P. (1990):**
Purulent otitis media: A silent source of sepsis in the pediatric intensive care unit. *Otolaryngol Head Neck Surg.*, 93: 330.
- Petros, A.; Bennett, D.; Valance, P. (1991):**
Effect of nitric oxide synthase inhibitors on hypotension in patients with septic shock. *Lancet*, 338: 1556-1558.
- Petros, A.; Lamb, G.; Leone, A. et al. (1995):**
Effect of a nitric oxide synthase inhibitor in humans with septic shock. *Cardiovasc. Res.*, 28: 34-39.
- Piccioli, A.; Chini, G.; Mannelli, M. et al. (1994):**
Bilateral massive adrenal hemorrhage due to sepsis: Report of two cases. *J. Endocrinol Invest.*, 17: 821.
- Pollock, M.; Alexander, S.; Clarke, N. et al. (1987):**
Evaluation of Oregon pediatric critical care outcome by hospital resource level. Presented at the National Conference on Pediatric Trauma, Boston.
- Pollock, M.M.; Fields, A.I.; Ruttiman, U.E. (1985):**
Distribution of cardiopulmonary variables in pediatric survivors and nonsurvivors of septic shock. *Crit Care Med.*, 13: 454.
- Pollock, M.M.; Fields, A.I.; Ruttiman, U.E. (198):**
Sequential cardiopulmonary variables of infants and children in septic shock. *Crit Care Med.*, 12: 554.
- Porte, D.Jr.; Robertson, R.P. (1983):**
Control of insulin secretion by catecholamines, stress, and the sympathetic nervous system. *Ped. Proc.*, 32: 1792.

- Powars, D.; Larsen, R.; Johnson, J. et al. (1993):**
Epidemic meningococemia and purpura fulminans with induced protein C deficiency. *Clin. Infect. Dis.*, 17: 254.
- Pralong, G.; Calandra, T.; Glauser, M.T. et al. (1989):**
Plasminogen activator inhibitor 1: A new pronostic marker in septic shock. *Thromb Haemost.*, 61: 459.
- Proulx, F.; Fayon, M.; Farrell, C.A. et al. (1996):**
Epidemiology of sepsis and multiple organ dysfunction syndrome in children. *Chest.*, 109: 1033.
- Preiser, J.C.; Zhang, H.; Wachel, D.; Boeynaems, J.M.; Buurman, W.; Vincent, J.L. (1994):**
Is endotoxin-induced hypotension related to nitric oxide formation.
Eur. Surg. Res., 26: 10-18.
- Rackow, E.C.; Kaufman, B.S.; Falk, J.L. et al. (1987):**
Hemodynamic response to fluid repletion in patients with septic shock: Evidence for early depression of cardiac performance.
Circ Shock, 22:11-22.
- Ramenotsky, M.; Luterman, A.; Quindlen, E. (1984):**
Maximum survival in pediatric trauma: The ideal system. *J. Trauma*, 24: 818.
- Ramenotsky, M.; Luterman, A.; Curreir, P.W. (1983):**
EMS for pediatrics: Optimum treatment or unnecessary delay?
J. Pediatr. Surg., 18: 498.
- Rand, M.J. (1992):**
Nitrgic transmission: nitric oxide as a mediator of non-adrenergic non-cholinergic neuro-effector transmission.
Clin. Exp. Pharmacol. Physiol., 19: 147-169.
- Reily, J.; Cunnion, R.; Burch-Whitman, C. et al. (1989):**
A circulating myocardial depressant substance is associated with cardiac dysfunction and peripheral hypoperfusion in patients with septic shock. *Chest*, 95: 1072-1080.

- Ress, D.D.; Cellek, S.; Palmer, R.M. (1990):**
Monuadazone prevents the induction by endotoxin synthase and the associated effects on insight into endotoxin shock. *Biochem. Biop.*, 173: 541-7.
- Richard, S.D.; Downy, Q. (1997):**
Clinical care and metabolic responses. *Critical Care*, 325: 335.
- Richmond, J.; Sibbald, W.; Linton, A. (1982):**
Patterns of urinary protein excretion in patients with sepsis. *Nephrology*, 31: 219-223.
- Rivard, G.E.; David, N.; Ferrell, C. et al. (1995):**
Treatment of purpura fulminans in meningococemia with protein C concentrate. *J. Pediatr.*, 126: 646.
- Roberts, J.D.; Polaner, D.M.; Lang, P.; Zpol, W.M. (1992):**
Inhaled nitric oxide in persistent pulmonary hypertension of the newborn. *Lancet*, 340: 818-819.
- Robertson, F.M.; Ofner, P.J.; Ciceri, D.P.; Becker, W.K.; Pruitt, B.A.Jr. (1994):**
Detrimental hemodynamic effects of nitric oxide synthase inhibition in septic shock. *Arch. Surg.*, 129: 149-155.
- Rogers, F.; Dunn, R.; Barrett, R. et al. (1985):**
Alterations of capillary flow during sepsis. *Circ. Shock.*, 15: 105-110.
- Roman, J.; Velasco, F.; Fernandez, F. et al. (1992):**
Protein C, protein S and C4b-binding protein in neonatal severe infection and septic shock. *J. Perinatal Med.*, 20: 111.
- Rosenthal, J.; Healey, T.; Ellis, R. et al. (1996):**
A two year follow up of neonates with presumed sepsis treated with recombinant human granulocyte colony stimulating factor during the first week of life. *J. Pediatr.*, 128: 135.

- Russel, J.; Ronco, J.; Lockhat, D. et al. (1990):**
Oxygen delivery and consumption and ventricular preload are greater in survivors than in nonsurvivors of the adult respiratory distress syndrome. *Am. Rev. Respir. Dis.*, 141: 659-665.
- Schafer, R.; Sheil, J.M. (1995):**
Superantigens and their role in recurrent disease. *Infect. Dis.*, 10: 369.
- Schmidt, HHHW, Walter, U.; (1994):**
NO at work.
Cell, 78: 919-925.
- Schroder, C.H.; Severijnen, R.S.; Potting, C.M. (1992):**
Continuous arteriovenous hemofiltration in a premature newborn as treatment of over hydration and hyperkalemia due to sepsis. *Eur. J. Pediatr. Surg.*, 2: 368.
- Schumer, W.; Dugapta, T.; Moss, G. et al. (1970):**
Effect of endotoxemia on liver cell mitochondria in man. *Ann. Surg.*, 171: 875-882.
- Seitz, R.; Wolf, M.; Ebbring, R. et al. (1989):**
The disturbances of hemostasis in septic shock: role of neutrophil elastase and thrombin, effects of antithrombin III and plasma substitution. *Eur. J. Haematol.*, 43: 22.
- Shamoon, H.M.; Hendler, R.; Sherwin, R.S. (1981):**
Synergetic interaction among anti insulin hormone in the pathogenesis of stress hypoglycemia in humans.
J. Clin. Endocrin. Metab., 52: 1235.
- Sheridan, R.L.; Weber, J.M.; Peterson, H.F. et al. (1995):**
Central venous catheter sepsis with weekly catheter change in paediatric burn patients: An analysis of 221 catheters. *Burns* 21: 127.
- Shapira, M.R.; Greenfield, S. (1987):**
The complete blood count and leukocyte differential count.
Ann. Intern. Med., 106: 65-74.

- Shoemaker, W.; Appel, P.; Kram, H. (1988):**
Tissue oxygen debt as a determinant of lethal and nonlethal postoperative organ failure. *Crit Care Med.*, 16: 1117-1120.
- Sibbald, W.J.; Paterson, W.; Holliday, R.L. et al. (1988):**
Pulmonary hypertension in sepsis: Measurement of pulmonary arterial diastolic-pulmonary wedge pressure gradient and the influence of passive and active factors. *Chest*, 72: 583-591.
- Siedel, J.S.; Hornbein, M.; Yoshuyama, K. (1994):**
Emergency medical services and the pediatric patients: Are the needs being met? *Pediatrics*, 73: 769.
- Simon, J.E.; Smookler, S.; Guy, B. (1989):**
A regionalized approach to pediatric emergency care. *Pediatr. Clin. North Am.*, 28: 677.
- Sorenson, J.; Kald, B.; tagesson, C. et al. (1994):**
Platelet-activating factor and phospholipase A2 in patients with septic shock and trauma. *Intensive Care Med.*, 20: 555, 1994.
- Souba, W.W.; Wilmore, D.W. (1983):**
Postoperative alterations of arteriovenous exchange of amino acids across the gastrointestinal tract. *Surgery*, 94: 342.
- Sprung, C. (1991):**
Definition of sepsis-have we reached a consensus? *Crit Care Med.*, 19: 849-851.
- Sprague, R.; Thiernemann, C.; Varve, J.R. (1992):**
Endogenous endothelium-derived relaxing factor opposes hypoxic pulmonary vasoconstriction and supports blood flow to hypoxic alveoli in anaesthetised rabbits. *Proc. Natl. Acad. USA*, 89: 8711-8715.
- Stamler, J.S. (1994):**
Redox signaling: nitrosylation and related target interactions of nitric oxide. *Cell*, 78: 931-936.

Steinhorn, D.M.; Green, T.P. (1991):

Severity of illness correlates with alterations in energy metabolism in the pediatric intensive care unit. Crit Care Med., 19: 1503.

Steven, M.; Hollenberg (1998):

Inhibition of nitric oxide synthesis in sepsis.

Apromising Strategy? Critic care Md. 26: 638-640.

Stratton, C.W. (1981):

Tracheostomy and endotracheal tube specimens. Infect. Control., 2: 162-164.

Stuber, F.; Petersen, M.; Bokelmann, F. et al. (1996):

A genomic polymorphism within the tumor necrosis factor locus influences plasma tumor necrosis factor-alpha concentrations and outcome of patients with severe sepsis. Crit Care Med., 24: 381.

Tendler, C.; Grossman, S.; Tenenbaum, J. (1989):

Medication dosages during pediatric emergencies: A simple and comprehensive guide. Pediatrics, 84: 731.

Teule, G.J.; den Hollander, W.; Bronsveld, W. et al. (1981):

Noninvasive detection of blood volume redistribution in canine endotoxin shock. Circ. Shock, 8: 627-634.

Teule, G.J.; den Hollander, W.; Bronsveld, W. et al. (1983):

Effect of volume loading and dopamine on hemodynamics and red-cell redistribution in canine endotoxin shock. Circ Shock, 10: 41-50.

Tischler, M.E.; Fagan, J.M. (1993):

Response to trauma of protein, amino acid and carbohydrate metabolism in injured rat skeletal muscle. Metabolism, 32: 853.

Torres, A.; El-Ebiary, M.; Padro, L. et al. (1994):

Validation of different techniques for the diagnosis of ventilator-associated pneumonia.

Am. J. Respir. Crit. Care Med., 149: 324.

- Townsend, M.; Hampton, W.; Haybron, D. et al. (1986):**
Effective organ blood flow and bioenergy status in murine peritonitis. *Surger*, 100: 205-212.
- Tuchschiidt, J.; Fried, J.; Astiz, M. et al. (1992):**
Elevation of cardiac output and oxygen delivery improves outcome in septic shock. *Chest*, 102: 216-220.
- Tuchschiidt, J.; Fried, J.; Swinney, R. et al. (1989):**
Early hemodynamic correlates of survival in patients with septic shock. *Crit Care Med.*, 17: 719-723.
- Vallance, P.; Moncada, S. (1994):**
Nitric oxide-from mediator to medicines.
J. R. Coll. Phys. Lond., 28: 209-219.
- Van Dervort, A.L.; Yan, L.; Madara, P.J. et al. (1994):**
Nitric oxide regulates endotoxin-induced TNF- α production by human neutrophils. *J. Immunol.*, 152: 4102-4109.
- Van Deuren, M.; Santman, F.W.; van Dalen, R. et al. (1992):**
Plasma and whole blood exchange in meningococcal sepsis. *Clin. Infect. Dis.*, 15: 424.
- Velecek, F.L.; Weiss, A.; DiMato, D. (1987):**
Traumatic death in children. *J. Pediatr. Surg.*, 12: 375.
- Vincent, J.L.; (1997):**
Clinical trials in sepsis: Where do we stand? *J. Crit Care*, 12: 3.
- Wagner, D.A.; Young, V.R.; Tannenbaum, S.R. (1983):**
Mammalian skrate biosynthesis: incorporation of $^{19}\text{NH}_3$ into nitrate is enhanced by endotoxin treatment. *Proc. Natl. Acad. Sci., USA*, 80: 4518-21.
- Weisel, R.; Vitto, L.; Dennis, R. et al. (1977):**
Myocardial depression during sepsis. *Am. J. Surg.*, 133: 512-521.
- Wilbert, C.; Davis, A.T.; Herman, J.J. (1985):**
Short-term holding room treatment of asthmatic children. *J. Pediatric.*, 106: 707.

- William, W.; Robert, R. (1997):**
Metabolic response to injury.
Critical care: 335: 340.
- Wilmore, D.W.; Goodwin, C.W.; Aulick, A.H. (1987):**
Effect of injury and infection on visceral metabolism and circulation.
Ann. Surg., 192-491.
- Wilson, M.L.; Weinstein, M.P. (1994):**
General principles in the laboratory detection of bacteria and fungemia. Clin. Lab. Med., 14: 69-82.
- Wilmore, D.W.; Goodwin, C.W.; Aulick, A.H. et al. (1980):**
Effect of injury and infection on visceral metabolism and circulation. Ann. Surg., 192: 491.
- Wolfe, R.R.; Herndon, D.N.; Jahoor, F. (1987):**
Effect of severe burn injury on substrate cycling by glucose and fatty acids.
N. Eng. J. Md., 317-403.
- Wong, H.R.; Carcillo, J.A.; Shan, N. (1996):**
Increased serum nitrate and nitrite levels in children with sepsis syndrome.
Critical Care Md., 835: 892.
- Yokokawa, K.; Mankus, R.; Saklayer, M. et al. (1995):**
Increased nitric oxide production in patients with hypotension during hemodialysis.
Am. Intern. Med., 123: 35-7.
- Yuan, Shi, M.S.; Hua-aiang, L.I.; Chikao-Shen, M.D.; Jlanghual-Wang, M.D.; Shtwen-Qin, M.S.; Ren-Llu, M.S. and Pan, J.K. (1993):**
Plasma nitric oxide levels in newborn infants with sepsis the Journal of Pediatric. 433-439.
- Zeller, W.P.; Goto, M.; Hurley, R.M. (1988):**
Endotoxin in newborn septic shock: Significance, metabolic, and cardiovascular changes. Ann. Clin. Lab. Sci., 18: 253.

Zenz, W.; Muntean, W.; Gallistl, S. et al. (1995):

Recombinant tissue plasminogen activator treatment in two infants with fulminant meningococemia. *Pediatrics.*, 96: 44.

Zingarelli, B.; Salzman, A.; Szabo, C. (1995):

Protective effects of nicotinamide against nitric oxide mediated vascular failure in endotoxic shock: Potential involvement of poly ADP ribosyl synthetase. *Shock*, 5: 258.

Zoritch, B. (1994):

Nitric oxide in asthma.

Arch. Dis. Child, 72: 259-262.