

References

- **Acland, K.M. ; Darvay, A. ; Griffin, C. ; Aali, S.A. and Russell-Jones, R. (1999):** Staphylococcal scalded skin syndrome in an adult associated with methicillin-resistant, *Staphylococcus aureus*. Br J Dermatol 140, pp. 518–520.
- **Aguita and Maria. (2009):** "Combating Bacterial Infection". LabNews.co.uk.http://www.labnews.co.uk/printer_friendly.php/2912/combating-bacterial-infection. Recieved on 2009-05-05.
- **Aldeyab, Mamoon A. ; Monnet, Dominique L. ; López-Lozano, José María ; Hughes , Carmel M. ; Scott, Michael G. ; Kearney, Mary P.; Magee, Fidelma, A. and McElnay, James C.(2008):** Modelling the impact of antibiotic use and infection control practices on the incidence of hospital-acquired methicillin-resistant *Staphylococcus aureus*: a time-series analysis. JAC Advance Access published online on May 7, 2008 Journal of Antimicrobial Chemotherapy, doi:10.1093/jac/dkn198.
- **Amyes, S.G.B.; Paton, R.H.; Miles, R.S. and Hood, J. (1992):** Ofloxacin as the standard 4- quinolone in sensitivity tests. Journal of Antimicrobial Chemotherapy. 30:101-102.
- **Anonymous. (2004):**National Nosocomial Infections Surveillance (NNIS) System Report; data summary from January 1992 through June 2004, issued October 2004. Am. J. Infect. Control 32:470-485.
- **Apfalter, P.; Assadian, O.; Kalczyk, A.; Lindenmann, V.; Makristathis, A. and Mustafa, S. (2002):** Performance of a

new chromogenic oxacillin resistance screen medium (Oxoid) in the detection and presumptive identification of methicillin-resistant *Staphylococcus aureus*. *Diagn Microbiol Infect Dis*. 44:209-211.

- **Arias ; Cesar, A . ; Murray and Barbara, E. (2009):** "Antibiotic-Resistant Bugs in the 21st Century — A Clinical Super-Challenge". *New England Journal of Medicine* 360 (5): 439–443. doi:10.1056/NEJMp0804651.
- **Arvidson, S. ; Janzon, L. and Lofdahl , S.(1990):** The role of the hemolysin gene (hld) in the agr-dependent regulation of exoprotein synthesis in *Staphylococcus aureus*, p. 419-431. In R. P. Novick (ed.), *Molecular biology of the staphylococci*. VCH Publishers Inc., New York, N.Y.
- **Baird, D. (1996):** *Staphylococcus: cluster-forming gram-positive cocci*. in Collee J. G.; Fraser A. G. ; Marmion B. P. and Simmons, A. (eds.) *Mackie & McCartney Practical Medical Microbiology*. Churchill Livingstone. P: 245-274.
- **Baird-Parker, A. C. (1979):** Methods for identifying staphylococci and micrococci. In: Skinner, F. A. and Lovelock, D. W. (eds) *Identification methods for microbiologists*, The society for Applied Microbiology Technical Series No. 14 Academic Press, London, pp 201-210.
- **Bannerman, T.L. (2003):** *Staphylococcus, Micrococcus and other catalase-positive cocci that grow aerobically*. In: P.R. Murray, E.J. Baron, J.H. Jorgensen, M.A. Pfaller and R.H. Tenover, Editors, *Manual of Clinical Microbiology* (8th edn), American Society for Microbiology, Washington. pp. 384–404.

- **Barber, M. and Rozwadowska-Dowzenko, M. (1948):** Infection by penicillin resistant staphylococci. *Lancet*. 252:641-4.
- **Bayston, R. ; Ashraf, W. and Smith, T. (2007):** "Triclosan resistance in methicillin-resistant *Staphylococcus aureus* expressed as small colony variants: a novel mode of evasion of susceptibility to antiseptics". *J. Antimicrob. Chemother.* 59 (5): 848–53. doi:10.1093/jac/dkm031. PMID 17337510.
- **Becker, K. ; Harmsen, D. ; Mellmann, A. ; Meier, C. ; Schumann, P. and Peters, G. (2004):** Development and evaluation of a quality-controlled ribosomal sequence database for 16S ribosomal DNA-based identification of *Staphylococcus* species. *J Clin Microbiol.* 42:4988–95
- **Belmonte, G. ; Cescatti, L. ; Ferrari, B. ; Nicolussi, T. ; Ropele, M. and Menestrina, G.(1987):** Pore-formation by *Staphylococcus aureus* alpha- toxin in lipid bilayers: temperature and toxin concentration dependence. *Eur. J. Biophys.* 14:349-358.
- **Nelson, J.M. ; Chiller, T.M. ; Powers, J.H. and Angulo, F.J. (2007):** "Fluoroquinolone-resistant *Campylobacter* species and the withdrawal of fluoroquinolones from use in poultry: a public health success story". *Clin. Infect. Dis.* 44 (7): 977–80. doi:10.1086/512369.PMID17342653.
- **Berger-Bachi, B . and Rohrer, S . (2002):** Factors influencing methicillin resistance in staphylococci. *Arch Microbiol.* 178:165-71.
- **Bhakdi, S. and Trantum-Jensen , J. (1991):** Alpha-toxin of *Staphylococcus aureus*. *Microbiol Rev* 55:733.

- **Binnicker, M.J. (2007):** Detection of *Coccidioides* Species in Clinical Specimens by Real-Time PCR.. J. Clin. Microbiol., 45: 173 - 178.
- **Birmingham, M.C.; Rayner, C.R. ; Meagher, A.K. ; Flavin, S.M. ; Batts, D.H. and Schentag, J.J. (2003):** "Linezolid for the treatment of multidrug-resistant, gram-positive infections: experience from a compassionate-use program". Clin. Infect. Dis. 36 (2): 159–68. doi:10.1086/345744. PMID 12522747.
- **Blanc, D. S.; Wenger, A. and Bille, J. (2003):** Evaluation of a novel medium for screening specimens from hospitalized patients to detect methicillin-resistant *Staphylococcus aureus*. J. Clin. Microbiol. 41:3499-3502.
- **Bohach, G. A. ; Fast, D. J. ; Nelson, R. D. and Schlievert, P. M.(1990):** Staphylococcal and streptococcal pyrogenic toxins involved in toxic shock syndrome and related illnesses. Crit. Rev. Microbiol. 17:251-272.
- **Bohach, G. A. ; Dinges, M. M. ; Mitchell, D. T. ; Ohlendorf, D. H. and Schlievert, P. M. (1997):** Exotoxins, p. 83-111. In Crossley, K. B. ;Archer, G. L. (ed.), The staphylococci in human disease. Churchill Livingstone, Inc., New York, N.Y.
- **Bohach, G. A. ; Jablonski, L. ; Roggiani, M. ; Sadler, I. ; Schlievert , P. M. ; Mitchell, D. and Ohlendorf, D. (1998):** Biological activity of pyrogenic toxins delivered at the mucosal surface, p. 170-172. In Arbuthnott, J. ; Furman, B. (ed.), European Conference on Toxic Shock Syndrome. International Congress and Symposium Series 229. Royal Society of Medicine Press Ltd, New York, N.Y.

- **Brakstad, O.G.; Aasbakk, K. and Maeland, J.A. (1992):** Detection of *Staphylococcus aureus* by Polymerase Chain Reaction Amplification of the nuc Gene. J Clin Microbiol. 30(7):1654-60.
- **Brakstad, O.G. and Maeland, J.A. (1997):** Mechanisms of methicillin resistance in staphylococci. 105(4):264-76.
- **Brumfitt, W. and Hamilton-Miller, J.M. (1994):** "Limitations of and indications for the use of co-trimoxazole". J Chemother 6 (1): 3–11. PMID 8071675.
- **Burke, (1922):** Notes on Gram stain with description of a new method. J. of Bacteriology 7:159.
- **Carleton, HA.; Diep, BA.; Charlebois, ED.; Sensabaugh, GF.; Perdreau-Remington, F.(2004):** Community-adapted methicillin-resistant *Staphylococcus aureus* (MRSA): population dynamics of an expanding community reservoir of MRSA. J Infect Dis.190:1730–1738. doi: 10.1086/425019.
- **Carter, A . P . ; Clemons, W.M. ; Brodersen, D . E . ; Morgan-Warren, R. J. ; Wimberly, B .T . and Ramakrishnan, V . (2000):** "Functional insights from the structure of the 30S ribosomal subunit and its interactions with antibiotics". Nature 407 (6802): 340–8. doi:10.1038/35030019. PMID 11014183.
- **Cauwelier, B.; Gordts, B.; Descheemaeker, P. and Van Landuyt, H. (2004):** Evaluation of a disc diffusion method with cefoxitin (30 µg) for detection of methicillin-resistant *Staphylococcus aureus*. Eur. J. Clin. Microbiol. Infect. Dis., 23: 389-92.
- **CDC (2005):** Department of Health and Human Services. Laboratory Detection of Oxacillin/Methicillin-resistant *Staphylococcus*

- aureus*. Available from: <http://www.cdc.gov/ncidod/hip/Lab/FactSheet/mrsa.htm>.
- **CDC (2006):** "Management of Multidrug-Resistant Organisms in Healthcare Settings.
 - **Centers for Disease Control and Prevention. (2007):** ",MRSA; Methicillin-Resistant *Staphylococcus aureus* in Healthcare Settings".
 - **Chambers, H . F. (1988):** Methicillin resistant staphylococci. Clin. Microbiol. Rev., 1:173-186.
 - **Chambers H . F. (1997):** Methicillin resistance in staphylococci: molecular and biochemical basis and clinical implications. Clin Microbiol. Rev., 10(4):781-91.
 - **Chambers, H. F. (2001):** The changing epidemiology of *Staphylococcus aureus*. Emerg. Infect. Dis. 7:178-182.
 - **Chambers, H. F.(2003):** Solving staphylococcal resistance to beta-lactams. Trends Microbiol. 11:145-148.
 - **Cheesbrough, M. (2000):** Culturing blood. In: District Laboratory Practice in Tropical Countries. Part 2. Cambridge University Press. P: 124-130.
 - **Chen, X.; Zehnbaauer, B.; Gnirke, A. and Kwok, P.Y. (1997):** Fluorescence energy transfer detection as a homogeneous DNA diagnostic method. Proc Natl Acad Sci U S A. 94(20):10756-61.
 - **Cherkaoui, A.; Renzi, G.; Francois, P. and Schrenzel, J. (2007):** Comparison of four chromogenic media for culture-based screening of methicillin-resistant *Staphylococcus aureus*. J Med Microbiol. 56(Pt 4):500-3.

- **Cheung, A. L. ; Koomey, J. M. ; Butler, C. A. ; Projan, S. J. and Fischetti, V. A. (1992):** Regulation of exoprotein expression in *Staphylococcus aureus* by a locus (sar) distinct from agr. Proc. Natl. Acad. Sci. USA. 89:6462-6466.
- **Clauditz, A. ; Resch, A. ; Wieland, K. P. ; Peschel, A. and Götz, F. (2006):** "Staphyloxanthin plays a role in the fitness of *Staphylococcus aureus* and its ability to cope with oxidative stress". Infection and immunity 74 (8): 4950–3. doi:10.1128/IAI.00204-06. PMID 16861688. PMC: 1539600.
- **Colakoglu, S.; Aliskan, H.; Senger, S.S.; Turunc, T.; Demiroglu, Y.Z.; Arslan, H. (2007):** Performance of MRSA ID chromogenic medium for detection of methicillin-resistant *Staphylococcus aureus* directly from blood cultures and clinical specimens. Diagn Microbiol Infect Dis. 2007 Nov; 59(3):319-23. Epub 2007 Oct 29.
- **Coleman, D. C. ; Arbuthnott, J. P. ; Pomeroy, H. M. and Birbeck, H. (1986):** Cloning and expression in *Escherichia coli* and *Staphylococcus aureus* of the beta-lysin determinant from *Staphylococcus aureus*: evidence that bacteriophage conversion of beta-lysin activity is caused by insertional inactivation of the beta-lysin determinant. Microb. Pathog. 1:549-564.
- **Colin, D. A. ; Mazurier, I. ; Sire, S. and Finck-Barbancon, V.(1994):** Interaction of the two components of leukocidin from *Staphylococcus aureus* with human polymorphonuclear leukocyte membranes: sequential binding and subsequent activation. Infect. Immun. 62:3184-3188.

- **Compernelle, V.; Verschraegen, G. and Claeys, G. (2007):** Combined Use of Pastorex Staph-Plus and Either of Two New Chromogenic Agars, MRSA ID and CHROMagar MRSA, for Detection of Methicillin-Resistant *Staphylococcus aureus*. J Clin Microbiol. 45 (1):154-8.
- **Cormican, M. G. and Jones, R. N. (1996):** Emerging resistance to antimicrobial agents in gram-positive bacteria. Enterococci, staphylococci and nonpneumococcal streptococci. Drugs 51:(Suppl. 1):6-12.
- **Cormican, M.G.; Wilke, W. and Barret, M. (1996):** Phenotypic detection of mec A positive Staph blood stream isolates. Diagn. Microbiol. Infect. Dis. 25:107-112.
- **Cosgrove, S.E.; Vigliani, G.A. and Campion, M. (2009):** "Initial low- dose gentamicin for *Staphylococcus aureus* bacteremia and endocarditis is nephrotoxic". Clin Infect Dis. 48 (6): 713–721.
- **Coulson Christopher, J. (1994):** "Bacterial RNA-Polymerase - Rifampin as Antimycobacterial." Molecular Mechanisms of Drug Action. 2nd ed. Bristol, PA: Taylor Francis,. pp. 40-41.
- **Cremer, A.W. and Grüneberg, R.N. (1988):** Assessment of a new test ("Staphylase") for the identification of *Staphylococcus aureus*. Med Lab Sci. 45(3):221–224.
- **Crum, N. F. ; Lee, R. U. ; Thornton, S. A. ; Stine, O. C. ; Wallace, M. R. Barrozo, C.; Keefer-Norris, A.; Judd, S.and Russell, K. L. (2006):** Fifteen-year study of the changing epidemiology of methicillin-resistant *Staphylococcus aureus*. Am. J. Med. 119:943-951.

- **Davies, A.; Perry, J.D. and Butterworth, L.A. (2004):** An evaluation of MRSA ID: a new chromogenic medium for the isolation and identification of methicillin-resistant *Staphylococcus aureus*. R2151, Pragues (République Tchèque) 2004 14th, ECCMID.
- **Davis, S.L.; McKinnon, P.S. and Hall, L.M. (2007):** Pharmacotherapy 27 (12): 1611–1618. PMID 18041881.
- **de San, N.; Denis, O.; Gasasira, M.F.; De Mendonca, R.; Nonhoff, C. and Struelens, M.J. (2007):** Controlled Evaluation of the IDI-MRSA Assay for Detection of Colonization by Methicillin-Resistant *Staphylococcus aureus* in Diverse Mucocutaneous Specimens. J Clin Microbiol. 45(4):1098-101.
- **de Sousa, M .A. and de Lencastre, H. (2004):** Bridges from hospitals to the laboratory: genetic portraits of methicillin-resistant *Staphylococcus aureus* clones. FEMS Immunol Med Microbiol. 40:101-111.
- **Deurenberg, R . H. ; Vink, C . ; Kalenic, S . ; Friedrich, A .W . ; Bruggeman, C . A . and Stobberingh, E .E. (2007):**The molecular evolution of methicillin-resistant *Staphylococcus aureus*. Clin Microbiol Infect. 13:222-35.
- **Diederer, B.; van Duijn, I.; Willemse, P.; van Keulen, P. and Kluytmans, J. (2005):** Performance of CHROMagar MRSA medium for detection of methicillin- resistant *Staphylococcus aureus*. J.Clin.Microbiol. 43:1925-1927.
- **Diederer, B.M. ; van Leest, M.L. ; Duijn, I. ; Willemse, P. ; van Keulen, P.H. and Kluytmans, J.A. (2006):** Performance of MRSA ID, a new chromogenic medium for detection of

- methicillin-resistant *Staphylococcus aureus*, J. Clin. Microbiol. 44, pp. 586–588.
- **Dinges, M. M ; Orwin, P. M and Schlievert, P. M.(2000):** Exotoxins of *Staphylococcus aureus*. Clin Microbiol Rev. 13:16–34.
 - **Domann, E.; Hossain, H.; Fussle, R. and Chakraborty, T.(2000):** Rapid and reliable detection of multiresistent *Staphylococcus aureus* (MRSA) by multiplex PCR. Dtsch. Med. Wochenschr. 19: 125(20):613-8.
 - **Donnio, P. Y. ; Oliveira, D. C. ; Faria, N. A. ; Wilhelm, N. ; Le Coustumier, A. and de Lencastre, H. (2005):** Partial excision of the chromosomal cassette containing the methicillin resistance determinant results in methicillin-susceptible *Staphylococcus aureus*. J. Clin. Microbiol. 43:4191-4193.
 - **Drancourt, M. and Raoult, D. (2002):** rpoB gene sequence-based identification of *Staphylococcus* species. J Clin Microbiol. 40:1333–8.
 - **Duckworth, G .J . ; Lothian, J . L . and Williams, J .D.(1988):** Methicillin- resistant *Staphylococcus aureus*: report of an outbreak in a London teaching hospital. J Hosp Infect. 11:1-15.
 - **Dufour, P.; Gillet, Y. and Bes, M. (2002):** Community-acquired methicillin-resistant *Staphylococcus aureus* infections in France: emergence of a single clone that produces Pantone-Valentine leukocidin. Clin Infect Dis. 35:819–24.
 - **Eady, E. A. and Cove, J. H. (2003):** Staphylococcal resistance revisited: community-acquired methicillin resistant

- Staphylococcus aureus*—an emerging problem for the management of skin and soft tissue infections. Curr. Opin. Infect. Dis. 16:103-124.
- **Earhart, C. A. ; Mitchell, D. T. ; Murray, D. L. ; Pinheiro, D. M. ; Matsumara, M. ; Schlievert, P. M. and Ohlendorf, D. H. (1998):** Structures of five mutants of toxic shock syndrome toxin-1 with reduced biological activity. Biochemistry 37:7194-7202.
 - **Easmon, CSF. and Adlam, C. (1983):** Staphylococci and staphylococcal infections. Vols 1 and 2 Academic Press, London.
 - **Elek, S. D., and Levy, E. (1950):** Distribution of hemolysins in pathogenic and non-pathogenic staphylococci. J. Pathol.Bacteriol. 62:541-554.
 - **Engels, w. and Kamp, M.A.F. (1981):** Staphylococcus and micrococcus. In:Topley & Wilson Principles of Bacteriology, Virology and Immunity. Tom-Parker, M. and Duerden, B.I.(eds) 8th edition ,vol.II, Edward Arnold, A. division of Hodder & Stoughton, London. P.172.
 - **Engemann, J. J. ; Carmeli, Y. ; Cosgrove, S. E. ; Fowler, V. G. ; Bronstein, M. Z. ; Trivette, S. L. ; Briggs, J. P. ; Sexton, D. J. and Kaye, K. S.(2003):** Adverse clinical and economic outcomes attributable to methicillin resistance among patients with *Staphylococcus aureus* surgical site infection. Clin. Infect. Dis. 36:592-598.
 - **Enright, M.C.; Day, N.P. and Davies, C.E. (2000):** Multilocus sequence typing for characterization of methicillin- resistant

- and methicillin –susceptible clones of *staphylococcus aureus*.
J Clin Microbiol. 38:1008-15.
- **Espy, M. J.; Uhl, J. R.; Sloan, L. M.; Buckwalter, S. P.; Jones, M. F. ; Vetter, E. A. ;Yao, J. D. C. ;Wengenack, N. L. ; Rosenblatt, J. E. ; Cockerill III, F. R. and Smith, T. F. (2006):** Real-Time PCR in Clinical Microbiology: Applications for Routine Laboratory Testing. Clinical Microbiology Reviews, p. 165-256, Vol. 19, No. 1 doi:10.1128/CMR.19.1.165-256.2006.
 - **Felton, A. ; Grandry, B. ; Lagrange, P.H. and Casin, I. (2002):**Evaluation of three techniques for detection of low-level methicillin-resistant *Staphylococcus aureus* (MRSA): a disc diffusion method with cefoxitin and moxalactam, the Vitek 2 System, and the MRSA-screen latex agglutination test. J. Clin. Microbiol. 40:2766-2771.
 - **Fischer , W. ; Rösel , P. and Koch , H. U. (1981):** Effect of alanine ester substitution and other structural features of lipoteichoic acids on their inhibitory activity against autolysins of *Staphylococcus aureus*. J. Bacteriol. 146:467-475.
 - **Fischer , W. (1988):** Physiology of lipoteichoic acids in bacteria. Adv. Microb. Physiol. 29:233-302.
 - **Fischer , W. (1994):** Lipoteichoic acids and lipoglycans. In Bacterial Cell Wall, pp. 199–215. Edited by J.-M. Ghuysen & R. Hakenbeck. Amsterdam: Elsevier.
 - **Forbes, B.A. (2007):** Bailey and Scott's Diagnostic Microbiology, 12th ed. C.V. Mosby Company, St. Louis, MO.
 - **Foster,T.J. and McDevitt,D. (1994):** Molecular basis of adherence of staphylococci to biomaterials. p. 31, In Bisno AL, Waldvogel FA

- (eds): Infections Associated with Indwelling Medical Devices, 2nd Edition. American Society for Microbiology, Washington, D.C.
- **Fournier, J. M.; Bouvet, A. and Mathieu, D. (1993):** New latex reagent using monoclonal antibodies to capsular polysaccharide for reliable identification of both oxacillin-susceptible and oxacillin-resistant *Staphylococcus aureus*. Journal of Clinical Microbiology. 31: 1342-1344.
 - **Fowler, V. G. ; Jr. ; Sakoulas, G. ; McIntyre, L. M. ; Meka, V. G. ; Arbeit, R. D. ; Cabell, C. H. ; Stryjewski, M. E. ; Eliopoulos, G. M. ; Reller, L. B. ; Corey, G. R. ; Jones, T. ; Lucindo, N. ; Yeaman, M. R. and Bayer, A. S. (2004):** Persistent bacteremia due to methicillin-resistant *Staphylococcus aureus* infection is associated with agr dysfunction and low-level in vitro resistance to thrombin-induced platelet microbiocidal protein. J. Infect. Dis. 190:1140-1149.
 - **Francois, P. and Schrenzel, J. (2008):** "Rapid Diagnosis and Typing of *Staphylococcus aureus*". Staphylococcus: Molecular Genetics. Caister Academic Press. ISBN 978-1-904455-29-5.
 - **Frebourg, N.B. ; Nouet, D. ; Lemee, L. and Iemeland, J.F. (1998):** Comparison of ATB Staph., rapid ATB Staph., Vitek and E-test method detection of oxacillin heterogeneous resistance in staphylococci possessing mec A gene. J. Clin. Microbiol. 36:52-57.
 - **Geary, C. and Stevens, M. (1991):** A rapid test to detect the most clinically significant *Staphylococcus* species. Medical Laboratory Sciences. 48:99-105. nicable Disease Report,

- Public Health Laboratory Service Communicable Diseases Surveillance Center, 61 Colindale Avenue, London 2: R25-29.
- **Gemmell, C. G.; King, S. L. and Sturrock, R. D. (1987):** Teichoic acid antibody measurement as an aid to the diagnosis of septic arthritis due to *Staphylococcus aureus*. Serodiagnosis and Immunotherapy. 1:201-207.
 - **Ghuysen, J.M. and Hakenbeck, R.(1994):** Bacterial cell wall.
 - **Gill, S.R. ; Founts, D. E. ; Archer, G. L. ; Mongodis, E. F. ; Deboy, R.T. ; Ravel, J. ; Paulsen, I.T. ; Kolonay, J.F. and Brinkac, L. (2005):** " Insights on evolution of virulence and resistance from the complete analysis of an early methicillin- resistant *staphylococcus aureus* strain and a biofilm producing methicillin- resistant staphylococcus epidermidis strain" Journal of bacteriology. 187 (7): 2426- 38.
 - **Gilligan, P. H.; Gage, P. A.; Welch, D. F.; Muszynski, M. J. and Wait, K. R. (1987):** Prevalence of thymidine-dependent *Staphylococcus aureus* in patients with cystic fibrosis. Journal of Clinical Microbiology. 25:1258-1261.
 - **Gouaux, J. E. ; Braha, O. ; Hobaugh, M. R. ; Song, L. ; Cheley, S. ; Shustak, C. and Bayley, H.(1994):** Subunit stoichiometry of Staphylococcal alpha-hemolysin in crystals and on membranes: a Heptameric transmembrane pore. Proc. Natl. Acad. Sci. USA 91:12828-12831.
 - **Gould, J.C. (1975):** Tests for sensitivity to antibiotics of the laboratory. In: Mackie & McKartney Practical Medical Microbiology. Duguid, J.P. ; Marmion, B.P. and Swan, R.H.A.(eds). 12th edition, vol.II, Churchill Livingstone, P. 199.

- **Gray, G. S. and Kehoe, M.(1984):** Primary sequence of the toxin gene α from *Staphylococcus aureus* Wood 46. Infect. Immun. 46:615-618.
- **Griffith, R.S. (1981):**"Introduction to vancomycin". Rev Infect Dis 3: S2004.
- **Grundmann, H . ; Aires-de-Sousa, M . ; Boyce, J . and Tiermersma, E. (2006):** Emergence and resurgence of methicillin-resistant *Staphylococcus aureus* as a public-health threat. Lancet. 368:874-85.
- **Gurran, C. ; Holliday, M. G. ; Perry, J. D. ; Ford, M. ; Morgan, S. and Orr, K. E. (2002):** A novel selective medium for the detection of methicillin-resistant *Staphylococcus aureus* enabling result reporting in under 24 h. J. Hosp. Infect. 52:148-151.
- **Hamilton-Miller, J. (1991):** "From foreign pharmacopoeias: 'new' antibiotics from old?". J Antimicrob Chemother 27 (6): 702–5. doi:10.1093/jac/27.6.702. PMID 1938680.
- **Harris, T.O. ; Grossman, D. ; Kappler, J.W. ; Marrack, P. ; Rich, R.R. and Betley, M.J. (1993):** Lack of complete correlation between emetic and T-cell-stimulatory activities of staphylococcal enterotoxins. Infect. Immun. 61, pp. 3175–3183.
- **Harshman, S. (1988):** Preparation and purification of staphylococcal alpha toxin. Meth. Enzymol., 165, 3-7.
- **Hartstein, A.I. ; Phelps, C.L. ; Kwok, R.Y.Y. and Mulligan, M.E. (1995):** In vivo stability and discriminatory power of methicillin-resistant *Staphylococcus aureus* typing by restriction endonuclease analysis of plasmid DNA compared

- with those of other molecular methods. J Clin Microbiol. 33:2022-6.
- **Hiramatsu, K . ; Hanaki, H . and Ino,T. (1997):** Methicillin-resistant *Staphylococcus aureus* clinical strains with reduced vancomycin susceptibility. J Antimicrob Chemother. 40:135-6.
 - **Hochkeppel, H. K. ; Braun, W. F. ; Vischer, W. ; Imm, A. ; Sutter, S. ; Staebli, U. ; Guggenheim, R. ; Kaplan, E. L. ; Boutonnier, A. and Fournier , J. M. (1987):** Serotyping and electron microscopy studies of *Staphylococcus aureus* clinical isolates with monoclonal antibodies to capsular polysaccharide types 5 and 8. J. Clin. Microbiol. 25:526-530.
 - **Holland, P.M.; Abramson, R.D.; Watson, R. and Gelfand, D.H. (1991):** Detection of specific polymerase chain reaction product by utilizing the 5'----3' exonuclease activity of *Thermus aquaticus* DNA polymerase. Proc Natl Acad Sci U S A. 88(16):7276-80.
 - **Howden, B.P. and Grayson, M.L. (2006):** "Dumb and dumber—the potential waste of a useful antistaphylococcal agent: emerging fusidic acid resistance in *Staphylococcus aureus*". Clin Infect Dis 42 (3): 394–400. doi:10.1086/499365.
 - **Idzik, D. ;Wojtyczka, R. ; Kepa,M. ; Wydmuch, Z. ; Pacha, J. ; Glab, S. ; Wroblewska, I. and Skowronek-Ciolek, O. (2000):** Evaluation of methicillin-resistance in *Staphylococcus aureus* by the agar disc diffusion method and PCR. Med Dosw Microbiol. 52(4):327-32.
 - **Inaam A. Al Obaida ; Udob, E.E. ; Jacobb, L.E. and Johnya, M. (1999):** Isolation and Characterization of Coagulase-Negative

- Methicillin-Resistant *Staphylococcus aureus* from Patients in an Intensive Care Unit. *Med Princ Pract.* 8:230-236 (DOI: 10.1159/000026097).
- **Ito, T.; Katayama, Y.; Asada, K.; Mori, N.; Tsutsumimoto, K.; Tiensasitorn, C. and Hiramatsu, K. (2001):** Structural comparison of three types of staphylococcal cassette chromosome mec integrated in the chromosome in methicillin-resistant *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 45:1323-1336.
 - **Ito, T . ; Okuma, K . ; Ma, X . X . ; Yuzawa, H . and Hiramatsu, K . (2003):** Insights on antibiotic resistance of *Staphylococcus aureus* from its whole genome: genomic island SCC. *Drug Resist.* 6:41- 52.
 - **Ito, T.; Ma, X. X.; Takeuchi, F.; Okuma, K.; Yuzawa, H. and Hiramatsu, K .(2004):** Novel type V staphylococcal cassette chromosome mec driven by a novel cassette chromosome recombinase, ccrC. *Antimicrob. Agents Chemother.* 48:2637-2651.
 - **Janknegt, R. (1997):** "The treatment of staphylococcal infections with special reference to pharmacokinetic, pharmacodynamic, and pharmacoeconomic considerations". *Pharmacy world & science* : PWS 19 (3): 133–41. doi:10.1023/A:1008609718457. PMID 9259029.
 - **Jawetz , E. ; Melnick , J. L. and Adelberg, E.A.(1989):** *Medical Microbiology.* Appleton & Lange, East Norwalk, CT.
 - **Jeffries, C. D. (1961):** Comparison of six physiological characteristics of staphylococci from laboratory specimens. *American Journal of Clinical Pathology.* 36:114-118.

- **Jek, V. H. ; Součkov, A. ; Koagul.za-pozitivn and stafylokoky.**
In: **Bedn, R. M. ; Fraňkov, V. ; Schindler, J. ; Souček, A.(1996):** editors. *Lab. mikrobiologie*. Praha: Marvil;. p.194-203.
- **Jevons, M . P. (1961):** Br Med J. 1:124–125.
- **Jonas, D.; Speck, M.; Daschner, F.D. and Grundmann, H. (2002):**
Rapid PCR-Based Identification of Methicillin-Resistant *Staphylococcus aureus* from Screening Swabs. J.Clin.Microbiol., 40(5): 1821-1823.
- **Jorgenson, J. H. and Kloos, W. E. (1987):** Staphylococcal Infections. In B. B. Wentworth (ed.), Diagnostic procedures for bacterial infections. 7th ed. American Public Health Association. Washington, D. C.
- **Jungk, J . ; Como-Sabetti, K. ; Stinchfield , P. ; Ackerman , P . and Harriman, K. (2007):** Epidemiology of Methicillin-Resistant *Staphylococcus aureus* at a Pediatric Healthcare System, 1991-2003. *Pediatr Infect Dis J.* 26:339-44.
- **Kamio, Y. ; Rahman, A. ; Nariya, H. ; Ozawa, T. and Izaki, K. (1993):** The two staphylococcal bi-component toxins, leukocidin and gamma- hemolysin, share one component in common. *FEBS Lett.* 321:15-18.
- **Karakawa, W. W. and Vann, W. F. (1982):** Capsular polysaccharides of *Staphylococcus aureus*. *Semin. Infect. Dis.* 4:285-293.
- **Kasbekar, N. (2006):** "Tigecycline: a new glycyclcycline antimicrobial agent.". *Am J Health Syst Pharm* 63 (13): 1235–43. doi:10.2146/ajhp050487. PMID 16790575.

- **Kasten Britta and Reski Ralf. (1997):** " β -lactam antibiotics inhibit chloroplast division in a moss (*Physcomitrella patens*) but not in tomato (*Lycopersicon esculentum*)". Journal of Plant Physiology 150 (1-2): 137–140.
<http://cat.inist.fr/?aModele=afficheN&cpsidt=2640663>.
- **Katayama, Y.; Ito, T. and Hiramatsu, K. (2000):** A new class of genetic element, *Staphylococcus* cassette chromosome *mec*, encodes methicillin resistance in *Staphylococcus aureus*. Antimicrob. Agents Chemother. 44:1549-1555.
- **Kenneth, T. (2005):** Edited from
<http://www.textbookofbacteriology.net>
- **Kirby, W .M. (1944):** Extraction of a high potent penicillin inactivator from penicillin resistant staphylococci. Science. 99:452-3.
- **Klein, E . ; Smith, D . L . and Laxminarayan, R . (2007):** "Hospitalizations and Deaths Caused by Methicillin-Resistant *Staphylococcus aureus*, United States, 1999–2005". Emerg Infect Dis. 13 (12): 1840-6.
- **Klevens, (2007):** " Invasive Methicillin- Resistant *Staphylococcus aureus* infections in the United States". JAMA. Retrieved on 10-31-2007.
- **Kloos, W.E. and Schleifer, K.H. (1975):** Simplified scheme for routine identification of human staphylococcus species. J. Clin. Microbiol. 1:82-88.
- **Kluytmans, J.; van Belkum, A. and Verbrugh, H. (1997):** "Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks". Clin. Microbiol. Rev. 10(3): 505–20. PMID 9227864. PMC:172932.

- **Kohli, R.M.; Walsh, C.T. and Burkart, M.D. (2002):** "Biomimetic synthesis and optimization of cyclic peptide antibiotics". *Nature* 418 (6898): 658–61. doi:10.1038/nature00907. PMID 12167866.
- **Kohne, D. E.; Steigerwalt, A. G. and Brenner, D. J. (1984):** Nucleic acid probe specific for members of the genus *Legionella*. *JID*.75(10):123-5
- **Kuehnert, M .J.; Kruszon-Moran, D.; Hill, H .A.; McQuillan, G.; McAllister, K. and Fosheim, G. (2006):** Prevalence of *Staphylococcus aureus* Nasal Colonization in the United States, 2001-2002. *JID*. 193(15):172-9.
- **Labischinski, H. (1992):** Consequences of the interaction of β -lactam antibiotics with penicillin binding proteins from sensitive and resistant *Staphylococcus aureus* strains. *Med. Microbiol. Immunol.* 181:241-65.
- **Ladhani, S. ; Joannou, C. L. ; Lochrie, D. P. ; Evans , R. W. and Poston, S. M. (1999):** Clinical, microbial, and biochemical aspects of the exfoliative toxins causing staphylococcal scalded-skin syndrome. *Clin. Microbiol. Rev.* 12:224-242.
- **Lally, R. T.; Ederer, M. N. and Woolfrey, B. F. (1985):** Evaluation of mannitol salt agar with oxacillin as a screening medium for methicillin-resistant *Staphylococcus aureus*. *J. Clin. Microbiol.* 22:501-504.
- **Larry Koreen ; Srinivas, V. ; Ramaswamy ; Edward, A. ; Graviss ; Steven Naidich ; James, M. ; Musser and Barry N. Kreiswirth.(2004):** spa Typing Method for Discriminating among *Staphylococcus aureus* Isolates: Implications for Use of a Single Marker To Detect Genetic Micro- and

Macrovariation. *Journal of Clinical Microbiology*, February 2004, p. 792-799, Vol. 42, No. 2 DOI: 10.1128/JCM.42.2.792-799.2004.

- **Lauderdale, T. L.; Kimberle, C. C. and Murray, P. R. (1999):** Reagents, stains and media. In: Patrick, R.M.; Boron, E.J.; Pfaller, M.A.; Tenover, F.C. and Tenover, R.H. (eds.) *Manual of Clinical Microbiology*. 7th ed. American Society for Microbiology. P: 1663-1707.
- **Lee, K.H. ; Fitton, J.E. and Wuthrich, K. (1987):** Nuclear magnetic resonance investigation of the conformation of α -haemolysin bound to dodecylphosphocholine micelles. *Biochim. Biophys. Acta* 911:144-153.
- **Lee, J. C. ; Liu, M. J. ; Parsonnet, J. and Arbeit, R. D. (1990):** Expression of type-8 capsular polysaccharide and production of toxic shock syndrome toxin-1 are associated among vaginal isolates of *Staphylococcus aureus*. *J. Clin. Microbiol.* 28:2612-2615.
- **Lee, C. Y.(1995):** Association of staphylococcal type-1 capsule-encoding genes with a discrete genetic element. *Gene* 167:115-119.
- **Lincosamides, Oxazolidinones, and Streptogramins". Merck Manual of Diagnosis and Therapy. Merck & Co., Inc.. November 2005.**
<http://www.merck.com/mmpe/print/sec14/ch170/ch170g.html>.
Retrieved on 2007-12-01.

- **Linda M. Weigel ; Don B. Clewell ; Steven R. Gill ; Nancye C. Clark ; Linda K. McDougal ; Susan E. Flannagan ; James F. Kolonay ; Jyoti Shetty ; George E. Killgore ; Fred C. Tenover. (2003):** Genetic Analysis of a High-Level Vancomycin-Resistant Isolate of *Staphylococcus aureus*. Science 28 November 2003. Vol. 302. no. 5650, pp. 1569 – 1571 DOI: 10.1126/science.1090956
- **Lindsay, J. A. ; Ruzin, A. ; Ross, H. F. ; Kurepina, N. and Novick, R. P. (1998):** The gene for toxic shock toxin is carried by a family of mobile pathogenicity islands in *Staphylococcus aureus*. Mol. Microbiol. 29:527- 543.
- **Livak, K.J.; Flood, S.J. ; Marmaro, J. ; Giusti, W. and Deetz, K.(1995):** Oligonucleotides with fluorescent dyes at opposite ends provide a quenched probe system useful for detecting PCR product and nucleic acid hybridization. PCR Methods Appl. 4(6):357-62.
- **Lowy, F. D. (1998):** *Staphylococcus aureus* infections. N. Engl. J. Med. 339:520-532.
- **Lyon, B . R and Skurray, R . (1987):** Antimicrobial resistance in *Staphylococcus aureus*: genetic basis. Microbiol Reviews 51:88.
- **Machov, I. ; Petr, š. P. ; Blažkov, E. ; Kňapov, L. ; Sledov, n.; genů, k. and duj.c.ch.(2007):** Pantónův-Valentinův leukocidin u kmenů *Staphylococcus aureus*. Epidemiol Mikrobiol Imunol. 56(2):88-93.
- **Maguire, G. P.; Arthur, A. D.; Boustead, P. J.; Dwyer, B. and Currie, B. J. (1998):** Clinical experience and outcomes of community-acquired and nosocomial methicillin-resistant

- Staphylococcus aureus* in a northern Australian hospital. J. Hosp. Infect. 38:273-281.
- **Manzella, J.P. (2001):** "Quinupristin-dalfopristin: a new antibiotic for severe gram-positive infections.", Am Fam Physician. 1;64(11):1863-6. PMID 11764864.
 - **Marples, R. R. and Reith, S. (1992):** Methicillin-resistant *Staphylococcus aureus* in England and Wales. Communicable Disease Report, Public Health Laboratory Service Communicable Diseases Surveillance Center, 61 Colindale Avenue, London 2: R25-29.
 - **Marrack, P. and Kappler, J. (1990):** The staphylococcal enterotoxins and their relatives. Science 248:705-711.
 - **Massanari, R.M. ;Pfaller, M.A. ;Wakefield, D.S. ;Hammons, G.T. ;McNutt, L.A. ; Woolson, R.F. and Helms, C.M. (1988):**Implications of acquired oxacillin resistance in the management and control of *Staphylococcus aureus* infections. J. Infect. Dis., 158(4):702-9.
 - **Matsuzaki, S. ; Yasuda, M. ; Nishikawa, H. ; Kuroda, M. ; Ujihara, T. ; Shuin, T. ; Shen, Y. ; Jin, Z. ; Fujimoto, S. ; Nasimuzzaman, M.D. ;Wakiguchi, H. ; Sugihara, S. ;Sugiura, T. ; Koda, S. ; Muraoka, A. and Imai, S. (2003):** "Experimental protection of mice against lethal *Staphylococcus aureus* infection by novel bacteriophage phi MR11". J. Infect. Dis. 187 (4): 613–24. doi:10.1086/374001. PMID 12599078.
 - **Matthew, P.; Johnson; Larisa, M.; Haupt; Lyn, R. and Griffiths. (2004):** Locked nucleic acid (LNA) single nucleotide polymorphism (SNP) genotype analysis and validation using

real-time PCR. Nucleic Acids Research, Vol. 32, No. 6 e55 © 2004 Oxford University Press.

- **Matthews, K. R ; Roberson, J. ; Gillespie, B.E. ; Luther, D.A. and Oliver, S.P. (1997):** "Identification and Differentiation of Coagulase-Negative *Staphylococcus aureus* by Polymerase Chain Reaction". Journal of Food Protection 60 (6): 686–8.
- **MayoClinic staff, (2010): MRSA infection, edited from <http://www.mayoclinic.com/health/mrsa/DS00735/DSECTION=causes>. In: Feb. 24, 2010.**
- **Megson, G. M; Law, D.and Ganguli, L. A. (1991):** Problems of thermonuclease detection for identifying *Staphylococcus aureus* in blood culture broths. Journal of Clinical Pathology. 44:772-774.
- **Melish, M. ; Fukunaga, C. and Murata, S. (1998):** TSST-1 dissemination in a vaginal model for TSS, p. 169. In J. Arbuthnott, and B. Furman (ed.), European Conference on Toxic Shock Syndrome. International Congress and Symposium Series 229. Royal Society of Medicine Press Ltd, New York, N.Y.
- **Mellor, I. R. ; Thomas, D. H. and Sansom, M. S. P. (1988):** Properties of ion channels formed by *Staphylococcus aureus* - toxin. Biochim. Biophys. Acta. 942:280-294.
- **Merlino, J.; Leroi, M.; Bradbury, R.; Veal, D. and Harbour, C. (2000):** New chromogenic identification and detection of *Staphylococcus aureus* and methicillin-resistant *S. aureus*. J. Clin. Microbiol. 38:2378-2380.
- **Meyers, B.R.; Kaplan, K. and Weinstein, L. (1969):** "Microbiological and pharmacological behavior of 7-

- chlorolincomycin". Appl Microbiol 17 (5): 653–7. PMID 4389137. Full text at PMC: 377774.
- **Mifsud, A.J. and Parker, S.C. (1987):** Staphaurex reappraised: role of protein A in false positive results and recommendations for use. J Clin Pathol. 40(7):808–809.
 - **Miles, R.S. and Amyes, S.G.B. (1996):** Laboratory control of antimicrobial therapy. In: Collee, J. G.; Fraser, A. G.; Marmion, B.P. and Simmons, A. (eds.) Mackie & McCartney Practical Medical Microbiology. Churchill Livingstone. P: 151-178.
 - **Mitchell, C. J.; Geary, C. and Stevens, M. (1991):** Detection of *Staphylococcus aureus* in blood cultures: evaluation of two-hour method. Medical Laboratory Sciences. 48:106-109.
 - **Modric, J.(2008):** MRSA Superbug Types: CA-MRSA, HA-MRSA in Y. 2008. last modified: November 10th. Received from <http://www.healthhype.com/mrsa-superbug.html>.
 - **Monecke, S. ; Slickers, P. ; Ellington, M.J. ; Kearns, A.M. and Ehricht, R. (2007):** High diversity of Pantón-Valentine leukocidin-positive, methicillin-susceptible isolates of *Staphylococcus aureus* and implications for the evolution of community-associated methicillin-resistant *S.aureus*. Clin Microbiol Infect. 13(12):1157-64.
 - **Mongkolrattanothai, K.; Boyle, S.; Kahana, M.D. and Daum, R.S. (2003):** Severe *Staphylococcus aureus* infections caused by clonally related community-associated methicillin-susceptible and methicillin-resistant isolates. Clin. Infect. Dis. 37 (8): 1050–8. doi:10.1086/378277. PMID 14523769.

- **Moran, G. J., A. ; Krishnadasan, R. J. ; Gorwitz, G. E. ; Fosheim, L. K.; McDougal, R. B.; Carey and Talan, D. A.(2006):** Methicillin-resistant *S. aureus* infections among patients in the emergency department. *N. Engl. J. Med.* 355:666-674.
- **Morrison, T.B.; Weis, J.J. and Wittwer, C.T. (1998):** Quantification of low-copy transcripts by continuous SYBR Green I monitoring during amplification. *Biotechniques.* 24(6):954-8, 960, 962.
- **Murakami, K.; Minamide, W.; Wada, K.; Nakamura, E.; Teraoka H. and Watanabe, S. (1991):** Identification of methicillin resistant strains of staphylococci by polymerase chain reaction. *J. Clin. Microbiol.* 29:2240-2244.
- **Nada, T. ; Ichiyama, S. ; Iinuma, Y. ; Inuzuka, K. ; Washida, H. ; Ohta, M. ; Shimokata, K. ; Kato, N. and Nakashima, N. (1996):** Types of methicillin- resistant *Staphylococcus aureus* associated with high mortality in patients with bacteremia. *Eur. J. Clin. Microbiol. Infect. Dis.* Apr. 15(4):340-343.
- **Nakatomi, Y. and Sugiyama, J. (1998):** *Microbiol. Immunol.*; 42:739-743.
- **National Nosocomial Infections Surveillance System. (2003):** National Nosocomial Infections Surveillance (NNIS) System report, data summary from January 1992 through June 2003, issued August 2003. *Am. J. Infect. Control* 31:481-498.
- **National Nosocomial Infections Surveillance System. (2004):** National Nosocomial Infections Surveillance (NNIS) System Report, data summary from January 1992 through June 2004, issued October 2004. *Am. J. Infect. Control* 32:470-485.

- **Neville, O. ; Billington, O. J. ; Kibbler, C. C. and Gillespie, S. H. (1991):** Methicillin resistant *Staphylococcus aureus* without clumping factor, protein A and DNase. *Lancet* 338:518.
- **Niwa, t. (2005):** *J. Microbiol. Methods*, 61, 251-60.
- **Noble, W .C . ; Virani, Z . and Cree, R .G. (1992):** Co-transfer of vancomycin and other resistance genes from *Enterococcus faecalis* NCTC 12201 to *Staphylococcus aureus*. *FEMS Microbiol Lett.* 72:195-8.
- **Noto, M. J. and Archer, G. L. (2006):** A subset of *Staphylococcus aureus* strains harboring staphylococcal cassette chromosome mec (SCCmec) type IV is deficient in CcrAB-mediated SCCmec excision. *Antimicrob. Agents Chemother.* 50:2782-2788.
- **Noto, Michael J.; Kreiswirth, Barry N.; Monk, Alastair B. and Archer, Gordon L. (2008):** Gene Acquisition at the Insertion Site for SCCmec, the Genomic Island Conferring Methicillin Resistance in *Staphylococcus aureus*. *J Bacteriol.* 2008 February; 190(4): 1276–1283. Published online 2007 December 14. doi: 10.1128/JB.01128-07.
- **Novak, S. M.; Hindler, J. and Bruckner, D. A. (1993):** Reliability of two novel methods, Alarm and E test, for detection of methicillin-resistant *Staphylococcus aureus*. *Journal of Clinical Microbiology* 31: 3056-3057.
- **Novick, R. ; Lindsay, J. ; Ruzin, A. and Ross , H. (1998):** Molecular genetics of the test element, a mobile pathogenicity island in *Staphylococcus aureus*., p. 109-112. In J. Arbuthnott, and B. Furman (ed.), *European Conference on Toxic Shock*

- Syndrome. International Congress and Symposium Series 229. Royal Society of Medicine Press Ltd, New York, N.Y.
- **Novick, R. P. (2006):** Staphylococcal pathogenesis and pathogenicity factors: genetics and regulation, p. 496-516. In V. A. Fischetti, R. P. Novick, J. J. Ferretti, D. A. Portnoy, and J. I. Rood (ed.), Gram-positive pathogens, 2nd ed. ASM Press, Washington, DC.26.
 - **Nsira, S.B.; Dupuis, M. and Leclercq, R. (2006):** Evaluation of MRSA Select, a new chromogenic medium for the detection of nasal carriage of methicillin-resistant *Staphylococcus aureus*. Antimicrob Agents Int. J. 27:561-4.
 - **Ogston, A. (1984):** "On Abscesses". Classics in Infectious Diseases". Rev Infect Dis 6 (1): 122–28. PMID 6369479.
 - **Okuma, K.; Iwakawa, K. and Turnidge, J. (2002):** Dissemination of new methicillin-resistant *Staphylococcus aureus* clones in the community". J Clin Microbiol. 40 (11): 4289–4294.
 - **Olive, D.M. and Bean, P. (1999):** Principles and Applications of methods of DNA-based typing of microbial organisms. J Clin Microbiol. 37: 1661-9.
 - **Oliveira, D.; Sanches, I. S.; Tamayo, M.; Ribeiro, G.; Maro, R.; Costa, D. and de Lencastre, H. (1998):** Virtually all MRSA infections in the largest Portuguese hospital are caused by two internationally spread multiresistant strains: the “Iberian” and the “Brazilian” clones of MRSA. Clin. Microbiol. Infect. 4:373-384.
 - **Oliveira, D. C., Milheirico, C. and de Lencastre, H. (2006):** Redefining a structural variant of staphylococcal cassette

chromosome mec, SCCmec type VI. Antimicrob. Agents Chemother. 50:3457-3459.

- **O'Reilly, M. ; Kreiswirth, B. N. and Foster, T. J. (1990):** Molecular analysis of a Non expressed toxin gene (hla) of clinical isolates of *Staphylococcus aureus*, p. 439-443. In R. P. Novick (ed.), Molecular biology of the staphylococci. VCH Publishers Inc, New York, N.Y.
- **Orwin, P.M. ; Fitzgerald, J. R. ; Leung, D.Y ; Gutierrez, J. A ; Bohach , G.A and Schlievert , P.M. (2003):** Characterization of *Staphylococcus aureus* enterotoxin L. Infect Immun. 71:2916–9.
- **Papakyriacou, H.; Vaz, D.; Simor, A.; Louie, M. and McGavin, M.J. (2000):** Molecular analysis of the accessory gene regulator (agr) locus and balance of virulence factor expression in epidemic methicillin-resistant *Staphylococcus aureus*, J Infect Dis 181. pp. 990–1000.
- **Parfitt, T. (2005):** "Georgia: an unlikely stronghold for bacteriophage therapy". Lancet 365 (9478): 2166–7. doi:10.1016/S0140-6736(05)66759-1. PMID 15986542.
- **Parsonnet, J. (1998):** Case definition of staphylococcal TSS: A proposed revision incorporating laboratory findings, p. 15. In J. Arbuthnott, and B. Furman (ed.), European Conference on Toxic Shock Syndrome. International Congress and Symposium Series 229. Royal Society of Medicine Press Ltd, New York, N.Y.
- **Passet, L.A. (1985):** The Staphylococci. In: John Wiley and Sons. Cohen, J.O. (eds). Inc. New York, London. P. 1-20.

- **Payne, M.C. ; Wood, H.F. ; Karakawa, W. and Gluck, L.(1966):**
A prospective study of staphylococcal colonization and infections in newborns and their families. Am J Epidemiol;82:305-16.
- **Peng, H. L. ; Novick, R. P. ; Kreiswirth, B. ; Kornblum, J. and Schlievert , P. (1988):** Cloning, characterization, and sequencing of an accessory gene regulator (agr) in *Staphylococcus aureus*. J. Bacteriol. 170:4365-4372.
- **Perry, J. D.; Davies, A. ; Butterworth, L. A.; Hopley, A. L. J.; Nicholson, A. and Gould, F. K. (2004):** Development and evaluation of a chromogenic agar medium for methicillin-resistant *Staphylococcus aureus*. J. Clin. Microbiol. 42 (10):4519-4523.
- **Pfizer, (a b c d e f g h i j k. (June 20, 2008). "ZYVOX (linezolid) Label Information"(PDF).**<http://www.fda.gov/cder/foi/label/2008/021130s016,021131s013,021132s014lbl.pdf>. Recieved on 2008-08-24.
- **Pirisi, A. (2000):** "Phage therapy—advantages over antibiotics?". Lancet 356 (9239): 1418. doi:10.1016/S0140-6736(05)74059-9. PMID 11052592.
- **Prasad, G. S. ; Radhakrishnan, R. ; Mitchell, D. T. ; Earhart, C. A. ; Dinges, M. ; Cook, W. J. ; Schlievert, P. M. and Ohlendorf, D. H. (1997):** Refined structures of three crystal forms of toxic shock syndrome toxin-1 and of a tetramutant with reduced activity. Protein Sci. 6:1220-1227.
- **Prevost, G. ; Cribier, B. ; Couppie, P. ; Petiau, P. ; Supersac, G. ; Finck- Barbancon,V. ; Monteil, H. and Piemont, Y.(1995):**

- Panton-Valentine leukocidin and gamma hemolysin from *Staphylococcus aureus* ATCC 49775 are encoded by distinct genetic loci and have different biological activities. *Infect. Immun.* 63:4121-4129.
- **Projan, S. J. ; Kornblum, J. ; Kreiswirth, B. ; Moghazeh, S. L. ;Eisner, W. and Novick, R. P. (1989):** Nucleotide sequence: the β –hemolysin gene of *Staphylococcus aureus*. *Nucleic Acids Res.* 17:3305.
 - **Projan, S. J. and Novick, R. P. (1997):** The molecular basis of pathogenicity, p. 55-81. In K. B. Crossley and G. L. Archer (ed.), *The staphylococci in human disease*. Churchill Livingstone, New York, N.Y.
 - **Recsei, P. ; Kreiswirth, B. ; O'Reilly, M. ; Schlievert, P. ; Gruss, A. and Novick, R. P. (1986):** Regulation of exoprotein gene expression in *Staphylococcus aureus* by agr. *Mol. Gen. Genet.* 202:58-61.
 - **Rice, L. B. (2006):** Antimicrobial resistance in gram-positive bacteria. *Am. J. Infect Control* 34:S11-S199, S64-S73.
 - **Riza, A. ;Yasar, N. ; Zeynep, C. K. ; Cihan, T. ; Fatma, K.(2008):** Comparison of polymerase chain reaction and conventional methods in detecting methicillin-resistant *Staphylococcus aureus*. *J Infect Developing Countries* 2008; 2(1):46-50.
 - **Rogers, H. J. ; Perkins, H. R. and Ward, J. B. (1980):** Microbial cell walls and membranes. Chapman and Hall, London, United Kingdom.
 - **Rogers, H . J. (1983):** Bacterial Cell Structure. American Society for Microbiology, Washington, D.C.

- **Ross, S. ; Rodroquez, W. ; Controni, G. and Khan, W.(1974):**
Staphylococcal susceptibility to penicillin G: changing pattern among community isolates. JAMA. 229:1075-7.
- **Rossney, A. S.; English, L. F. and Keane, C. T. (1990):** Coagulase testing compared with commercial kits for routinely identifying *Staphylococcus aureus*. Journal of Clinical Pathology. 43:246-252.
- **Rubin, R. J. ; Harrington, C. A. ; Poon, A. ; Dietrich, K. ; Greene, J. A. & Moiduddin, A. (1999):**The economic impact of *Staphylococcus aureus* infection in New York City hospitals. Emerg Infect Dis 5, 9–17.
- **Sadaka, S. M.; El-Ghazzawy, E. F.; Harfoush, R. A. and Meheissen, M.A. (2009):** Evaluation of different methods for the rapid diagnosis of methicillin-resistance in *Staphylococcus aureus*. African Journal of Microbiology Research Vol. 3 (2) pp. 049-055 February, 2009.
- **Sadoyama, G. and Gontijo-Filho, P.P. (2000):** Risk factors for methicillin-resistant and sensitive *Staphylococcus aureus* infection in Brazilllian University Hospital. Braz. J. Infect. Dis. 4(3):135-143.
- **Safdar, N.; Narans, L.; Gordon, B. and Maki, D. G. (2003):** Comparison of culture screening methods for detection of nasal carriage of methicillin-resistant *Staphylococcus aureus*: a prospective study comparing 32 methods. J. Clin. Microbiol. 41:3163-3166.
- **Sails, A.D. (2009):** "Applications in Clinical Microbiology". Real-Time PCR: Current Technology and Applications. Caister Academic Press. ISBN 978-1-904455-39-4.

- **Sakon, J . ; Liao, H . H . ; Kanikula, A . M . ; Benning, M . M . ; Rayment, I . and Holden, H . M . (1993):** "Molecular structure of kanamycin nucleotidyltransferase determined to 3.0-Å resolution". *Biochemistry* 32 (45): 11977–84. PMID 8218273.
- **Salton, M. R. J. (1994):** The bacterial cell envelope—a historical perspective, p. 1-22. In J.-M. Ghuysen, and R. Hakenbeck (ed.), *Bacterial cell wall*. Elsevier Science, Amsterdam, The Netherlands.
- **Sanches, I.S . ; Saravaiva, Z .C . ; Tendeiro, T .C . ; Serra, J .M . ; Dias , D .C . and de Lencastre, H. (1998):** Extensive intra-hospital spread of a methicillin-resistant *Staphylococcus aureus* clone. *Int J Infect Dis.* 3:26–31.
- **Sanford , M. D. ; Widmer , A.F. ; Bale, M. J. ; Jones , R. N. and Wenzel, R. P. (1994):** Efficient detection and long-term persistence of the carriage of methicillin-resistant *Staphylococcus aureus*. *Clin Infect Dis.* 19:1123-8.
- **Sau, S. ; Bhasin, N. ; Wann, E. R. ; Lee, J. C. ; Foster, T. J. and Lee, C. Y. (1997):** The *Staphylococcus aureus* allelic genetic loci for serotype 5 and 8 capsule expression contain the type-specific genes flanked by common genes. *Microbiology* 143:2395-2405.
- **Savas, L.; Duran, N.; Onlen, Y.; Savas, N. and Erayman, M. (2005):** Prospective analysis of antibiotic susceptibility patterns of MRSA in a Turkish University Hospital. *Turk. J. Med. Sci.*, 35: 323-327.
- **Schito, G.C. (2006):** "The importance of the development of antibiotic resistance in *Staphylococcus aureus*". *Clin*

- Microbiol Infect 12 Suppl 1: 3–8. doi:10.1111/j.1469-0691.2006.01343.x. PMID 16445718.
- **Schwalbe, R . S .; Stapleton, J .T . and Gilligan, P .H. (1987):** Emergence of vancomycin resistance in coagulase-negative staphylococci. N Engl. J. Med. 316:927-31.
 - **Seifert, H. ;Wisplinghoff, H. ;Schnabel, P. and von Eiff, C. (2003):** Small colony variants of *Staphylococcus aureus* and pacemaker-related infection. Emerg Infect Dis. 9:1316–8.
 - **Sensi, P.; Margalith, P. and Timbal, M.T. (1959):** "Rifomycin, a new antibiotic-preliminary report". Farmaco Ed Sci 14: 146–147.
 - **Shockman, G. D. and Höltje, J.-V. (1994):** Microbial peptidoglycan (murein) hydrolases, p. 131-166. In J.-M.
 - **Simor, A. E.; Goodfellow, J.; Louie, L. and Louie, M. (2001):** Evaluation of a new medium, oxacillin resistance screening agar base, for the detection of methicillin-resistant *Staphylococcus aureus* from clinical specimens. J. Clin. Microbiol. 39:3422 .
 - **Skov, R. ; Smyth, R. ; Clausen, M. ; Larsen, A. R. ; Frimodt-Möller, N. ; Olsson-Liljequist, D. and Kahlmeter G. (2003):** Evaluation of a cefoxitin 30 µg disc on IsoSensitest agar for detection of methicillin-resistant *Staphylococcus aureus*. J. Antimicrob. Chemother. 52:204-207.
 - **Skov, R. ; Smyth, R. ; Larsen, A. R. ; Frimodt-Möller, N. and Kahlmeter, G. (2005):** Evaluation of cefoxitin 5 and 10 µg discs for the detection of methicillin resistance in staphylococci. J. Antimicrob. Chemother. 55:157-161.

- **Skulnick, M.; Simor, A.E.; Gregson, D.; Patel, M. and Small, G.W. (1992):** Evaluation of commercial and standard methodology for determination of oxacillin susceptibility in *Staphylococcus aureus*. J. Clin. Microbiol., 30: 1985-1988.
- **Slatore, C.G. (2004):** "Sulfonamide hypersensitivity". Immunology and Allergy Clinics of North America 24 (3): 477–90,vii. doi:10.1016/j.iac.2004.03.011.
- **Smyth, R.W. and Kahlmeter, G. (2005):** Mannitol Salt Agar-Cefoxitin Combination as a Screening Medium for Methicillin-Resistant *Staphylococcus aureus*. J Clin Microbiol 43: 3797–3799.
- **Solomon, J. J and San Clemente, C. L.(1963):** Some physiological characteristics of two sets of phage-propagating strains of *Staphylococcus aureus*. Appl. Microbiol. 11:36-41.
- **Soriano, A. ; Martinez, J.A. ; Mensa, J. ; Marco, F. ; Almela, M. and Moreno-Martinez, A. (2000):** Pathogenic significance of methicillin resistance for patients with *Staphylococcus aureus* bacteremia. Clin Infect Dis. 30:368-73.
- **Spreitzer, H. (2007):** "Neue Wirkstoffe - Incyclinid" (in German). Österreichische Apothekerzeitung (14/2007): 655.
- **Stamper, P.D.; Cai, M.; Howard, T.; Speser, S. and Carroll, K.C. (2007):** Clinical Validation of the Molecular BD GeneOhm StaphSR Assay for Direct Detection of *Staphylococcus aureus* and Methicillin-Resistant *Staphylococcus aureus* in Positive Blood Cultures. J Clin Microbiol. 45(7):2191-6.
- **Stenhem, Mikael ; Örtqvist, Åke; Ringberg, Håkan ; Larsson, Leif ; Olsson-Liljequist, Barbro ; Hæggman, Sara ; Kalin, Mats and Ekdahl, Karl (2010):** Imported Methicillin-

Resistant *Staphylococcus aureus*, Sweden. EID Journal Home
Volume 16, Number 2-February 2010.

- **Stoakes, L. ; Reyes, R. ; Daniel, J. ; Lennox, G. ; John, M.A. ; Lannigan, R. and Hussain, Z. (2006):** Prospective comparison of a new chromogenic medium, MRSASelect, to CHROMagar MRSA and mannitol–salt medium supplemented with oxacillin or ceftiofur for detection of methicillin-resistant *Staphylococcus aureus*, J. Clin. Microbiol. pp. 637–639. Vol. 44, No. 2
- **Sugai , M. ; Yamada, S. ; Nakashima, S. ; Komatsuzawa, H. ; Matsumoto, A. ; Oshida , T. and Suganaka, H.(1997):** Localized perforation of the cell wall by a major autolysin: atl gene products and the onset of penicillin-induced lysis of *Staphylococcus aureus*. J. Bacteriol. 179:2958-2962.
- **Suggs, A. H.; Maranan, M. C.; Boyle-Vavra, A. and Daum, R. S. (1999):** Methicillin-resistant and borderline methicillin-resistant asymptomatic *Staphylococcus aureus* colonization in children without identifiable risk factors. Pediatr. Infect. Dis. J. 18:410-414.
- **Sundin, D.P.; Sandoval, R. and Molitoris, B.A. (2001):** Gentamicin Inhibits Renal Protein and Phospholipid Metabolism in Rats: Implications Involving Intracellular Trafficking. J Am Soc Nephrol 12:114-123.
- **Swaminathan, S. ; Furey, W. ; Pletcher, J. and Sax, M . (1992):** Crystal structure of staphylococcal enterotoxin B, a superantigen. Nature (London) 359:801-806.

- **Swenson, J.M. and Tenover, F.C. (2005):** Results of disc diffusion testing with cefoxitin correlate with presence of *mecA* in *Staphylococcus* spp. *J.Clin. Microbiol.* 43: 3818-3823.
- **Swinscow, T.D.V. (1997):** Statistics at Square One. BMJ Publishing Group. Engemann, J. J. ; Carmeli, Y. ; Cosgrove, S. E. ; Fowler, V. G. ; Bronstein, M. Z. ; Trivette, S. L. ; Briggs, J. P. ; Sexton, D. J. and Kaye, K. S. Adverse clinical and economic outcomes attributable to methicillin resistance among patients with *Staphylococcus aureus* surgical site infection. *Clin. Infect. Dis.* 36:592-598.
- **Tenover Fred, C. ; Robert Arbeit ; Gordon Archer ; James Biddle ; Sean Byrne ; Richard Goering ; Gary Hancock ; Ann Hébert, G. ; Bertha Hill ; Richard Hollis ; William R. Jarvis ; Barry Kreiswirth ; William Eisner ; Joel Maslow ; Linda K. MacDougall ; Michael Miller, J. ; Maury Mulligan and Michael A. Pfaller. (1994):** Comparison of Traditional and Molecular methods of typing isolates of *Staphylococcus aureus*. *JOURNAL OF CLINICAL MICROBIOLOGY* Feb. P.407-415 vol.32, No.2.
- **Thompson, R .L , ; Cabezudo, I . and Wenzel, R .P. (1982):** Epidemiology of nosocomial infections caused by methicillin-resistant *Staphylococcus aureus*. *Ann Intern Med.* 97:309-17.
- **Thornsberry, C. and McDougal, L. K. (1983):** Successful use of broth microdilution in susceptibility tests for methicillin-resistant (heteroresistant) staphylococci. *Journal of Clinical Microbiology.* 18:1084-1091.

- **Toku, Y.; Shoji, S.; Satoh, K.; Watanabe, A. and Motomiye, M. (1991):** Detection of methicillin resistance *staphylococcus aureus* using polymerase chain reaction amplification. Tohokue. J. Exp. Med. P:31-37.
- **Tomasz, A.; Drugeon, H.B.; de Lencastre, H.M.; Jabes, D.; McDougall, L. and Bille, J. (1989):** New mechanism for methicillin resistance in *Staphylococcus aureus*:clinical isolates that lack the PBP2a gene and contain normal penicillin-binding protein with modified penicillin-binding capacity. Antimicrob. Agents Chemother., 33:1869-1874.
- **Trindade, P.A.; Mc Culloch, J.A.; Oliveira, G. A. and Mamizuka E.M. (2003):** Molecular Techniques For MRSA Typing: Current Issues and Perspectives. The Brazilian Journal of Infectious Diseases. 7(1):32-43.
- **Valeva, A. ; Palmer, M. and Bhakdi , S.(1997):** Staphylococcal exotoxin:formation of the heptameric pore is cooperative and proceeds through multiple intermediate stages. Biochemistry 36:13298-13304.
- **Van Enk, R. A. and Thompson, K. D. (1992):** Use of a primary isolation medium for recovery of methicilin-resistant *Staphylococcus aureus*. Journal of Clinical Microbiology. 30:504-505.
- **van Hal, S.J. ; Stark, D. ;Lockwood, B. ; Marriott, D. and Harkness, J. (2007):** Methicillin-Resistant *Staphylococcus aureus* (MRSA) Detection:Comparison of Two Molecular Methods (IDI-MRSA PCR Assay and GenoType MRSA Direct PCR Assay) with Three Selective MRSA Agars (MRSA ID, MRSASelect, and CHROMagar MRSA) for Use

- with Infection-Control Swabs. J Clin Microbiol. 45(8):2486-90.
- **van Loo, I.H. ; van Dijk, S. ; Verbakel-Schelle, I. and Buiting, A.G.(2007):** Evaluation of a chromogenic agar (MRSASelect) for the detection of methicillin-resistant *Staphylococcus aureus* with clinical samples in The Netherlands. J Med Microbiol. 56(Pt 4):491-4.
 - **Vandenesch, F. ; Naimi, T. and Enright, M. C. (2003):** Community- acquired methicillin-resistant *Staphylococcus aureus* carrying Panton- Valentine leukocidin genes: worldwide emergence. Emerging Infectious Diseases. 9, 978–84.
 - **Velasco, D.; Tomas, M. M.; Cartelle, M. and Beceiro, A. (2005):** Evaluation of different methods for detecting methicillin (oxacillin) resistance in *Staphylococcus aureus*. J. Clin. Microbiol.55:379-382.
 - **Wadlvogel, F.A. (2000):** *Staphylococcus aureus* (including staphylococcal toxic shock). In: Mandell GL, Bennett JE, Dolin R, editors. Principles and practice of infectious diseases. 5th ed. Philadelphia: Churchill Livingstone. p.2072-3 .
 - **Wang, Jann-Tay ; Fang, Chi-Tai ; Chen, Yee-Chun ; Wu, Chia-Ling ; Chen, Mei-Ling and Chang, Shan-Chwen.(2007):** Staphylococcal Cassette Chromosome mec in MRSA, Taiwan. Emerging Infectious Diseases Journal, Volume 13, Number 3–March 2007.

- **Watt, B. ; Rayner, A. and Harris, G. (1993):** Modern Methods in mycobacteriology. Reviews in Medical Microbiology 4: 97-105.
- **Weller, T.M.A.(2000):** Methicillin –resistant *Staphylococcus aureus* typing methods: which should be the international standard? J Hosp Infect. 44:160-72.
- **Wilkerson, M.; McAllister, S.; Miller, J. M.; Heiter, B.J. and Bourbeau, P .P. (1997):** Comparison of five agglutination tests for identification of *Staphylococcus aureus*. J Clin Microbiol. 35:148–151.
- **Williams, R.E. and Harber, G.J. (1964):** Determination of coagulase and alpha. Hemolysin production by staphylococci. Brit. J. Eup. Path. 27:72.
- **Woodin, A. M.(1959):** Fractionation of the two components of leukocidin from *Staphylococcus aureus*. Biochem. J. 73:225-237.
- **Woodin, A. M. (1960):** Purification of the two components of leukocidin from *Staphylococcus aureus*. Biochem. J. 75:158-165.
- **Wright, H. T. and Reynolds, K.A. (2007):** Antibacterial targets in fatty acid biosynthesis. Current Opinion in Microbiology doi:10.1016/j.mib.2007.07.001.

- **York, M.K.; Gibbs, L.; Chehab, F. and Brooks, G.F. (1996):**
Comparison of PCR detection of mec A with standard susceptibility testing methods to determine methicillin resistance in coagulase negative staphylococci. J. Clin. Microbiol. 34:249-253.

- **Zetola, N. ; Francis, J.S. ; Nuermberger,E.L. ; Bishai,W.R.(2005):**
Community-acquired meticillin-resistant *Staphylococcus aureus*: an emerging threat. The Lancet infectious diseases. 5:275–286. doi: 10.1016/S1473-3099(05)70112-2.