

REFERENCES

AbdEl-Aziz, K. B .and Mustafa, H. N. (1989):

Toxic effects of pireprofen on chromosome and blood criteria in mice.

Vet. Med. J. Giza, 37 (1):33-45.

Avers, C.J. (1980):

Genetics.

Chapter 11. PP. 417 – 418. D.

Van Nastrand and Company, New York.

Aviado, D. M. (1972):

Pharmacologic principals of medical practice

A text book on pharmacology and therapeutics for students and practitioners of medicine pharmacy.

Chapter 53, P, 830-831

The Williams &Wilkins company/Baltimore.

Ayala, F. J. and Kiger, J. A., Jr. (1984):

Modern Genetics.

Second Edition, the Benjamin/ Cummings, London. Chapter21, PP. 688 – 729.

Balfour J.A (1996):

Lornoxicam: a review of its pharmacology and therapeutic potential in the management of painful and inflammatory conditions.

Drugs, 51: 639-57.

Bertram, G. Katzung (1997):

Basic and clinical pharmacology.

Text book.17th Edition.

Bianchi, -M, Panerai, -A- E (2002):

Effects of lornoxicam, piroxicam, and meloxicam in a model of thermal hindpaw hyperalgesia induced by formaline injection in rat tail. Pharmacol-Res. 45(2): 101-5

Brogger, A (1974):

Chromosomes damage in human mitotic cells after *in vivo* and *In vitro* exposure to mutagens.

Mutation Res. 23:353

Craig, C. R. and Stitzel, R.E., (1986):

Modern pharmacology, second edition .1029-1030.

Little, Brown and Company

Boston/Toronto.

Crippa, L., Klein, D. and Linder, A. (1976):

Cytogenetic and statistical study of a group of patients treated with Butazolidine (Phenylbutazone)

J. Genet.Hum.24 (1):1-13.

De-Hondt, H.A., El-Laithy, A.F. and Ali, S.M. (1981):

Biochemical and chromosomal studies on rats injected with captan Egypt. J. Genet. Cytol, 11: 245 – 253.

El-Nabi, E.S.(1981):

Physiological and karyyological studies on the effect of organophosphorous insecticide on male albino mice.

M.Sc.Thesis Zoology, Menoufia Univ.Faculty of Science, 180.

Evan, H.J.(1977):

Molecular mechanisms in the induction of chromosome aberration.

Scott, D.Bridges, B.A. and Sobier / North Holland,
Amsterdam, 57-74.

Ferreira, S.H.(1972):

Nature New Biol., 240:200-203.

Ferreira, S.H. and Vane, J.R.(1974)

A.Rev.Pharmac., 14: 57-73.

Fraser, G. and Mayo O.(1975):

Textbook of human genetic chapter 6.

Blackwell scientific publications

(Oxford, London, Edinburgh, Melbourne).

Gebhart, E. and Wissmuller, H.F. (1973):

Investigation on the effect of phenylbutazone on chromosomes and mitosis in the bone marrow of rats.

Mutation Research, 17, 283-286.

Goodwin, J.S. (1984):

Mechanism of action of None steroidal anti-inflammatory agents.

The American Journal of Medicine, 13:57-64.

Hartl, D.L., Ferifelder, D. and Snyder, L.A. (1987) :

Basic Genetics.

Chapter 6, PP. 155 – 178. Jones and Bartlett, Boston.

Khater, (1991):

Cytogenic study on the effect of some drugs on albino rat

M.Sc. Thesis Zoology, Benha Univ. Faculty of Science, p. 95-96.

Kihlman, B.A., Anerson, H.C. and Nataragon, A.T. (1977):

Molecular mechanism in the production of chromosomal aberrations : studies with the 50 bromodeoxyuridine.

Labelling method, in dela chapella, A. and Sorsa, M. (eds)

Chromosomes today. Vol. 6, El Sevier.

North Holland, Amsterdam, 287-296.

Klein,G.and Wattawa,A.(1975):

Der Einflub,sogenonnter basis therapeutika und symptomatisch
Wirksomer antirhematika ouf enzyme der DNA-synthesis und
DNA-reparatur.

Acata Med.Austriaca,J.2:153-156.

Kullich ,W.and Klien ,G.(1986):

Investigation of the influence of none-steroidal anti-rheumatic drugs
on the rates of sister chromatid exchange.

Mutation research, 131-134.

Leonard, A. (1976):

Heritable chromosome aberrations in mammals after exposure to
Chemicals.

Rad.And Environm.Biophys.13:1-8.

Loscher, w. and Blazaki, d. (1986):

Effect of NSAIDs on fertility of male rats

J. reprod .fert.69: 65-73.

Court Brown ,W.M.,Jacobs,P.A,Buckton ,K.E,

Loscher, w., luttgenau, tl. , schlegel, w. and kruiger, s. (1988):

Pharmacokinetics of NSAIDs in male rabbits after acute and chronic administration and effects of chronic treatment on seminal PGs, sperm quality and fertility
J. reprod. fert. 82: 353-364.

Mastuda, k. ohnishi, k., misaka, e. and yamazaki, m. (1983):

Decrease of urinary PG E2 and PG E2alpha excretion by NSAIDs in rats) biochem. Pharmacol, 32, 1347-1352.
Cited by Losher, w. et al. (1988).

Mauer,E.F.(1955):

The toxic effects of phenylbutazone(Butazidine)
New England Journal of Medicine,253:404-410.

Muller,D.and Strasser,F.F.(1971):

Comparative studies on the Chinese Hamster bone marrow after treatment with phenylbutazone and cyclophosphamide.
Mutation Research,13:337-382.

Neal, M, (1992):

in (Medical Pharmacology at a glance) Second edition.,
Neal, M. (ed.) Blackwell Scientific publications, P.66.

O' Brien ,W.M.(1983):

Pharmacology of NSAIDs.

The American journal of Medicine.

Practical review for clinicians,31: 32-39.

Pai,A.C.(1985):

Foundations of genetics .

A Science for Society second edition.Chapter six.

Mc Graw,Hill Book company,New York,Hamburg,New Delhi.

Perpignano,G.,Cacace,E.,Matulli,C.and Ruggiero,V.(2003):

Sicurezza terapeutica di lornoxicam in soggetti G-6-PDH carenti.

(Safety of lornoxicam in G-6-PDH deficiency)

Reumatismo. 55(1):45-7

Phillips, D.M.(1974):

(Sprmatogenesis). Academic Press.Inc.fifth Avenue,New York.

U.K.Edition publishd by Academic Press.Inc,(London)LTD.

Rathenberg,R.and Muller,D. (1972):

Comparative cytogenetic studies of the influences of
phenylbutazone and cyclophosphamide on spermatogenesis in the
mouse.

Agents and actions,2(4):180-185.

Schorderet, M. (1976):

J.Exp.Med., 126:171. Quated from: Arrigoni-Martelli, E. (1977 b).

Skjodt N.M.and Davies, N.M.(1998):

Clinical pharmacokinetics of Lornoxicam:a short halfe-life
Oxicam .clin pharmacokinet. 34:421-428.

Smith,A.L.,Lodge,J.P.and Pandleink,R.P.(1979):

In Vitro effects of phenylbutazone on equine chromosome.
Veterinary Record, 104 (9):195-196.

Snedecor, G.W. (1946):

Statistical methods.
Iowa state college press, Amer. Iowa.4 th Edition.

Sorsa,M.(1992):

Human cytogenetic damage as predictor of cancer risk.In
Vainiq,H.,Magee, P.,Gregore,D.and Cheal,A.(Editor):mechanisms of
carcinogenesis in risk identification,4th ed.,Macmillan publishing
CO., New York,pp.543.

Staunstrup H. (1999):

Efficacy and tolerability of lornoxicam versus tramadol in
postoperative pain,J clin pharmacol ,39:834-841.

Stevenson,A.C.;Bedford,J.;Hill,A.G.S. and Hill,B.F.A.(1971):

Chromosomal studies in patients taking phenylbutazone.

Ann. Rheum. Dis., 30 : 487 – 500 .

Stoian,V.and Raicu,P.(1975):

Morphological characteristics and the nature of gaps (achromatic lesions)induced by 4-Aminouracil in chromosomes of Vicia faba.

Mutation research.,217-200.

El-Sevier Scientific publishing company,

Amsterdam-printed in th Netherlands.

Vormittage,W.(1975):

Induktian chromosomaler aberrationen duch.

Antirheumatische therapie.

Acta.Med.Austeria,J.2:148-153.

Walker,S.,Evans,D.A.P.,Benn,P.A.,Litter,T.R.and Holliday,D.C.(1975):

Phenylbutazone and chromosomal damage.

Annals Rheumatoid disease,34:409-415.

Wagner, R.P., Judd, B.H., Sanders, B.G. and Richardson,

R.H. (1980) :

Introduction to modern genetics .

John. Willy and Sons. New York.

Wissmuller, V.H.F.(1971):

Untersuchungen über die cytogenetische Wirkung von
Phenylbutazone an menschlichen Lymphocyten in vitro.
Arzneim-Forsch(Drug Res.) Jahrgang 21(11), 1938-1950.

Wyrobek, A.J. and Burce, W.R.(1975):

Chemical introduction of sperm abnormalities in mice.
Proc.Nat.Acad.Sci.USA, 72(11):4425-4429

Yosida, T.M. and Amano, K. (1965) :

Autosomal Polymorphism in laboratory bred and wild
Norway rats. Rattus norvegicus found in Misima
Chromosoma. 16: 658 – 667.

Zdzienicka, M., Hyniewicz, M. and Pienkowska, M.(1982):

Thiram induced sperm head abnormalities in mice.
Mutation, Res. 102:261-265.

Zowail, M.E.M. and Abu-Saba, M.A.(1992):

Cytogenetic evaluation of Feldene (piroxicam)
The Egyptian Society of Histology and Cytology (Cairo)
15:

Zubay, G. (1987) :

Genetics.

Chapter 9, PP. 308 – 357. The Benjamin/ Cummings, Menlo
Park, California.