

REFERENCES

- Agnese, J. F.; Adepo-Gourene, B.; Abban, E. K.; Fermon, Y. (1997):** Genetic differentiation among natural populations of the Nile tilapia *Oreochromis niloticus* (Teleostei, Cichlidae). **Heredity, (79): 88-96.**
- Ahmed, M. M.; Ali, B. A. and EL-Zaeem, S. Y. (2004):** Application of RAPD markers in fish: Part I- some genera (*Tilapia*, *Sarotherodon* and *Oreochromis*) and species (*Oreochromis aureus* and *Oreochromis niloticus*) of Tilapia. **Inter. J. Biotech., 6 (1): 86-93.**
- Albertson, R. C. and Kocher, T. D. (2001):** Assessing morphological differences in an adaptive trait: A landmark-based morphometric approach to Jaw morphology in lake Malawi Cichlid fish. **J. Exp. Zool., 282: 385-403.**
- Albertson, R. C.; Markert, J. A.; Danley, P. D. and Kocher, T. D. (1999):** Phylogeny of a rapidly evolving clade: The cichlid fishes of Lake Malawi, East Africa. **Proc. Nat. Acad. Sci. USA, 96: 5107-5110.**
- Albertson, R. C.; Streelman, J. T. and Kocher, T. D. (2003a):** Directional selection has shaped the oral jaws of Lake Malawi cichlid fishes. **Proc. Natl. Acad. Sci. USA, 100: 5252-5257.**
- Albertson, R. C.; Streelman, J. T. and Kocher, T. D. (2003b):** Genetic basis of adaptive shape differences in the cichlid head. **J. Heredity, 94 (4): 291-301.**

- Alrubaian, J.; Lecaude, S.; Barba, J.; Szynskie, L.; Jacobs, N.; Bauer, D.; Brown, C.; Kaminer, I.; Bagrosky, B. and Dores, R. M. (2006):** Trends in the evolution of the prodynorphin gene in teleosts: cloning of eel and tilapia prodynorphin cDNAs. **Peptides**, 27 (4): 797-804.
- Al-Zahaby, A. S.; El-Assy, N. S.; El-Attar, A. E. and Awwad, G. S. (1990):** Comparative developmental studies on the lateral canal system in *Cyprinus carpio* and *Gambusia affinis affinis*. **Quarter Univ. Sci. Bull.**, 10.
- Anderson, J. D.; Karel, W. J.; Anderson, K. A. and Roperfoo, P. A. (2008):** Genetic Assessment of Sheepshead stock structure in the Northern Gulf of Mexico: Morphological divergence in the face of gene flow. **North Amer. J. Fish. Manag.**, 28 (2): 592-606.
- Anene, A. (1999):** Morphometric and meristic description of *Tilapia mariae* (Boulenger, 1901) and *Tilapia zillii* (Gervais, 1848) from the Umuoseriche Lake in the freshwater reaches of the Niger delta floodplains. **Acta Hydrobiologica**, 41: 211-218.
- Argenton, F.; Ramoz, N.; Charlet, N.; Bernardint, S.; Colombo, L. and Bortolussi, M. (1996):** Mechanisms of transcriptional activation of the promoter of the rainbow trout prolactin gene by GHF1/Pit1 and glucocorticoid. **Biochem. Biophys. Res. Commun.**, 234: 57-66.
- Arnason, U.; Gullberg, A.; Janke, A.; Joss, J. and Elmerot, C. (2004):** Mitogenomic analyses of deep gnathostome divergences: A fish is a fish. **Gene**, 333: 61-70.

- Astola, A.; Ortiz, M.; Calduch-Giner, J. A.; Perez-sanchez, J. And Valdivia, M. M. (2003):** Isolation of *Sparus auratus* prolactin gene and activity of the *cis*-acting regulatory elements. **Gen. Comp. Endocrinol., 134: 57-61.**
- Awwad, M. H. (1995):** Small-subunit ribosomal RNA gene phylogeny of the opportunistic human pathogen *Acanthamodeba polyphaga*. **Ph. D. Thesis, Faculty of Science, Benha University, Egypt, 142 pp.**
- Awwad, M. H. and Morsy, G. H. (2001):** Small subunit Ribosomal RNA gene identification of the human pathogen *Schistosoma* sp. in Egypt. **Egypt. J. Aquat. Biol. & Fish., 5 (4): 281-294.**
- Azab, M. S. (2003):** Studies on some molecular biological aspects of Tilapia fish in Egypt. **M. Sc. Thesis, Faculty of Science, Benha University, Egypt, 166 pp.**
- Balarin, J. D. and Hatton, J. P. (1997):** Tilapia, a guide to their biology and culture in Africa. **Unit of Aquatic Pathobiology, University of Sterling, Scotland, U. K., 174 pp.**
- Baumsteiger, J. and Kerby, J. L. (2009):** Effectiveness of salmon Garcaass tissue for use in DNA extraction and amplification in conservation Genetic studies. **North Amer. J. Fish. Manag., 29 (1): 40-49.**
- Beacham, T. D. and Murray, C. B. (1986):** The effect of spawning time and incubation temperature in meristic variation in chum salmon (*Oncorhynchus keta*). **Can. J. Zool., 64: 45-48.**

- Beacham, T. D.; Candy, J. R.; McIntosh, B.; Connachie, C. M.; Tabata, A.; Miller, K. M. and Withler, R. E. (2005):** DNA-Level Variation of Sockeye Salmon in Southeast Alaska and the Nass and Skeena Rivers, British Columbia, with Applications to Stock Identification. **North Amer. J. Fish. Manag.**, 25 (3): 763-776.
- Beacham, T. D.; Wetklo, M. and Wallace, C. (2008):** The application of microsatellites for stock identification of Yukon River Chinook salmon. **North Amer. J. Fish. Manag.**, 28 (1): 283-295.
- Beccari, E. and Mazzetti, P. (1987):** The nucleotide sequence of the ribosomal protein L14 gene of *Xenopus laevis*. **Nucleic Acids Res.**, 15: 1870-1872.
- Berridge, B. R.; Bercovier, H. and Frelief, P. F. (2000):** *Streptococcus agalactiae* and *Streptococcus difficile* 16S-23S intergenic rDNA: genetic homogeneity and species-specific PCR. **Veterin Microbiol.**, 78: 165-173.
- Beveridge, M. C. M. and McAndrew, B. J. (2000):** "Tilapias: Biology & Exploitation". **Kluwer Academic Dordrecht**, 532 pp
- Blache, J. and Miton, F. (1960):** Poissons nouveaux du bassin du Tchad et du bassin adjacent du Mayo Kebbi. (Suite et fin). III. Cyprinodontidae Cichlidae. **Bull. Mus. Nat. Hist. Nat. (Sér. 2)**, 32 (3): 214-218.

- Bosworth, A. and Farrell, J. M. (2006):** Genetic Divergence among Northern Pike from Spawning Locations in the Upper St. Lawrence River. **North Amer. J. Fish. Manag.**, **26** (3): 676-684.
- Bouck, A. and Vision, T. (2007):** The molecular ecologist's guide to expressed sequence tags. **Mol. Ecol.**, **16** (5): 907-924.
- Boulenger, G. A. (1899):** (June) A revision of the African and Syrian fishes of the family Cichlidae. Part II. **Proc. Zool. Soc. Lond. (pt. 1):** 98-143.
- Bouvet, P.; Diaz, J. J.; Kindbeiter, K.; Madjar, J. J. and Amalric, F. (1998):** Nucleolin interacts with several ribosomal proteins through its RGG domain. **J. Biol. Chem.**, **273**: 19025-19029.
- B-Rao, C. and Majumdar, K. C. (1998):** Multivariate map representation of phylogenetic relationships: application to tilapiine fish. **J. Fish Biol.**, **25** (6): 1199-1217.
- Bruet, B. J. F.; Song, J.; Boyce, M. C. and Ortiz, C. (2008):** Materials design principles of ancient fish armour. **Nat. Mat.**, **7** (9): 1-9.
- Calderaro, A.; Piccolo, G.; Perandin, F.; Gorrini, C.; Peruzzi, S.; Zuelli, C.; Ricci, L.; Manca, N.; Dettori, G.; Chezzi, C. And Snounov, G. (2007):** Genetic polymorphisms influence *Plasmodium ovale* PCR Detection Accuracy. **J. Clin. Microbiol.**, **45**: 1624-1627.

- Carolsfeld, J.; Powell, J. F. F.; Park, M.; Fischer, W. H.; Craig, A. G.; Chang, J. P.; Rivier J. E. and Sherwood, N. M. (2000):** Primary structure and function of three gonadotropin-releasing hormones, including a novel forms, from an ancient teleost, herring. **Endocrinology**, **141 (2): 505-512.**
- Carpio, Y.; Acosta, J.; Morales, A.; Herrera, F., Gonzalez, L. J. and Estrada, M. P. (2006):** Cloning, expression and growth promoting action of Red tilapia (*Oreochromis* sp.) Neuropeptide Y. **Peptides**, **27 (4): 710-718.**
- Carrera, E.; Garcia, T.; Cespedes, A.; Gonzales, I.; Fernandez, A.; Asensio, L. M.; Hernandez, P. E. and Martin, R. (2000):** Differentiation of smoked *Salmo salar*, *Oncorhynchus mykiss* and *Brama rail* using the nucleolar marker 5S rDNA.. **Int. J. Food Sci. Tech.**, **35: 401-406.**
- Céspedes, A.; Garcia, T.; Carrera, E.; Gonzalez, I.; Sanz, B.; Hernandez, P. E. And Martin, R. (1998):** Identification of Flatfish Species Using Polymerase Chain Reaction (PCR) Amplification and Restriction Analysis of the Cytochrome *b* Gene. **J. Food Sci.**, **63(2): 206-209.**
- Céspedes, A.; Garcia, T.; Carrera, E.; Gonzalez, I.; Fernandez, A.; Hernandez, P. E. and Martin, R. (1999):** Identification of sole (*Solea solea*) and greenlanda halibut (*Reinhardtius hippoglossoides*) by PCR amplification of the 5S rDNA gene. **J. Agric. Food Chem.**, **47: 1046-1050.**
- Chen, Y.; Conway, C.; Keeler-Foster, C.; Hamman, R. and Meisner, S. (2009):** Genetic characterization of variation in

- Captive and Wild Woundfin. **North Amer. J. Fish. Manag.**, **29** (4): 43-849.
- Chow, S. and Inoue, S. (1993):** Intra-and interspecific restriction fragment length polymorphism in mitochondrial genes of *Thunnus tuna* species. **Bull. Nat. Res. Inst. Far Seas Fish.**, **30**: 207-224.
- Chow, S.; Clarke, M. E. and Walsh, P. J. (1993):** PCR-RFLP analysis on thirteen western Atlantic snappers (Subfamily Lutjaninae): A simple method for species and stock identification. **Fish. Bull.**, **91**: 619-627.
- Chu, S.; Weng, C.; Hsiao, C.; Hwang, P.; Chen, Y.; Ho, J. and Lee, S. (2006):** Profile analysis of expressed sequence tags derived from the ovary of tilapia, *Oreochromis mossambicus*. **Aquaculture**, **251** (2-4): 537-548.
- Colomba, M. S.; Vitturi, R. Castriota, L.; Bertoni, R. And Libertini, A. (2002):** Fish mapping of 185-285 and 5S ribosomal DNA, (GATA) and (TTAGGG) telomeric repeats in the peri winkle *Melarhaphe neritoides* (Prosobranchia, Gastropoda, Caenogastropoda). **Heredity**, **88** (5): 381-384.
- Cooper, J. E. (1981):** Development and replacement order of pharyngeal teeth in the golden shiner, *Notemigonus crysoleucas*. **Ohio, J. Sci.**, **81**(1): 14-18.
- Daget, J. (1954):** Les poisons du Niger Supérieur Mem. **Inst. Franc. Afr. Noire**, **36**: 1-391.

- De Rosa Sntos, L. V.; Foresti, F.; Wasko, A. P.; Oliveira, C. And Martins, C. (2006):** Nucleotide sequences, genomic orgnaization and chromosome localization of 5S rDNA in two species of Curimatidae (Teleostei, characifores). **Genet. Mol. Biol., 29 (2): 15-57.**
- Desdevises, Y.; Jovelín, R.; Jousson, O. and Morand, S. (2000):** Comparison of ribosomal DNA sequences of *Lamellodiscus* spp. (Monogenea, Diplectanidae) parasitizing *Pagellus* (Sparidae, Teleostei) in the North Mediterranean sea: species divergence and coevolutionary interactions. **J. Parasitol., 30: 741-746.**
- El-Serafy, S. S.; Abdel-Hamide, N. H.; Awwad, M. H. and Azab, M. S. (2007):** Comparative study on electrophoretic protein pattern of tilapia species in the River Nile, Egypt. **Proc. 7. Internat. Symp. Tilapia Aquacul. (ISTA7), Arizona University, 281-292.**
- El-Serafy, S. S.; Abdel-Hamide, N. H.; Awwad, M. H. and Azab, M. S. (2007):** DNA riboprinting analysis of *Tilapia* species and their hybrids using restriction fragment length polymorphisms of the small subunit ribosomal DNA. **Aquacul. Res., 38: 295-303.**
- El-Serafy, S. S.; Al-Zahaby, A. S.; Zowail, M. E.; Labib, W. and Badawy, E. A. (1992):** Comparative cytogenetic studies on two *Tilapia* species, from different localities. **Bull. Fac. Sci. Zagazig Univ., 14 (2).**

- El-Serafy, S. S.; Awwad, M. H.; Abdel-Hamide, N. H. and Azab, M. S. (2003):** Using restriction fragment length polymorphisms (RFLPs) of the small-subunit ribosomal DNA as a tool for identification of *Tilapia* spp. **Egypt. J. Aquat. Biol. & Fisheries**, 7 (4): 465-282.
- Esmaeili, H. R. and Gholami, Z. (2007):** Investigation on the surface ultrastructure of scale of Geno tooth-carp, *Aphanius ginaonis* (Holly, 1929) (Actinopterygii: Cyprinodontidae) using scanning electron microscope. **Iran. J. Boil.**, 20: 307-314.
- Esmaeili, H. R.; Ansari, T. H. and Teimory, A. (2007):** Scale structure of a cyprinid fish, *Capoeta damascina* (Valenciennes in Cuvier and Valenciennes, 1842) using scanning electron microscope (SEM). **Iran. J. Sci. & Technol. Trans.**, 31 (A3).
- Esmaeili, H. R.; Baghbani, S.; Zareian, H. and Shahryari, F. (2009):** Scale morphology of tank goby *Glossogobius giuris* (Hamilton-buchanan, 1822) (Perciformes: Gobiidae) using scanning electron microscope. **J. Boil. Sci.**, 9(8): 899-903.
- Falk, T. M.; Abban, E. K.; Oberst, S.; Villwock, W.; Pullin, R. S. V. and Renwranztz, L. (1996):** A Biochemical laboratory manual for species characterization of some Tilapine fishes. **ICLARM Educ. Ser.**, 17: 93 pp.
- FAO (1980):** Report on the adhoc consultation on aquaculture research. **FAO Fish. Rep.**, 238: 26 pp.

- FAO (1984):** News feature, Reshaping world fisheries. **Food Agric. Organ. Un. Nat.**, 191-193.
- Farahmand, H.; Azizur Rahman, M.; Sohm, F.; Hwang, G. and Maclean, N. (2003):** Isolation and expression of tilapia (*Oreochromis niloticus*) serine 8-type GnRH coding and regulatory sequences. **Gene**, 304: 97-106.
- Felsenstein, J. (1989):** Phylogenies from molecular sequences: inference and reliability. **Ann. Rev. Genet.**, 22: 521-565.
- Fernandez, A.; Garcia, T.; Asensio, L.; Rodreguez, M. A.; Gonzalez, I. Hernandez, P. E. and Martin, R. (2001):** PCR-RFLP analysis of the internal Transcribed Spacer (ITS) Region for Identification of 3 Clam Species. **J. Food Sci.**, 66(5): 657-661.
- Fishelson, L. (1984):** A comparative study of ridge-mazes on surface epithelial cell-membranes of fish scales (Pisces, Teleostei). **Zoomorphology**, 104: 231-238.
- Fitch, D. H. A.; Bugaj-Gaweda, B. and Emmons, S. W. (1995):** 18 Sribosomal RNA Gene Phylogeny for some Rhabditidae related to Caenorhabditis. **Mol. Biol. Evol.**, 12 (2): 346-358.
- Flannery, B. G.; Wenburg, J. K. and Gharrett, A. J. (2007):** Variation of Amplified Fragment Length Polymorphisms in Yukon River Chum Salmon: Population Structure and Application to Mixed-Stock Analysis. **Trans. Amer. Fish. Soc.**, 136 (4): 911-925.

- Franck, J. P. C.; Kornfield, I. and Wright, J. M. (1994):** The utility of SATA satellite DNA sequences for inferring phylogenetic relationships among the three major genera of tilapiine fishes. **Mol. Phyl. Evol.**, 3: 10-16.
- Fry, N. K.; Rowbotham, T. J., Saunders, N. A. and Embley, T. M. (1991):** Direct amplification and sequencing of the 16S ribosomal DNA of an intracellular legionella species recovered by amoebal enrichment from the sputum of a patient with pneumonia. **Fems microbiology letters**, 67 (2): 16-18.
- Fryer, G. and Iles, T. D. (1972):** The cichlid fishes of the great Lakes of Africa: their biology and evaluation. **T. F. H. Publ., Neptune City, New Jersey. Also Published by Oliver and Boyd, Edinburgh.**
- Gabr, S. A. and Sakr, S. A. (1992):** A scanning electron microscopic study on the scales of *Tilapia nilotica*. **J. Egypt. Ger. Soc. Zool.**, 7(c): 41-48.
- GAF (1997):** General authority of the development of fish resources. **Data on fish production in Egypt, Cairo, Geoarchaeology**, 18 (3): 395-402.
- Garclá-Varela, M.; Cummings, M. P.; Pérez-once de léon, G.; Gardner, S. L. and Laclette, J. P. (2002):** Phylogenetic analysis based on 18S ribosomal RNA gene sequences supports the existence of class polyacanthocephala (Acanthocephala). **Mol. Phylogen. Evol.**, 23 (2): 288-292.

- Gervais, F. L. P (1848):** Sur les animaux vertébrés de l' Algérie, Envisagés sous le double rapport de la géographie zoologique et de la domestication. **Ann. Sci. Nat., Paris (Zool.) (Sér. 3), 10: 202-208.**
- Gibert, Y.; Bernard, L. and Laudet, V. (2007):** Crucial roles of retinoic acid signaling during teeth induction in teleost fish. **Europ. Cell Mat., 14 (2): p6.**
- Ginisty, H.; Amalric, F. and Bouvet, P. (1998):** Nucleolin functions in the first step of ribosomal RNA processing. **EMBO J., 17: 1476-1486.**
- Ginisty, H.; Sicard, B.; Roger, P. and Bouvet, P. (1999):** Structure and functions of nucleolin. **J. Cell Sci., 112: 761-772.**
- Giribet, G.; Carranzo, S.; Riutort, M.; Baguna, J. and Riera, C. (1999):** Internal phylogeny of chilopoda (Myriapoda, Arthropoda) using complete 18S rDNA and partial 28S rDNA sequences. **Phil. Trans. Roy. Soc. Lond., 354: 215-222.**
- Goldstein, R. M. (2009):** Interpreting stream physical characteristics in index of biotic integrity classifications: General similarities and Specific differences. **North Amer. J. Fish. Manag., 29 (1): 151-162.**
- Gregory, T. R. (2001):** Animal Genome size Database. <http://www.genomesize.com>.
- Günther, A. (1859):** Catalogue of the fishes in the British Museum, 1: I-XX.

- Günther, A. (1862): Catalogue of the fishes in the British Museum, 4: I-XII +1-534 pp.**
- Günther, A. (1864):** Description of a labyrinthibranchiate fish from the Nile. **Ann. Mag. Nat. Hist. (Ser.3), 13 (75): 211 pp.**
- Gur, G.; Rosenfoeld, H.; Melamed, P.; Meiri, I., Elizur, A. and Yaron, Z. (2001):** *Tilapia* glycoprotein hormone α subunit: cDNA cloning and hypothalamic regulation. **Mol. Cell. Endocrinol., 182: 49-60.**
- Gyllensten, U. (1985):** The genetic structure of fish: differences in the intraspecific distribution of biochemical genetic variation between marine, anadromous, and freshwater species. **J. Fish Biol., 26: 691-699.**
- Habicht, C.; Seeb, L. W. and Seeb, J. E. (2007):** Genetic and Ecological Divergence Defines Population Structure of Sockeye Salmon Populations Returning to Bristol Bay, Alaska, and Provides a Tool for Admixture Analysis. **Trans. Amer. Fish. Soc., 136 (1): 82-94.**
- Hamilton, L. C.; Macpherson, G. R. and Wright, J. M. (2000):** Expressed sequence tags derived from brain tissue of *Oreochromis niloticus*. **J. Fish Biol., 56: 219-222.**
- Hanfling, B. and Brandl, R. (2000):** Phylogenetics of European cyprinids: insights from allozymes. **J. Fish Biol., 57: 265-276.**
- Harabawy, A. S. A. (2002):** Biological and taxonomic studies on some fish species of the genus *Lethrinus* (Family

- Lethrinidae) from the Red Sea, Egypt and the genus *Abramis* (Family Cyprinidae) from the Baltic Drainage. **Ph. D. Thesis, Faculty of science, Assiut University, Egypt, 294 pp.**
- Hariharan, N. and Perry, R. P. (1990):** Functional dissection of a mouse ribosomal protein promoter significance of the polypyrimidine initiator and an element in the TATA box region. **Proc. Natl. Acad. Sci. USA, 87: 1526-1530.**
- Haroun, M. R. (1999):** Cross-breeding Egyptian races of tilapias to produce more productive hybrids. **Ph. D. Thesis, Faculty of Science, Tanta University, Egypt, 193 pp.**
- Hassenkam, T.; Fantner, G. E.; Cutroni, J. A.; Weaver, J.; Morse, D. and Hansmaa, P. K. (2004):** High-resolution AFM imaging of intact and fractured trabecular bone. **Bone, 35: 4-10.**
- Hauser, L. and Seeb, J. E. (2009):** Advances in molecular technology and their impact on fisheries genetics. **Fish and Fisheries, 9 (4): 473-486.**
- Henry, S. D.; Barkley, S. W. and Johnson, R. L. (2005):** Exploitation of Nile Tilapia in a Closed-System Public Fishing Reservoir in Northern Arkansas. **North Amer. J. Fish. Manag., 25 (3): 853-860.**
- Hoening, J. M.; Latour, R. J. and Olney, J. E. (2008):** Estimating stock composition of anadromous fishes from Mark-

- Recovery data: Possible application to American Shad.
North Amer. J. Fish. Manag., 28 (2): 507-515.
- Hsieh, S. L.; Chang, H. T.; Wu, C. H. and Kuo, C. M. (2004):**
Cloning, Tissue distribution and hormonal regulation of
stearoyl-CoA desaturase in tilapia, *Oreochromis mossambicus*.
Aquaculture, 230: 527-546.
- Hudelot, C.; Gowri-Shankar, V.; Jow, H.; Rattray, M. and Higgs,
P.O. (2003):** RNA-based phylogenetic methods application
to mammalian mitochondrial RNA sequences. **Mol. Phyl.
Evol.**, 28: 241-252.
- Hugo, E. R.; Stewart, V. J.; Gast, R. J. and Byers, T. J. (1992):**
Purification of amoeba mtDNA using the UNSET procedure.
In: "Protocols in Protozoology", J. J. Lee & A. T. Soldo,
eds Soc. Protozoologists, Lawrence, KA., D 7.1 - D 7.2.
- Huysseune, A. and Sire, J. Y. (1998):** Evolution of patterns and
processes in teeth and tooth-related tissues in non-mammalian
vertebrates. **Eur. J. Oral. Sci.**, 106 (1): 437-481.
- Hwang, G.; Rahman, M. A.; Abdul Razak, S; Sohm, F.;
Farahmand, H.; Smith, A.; Brooks, C. and Maclean, N.
(2002):** Isolation and characterization of tilapia β -actin
promoter and comparison of its activity with carp- β -actin
promoter. **Biochem. Biophys. Acta**, 1625: 11-18.
- Ikoma, T.; Kobayashi, H.; Tanaka, J.; Walsh, D. and Mann, S.
(2003):** Microstructure, mechanical, and biomimetic

-
-
- properties of fish scales from *Pagrus major*. **J. Struc. Biol.**, **142**: 327–333.
- Il'inskii, A. V.; Silva-Andrade, F.; Shadrin1, E. B.; Samoilov, V. O. and Orbeli, A. L. (2006):** Biological structures as photonic objects. **Biophysics**, **51(4)**: 664–667.
- Jawad, L. A. (2005):** Comparative morphology of scales of four teleost fishes from Sudan and Yemen. **J. Nat. History**, **39 (28)**: 2643-2660.
- Jawad, L. A. and Al-Jufaili, S. M. (2007):** Scale morphology of greter lizard fish *Saurida tumbil* (Bloch, 1795) (Pisces: Synodontidae). **J. Fish Boil.**, **70**: 1185-1212.
- Jiao, Y.; Reid, K.; Nudds, T. and Smith, E. (2009):** Graphical evaluation of fishery status using a likelihood inference Aproach. **North Amer. J. Fish. Manag.**, **29 (4)**: 1106-1118.
- Jin, L.; Cho, J.; Seong, K.; Park, J.; Kong, I. and Hong, Y. (2006):** 18S rRNA gene sequences and random amplified polymorphic DNA used in discriminating Manchurian Trout from other freshwater salamonids. **Fish. Sci.**, **72 (4)**: 903-905.
- Johal, M. S. and Agarwal, T. (1997):** Scale structure of *Oreochromis mossambicus* (Peters). **Res. Bull. Panjab University**, **47(1-4)**: 41-49.
- Johnson, A. M.; Fielke, R.; Christy, P. E.; Robinson, B. and Baverstock, P. E. (1990):** Small subunit ribosomal RNA

- evolution in the genus *Acanthamoeba*. **J. Gen. Microbiol.**, **136**: 1689-1698.
- Jolicoeur, P. and Moismann, J. (1960):** Size and shape variation in the painted turtle. A principal component analysis. **Evolution**, **24**: 339-354.
- Ju, Z.; Karsi, A.; Kocabas, A.; Patterson, A.; Li, P.; Cao, D.; Dunham, R. and Liu, Z. (2000):** Transcriptome analysis of channel catfish (*Ictalurus punctatus*): genes and expression profile from the brain. **Gene**, **261**: 373-382.
- Kaiya, H.; Kojima, M.; Hosoda, H.; Riley, L. G.; Hirano, T.; Grau, E. G. and Kangawa, K. (2003):** Identification of tilapia ghrelin and its effects on growth hormone and prolactin release in the tilapia, *Oreochromis mossambicus*. **Comp. Biochem. Physiol.**, **135**: 421-422.
- Kaneko, A.; Kinoshita, S.; Yoshinaga, T.; Tsukamoto, K. and Watabe, S. (2002):** Changes in expression patterns of stress protein genes during population growth of the rotifer *Brachionus plicatilis* GEN. **Fish. Sci.**, **68 (6)**: 13-17.
- Kaur, N. and Dua, A. (2004):** Species specificity as evidenced by scanning electron microscopy of fish scales. **Current Sci.**, **87(5)**: 692-696.
- Kawamoto, F.; Miyake, H.; Kimura, M.; Nguyen, T. D.; Liu, O.; Zhou, M.; Le, D. D.; Kawai, S.; Isomura, S. And Wataya, Y. (1996):** Sequence variation in the 18S rDNA gene, a target

for PCR-based malaria diagnosis, in *Plasmodium ovale* from southern Vietnam. **J. Clin. Microbiol.**, 34 (9): 2287-2289.

Kenndy, A. J.; Daugherty, D. J. and Sutton, T. M. (2007): Population characteristics of shovelnose sturgeon in the upper Wabash River, Indiana. **North Amer. J. Fish. Manag.**, 27 (1): 52-62.

Kessing, B.; Croom, H.; Martin, A.; McIntosh, C.; McMillan, W.O. and Palumbi, S. (1989): The Simple Tool's Guide to PCR, University of Hawaii, Honolulu, Hawaii, p 23.

Khaw, H. L.; Bovenhuis, H.; Ponzoni, R. W.; Rezk, M. A.; Charo-Karisa, H. and Komen, H. (2009): Genetic analysis of Nile tilapia (*Oreochromis niloticus*) selection line reared in two input environments. **Aquaculture**, 294 (1-2): 37-42.

Kim, S. J.; Ogasawara, K.; Park, J. G.; Takemura, A. and Nakamura, M. (2002): Sequences and expression of androgen receptor and estrogen receptor gene in the sex types of protogynous wrasse, *Halichoeres trimaculatus*. **Gen. Comp. Endocrinol.**, 127: 165-173.

Kjer, K. M. (2004): Aligned 18S and insect phylogeny. **Sys. Biol.**, 53: 506-514.

Kjer, K. M.; Carie, F. L.; Litman, J. and Ware, J. (2006): A molecular phylogeny of Hexapoda. **Arthropod sys. Phyl.**, 64: 35-44.

Koblmüller, S., Sturmbauer, C., Verheyen, E. Meyer, A. And Salzburger, W. (2006): Mitochondrial phylogeny and

phylogeography of East African squeaker catfishes (*Siluriformes: synodontis*). **BMC Evol. Biol.**, **6** (49): 2148-2149.

Kocher, T. D. (2004): Adaptive evolution and explosive speciation: the cichlid fish model. **Nat. Rev. Gen.**, **5**: 288-298.

Kocher, T. D.; Lee, W. J.; Sobolewska, H.; Penman, D. and McAndrew, B. (1998): A genetic linkage map of a cichlid fish, the tilapia (*Oreochromis niloticus*). **Genetics**, **148** (3): 1225-1232.

Kondo, M.; Froschauer, A.; Kitano, A.; Nanda, I.; Hormung, V.; Volff, J.; Asakawa, S.; Mitani, H.; Naruse, K.; Tanaka, M.; Schmid, M.; Shimizu, N.; Scharl, M. and Shima, A. (2002): Molecular cloning and characterization of DMRT genes from the *Medaka oryzias latipes* and the platyfish *Xiphophorus maculatus*. **Gene**, **295**: 213-222.

Kraakman, L. S.; Mager, W. H.; Grooftjans, J. J. and Planta, R. J. (1991): Functional analysis of the promoter of the gene encoding the acidic ribosomal protein L45 in yeast. **Biochim. Biophys. Acta**, **1090**: 204-210.

Krieger, J. and Fuerst, P. A. (2002): Evidence of multiple alleles of the nuclear 18S ribosomal RNA gene in sturgeon (family: Acipouseridae). **J. Appl. Ichthyol.**, **18** (4-6): 290-297.

Krieger, J. and Fuerst, P. A. (2004): Characterization of nuclear 18S rRNA gene sequence diversity and expression in an

- individual lake sturgeon (*Acipenser fulvescens*). **J. Appl. Ichthyol.**, 20 (6): 433-439.
- Kumar, R.; Kushwaha, B.; Nagpure, N. S.; Behera, B. K.; Srivastava, S. K. and Lakra, W. S. (2009):** Physical mapping of rRNA gene in endangered fish *Osteobrama belangeri* (Valenciennes, 1844) (Family: Cyprinidae). **Indian J. Exp. Biol.**, 47: 597-601.
- Lagler, K. E.; Bardach, J. E.; Millet, R. and Passino, P. R. M. (1977):** Ichthyology. **John Wiley and Sons, Inc., New York**, 528 pp.
- Lanzing, W. J. R. and Higginbotham, D. R. (1976):** Scanning microscopy of pharyngeal and oral teeth of the teleost *Tilapia mossambica*. **J. Hydrobiol.**, 48 (2): 137-139.
- Ledee, D. R.; Booton, G. C.; Awwad, M. H.; Sharma, S.; Aggarwal, R. K.; Niszl, I. A.; Markus, M. B.; Fuerst, P. A. and Byers, T. J. (2003):** Advantages of using mitochondrial 16S rDNA sequences to classify clinical isolates of *Acanthamoeba*. **Invest. Ophthalmol. Vis. Sci.**, 44 (3): 1142-49.
- Leitner, J. and Bulak, J. (2008):** Performance differences between two endemic stocks of Largemouth Bass in South Carolina. **North American J. Fish. Manag.**, 28 (2): 516-522.
- Levavi-Sivan, B.; Aizen, J. and Avitan, A. (2005):** Cloning, characterization and expression of the D₂ dopamine receptor from the tilapia pituitary. **Mol. Cell. Endocrinol.**, 236: 17-30.

- Lévêque, C. and Paugy, D. (1984):** Guides des poisons d'eau douce de la zone du programme de lutte contre l'onchocercose en Afrique de l'Ouest. **Convention ORSTOM/OMS: 393 pp.**
- Linnaeus, C. (1758):** (1 Jan.) *Systema Naturae*, Ed X. (*Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. (Tomus I. Editio decima, reformata) Holmiae, 1: i-ii + 1-824.*
- Lippitsch, E. (1990):** Scale morphology and squamation patterns in Cichlids (Teleostei, Perciformes): A comparative study. **J. Fish Boil., 37: 265-291.**
- Lippitsch, E. (1992):** Squamation and scale character stability in Cichlids, examined in *Sarotherodon galilaeus* (Linnaeus, 1758) (Perciformes, Cichlidae). **J. Fish Boil., 41: 355-362.**
- Lippitsch, E. (1993):** A phyletic study on *Lacustrine haplochromine* fishes (Perciformes, Cichlidae) of East Africa, based on scale and Squamation characters. **J. Fish Boil., 42: 903-946.**
- López-Piñón, M. J.; Freire, R.; Insua, A. and Mendez, J. (2008):** Sequence characterization and phylogenetic analysis of the 5Sribosomal DNA in some scallopus (Bivalvia: Pectinidae), **Heriditas, 145 (1): 9-19.**
- Lovshin, L. L. (1982):** Tilapia hybridization, P.279-308. In R. S. V. Pullin and R. H. Lowe-McConnell, eds. *The Biology and Culture of Tilapias. ICLARM Conf. Proc., 7. Manila, Philippines, p279-308.*

- Majumdar, K. C. and McAndrew, B. J. (1986):** Relative DNA content of somatic nuclei and chromosomal studies in three genera, tilapia, *Sarotherodon*, and *Oreochromis* of the tribe tilapiini (Pisces, Cichlidae). **Genetica**, **68**: 175-188.
- Mallatt, J. and Giribet, G. (2006):** Further use of nearly complete 28S and 18S rRNA genes to classify Ecdysozoa: 37 more orthopods and a kinorhynch. **Mol. Phyl. Evol.**, **40**: 772-794.
- Mallatt, J. and Sullivan, J. (1998):** 28S and 18S rDNA sequences support the monophyly of lampreys and hagfishes. **Mol. Biol. Evol.**, **15**: 1706-1718.
- Mallatt, J. and Winchell, C. J. (2007):** Ribosomal RNA genes and deuterostome phylogeny revisited: More cyclostomes, elasmobranchs, reptiles and a brittle star. **Mol. Phyl. Evol.**, **43**: 1005-1022.
- Maltagliati, F. (1998):** A preliminary investigation of allozyme genetic variation and population geographical structure in *Aphanius fasciatus* from Italian brackish-water habitats. **J. Fish Biol.**, **52**: 1130-1140.
- Mansour, M.; Wright, J. R. and Pohajdak, B. (1998):** Cloning, sequencing and characterization of the tilapia insulin gene. **Comp. Biochem. Physiol.**, **121**: 291-297.
- Manylov, O. G.; Vladychenskaya, N. S.; Milyutina, I. A.; Kedrova, O. S.; Korokhov, N. P.; Dvoryanchikov, G. A.; Aleshin, V. V. and Petrov, N. B. (2004):** Analysis of 18

SrRNA gene sequences suggests significant molecular differences between Macrodasyda and chaetonotida (Gastrotricha). **Mol. Phyl. Evol.**, **30**: 850-854.

Marques, J. F.; Teixeira, C. M. and Cabral, H. (2006): Differentiation of commercially important flatfish populations along the Portuguese coast: Evidence from morphology and parasitology. **Fish. Res.**, **81** (2-3): 293-305.

Martins, C.; Oliverira, C.; Wasko, A. P. and Wright, J. M. (2004): Physical mapping of the Nile tilapia (*Oreochromis niloticus*) genome by fluorescent in situ hybridization of repetitive DNAs to metaphase chromosomes-A review. **Aquaculture**, **231** (2): 37-49.

Martins, C.; Wasko, A. F.; Oliveira, C. and Wright, J. M. (2000): Nucleotide seauence of 5S rDNA and localization of the ribosomal DNA genes to metaphase chromosomes of the Tilapiine cichlid fish, *Oreochromis niloticus*. **Hereditas**, **133**: 39-46.

Martins, C.; Wasko, A. P.; Oliveira, C.; Porto-Foresti, F.; Parise-Maltempi, P. P.; Wright, J. M. and Foresti, F. (2002): Dynamics of 5 Sr DNA in the Tilapia (*Oreochromis niloticus*) genome: repeat units, inverted sequences, pseudogenes and chromosome loci. **Cytogenet. Genome Res.**, **98**: 78-85.

Matejusova, I.; Gelnar, M.; McBeath, A. J. A.; Collins, C. M. and Cunningham, C. O. (2001): Molecular markers for gyroductylids (Gyrodactylidae: Monogenea) from five fish families (Teleostei). **J. Parasitol.**, **31**: 738-745.

- McAndrew, B. J. and Majumdar, K. C. (1983):** Tilapia stock identification using electrophoretic markers. **Aquaculture**, **30**: 249-261.
- McAndrew, B. J. and Majumdar, K. C. (1984):** Evolutionary relationships within three tilapiine genera. **Zool. J. Linn. Soc.**, **80**: 421-435.
- Mekkawy, I. A. A. (1987):** Taxonomic studies on some Nile Mormyriiform fishes. **Ph. D. Thesis, Faculty of Science, Assiut University, Egypt.**
- Mekkawy, I. A. A.; Mohamad, S. H.; Abass, F. F. and Okasha, S. A. (1994):** Some biological aspects of *Oreochromis niloticus* (Linnaeus, 1758) from Lake Nasser, Egypt and the effect of Lake Impoundment. **Bull. Faculty of Science, Assiut University**, **23 (2)**: 101-142.
- Mekkawy, I. A. A.; Saber, S. A.; Shehata, S. M. A. and Osman, A. G. M. (2002):** Morphometrics and meristics of four fish species of genus *Epinephelus* (Family: Serranidae) from the Red Sea, Egypt. **Bull. Faculty of Science, Assiut University**, **31 (1)**: 21-41.
- Mekkawy, I. A. A.; Shehata, S. M. A.; Saber, S. A. and Osman, A. G. M. (1999):** Scale characteristics of five species of genus *Epinephelus* (Family: Serranidae) from the Red Sea, Egypt. **J. Egypt. Ger. Soc. Zool.**, **30(B)**: 71-102.

- Meyer, A. and Wilson, A. C. (1990):** Origin of the tetrapods inferred from their mitochondrial DNA affiliation to lungfish. **J. Mol. Evol.**, **31**: 359-364.
- Meyer, A. and Zardoya, R. (2003):** Recent advances in the (molecular) phylogeny of vertebrates. **Ann. Rev. Ecol. Evol. Sys.**, **34**: 311-335.
- Molina, A.; Iyengar, A.; Marins, L. F.; Biemar, F.; Hanley, S.; Maclean, N.; Smith, T. J.; Martial, J. A. and Muller, M. (2001):** Gene structure and promoter function of a teleost ribosomal protein: A tilapia (*Oreochromis mossambicus*) 118 gene. **Biochim. Biophys. Acta**, **1520**: 195-202.
- Morals, R.; Herrera, M. T.; Arenal, A.; Cruz, A.; Hernández, O.; Pimentel, R.; Guillén, I.; Martinez, R. and Estrada, M. P. (2001):** Tilapia chromosomal growth hormone gene expression accelerates growth in transgenic zebrafish (*Danio rerio*). **Elec. J. Biotech.**, **4** (2): 17-58.
- Mullis, K. and Faloona, F. (1987):** Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. **Methods enzymol.**, **155**: 335-350.
- Musilová, Z.; Rican, O.; Janko, K. and Novak, J. (2008):** Molecular phylogeny and Biogeography of the Neotropical cichlid fish tribe Cichlasomatini (Teleostei: Cichlidae: Cichlasomatinae). **Mol. Phyl. Evol.**, **46**: 659-672.
- Nagi, S.; Tichy, H.; Mayer, W. E.; Samonte, I. E.; McAndrew, B. J. and Klein, J. (2001):** Classification and phylogenetic

relationships of African tilapiine fishes inferred from mitochondrial DNA sequences. **Mol. Phyl. Evol.**, **20** (3): 361-374.

Naylor, R. L.; Goldberg, R. J.; Primavera, J. H.; Kautsky, N.; Beveridge, M. C.; Clay, J.; Folke, C.; Lubchenco, J.; Mooney, H. and Troell, M. (2001): Effects of aquaculture on world fish supplies. **Nature**, **405**: 1017-1024.

Nieddu, M.; Pichiri, G.; Coni, P.; Salvadori, S.; Deiana, A. M. and Mezzanote, R. (1998): A comparative analysis of European and American eel (*Anguilla anguilla* and *Anguilla rostrata*) genomic DNA: 5S rDNA polymorphism permits the distinction between the two populations. **Genome**, **41**: 728-732.

Nomura, M. (1999): Regulation of ribosome biosynthesis in *Escherichia coli* and *Saccharomyces cerevisiae*: diversity and common principles. **J. Bacteriol.**, **181**: 6857-6864.

Obady, Y. H. A. (2003): Biological and taxonomic studies on some species of genus *Siganus* (Family: Siganidae) from the Red Sea, Egypt. **M. Sc. Thesis, Faculty of Science, Assiut University, Egypt**, 286 pp.

Oberst, S.; Abban, E. K. and Villwock, W. (1996): Biochemical and immunological markers for the discrimination of three Tilapia species: *T. zillii* Gervais *T. guineensis* Bleeker and *T. dageti* Thys v. d. Audenaerde (Pisces: Cichlidae) from West Africa. **Aquacul. Res.**, **27**: 235-244.

- Oberst, S.; Abban, E. K.; Villwock, W. and Renwranzt, L. (1993):** Tilapia plasma proteins and muscle parvalbumins as markers for species identification. **Ger. J. Appl. Zool., 4: 487-501.**
- Oberst, S.; Villwock, W. and Renwarntz, L. (1992):** Western blot analysis of plasma components of three Tilapia speices, *Oreochromis aureus*, *O. niloticus* and *Sarotherodon galilaeus*. **J. Appl. Ichthyol., 8: 278-292.**
- Omoniyi, I. T. and Agbon, A. O. (2008):** Morphometric variations in *Sarotherodon melanotheron* (Pisces: Cichlidae) from brackish and fresh water habitats in South-western Nigeria. **West African J. Appl. Ecol., 12: 1-5.**
- Pante, M. J. R.; Lester, L. J. and Pullin, R. S. V. (1988):** A preliminary study on the use of canonical discriminate analysis of morphometric and meristic characters to identify cultured tilapia, **In: R. S. V. Pullin, T. Bhukaswan, K. Tonguthal and J. L. Maclean, 1-13.**
- Parthar, I. S.; Sato, H. and Sakuma, Y. (2003):** Ghrelin gene in Cichlid fish is modulated by sex and development. **Biochem. Biophys. Res. Commun., 305: 169-175.**
- Pendás, A. M.; Moran, P.; Martinez, J. L. and Garcia-Vasquez, E. (1995):** Applications of 5S rDNA in atlantic salmon, brow trout, and in Atlantic salmon x brown trout hybrid identification. **Mol. Ecol., 4: 275-276.**
- Poncelet, A. C.; Levavi-Sivan, B.; Muller, M.; Yaron, Z.; Martial, J. A. and Belayew, A. (1996):** The tilapia prolactin I gene:

evolutionary conservation of the regulatory elements directing pituitary-specific expression. **DNA Cell Biol.**, **15**: 679-692.

Ponzoni, R. W.; Nguyen, N. H. and Khaw, H. L. (2007): Investment appraisal of genetic improvement programs in Nile tilapia (*Oreochromis niloticus*). **Aquaculture**, **269** (1-4): 187-199.

Puder, M.; Barnard, G. F.; Staniunas, R. J.; Steele Jr., G. D. and Chen, L. B. (1993): Nucleotide and deduced amino acid sequence of human ribosomal protein L18. **Biochim. Biophys. Acta**, **1216**: 134-136.

Pullin, R. S. V. (1985): Tilapia "Everyman's fish". **Biologist.**, **32**: 84-88.

Pullin, R. S. V. (1996): World tilapia culture and its future prospects. In R. S. V. Pullin, J. Lazard, M. Legendre, J. B. Amon Kothias and D. Pauly (eds.). **The 3. Internat. Symp. Tilapia Aquacul. ICLARM Conf. proc.**, 41 pp.

Quackenbush, J.; Liang, F.; Holt, I.; Pertea, G. and Upton, J. (2000): The TIGR gene indices: reconstruction and representation of expressed gene sequences. **Nucleic Acids Res.**, **28**: 41-45.

Quezada, C.; Navarro, C.; Martin, R. S.; Alvarez, M.; Molina, A. and Ines vera, M. (2006): Genomic organization of ucleolin gene in carp fish: Evidence for several genes. **Biol. Res.**, **39**: 353-365.

- Quinterio, J.; Sotelo, C. G.; Rehbein, H.; Pryde, S. E.; Medina, I.; PerezMartin, R. I.; ReyMendez, M. and Mackie, I. M. (1998):** use of mtDNA direct polymerase chain reaction (PCR) sequencing and PCR-restriction fragment length polymorphism methodologies in species identification of canned tuna. **J. Agric. Food. Chem., 46: 1662-1669.**
- Radtke, R. L.; Collins, M. and Dean, J. M. (1982):** Morphology of the otoliths of the atlantic blue marlin (*Makaira nigricans*) and their possible use in age estimation. **Bull. Marine Sci., 32(2): 498-503.**
- Rajavarthini, P. B; Arunkumar, R. I. and Michael, R. D. (2000):** Partial characterization of serum immunoglobulins of *Oreochromis mossambicus*. **Ind. J. Exp. Biol., 38 (6): 549-553.**
- Ram, J. L.; Ram, M. L. and Baidoum, F. F. (1996):** Authentication of canned tuna and bonito by sequence and restriction site analysis of polymerase chain reaction products of mitochondrial DNA. **J. Agric. Food Chem., 44: 2460-2467.**
- Reddy, G. M. (2006):** Scale structure of a cobitid fish, *Cobitis linea* (Heckle, 1849) using different modes of SEM. **Current Sci., 91(11): 1464-1466.**
- Riley, D. E.; Samadpour, M. and Krieger, J. N. (1991):** Detection of variable DNA repeat in diverse eukaryotic microorganisms by a single set of polymerase chain reaction primers. **J. Clin. Microbiol., 29 (12): 2746-2751.**

- Rishi, K. K. and Jain, M. (1998):** Effect of toxicity of cadmium on scale morphology in *Cyprinus carpio* (Cyprinidae). **Bull. Environ. Contam. Toxicol.**, **60**: 323-328.
- Roger, B.; Moisand, A.; Amalric, F. and Bouvet, P. (2002):** Repression of RNA polymerase I transcription by nucleolin is independent of the RNA sequence that is transcribed. **J. Biol. Chem.**, **277**: 10209-10219.
- Roger, B.; Moisand, A.; Amalric, F. and Bouvet, P. (2003):** Nucleolin provides a link between RNA Polymerase I Transcription and pre-ribosome assembly. **Chromosoma**, **111**: 399-407.
- Rognon, X. and Guyomard, R. (2003):** Large extent of mitochondrial DNA transfer from *Oreochromis aureus* to *Oreochromis niloticus* in West Africa. **Mol. Ecol.**, **12** (2): p 435.
- Roldan, M. I.; Perrotta, R. G.; Cortey, M. and Pla, C. (2000):** Molecular and morphologic approaches to discrimination of variability patterns in chub mackerel, *Scomber japonicus*. **J. Exp. Mar. Biol. Ecol.**, **253**: 63-74.
- Rosenfeld, H.; Levavi-Sivan, B.; Gur, G.; Melamed, P.; Meiri, I., Yaron, Z. and Elizur, A. (2001):** Characterization of tilapia FSH β -gene and analysis of its 5' flanking region. **Comp. Biochem. Physiol. Part B**, **129**: 389-398.
- Rutten, M. J. M.; Komen, H.; Deerenberg, R. M.; Siwek, M. and Bovenhuis, H. (2004):** Genetic characterization of four

strains of Nile tilapia (*Oreochromis niloticus*) using microsatellite markers. **Animal Gen.**, 35: 93-97.

Saiki, R. K.; Scharf, S. J.; Faloona, F.; Mullis, K. B.; Horn, G. T.; Erlich, H. A. and Arnheim, N. (1985): Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. **Science**, 230: 1350-1354.

Salama, M. A. M. (2001): Experimental studies on the cytogenetic of a certain fresh water fish. **Ph. D. Thesis, Faculty of Science, Zagazig University, Egypt, 162 pp.**

Salini, J. P.; Milton, D. A.; Rahman, M. J. and Hussain, M. G. (2004): Allozyme and morphological variation throughout the geographic range of the tropical shad, hilsa *Tenualasa ilisha*. **Fish. Res.**, 66: 53-69.

Sanger, F.; Nicklen, S. and Coulson, A. R. (1977): DNA sequencing with chain-terminating inhibitors. **Proc. Natl. Acad. Sci. USA**, 74 (12): 5463–5467.

Santini, S.; Boore, J. L. and Meyer, A. (2003): Evolutionary conservation of regulatory elements in vertebrate Hox gene clusters. **Genome Res.**, 13 (6): 1111-1122.

Savill, N. J.; Hoyle, D. C. and Higgs, P. G. (2001): RNA sequence evolution with secondary structure constraints: Comparison of substitution rate models using maximum likelihood methods. **Genetics**, 157: 399-411.

- Seehausen, O. (2006):** African cichlid fish: a model system in adaptive radiation research. **Proc. Biol. Sci.**, **273** (1597): 1987-1998.
- Seyoum, S. (1989):** Stock identification and the evolutionary relationship of the Genera *Oreochromis*, *Sarotherodon* and *Tilapia* (Pisces: Cichlidae) using Allozyme analysis and restriction endonuclease analysis of mitochondrial DNA. **Ph. D. Thesis, University of Waterloo, Waterloo, Ontario.**
- Seyoum, S. (1990):** Stock identification and the evolutionary relationships of the tilapiine fishes of the genera *Oreochromis*, *Sarotherodon* and *Tilapia* (Pisces: Cichlidae) using allozyme analysis and restriction endonuclease analysis of mitochondrial DNA. **Dissert. Internat. Sci. Eng.**, **50**: 1-12.
- Seyoum, S. and Kornfield, I. (1992):** Identification of the subspecies of *Oreochromis niloticus* (Pisces: Cichlidae) using restriction endonuclease analysis of mitochondrial DNA. **Aquaculture**, **102**: 29-42.
- Shahin, A. A. B. (1999):** Phylogenetic Relationship between the Characid Nile fish *Alestes dentex*, Cyprinid *Barbus bynni*, *Labeo niloticus*, and the introduced grass carp *Ctenopharyngodon idella* elucidated by protein electrophoresis. **J. Egypt Ger. Sci. Zool.**, **29 (B)**: 21-42.
- Sharaf-Eldeen, K. and Abdel-Hamde, N. (2002):** Sublethal Effect of Copper Sulfate, Malathion and Paraquat on Protein Pattern

-
- of *Oreochromis niloticus*. **Egypt. J. Aquat. Biol. & Fish.**, **6 (2): 167-182.**
- Shiue, Y.; Wang, L.; Chao, T; Lin, C. and Tsai, C. (2004):** EST-based identification of genes expressed in the hypothalamus of adult tilapia, *Oreochromis mossambicus*. **Biochem. Biophys. Res. Commun.**, **316: 523-527.**
- Sire, J. (1988):** Evidence that mineralized spherules are involved in the formation of the superficial layer of the elasmoid scale in cichlids *Cichlasoma octofasciatum* and *Hemichromis bimaculatus* (Pisces, Teleostei): an epidermal active participation? **Cell Tissue Res.**, **253: 165-172.**
- Sire, J. Y. (1986):** Ontogenic development of surface ornamentation in the scales of *Hemichromis bimaculatus* (Cichlidae). **J. Fish Boil.**, **28: 713-724.**
- Smith, A. (1840):** "Illustrations to the Zoology of South Africa, consisting chiefly of figures and descriptions of the objects of natural history collected during an expedition into the interior of South Africa. **Fish. Lond., Pt. 5: 1834-1836.**
- Smith, C. T.; Elfstrom, C. M.; Seeb, L. W. and Seeb, J. E. (2005):** Use of sequence data from rainbow trout and Atlantic salmon for SNP detection in Pacific Salmon. **Mol. Ecol.**, **14 (13): 4193-4203.**
- Smith, R. and McVeagh, M. (2000):** Allozyme and microsatellite DNA markers of toothfish population structure in the Southern Ocean. **J. Fish Biol.**, **57 (A): 72-83.**

- Sogin, M. L. (1989):** Evolution of eukaryotic microorganisms and their small subunit ribosomal RNAs. **Amer. Zool., 29: 487-499.**
- Sogin, M. L. (1991):** The phylogenetic significance of sequence diversity and length variation in eukaryotic small subunit ribosomal RNA coding regions. **In L. Warren and H. Koprowski, eds. New Perspectives on evolution. Wiley-Liss, New York, 175-188.**
- Song, J.; Bruet, B. J. F.; Kalai, I.; Pelled, G.; Gazit, D.; Boyce, M. C. and Ortiz, C. (2007):** The graded structure and mechanical properties of the natural armor of *Polypterus senegalus*. **Mat. Res. Society, accepted as an oral presentation Symposium AA7.10, Boston, Mass.**
- Souza, I. L.; Santos-Silva, L. K.; Venere, P. C. and Moreira-Filho, O. (2008):** Molecular cytogenetics of *Salminus* fish (characiformes) based on 5S and 18S RNA genes hybridization, fluorochrome staining and C-banding. **Micron, 39 (7): 1036-1041.**
- Steindachner, F. (1864):** Ichthyologische Notizen. **Anzeiger, K. Akad. Wiss. Math. Naturwiss. Cl., 4 (5): 37-38.**
- Steindachner, F. (1870):** Ichthyologische Notizen (X). **Sitzungsber, Akad. Wiss. Wien, 623-642.**
- Stepien, C. A.; Hubers, A. N. and Skidmore, J. L. (1999):** Diagnostic genetic markers and evolutionary relationship among invasive dreissenoid and corbiculoid bivalves in North

- America: Phylogenetic signal from mitochondrial 16S rDNA. **Mol. Phyl. Evol.**, 13 (1): 31-49.
- Stiassny, M. L. J. (1991):** Phylogenetic intrarelationships of the family cichlidae: an overview, in **Cichlid fishes: behavior, Ecology and Evolution**, Edited by M. H. A. Keenleyside. Chapman & Hall. London, 1-35.
- Stiassny, M. L. J.; Schliewen, U. K. and Dominey, W. J. (1992):** A new species flock of cichlid fishes from Lake Bermin, Cameroon with a description of eight new species of *Tliapia* (Labroidei: Cichlidae). **Ichthyol. Explorat. Freshwaters**, 3: 311-346.
- Stothard, D.; Schroeder-Diedrich, J. M.; Awwad, M. H.; Gast, R. J.; Ledee, D. R.; Rodriguez-Zaragoza, S.; Dean, C. L.; Fuerst, P. A. and Byers, T. J. (1998):** The Evolution History of the *Acanthamoeba* and the Identification of Eight New 18S rRNA Gene Sequence Types. **J. Eukaryot. Microbiol.**, 45 (1): 45 – 54.
- Streelman, J. T.; Webb, J. F.; Albertson, R. C. and Kocher, T. D. (2003a):** The cusp of evolution and development: Model of cichlid tooth shape diversity. **Evol. Devel.**, 5: 600-608.
- Sun, P. L.; Hawkins, W. E.; Overstreet, R. M. and Brown-Peterson, N. J. (2009):** Morphological deformities as biomarkers in fish from Contaminated Rivers in Taiwan. **Int. J. Environ. Res. Public Health**, 6: 2307-2331.

- Suzuki, H.; Sakurai, S. and Matsuda, Y. (1996):** Rat rDNA spacer sequences and chromosomal assignment of the genes to the extreme terminal region of chromosome 19. **Cytogenet. Cell. Gen.**, 72: 1-4.
- Telford, M. J.; Wise, M. J. and Gowri-Shankar, V. (2005):** Consideration of RNA secondary structure significantly improves likelihood based estimates of phylogeny examples from the Bilateria. **Mol. Biol. Evol.**, 22: 1129-1136.
- Teugels, G. G. and Thys van den Audenaerde, D. F. E. (1992):** Cichlidae, P. 714-779. In C. Lévêque, D. Paugy and G. G. Teugels (eds.) Faune des poissons d'eaux douces et saumâtres de L'Afrique de L'Ouest. **Faune tropicale XXVIII. ORSTOM, Paris-MRAC, Tervuren. 902 pp.**
- Thys van den Audenaerde, D. E. F. (1970):** An annotated bibliography of Tilapia (Pisces: Cichlidae). **Mus. Roy. Afr. Cent. Doc. Zool.**, 14: 406 pp.
- Tiano, T. J.; Willis, C. M.; Noble, A. A.; Luttenton, M. R. and Nikitin, A. G. (2007):** Genetic Identification of Hatchery Stocks of Brown Trout Using Mitochondrial DNA Polymorphism. **North Amer. J. Fish. Manag.**, 27: 965-970.
- Trewavas, E. (1965):** *Tilapia aurea* (Steindachner) and the status of *Tilapia niloticus exul*, *T. monody* and *T. lemassoni*. **Isr. J. Zool.**, 14: 258-276.

- Trewavas, E. (1982):** Generic grouping of Tilapiini used in aquaculture. **Aquaculture**, 27: 79-81.
- Trewavas, E. (1984):** Tilapia fishes of the genera *Sarotherodon*, *Oreochromis* and *Danakilia*. **Brit. Mus. (Nat. Hist.) Lond.**, 600 pp.
- Tuisku, F. and Hildebrand, C. (1994):** Evidence for a neural influence on tooth germ generation in a polyphyodont species. **Devel. Biol.**, 165 (1): 1-9.
- Tuisku, F. and Hildebrand, C. (1996):** Occurrence of axons with certain immunohistochemical markers in teleost gingiva and teeth. **Brain Res.**, 729 (1): 137-141.
- Van der Salm, A. L.; Metz, J. R.; Bonga, S. E. W. and Flik, G. (2005):** Alpha-MSH, the melanocortin-1 receptor and background adaptation in the Mozambique tilapia, *Oreochromis mossambicus*. **Gen. Comp. Endocrinol.**, 144: 140-149.
- Vari, R. P. (2003):** Curimatidae. In: R. E. Reis, S. O. Kullander and C. J. Ferraris (eds) Check list of the Freshwater Fishes of South America. **Edipucrs, Porto Alegre**, 735 pp.
- Vera cruz, E. M.; Brown, C. L.; Luckenbach, J. A.; Picha, M. E; Bolivar, R. B. and Broski, R. J. (2006):** Insulin-like growth factor-I cDNA cloning, gene expression and potential use as a growth rate indicator in Nile tilapia, *Oreochromis niloticus*, **Aquaculture**, 251 (2-4): 585-595.

- Vera, M. I.; Molina, A.; Pinto, R.; Reyes, M.; Alvarez, M.; Krauskopf, E.; Quezada, C.; Toores, J. and Krauskoff, M. (2003): Genomic organization of the rDNA cistron of the teleost fish *Cyprinus carpio*. **Biol. Res.**, **36**: 241-51.
- Vodkin, M. H.; Howe, D. K.; Visvesvara, G. S. and McLaughlin, G. L. (1992): Identification of *Acanthamoeba* at the generic and specific levels using the polymerase chain reaction. **J. Protozool.**, **39** (3): 378-385.
- Vuorinen, J. A.; Bodaly, R. A. and Reist, J. D. (1993): Genetic and morphological differentiation between dwarf and normal size forms of Lake White fish (*Coregonus clupea formis*). In Como Lake, Ontario. **Can. J. Fish. Aquat. Sci.**, **50**: 210-216.
- Wadie, W. F.; Rizkalla, S. A.; Al-Zahaby, S. A. and El-Serafy, S. S. (1999): Biometric studies on two species of *Pagellus* (Family: Sparidae) In the Egyptian Mediterranean waters. **Egypt. J. Aquat. Biol. & Fish.**, **3** (3): 1-29.
- Wainwright, P. O.; Hinkle, G.; Sogin, M. L. and Stickel, S. L. (1993): Monophyletic origins of the metazoa: An evolutionary link with fungi. **Science**, **260**: 340-342.
- Wang, D.; Kobayashi, T.; Zhou, L. and Nagahama, Y. (2004): Molecular cloning and gene expression of fox 12 in the Nile tilapia, *Oreochromis niloticus*. **Biochem. Biophys. Res. Commun.**, **320**: 83-89.
- Welsh, A.; Hill, T.; Quinlan, H.; Robinson, C. and May, B. (2008): Genetic assessment of lake sturgeon population structure in

- the Laurentian Great Lakes. **North Amer. J. Fish. Manag.**, 28 (2): 572-591.
- White, M. M. (2000):** Genetic Variation in White Bass. **Trans. Amer. Fish. Soc.**, 129: 879-885.
- Winkler, C. and Moon, R. T. (2001):** Zebrafish *mdk2*, a novel secreted midkine, participates in posterior neurogenesis. **Devel. Biol.**, 229: 102-118.
- Xia, X.; Xie, Z. and Kjer, K. M. (2003):** 18S ribosomal rRNA and tetrapod phylogeny. **Sys. Biol.**, 52: 283-295.
- Yapi-Gnaoré, V. (2001):** Fish genetics research in Côte d'Ivoire. In **M. V. Gupta and B. O. Acosta (eds.)**, 25-28.
- Zanke, C.; Borisjuk, N.; Ruoss, B. and Schilderentschler, L. (1995):** A specific oligonucleotide of the 5S rDNA spacer and species specific elements identify symmetrical somatic hybrids between *Salanum tuberosum* and *S. pinnatisectum*. **Theor. Appl. Gen.**, 90: 720-726.
- Zrzavy, J. (2003):** Gastrotricha and Metazoan Phylogeny. **Zool. Scripta**, 32: 61-81.