

REFERENCES

Abd-Elmoneim, E.; Sulieman, M.O.; ElBoshra and El Amin, A. (2007): Nutritive value of clove (*Syzygium aromaticum*) and detection of antimicrobial effect of its bud oil. J. of Microbiology 2 (3): 266-271.

Abd-Elsamea, A. (2008): Studies on the resistance and susceptibility of bacteria implicated in urinary tract infection to antimicrobial drugs. Master Degree, Zagazig University. Faculty of Science. Botany Department.

ACCM, (2008): American council collaborative medicine. Antimicrobial resistance in the treatment of urinary tract infections is a major health problem.

Adedeji, B.A.M. and Abdulkadir, O.A. (2009): Etiology and antimicrobial resistance pattern of bacterial agents of urinary tract infections in students of tertiary institutions in Yola Metropolis. Department of Microbiology, Federal University of Technology, P.M.B. 2076. Advances in Biological Research 3 (3-4): 67-70.

Adeyemo, A.A.; Gbadegesin, R.A.; Onyemenem, T.N. and Ekweozor, C.C. (1994): Urinary pathogens and antimicrobial sensitivity patterns in children in Ibadan, Nigeria. Ann Trop Paediatr; 14: 271-274.

Adjei, O. and Opoku, C. (2004): Urinary tract infections in African infants. Int. J. Antimicrob Agents., 24 S1:32-34.

Agace, W.W.; Hedges, S.R.; Ceska, M. and Svanborg, C. (1993): Interleukin-8 and the neutrophil response to mucosal Gram-negative infection. J Clin. Invest; 92(2):780-5.

Ahmed, F.M. (2004): Antimicrobial studies of some essential oils on some bacteria; Master Degree; Botany Department; Faculty of Science.

Al-Haddad, A.M. (2005): Urinary tract infection among pregnant women in Al-Mukalla district, Yemen Eastern Mediterranean Health Journal, Vol. 11, No. 3. PP: 505-510.

Alonso, J. (2004): Tratado de fitofarmacos nutraceuticos. Barcelona: Corpus; p: 927-30 (633.8 ALO).

Ana, C.; Ronald, N. and Kelley, A. (2000): Activity and spectrum of 22 antimicrobial agents tested against urinary tract infection pathogens in hospitalized patients in Latin America. J. Antimicrob. Chemother.; 45: 295-303.

Andrade, S.S.; Sader, H.S.; Jones, R.N. and Pereira, A.S. (2006): Increased resistance to first-line agents among bacterial pathogens isolated from urinary tract infections in Latin America. Mem. Inst. Oswaldo. Cruz, Rio de Janeiro, Vol. 101 (7): 741-748.

Andriole, V.T. (2002): Asymptomatic bacteriuria in patients with diabetes-enemy or innocent visitor? N Engl J Med; 2002; 347:1617-1618.

Araki, W.B. and Tomasi, T.B. (1999): Quantitative relationships in human serum and non-vascular fluids; T.B.Jr. Gamma Globulins Science; 142; 1080.

Aridogan, B.C.; Baydar, H.; Kaya, S.; Demirci, M.; Ozbasar, D. and Mumcu, E. (2002): Antimicrobial activity and chemical composition of some essential oils. J. of Arch. Pharm. Res., 25: 860-864.

Asscher, N.; Legakis, N.J.; Papavassilion, J. and Sifalakis, J. (1999): Sterilization of urological instruments infected with *Pseudomonase aeroginosa*; Dept. Microbiol. Fac. Med.; Univ. Athens. Actampicrobiol Hellen (285-291).

Ayoola, G.A.; Lawore, F.M.; Adelowotan, T.; Aibinu, I.E.; Adenipekun, E.; Coker, HAB. and Odugbemi, T.O. (2008): Chemical analysis and

antimicrobial activity of the essential oil of *Syzigium aromaticum* (clove); Faculty of Pharmacy and Medicine, University of Lagos, Nigeria; vol.(2) pp. 162-166.

Barisic, Z.; Babic-Ereeg, A. and Borzic, E. (2003): Urinary tract infections in South Croatia: Etiology and antimicrobial resistance. International J Antimicrob Agents; 22: S61-4.

Bariska, J.; Valko, K.; Takacs-Novak, K. and Kalasz, H. (1999): Planar chromatography, J. Planar Chromatogr. 12: 46-50.

Baron, E.J. and Finegold, S.M. (1990): General microbiology; Baily and Scotts Diagnostic Microbiology. 8th. Ed. Mosby Co. St. Louis. Balitimore.

Barr, S.T. (1997): Hereditary interstitial nephritis. In: Biology of pyelonephritis; edited by B.I. Nas and Boston; Bittle; Browen and Marry; pp. 427-432.

Barry, H.C.; Ebell, M.H. and Hickner, J. (1997): Evaluation of suspected urinary tract infection in ambulatory women: a cost-utility analysis of office-based strategies. J Fam Pract; 44: 49-60.

Barry, L.B. and Thrnsberry, Y. (1987): Inhibition of growth and aflatoxin production by cinnamon and clove oils, cinnamic aldehyde and eugenol. Food Sci. 42:1107-1109.

Bartman, T.; Frank, R. and Goldenring, J.M. (2001): Newborn circumcision and urinary tract infections pediatrics; 107(1): 210b-214.

Bauer, A.W.; Kirby, W.M.; Sherris, J.C. and Turk, M. (1986): Antibiotics susceptibility testing by a standardized single disk method Am. J of C1. Pathology, 45:493-496.

Bent, S.; Nallamothu, B.K. and Simel, D.L. (2002): Does this woman have an acute uncomplicated urinary tract infection? JAMA; 287: 2701-2710.

Boon, E.M.; Downs, A. and Marcey, D. (2007): Catalase: H_2O_2 : H_2O_2 oxidoreductase. Catalase structural tutorial text. Retrieved on 2007-02-11.

Boyko, E.J.; Fih, S.D. and Scholes, D. (2002): Diabetes and risk of acute urinary tract infection among postmenopausal women. Diabetes care; 25: 1778-1783.

Brenner, R.M. and DeNavasquez, S.J. (1997): Experimental pyelonephritis in the mouse produced by *Escherichia coli*; *Pseudomonas aeruginosa* and *Proteus mirabilis*. J. Path. Bact.; 87: 79-89.

Bruce, G. and Galloway, N. (1998): Diagnostic and therapeutic considerations in catheters associated bacteruria in hospital/chronic care facility. In: Cunha B, ed. Urinary tract infections: current issue in diagnosis and treatment. Antibiotics for clinicians; 2(suppl 2): 11-16.

Buffenmyer, M.; Benesuva, D. and Rossmann, P. (1986): Pathogenicity of various strains of *Escherichia coli* for the intact rabbit kidney and the effect of repeated passage of renal tissue progress in pyelonephritis Philadelphia; S.A.; Davis Co.; pp. 155-196.

Burrows, M.O.; (1963): Role of ureter in pathogenesis of ascending pyelonephritis. JAWA; 168: 1983.

Burt, S.A. and Reinders, R.D. (2003): Antibacterial activity of selected plant essential oils against *E. coli* 0157: 117. Lett. applied microbiol., 363: 162-167.

Byrne, D.E.; Sedor, J.F. and Soroush, M. (2001): Interaction of bladder glycoprotein GP51 with uropathogenic bacteria. J Urol; 165(4): 1342-6.

Chessbrough, M. (1993): Collection, transport and examination of specimens. In: Chessbrough, M. (ed). Medical laboratory Manual for Wort- Heinemann Ltd., Oxford.

Chessbrough, M. (2000): Biochemical test to identify bacteria. In Cheesbrough, M. (ed). Laboratory Practice in Tropical Countries, Vol. 2:63-70. Cambridge University Press, Uk.

Choi, S.H.; Woo, J.H.; Jeong, J.Y.; Kim, N.J.; Kim, M.N.; Kim, Y.S. and Ryu, J. (2006): Clinical significance of *Staphylococcus saprophyticus* identified on blood culture in a tertiary care hospital. Diagn. Microbiol. Infect. Dis. 56:337-339.

Cieslikowski, T.; Gradecka, D. and Mielczarek, M. (2003): Tandem tetramer-based microsatellite fingerprinting for typing of *Proteus mirabilis* strains. Journal of clinical microbiology; 41: 1673-1680.

Coia, D. (2006): Guidance for the control and prevention of methicillin-resistant *Staphylococcus aureus* in health care facilities. Journal of hospital infection. 66, (S1) 1-44.

Collee, J.G. and Marr, W. (1989): Culture containers and culture media. In practical medical microbiology 13th ed. Vol. (2). Collee, J.G.; Duguid, J.P.; Fraser, A.G. and Marmion, B.P. (eds); Churchill. Livingstone. New York. Pp. 100-120.

Colodner, R.; Ken-Dror, S.; Kavenshtock, B.; Chazan, B. and Raz, R. (2006): Epidemiology and clinical characteristics of patients with *Staphylococcus saprophyticus* bacteriuria in Israel. Infection 34:278-281.

Conte, J.E. and Barriere, S.L. (1981): A manual of antibiotics and infectious disease. 4th Ed. LEA and FEBIGER Philadelphia.

Cornelis, P. (2008): *Pseudomonas: Genomics and Molecular Biology*, 1st ed., Caister Academic Press. ISBN 978-1-904455-19-6.

Cosentino, S.; Tuberoso, C.; Pisano, B.; Satta, M.; Mascia, V.; Arzedi, E. and Palmas, F. (1999): In-vitro antimicrobial activity and chemical composition of *Sardinian thymus* essential oils. Letters in Applied Bacteria. 29 (2): 130.

Duerden, B.I.; Reid, T.M.S.; Jewsbury, J.M. and Turk, D.C. (1990): A new short book of medical parasitic infection, ELBS Publishers; p. 576-581.

Daniel, J.G.; and David, W. (2004): Urinary tract infection; Canadian Pharmacists Association.

Dalia, A. (2005): Studies on some microorganisms causing urinary tract infection in diabetics. Master Degree, Zagazig University. Benha Branch. Faculty of Science. Botany Department.

Dan B.; Steven C.; Erich S. and Andrew G. (2004): Chinese herbal medicine: Materia Medica, Third Edition.

Darnton, N.C.; Turner, L.; Rojevsky, S. and Berg, H.C. (2007): On torque and tumbling in swimming *Escherichia coli*. J Bacteriol.; 189(5):1756-64.

Dauripandiyan, V.; Ayyanar, M. and Ignacimuthu, S. (2006): Antimicrobial activity of some ethnomedicinal plants used by Paliyar tribe from Tamil Nadu, India, doi: 10.1186/1472-6882-6-35.

Delzell, J.E. and Lefevre, M.L. (2000): Urinary tract infection in pregnancy. Am. Fam. Phys. 61:713-72.

Deng, F.M.; Ding, M. and Lavker, R.M. (2001): Urothelial function reconsidered: A role in urinary protein secretion. Proceedings of the national academy of sciences of the United States of America; 98(1): 154-159.

Duguid, D.M. and Staney, T.A. (1994): Bacteriologic localization patterns in bacterial. Prostatitis and Invest. Urol.; 5: 412.

Edward, M.F. and Little, J.D. (1997): Ureteral obstruction in infancy a study of 74 cases. J. A.M.A.; 92: 544.

Egorov, N.S. (1985): Antibiotics a scientific approach. MIR publishers, Moscow.

El-Mishad, J. (1991): Microorganisms isolated from urinary tract infections and their beta lactamase production and evaluation of clinical efficacy of sulperazone. Hinyokika Kivo., 34:8:1503-1514.

Elujoba, A.A.; Odelleye, O.M. and Ogunyemi, C.M. (2005): Traditional medicine development for medical and dental primary health care delivery system in Africa. Afr. J. Trad. CAM 2: 46-61.

Evandro, L.S.; Tania, L.M.S.; Edeltrudes, O.L.; Jose, M.B.F. and Marcia, O.M.M.; (2008): Interference of heating on the antimicrobial activity and chemical composition of *Origanum vulgare* L. (Lamiaceae) essential oil. Cienc. Tecnol. Aliment. Campinas, 28(2): 418:422.

Farag, M.C. and Juven, M.A. (2003): Spasmolytic and antimicrobial activity of essential oil from *Thymus leptophyllus* and *T. Webbii* anus. Fitoterapia.; 60(5): 469-473.

Farrell, D. and Qlobinson, M. (1972): Nutrient and blood agar J. Appl. Bact., 25: 625-630.

Fihn, S.D.; Boyko, E.J. and Chen, C.L. (1998): Use of spermicide-coated condoms and other risk factors for urinary tract infection caused by *Staphylococcus saprophyticus*. Arch Intern Med; 158: 281-287.

Firas, A. (2007): Synergistic antibacterial activity between *Thymus vulgaris* and *Pimpinella anisum* essential oils and methanol extracts. Department of Biology, College of Education, University of Mosul, Mosul, Iraq. Vol. 116, Issue 3, 403:406.

Foxman, B. (2002): Epidemiology of urinary tract infections: incidence, morbidity, and economic costs. Am J Med; 113 Suppl 1A:5S:13S.

Gales, A.C.; Jones, R.N. and Gordon, K.A. (2000): Activity and spectrum of 22 antimicrobial agents tested against urinary tract infection pathogens in hospitalized patients in Latin America: report from the second year of the sentry antimicrobial surveillance program (1998). J Antimicrob Chemother; 45: 295-303.

Gales, A.C.; Jones, R.N.; Turnidge, J.; Rennie, R. and Ramphal, R. (2001): Characterization of *Pseudomonas aeruginosa* isolates: Occurrence rates, antimicrobial susceptibility patterns, and molecular typing in the global sentry antimicrobial surveillance program. Clinical infection diseases; 32:S146-S155.

Gangoue-Pieboji, J.; Pegnyemb, D.E. and Niyitegeka, D. (2006): The in-vitro antimicrobial activities of some medicinal plants from Cameroon. Ann. Trop. Med. Parasitol; 100:237-243.

Gary, K. (2007): Gram stain and capsule stain. BIOL 230MICROBIOLOGY LABORATORY MANUAL.

Gebre-Selassie, S. (1998): Asymptomatic bacteriuria in pregnancy: epidemiology, clinical and microbiological approach. Ethiop. Med. J., 36:185-192.

Geerlings, S.E.; Stolk, R.P. and Camps, M.J.L. (2000): Risk factors for symptomatic urinary tract infection in women with diabetes. *Diabetes care*; 23: 1737-1741.

Gerberding, J.L. (2002): Hospital-onset infections: A patient safety issue. *Ann Intern Med*; 137(8): 665-670.

Ghosh, R.; Nadiminty, N.; Fitzpatrick, J.E.; Alworth, W.L.; Slaga, T.J. and Kumar, A.P. (2005): *J. Biol. Chem.*, 280, 5812.

Glisic, S.B.; Milojevic, S.Z.; Dimitrijevic, S.I.; Orlovic, A.M. and Skala, D.U. (2007): Antimicrobial activity of the essential oil and different fractions of *Juniperus communis* L. and a comparison with some commercial antibiotics. *J. Serb. Chem. Soc.*; 72 (4):311-320.

Gordon, M.; Fergal, O. and John, P.M. (2008): Signal-mediated interactions between *Pseudomonas aeruginosa* and *candida albicans*. Department of Microbiology and Biomerit Research Center, Bioscience Institute, University College Cork, Ireland. *Med. Microbiol* 57, 563-569.

Gosling, R.R. and Little, P.J. (1995): Renal infection and renal scarring edited by Kincaid Smith and Faircliff K.F.; Melbourne; Mace Publishing Co.; pp. 81.

Guentzel, M.N. (1996): *Escherichia, Klebsiella, Enterobacter, Serratia, Citrobacter, and Proteus*. In: Barron's Medical Microbiology (Barron S., eds.) (4th ed. ed.). Univ of Texas Medical Branch. (via NCBI Bookshelf) ISBN 0-9631172-1-1.

Gupta, K.; Hooton, T.M. and Roberts, P.L. (2001): Patient-initiated treatment of uncomplicated recurrent urinary tract infections in young women. *Ann Intern Med*; 135: 9-16.

Gupta, N.; Kundra, S.; Sharma, A. and Vikas Gautam, V. (2007): Antimicrobial susceptibility of uropathogens in india. J. Infect. Dis. Antimicrob. Agents, 24:13-18.

Harrington, R.D. and Hooton, T.M. (2000): Urinary tract infection risk factors and gender. J Gend Specif Med; 3: 27-34.

Harvy, S. (2003): Urinary tract infection. Areview A.D.A.M.pp: 1-19.

Hassan, H.A.; El-Mofty, M. and Nafiss, S.E. (2004): Distribution of pap A and hly genes in uropathogenic *Escherchia coli* isolated in Alexandria. The 12th International Conference of Egyptian Society for Medical Microbiology (ESSM) booklet; 92.

Hazes, B.; Sastry, P.A. and Hayakawa, K. (2000): Crystal structure of *Pseudomonas aeruginosa* PAK pilin suggests a main-chain-dominated mode of receptor binding. J Mol Biol; 299(4): 1005-17.

Hersch, M.P.; Leanos, B.E.; Solorzano, F. (2005): Antibacterial effects of commercial essential oils over locally prevalent pathogenic strains in Mexico. PubMed, 76 (5):453-7.

Holt, J.G.; Krieg, N.R.; Sneath, P.H.; Staley, J.T.; Williams, T. and Hensyl, W.R. (1994): "Bergy's manual of determination bacteriological" 9th ed Williams and Wilkins, Baltimore, Meryland.

Hooton, T.M. and Stamm, W.E. (1997): Diagnosis and treatment of uncomplicated urinary tract infection. Infect Dis Clin North Am 11:551 [PMID 9378923].

Hooton, T.M.; Scholes, D. and Stapheleton, A.E. (2000): A prospective study of asymptomatic bacteriuria in sexually active young women; 343: 992-997.

Hovelius, B. and Mardh, P.A. (1984): *Staphylococcus saprophyticus* as a common cause of urinary tract infections. Rev. Infect. Dis. 6:328-337.

Howes, D.S. and Remer, E.E. (2002): Male urologic infections. In: Harwood-Nuss Al, eds. Clinical practice of emergency medicine. 3rd ed. Philadelphia: Lippincott Williams and Wilkins; 267-272.

Hsieh, J. and Liu, G.D. (1979): Isolation, purification and antibiotic activity of o-methoxy cinnamaldhyde from cinnamon. Pharma Cognosy, phyto chemistry, Medicinal plants. 2nd Edition. Lavoisier publishing intercept. Paris Pg. 545-546.

Huang, S.; Contronic, G. and Khan, W. (1999): Resistance of *shigella* to ampecillin and other antibiotics. Clinical and epidemiological application. JAMA; 221: 45-47.

Iroha, I.R.; Adikwu, M.U.; Esimone, C.O.; Aibinu, I. and Amadi, E.S. (2009): Extended spectrum Beta- L actamase (EBSL) in *E.coli* isolated from a tertiary hospital in Enugu state, Nigeria. Pak J Med Sci; Vol. 25 No. 2; 279-282.

Jackie, R. (2005): Oxidase test, Richland College 12800 Abrams RD Dallas, TX, 75243 US 972.238.6100

Jacques, F.A. (1980): The disc susceptibility test in; Antibiotic in laboratory medicine by Victor – Lorian M. d; Ed Williams and Wilkins Baltimore/London.

Jansen, A.M.; Lockatell, C.V. and Johnson, D.E. (2003): Visualization of *Proteus mirabilis* morphotypes in the urinary tract: the elongated swarmer cell is rarely observed in ascending urinary tract infection. Infection and Immunity; 7(1): 63607-3613.

Javor, N. and Ivemark, B.I. (2000): Renal dysplasia and pyelonephritis in infants and children. Part I Arch. Path.; 66: 255-263.

Jones, A.L. (1999): The pathogenesis of experimental enterococcal pyelonephritis in mice. *J. Infect. Dis.*; 108: 181-188.

Joyce, E.C.B.; Rebeca, P.M.; Lidiane, N.B.; Luiz, C. and Ary, F.J. (2006): Synergism between plant extract and antimicrobial drugs used on *Staphylococcus aureus* diseases. *Mem. Inst. Oswaldo. Cruz, Rio de Janeiro*, Vol. 101 (4): 387-390.

Kalemba, D. and Kunicka, A. (2003): Antimicrobial antifungal properties of essential oils. *Current medicinal chemistry*, volume 10, number 10, pp. 813-829(17).

Kalpana G.; Walter, E. and Stamm, W. (2005): Urinary tract infections. *Medscape*; 505095.

Karen, W. and Edzard, E. (2003): Herbal medicines for treatment of bacterial infections. *J Antimicrob Chemotherapy*; 51,241-246.doi: 10.1093/jac/dkg087.

Kassis, W. (1997): Antimicrobial resistance of *salmonella sp.* isolates in a Spanish hospitals. *Lancet*; 2; 1059; cited from *Br. J. Urol.*; 36: 360-363.

Kathleen, H. (2006): Quick reference for basic SPSS procedures. Department of Psychology, Muhlenberg College.

Koch, (1942): Mannitol salt agar. *Zentralbl. Bakteriologie. Parasitenkd. Abt. I Orig.* 149:122.

Koneman, E.W.; Allen, S.D.; Dowell, V.R. and Cho, Y. (1983): Colour atlas and text book of diagnostic microbiology. 2th ed. J. P. Lippincott Company; N. Y.; London. Pp. 57-125.

Krieg, and Holt, (1984): *Bergey's manual of systematic bacteriology*, Vol. 1. Williams & Wilkins, Baltimore, Md.

Krieg, and Noel, (1984): Bergey's manual of systematic bacteriology, volume 1. Baltimore: Williams & Wilkins. ISBN 0683041088.

Kunin, C.M. (1994): Urinary tract infections in females; Clin. Infect. Dis.; 18(1):1-10; quiz 11-12.

Kunin, C. (1997): Urinary tract infections. 5 ed. Baltimore: Williams & Wilkins; P301-4

Kunin, C.M. (2001): Nosocomial urinary tract infections and the indwelling catheter: What is new and what is true? Chest; 120: 10-12.

Laemmli, V.K. (1973): Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature (London), 227: 680-685.

Lantz, R.C.; Chen, G.J.; Sarihan, M.; Solyom, A.M.; Jolad, S.D. and Timmermann, B.N. (2007): The effect of extracts from ginger rhizome on inflammatory mediator production. Phytomedicine 14 (2-3): 123-128.

Larcombe, J. (1999): Urinary tract infection in children. BMJ; 319(7218):1173-1175.

Lau, W.S. (1999): Infrared characterization for microelectronics. World Scientific.

Lau, S. and Adetubi, M. (2001): Pyelonephritis its relation to vascular lesions and to arterial hypertension. Medicine (Ball) 18: 211.

Lau, S.M.; Peng, M.Y. and Chang, F.Y. (2004): Among pathogens of both bacteremic and non-bacteremic community-acquired urinary tract infection.

Lawson, V.C. and Wood, J.C. (2003): Influence of different media and bloods on results of diffusion antibiotic susceptibility tests. Antimicrob. agents chemother; 1:116-122.

Lerner, G.R. (1994): Urinary tract infections in children *Pediatr. Ann.*, 23; 463-473.

Lepper, F.M.; Grusa, E. and Reichl, H. (2002): Consumption of imipenem correlates with B- Lactam resistance in *Pseudomonas aeruginosa*. *Antimicrobial agents and chemotherapy*; 46(9): 2920-2925.

Li, X.; Lockatell, C.V. and Johnson, D.E. (2004): Development of an intranasal vaccine to prevent urinary tract infection by *Proteus mirabilis*. *Infect. Immun.*; 72(1) 66-75.

Lowry, S.J. (1975): Diarrheal disease: Current concepts and future challenges. Enterodherent *E. coli*: A heterogeneous group of *E. coli* implicated as diarrheal pathogens. *Trans. R. Soc. Trop. Med. Hyg.* 87: 549-553.

Mackey, J.P. and Sandys, J.H. (1966): A trial of cystine-lactose electrolyte deficient medium for urinary bacteriology. *Br. Med. J.* 1:1173.

Madigan, M. and Martinko, J. (2005): Brock biology of microorganisms (11th ed. ed.). Prentice Hall. ISBN 0131443291.

Mady, S. and Helmi, H. (2003): Urinary tract infection pathogens and antibiotics sensitivity patterns in hospitalized infants and children in Al-Jahra Hospital, Kuwait. *Egyptian Journal of Medical Microbiology*; 12(2): 415-420.

Maki, G.D. (2001): Engineering out the risk of infection with urinary catheters. *Emerg. Infect. Dis.*; 7(2).

Malowany, L.; Smith, D.J. and Klavias, J.V. (1989): Renal papillary necrosis induced by heterologous serum. *Arch. Pathol.*; 78: 108.

Manges, A.R.; Natarajan, P.; Solberg, O.D.; Dietrich, P.S. and Riley, L.W. (2006): The changing prevalence of drug-resistant *Escherichia coli* clonal

groups in a community: evidence for community outbreaks of urinary tract infections. *Epidemiol infect.* ; 134:425-31. doi: 10.1017/S0950268805005005.

Marina, P.; Gabriella, P.; Monica, L.; Maria E. M.; Maria G. T.; Giuseppe P.; Davide F.; Maria A.D.; Giovanna D.; Mirco P. and Carla R. (2007): Antiproliferative and pro-apoptotic activity of eugenol-related biphenyls on malignant melanoma cells.

Maslin, M.M. and Pascule, J.V. (2000): Thiopropanal S-oxide: alachrymatory factor in onion. *J. Agric food Chern.* 19: 269-272.

Matsuyama, T.; Takagi, Y.; Nakagawa, Y.; Itoh, H.; Wakita, J. and Matsushita, M. (2000): Dynamic aspects of the structured cell population in a swarming colony of *Proteus mirabilis*. *J. Bacteriol.* 182 (2): 385–93. PMID 10629184. PMC: 94287.

Mazzulli, T. (2001): Susceptibility of community Gram-negative urinary tract isolates to mecillinam and other oral agents. *Can J Infect Dis*; 12(5):289-292.

Mehnert-Kay, S.A. (2005): Diagnosis and management of uncomplicated urinary tract infections. *Am. Fam. Physician.*, 72:451-456.

Melendez, P.A. and Capriles, V.A., (2006): Antibacterial properties of tropical plants from Puerto Rico. *Vol. 13, (4):* 272-276.

Meyer, J.M.; Geoffroy, V.A. and Baida, N. (2002): Siderophore typing, a powerful tool for the identification of fluorescent and nonfluorescent *Pseudomonads*". *Appl. Environ. Microbiol.* 68 (6): 2745–53. doi:10.1128/AEM.68.6.2745-2753.2002. PMID 12039729. Retrieved on 2007-05-27.

Moore, K.N.; Day, R.A. and Albers, M. (2002): Pathogenesis of urinary tract infections. *J Clin Nurs*; 11(5): 568-74.

Monica, C. (1985): Medical laboratory manual for tropical countries. ELBS. Ed. English Language Book Society; tropical health. Technology. Buttrworth. Vol. II. Enteric Gram-negative rods. Chap. 42 : 250-255.

Moreno, S.; Scheyer, T.; Romano, C.S. and Vojnov, A.A. (2006): Antioxidant and antimicrobial activities of rosemary extracts linked to their polyphenol composition. Free Radical Research, 40 (2): 223-231.

Mounnissamy, V.M.; Kavimani, S. and Gunasegaran, R. (2002): Antibacterial activity of gossypetin isolated from *hibiscus sabdariffa*; Pharmacy Division, Mother Theresa Institute of Health Sciences, Govt. Pharmacy, Gorimedu, Pondichery; 99(3): 81-2

Muckenstrum, B.; Foechterlen, D.; Reduron, J.P.; Danton, P. and Hildenbrand M. (1997): Phytochemical and chemotaxonomic studies of *Foeniculum vulgare*. Biochem. Syst. Ecol. 25, 353: 358.

Muhammad, G.; Sajid, M.; Nasir, M.; Nyla, J.; Rehana, K.; Kokab, J. and Gulshan, A. (2008): Composition and antimicrobial properties of essential oil of *Foeniculum vulgare*. African Journal of Biotechnology Vol. 7(24), pp. 4364:4368.

NCCLS, (1998): National Committee for Clinical Laboratory Standard Recommendations

NCCLs (2001): Approved guideline, 2nd ed. Routine urinalysis and collection, transport and preservation of urine specimens. Document GP16-A2, Nov. 2001.

Neu, H.C. (1989): Overview of mechanisms of bacterial resistance. DIAGM. Microbiol. Inf. Dis., 12:1095.

Nicolle, L.E. (1999): Urinary infections in the elderly: symptomatic or asymptomatic. Int J Antimicrob Agents; 11: 265-268.

Nicolle, L.E. (2001): Epidemiology of urinary tract infection. *Infect. Med*; 18(3): 153-162.

Nicolle, L.E. (2002): Resistant pathogens in urinary tract infection. *J Am Geriatr Soc*; 50(7suppl):S230-S235.

Norinder, B.S.; Norrby, R.; Palmgren, A.; Hollenberg, S.; Eriksson, U. and Nord, C.E. (2006): Microflora changes with norfloxacin and pivmecillinam in women with recurrent urinary tract infection. *Antimicrobial agents and chemotherapy*. PP: 1528-1530.

Nwakoby, I.E.; Reddy, K. and Patel, P. (2001): Fast-mediated apoptosis of neutrophils in sera of patients with infection. *Infection and Immunity*; 69(5): 3343-3349.

O'Donnell and Abrutya, (2001): Upper tract obstruction. In: *Comprehensive urology*, 1st ed. Ch 23. P: 33-346. Weiss, R.M.; George, N.J. and O'Reilly, P.H. Mosby International Limited London.

Orenstein, R. and Wong, E.S. (1999): Urinary tract infections in adults; *Am Family Phys*; 59(5):1225-1234.

Osman, S. and Musa, O. (2003): Antibacterial activity of Turkish spice hydrosols. *J. of Food control*. 14: 141-143.

Panton, J.C. and Panton, A.W. (1998): Pathogenesis and diagnosis of shiga toxin-producing *Escherichia coli* infections. *Clin. Microbiol. Rev.* 11 (3): 450–79. PMID 9665978. PMC: 88891.

Park, N; Juven, B.J.; Shaaya, E.; Menasherov, M.; Nitzan, R.; Weiss, H. and Ravid, U. (1984): Inhibitory effect of oregano and thyme essential oils on moulds and food borne bacteria letters in *Applied Microbiology*; 11 (1): 33-37.

Pastore, L.M.; Savitz, D.A. and Thorp, M. (1999): Predictors of symptomatic urinary tract infection after 20 weeks' gestation. *J Perinatol.*, 19:488-493.

Patra, M.; Shahi, S.K.; Midgely, G. and Dikshit, A. (2002): Utilization of essential oil as natural antifungal against nail infective fungi. *Flavour Fragr. J.* 17: 91-94.

Patricia, S. (2006): Department of cell biology and molecular genetics
University of Maryland College Park, MD 20742 USA

Podschun, R. and Ullmann, U. (1998): *Klebsiella* spp. as nosocomial pathogens: epidemiology, taxonomy, typing methods and pathogenicity factors". *Clin Microbiol Rev* 11 (4): 589–603. PMID 9767057.

Postgate, J. (1998): Nitrogen fixation, 3rd ed. Cambridge University Press.

Rajesh, A. (2009): Answer.com. Director, Urology and Renal transplantation.
Fortis Hospitals. New Delhi.

Raphael, M.M. and Patrick O. E. (2006): Susceptibility of common urinary isolates to the commonly used antibiotics in a tertiary hospital in southern Nigeria. Department of Medical Microbiology, University of Benin Teaching Hospital, Benin City, Nigeria.

Rebecca, B. (2005): Blood agar plates and hemolysis: *Streptococcus* and Other catalase negative Gram-positive cocci University of Utah, Department of Pathology, Salt Lake City, UT 84132. USA.

Ress, T.; Minney, Y.; Niskikira, K. and Mastuoka, S. (2000): Bacteria isolated from UTI and their drug sensitivity; statistical studies from 1966 to 1970 (Japanese); *Cent. Clin. Lab.; Univ. Hosp.; Yokohama Jap. Clin. Urol.*; 27/2 (155-162).

Robert, C.R.; Valkenburg, H.A.; Haverkorn, M.J. and Goslings, W.H. (1991): A prospective study of urinary tract infection in a Dutch general practice. *Lancet* 2: 674-7.

Ronald, A. (2002): The etiology of urinary tract infection: traditional and emerging pathogens. *Am J Med*; 113(suppl 1A): 14S-19S.

Ronald, A. and Ludwig, E. (2001): Urinary tract infections in adults with diabetes. *Int. J. Antimicrob. Agents* 17:287-292. Vol. 5 (11), pp. 1067-1071, 2 June 2006 Extended spectrum Beta - Lactamase (EBSL) in *E.coli* isolated from a tertiary hospital in Enugu state, Nigeria

Ronald, A.R. and Pattulo, M.S. (1991): The natural history of urinary infection in adults. *Med. Clin. North Am.*, 75:299-312.

Ruz, E.D.; Leoni, G.; Eugeni, A. and Rafae, L. (2000): Epidemiology and risk factors for urinary tract infections in patients with spinal cord injury. *Journal of Urology*; 164(4): 1285-1289.

Ryan, K.J. and Ray, C.G. (2004): Sherris medical microbiology, 4th ed., McGraw Hill. ISBN 0838585299.

Sabahat, S. and Perween, T. (2005): Antibacterial activities of *Mentha piperita*, *Pisum sativum* and *Momordica charantia*. *Pak. J. Bot.*, 37(4): 997:1001.

Savas, L.; Guvel, S.; Onlen, Y.; Savas3, N. and Duran, N. (2006): Nosocomial urinary tract infections: Micro-organisms, antibiotic sensitivities and risk factors. *West Indian Med J.*, 55 (3): 188.

Saxena D.; Amin, M. and Khanna, S. (1996): Modulation of protein profiles in *Rhizobium* sp. under salt stress. *Can. J. Microbiol.* Val. 42.

Schneider, P.F. and Riley, T.V. (1996): *Staphylococcus saprophyticus* urinary tract infections: epidemiological data from Western Australia. Eur. J. Epidemiol. 12:51-54.

Schaeffer, H. and Anthony, J. (2000): Urinary tract infection. Journal of urology; 163(1):384.

Schaeffer, A.J.; Rajan, N. and Cao, Q. (2001): Host pathogenesis in urinary tract infections. Int. J Antimicrob. Age; 17:245-251.

Schaeffer and Anthony, J. (2003): Infection and inflammation of the genitourinary tract. Journal of Urology; 170(4): 1468-1475.

Shortliffe, L.M. and McCue, J.D. (2002): Urinary tract infection at the age extremes: Pediatrics and geriatrics. Am. J Med; 113: 55S-66S.

Shymaa, A. (2004): Identification and susceptibility testing of non-lactose fermenting Gram-negative bacilli in urinary tract infection. Master Degree, Zagazig University, Faculty of Medicine, Microbiology and Immunology Department.

Silverstein, R.M.; Bassler, G.C. and Morrill, T.C. (1991): Spectrometric identification of organic compounds, 5th Ed., Wiley.

Smaill, F. (2001): Antibiotics for asymptomatic bacteriuria in pregnancy. Cochrane database Syst. Rev. (2):CD000490 [PMID 11405965].

Spach, D.H.; Stapleton, A.E. and Stamm, W.E. (1992): Lack of circumcision increases the risk of urinary tract infection in young men. JAMA 267:679 [PMID 1472171].

Sobel, J.D. and Kaye, D. (2000): Urinary tract infections. In: Mandell G.L.; Bennett J.E. and Dolin R., eds. Principles and practice of infectious diseases. 5th ed. Philadelphia: Churchill Livingstone; 773-805.

Sonnenwirth, A.C. (1980): Gradwohl's clinical laboratory methods and diagnosis Vol. 2; 8th ed. Sonnenwirth, A.C. and Jarett, L. (eds). The CV Mosby Co. London; Toronto.

Stamm, W.E. and Hooton, T.M. (1993): Management of urinary tract infections in adults. N Engl J Med; 329(18):1328-34.

Stamm, W.E. (1998): Urinary tract infections and pyelonephritides. In: Harrison's principles of internal medicine, 14th ed. Ch. 131; 1: 817-827. Fauci A.S.; Braunwald, E.; Isselbacher, K.J., eds. McGraw-Hill, New York.

Stamm, W. (2001): Urinary tract infections and pyelonephritis. In: Braunwald E., Fauci A., Kasper D.L., eds. Harrison's Principles of Internal Medicine. 15th ed. New York: McGraw-Hill Medical Publishing Division; 1620-1629.

Sue, G.D.; Ciaramonte, J.S. and Carofalo, M.M. (2001): Effects of vapors of aromatic chemicals on fungi. J. Amer. Pharm. Assoc.; 50:665-668.

Suslow, T.V.; Schroth, M.N. and Isaka, M. (1982): Application of a rapid method for Gram differentiation of plant pathogenic and saprophytic bacteria without staining. Phytopathology 72: 917-918.

Sussman, M. (1998): Urinary tract infections; 601-621. Microbial infections 9th ed, Topley & Wilson's Microbiology and Collier, L; Balows, A. and Sussman, M. (eds.) Vol. 3 (Bacterial infections), Hausler, W. J., and Sussman, M. (eds.). Oxford University Press.

Svanborg, C.; Godaly, G. and Hedlund, M. (1999): Cytokine responses during mucosal infections: role in disease pathogenesis and host defence. Curr Opin Microbiol; 2(1):99-105.

Szlyk, G.R. and Williams, (2003): Incidence of new renal parenchymal inflammatory changes following breakthrough urinary tract infection in patients

with vesicoureteral reflux treated with antibiotic prophylaxis: Evaluation by M technetium dimercapto-succinic acid renal scan. Journal of urology; 170(4): 1566-1569.

Talan, D.A.; Hoey, R. and Probst, F. (2003): Urinary tract infection. N Engl J Med; 349: 1674-1676.

Talbot, J.P. and Levine, M.M. (1988): Bacterial diarrheas; p.697-752. In AZ. Kapikian (ed); Viral infection of gastrointestinal tract; 2nd ed. Marcel Dekker. Inc.; New York; NY.

Tambyah, P.A. and Maki, D.G. (2000): Catheter-associated urinary tract infection is rarely symptomatic: A prospective study of 1497 catheterized patients. Arch intern Med; 160: 678-82.

Tepe, B.; Daferera, D.; Sokmen, A.; Sokmen, M. and Polissiou, M. (2005): Antimicrobial and antioxidant activities of the essential oil and various extracts of *Salvia tomentosa* (Lamiaceae). Food Chemistry, 90, 333-340.

Toama, M.A.; El-Kersh, T.A. and Ahmady, A.M. (1983): Antibiotic-resistance pattern and resistance curing of *E. coli* locally isolated. Bulletin of Faculty of Pharmacy; Cairo University; Vol. XXII No. 1: 129-146.

Trapp, S. and Croteau, R. (2001): Genomic organization of plant terpene synthases and molecular evolutionary implications. Annu. Rev. Plant. Physiol. Plant. Mol. Biol. 52,689.

Trautner, B.W. and Darouiche, R.O. (2004): Catheter-associated infections: pathogenesis affects prevention. Arch Intern Med; 164: 842-850.

Turnidge, J.; Bell, J. and Biedenbach, D.J. (2002): Pathogen occurrence and antimicrobial resistance trends among urinary tract infection isolates in the

Asia-Western Pacific Region: report from the SENTRY antimicrobial surveillance program, 1998-1999. *Int J Antimicrob Agent*; 20(1):10-17.

Vejlsgaard, R.H. (1999): Susceptibility of the kidney to experimental infection. In: O'Grady, F. and Brumfitt, W.; editors: urinary tract infection; London; Oxford; University Press; p. 24-36.

Walsh, T.J. (1993): Randomized double-blinded trial of rifampin with either novobiocin or trimethoprim-sulfamethoxazole against methicillin-resistant *Staphylococcus aureus* colonization: Prevention of antimicrobial resistance and effect of host factors on outcome. *Antimicrobial agents and chemotherapy*, Vol .37 No 6 p. 1334–1342

Warren, J.W. (1996): Clinical presentation and epidemiology of urinary tract infections. P.3-27. In H.L.T. Mobley and J.W. Warren (ed.), *Urinary Tract Infections. Molecular pathogenesis and clinical management*. American Society of Microbiology, Washington, D.C.

Weber, B. and Rocha, H. (1999): Bacteriuria in apparently healthy young adults.; Dep. Of Med. Hosp. Proi Edgard Santos; Univ of Bahia. Salvador. *J. Uro. (Baltimore)* 102/3: 342-344.

Well, B. and Stood, R.J. (1989): The genus *shigella*. In *Bergey's manual of systematic bacteriology*. Vol. I. 8th ed. Krieg, N.R.; Holt, J.G. and others (eds) Wilkins. Hong Kong., Pp. 423-427.

Wilson, F.; Match, J.A.; Miller, B.R. and Minman, F. (1984): A careful study. Relation of chronic pyelonephritis with diabetes. *J. Urol.*; 90:96.

Wilson, J.W. (1994): Non-obstructive *Escherchia coli* pyelonephritis; in the rat. *Proc. Sec. Exp. Biol. Med.*; 104: 116-120.

Wilson, S.C.; Sorhati, I.; Caspo, G. and Balazs, (1994): The relationship between diabetes mellitus and infection of the urinary tract amer. J. Med. Sci.; 240: 48-141.

Winsbury, S.M. and White, F.P. (1983): Herbal medicines: A guide for health care professionals. The pharmaceutical press: London. Pp 256-257.

Wright, S.W.; Wrenn, K.D. and Haynes, M. (2000): Prevalence and risk factors for multidrug resistant uropathogens in ED patients. Am J Emerg Med; 18:143-146.

Yu, Z.; Wu, H. and Ding, J. (1998): The volatile chemical components of fresh *Zingiber officinale*. Acta Botanica Yunnanica. 20: 113-118.

Zotti, C.M.; Messori Ioli, G. and Charrier, L. (2004): Hospital-acquired infections in Italy: A region wide prevalence study. J Hosp Infect; 56(2): 142-9.

Zunino, P.; Geymonate, L. and Allen, A. (2000): Virulence of a *Proteus mirabilis* ATF isogenic mutant is not impaired in a mouse model of ascending urinary tract infection. FEMS-Immunology and Medical Microbiology; 29: 137-144.
