
References

- Aal, A.A.A.; Abou Eisha, A.M. and El-Sheary, M.W. (1999):** Prevalence of fascioliasis among man and animals in Ismalia province. Assuit. Vet. Med. J., 41: 141-152.
- Abdo-Rabo, T.M.A. (1991):** Incidence and seasonal prevalence of trematodes infestation among cattle, buffaloes and sheep in Kafr-El-Sheikh province. J. Egypt. Med. Ass., 51(1-2): 561-570.
- Ahmadi, M.R.; Nazifi, S.; Ghaisari, H.R. (2006):** Comparison of hormonal changes of estrus cycle with cytology of cervical mucosa and hematological parameters in dairy heifers. Comp. Clin. Pathol., 15: 94-97.
- Ahmed, W.A.; Nabil, G.M.; El-khadrawy, H.H.; Hanafi, E.M. and Adel-Moez, S.I. (2006):** Monitoring progesterone level and markers of oxidative stress in blood of buffalo-cows with impaired fertility. Egypt. J. Biophys. Biomed. Engineering, 7: 71-83.
- Ahmed, W.M. (2006):** Adverse conditions affecting ovarian activity in large farm animals. Proceeding of the 3rd International Conference of Vet. Res. Div., NRC., Cairo, Egypt, 251-253.
- Ahmed, W.M. (2007):** Overview on some factors negatively affecting ovarian activity in large farm animals. Global Vet., 1: 53-66.
- Ahmedullah, F.; Akbor, M.; Haider, M.G.; Hossain, M.M.; Khan, M.A.H.N.A.; Hossain, M.I. and Shanta, I.S. (2007):** Pathological investigation of liver of the slaughtered buffaloes in Barisal district. Bangl. J. Vet. Med., 5 (1-2): 81-85.
- Ali, H.; Ai, L.; Song, H.Q.; Ali, S.; Lin, R.Q.; Seyni, B.; Issa, G. and Zhu, X.Q. (2008):** Genetic characterization of *Fasciola* samples from different host species and geographical localities revealed the existence of *F. hepatica* and *F. gigantica* in Niger. Parasitol. Res. 102(5): 1021-1024.
- Almazán, C.; Avila, G.; Quiroz, H.; Ibarra, F. and Ochoac, P. (2001):** Effect of parasite burden on the detection of *Fasciola hepatica* antigens in sera and feces of experimentally infected sheep. Vet. Parasitol., 97: 101-112.

- Alvarez-Sanchez, M.A.; Mainar-Jaime, R.C.; Perez-Garcia, J. and Rojo-Vazquez, F.A. (2006):** Resistance of *Fasciola hepatica* to Triclabendazole and albendazole in sheep in Spain. Vet. Rec., 159: 424-425.
- Aly, M.E.M. (1993):** Some biochemical and serological studies on gastrointestinal helminth infections in cattle and buffaloes in Dakahlia Governorate. Ph.D. Thesis, Vet. Med., Cairo Univ.
- Anderson, P.H.; Berrett, S.; Brush, P.J.; Hebert, C.N.; Parfitt, J.W. and Patterson, D.S. (1977):** Biochemical indicators of liver injury in calves with experimental fascioliasis. Vet. Rec., 100: 43-45.
- Anonymous (2003):** Fascioliasis causes losses in cattle, sheep and camelids. Vet. Rec., 152: 189-192.
- Anwer, H.A.; Hayat, C.S. and Amir, M.I. (1996):** Prevalence of gastrointestinal helminthiasis and comparative efficacy of anthelmintics in parasitized buffalo calves. Pakistan. Vet. J., 16: 160-163.
- Aris, M.; Morrondo, P.; Hillyer, G.V.; Sánchez-Andrade, R.; Suárez, J.L.; Lomba, C.; Pedreira, J.; Díez-Baños, P. and Paz-Silva, A. (2007):** Immunodiagnosis of current fasciolosis in sheep naturally exposed to *Fasciola hepatica* by using a 2.9 kDa recombinant protein. Vet. Parasitol., 146: 46-49.
- Arosh, J. A.; Kathiresan, D.; Devannathan, T. G.; Rajasundaram, C. R. and Rajasekaran, J. (1998):** Blood biochemical profile in normal cyclical and anestrus cows. Ind. J. Anim. Sci., 68: 1154-56.
- Association of Official Analytical Chemists (AOAC) (1990):** Official Methods of Analysis (13th Ed.), Washington, D.C. U.S.A.
- Atkins, J.A.; Busch, D.C.; Bader, J.F.; Keisler, D.H.; Patterson, D.J.; Lucy, M.C. and Smith, M.F. (2008):** Gonadotropin-releasing hormone induced ovulation and luteinizing hormone release in beef heifers: effect of day of the cycle. J. Anim. Sci., 86: 83-93.
- Ayob, A.A.M. (1983):** The interpretation of different tests used for estimation of parasitic state of *Fasciola gigantica* in Gharbia Governorate. M. Vet. Sci. Thesis, Cairo Univ., Egypt.
- Barakat, A.M.; Hanafi, E.M.; Sabra, H.A.; Zaabal, M.M. and Ahmed, W.M. (2001):** Effect of parasitic infection on ovarian activity in native Egyptian cows and ewes with special reference to changes in blood constituents and immunogenetic markers. Zagazig. Vet. J., 29: 121-136.

- Baratta, M.; Grasselli, F. and Tamanini, C. (1994):** Effects of gonadal steroids on tonic luteinizing hormone (LH) release and luteinizing hormone releasing hormone-induced LH release from bovine pituitary cells cultured *in vitro*. Biol. Reprod., 50: 1320–1327.
- Bartolome, J.A.; van Leeuwen, J.J.J.; Thieme, M.; Sa’filho, O.G.; Melendez, P.; Archbald, L.F. and Thatcher W.W. (2009):** Synchronization and resynchronization of inseminations in lactating dairy cows with the CIDR insert and the Ovsynch protocol. Theriogenol., 72: 869-878.
- Batzar, F. (1980):** Hormonal evaluation of early pregnancy. Fertil. Steril., 34: 1-13.
- Bauman, J. (1981):** Basal body temperature: unreliable method of ovulation detection. Fertil. Steril., 36: 729-733.
- Behm, C.A. and Sangester, N.C. (1998):** Pathology, pathophysiology and clinical aspects. In: Fasciolosis (Ed. Dalton, J.P.), CAB Inter. Publishing, Oxon, pp. 185-224.
- Belfield, A. and Goldberg, D.M. (1971):** Revised assay for serum phenyl phosphatase activity using 4-amino-antipyrine. Enzyme, 12(5): 561-573.
- Bello, N.M.; Steibel, J.P. and Pursley, J.R. (2006):** Optimizing ovulation to first GnRH improved outcomes to each hormonal injection of Ovsynch in lactating dairy cows. J. Dairy Sci., 89: 3413-3424.
- Broadbent, P.J.; Tregaskes, L.D.; Dolman, D.F.; Franklin, M.F., and Jones, R.L. (1993):** Synchronization of estrus in embryo transfer recipients after using a combination of PRID or CIDR-B plus PGF2 α . Theriogenol., 39: 1055–1065.
- Bulgin, M.S.; Anderson, B.C.; Hall, R.F. and Lang, B.Z. (1984):** Serum gamma glutamyl transpeptidase activity in cattle with induced fascioliasis. Res. Vet. Sci., 37: 167-171.
- Burt, A.D. and James O.F.W. (1994):** Pathophysiology of Liver. In: Pathology of Liver. Churchill Livingstone. Macsween, R.N.M.; Peter, P.A.; Peter, J.S.; Alistair, D.; Burt, A.D. and Brevard, C.P. (eds.), Edinburgh, London, Madrid, Melbourne, New York and Tokyo, pp: 57.

- Chauvin, A.; Moreau, E. and Boulard, C. (2001):** Responses of *Fasciola hepatica* infected sheep to various infection levels. Vet. Res., 32: 87-92.
- Chen, M. G. and Mott, K.E. (1990):** Progress in morbidity due to *Fasciola hepatica* infection. Trop. Dis. Bull., 87: 1–37.
- Clery, D.; Torgerson, P. and Mulcahy, G. (1996):** Immune responses of chronically infected adult cattle to *Fasciola hepatica*. Vet. Parasitol., 62: 71-82.
- Colazo, M.G.; Gordon, M.B.; Rajamahendran, R.; Mapletoft, R.J. and Ambrose, D.J. (2009):** Pregnancy rates to timed artificial insemination in dairy cows treated with gonadotropin-releasing hormone or porcine luteinizing hormone. Theriogenol., 72: 262–270.
- Colazo, M.G.; Small, J.A.; Ward, D.R.; Erickson, N.E.; Kastelic, J.P. and Mapletoft, R.J. (2004):** The effect of presynchronization on pregnancy rate to fixed-time AI in beef heifers subjected to a Cosynch protocol. Reprod. Fertil. Dev., 16: 128.
- Curtale, F.; Hassanein, Y.A. and Savioli, L. (2005):** Control of human fascioliasis by selective chematography: design, cost and effect of the first public health, school-based intervention implemented in endemic areas of the Nile Delta, Egypt. Trans. R. Soc. Trop. Med. Hyg., 99(8): 599-609.
- Dalton, J.P. (1999):** Fasciolosis. New York: CAB International Publishing, Cambridge University Press, pp. 113-149.
- Değer, Y.; Ertekín, A. Değer, S. and Mert, H. (2008):** Lipid peroxidation and antioxidant potential of sheep liver infected naturally with Distomatosis. Türk. Parazitol. Derg., 32 (1): 23-26.
- Díaz-Torga, G.S.; Mejía, M.E.; González-Iglesias, A.; Formía, N.; Becú-Villalobos, D. and Lacau-Mengido, I.M. (2001):** Metabolic cues for puberty onset in free grazing Holstein heifers naturally infected with nematodes. Theriogenol., 56: 111-122.
- Doumas, B.T.; Watson, W.A. and Biggs, H.G. (1971):** Albumin standards and the measurement of serum albumin with bromcresol green. Clin. Chem. Acta, 31(1): 87-96.
- Dreux, C. (1977):** Selected method. Analysis of human serum: assay of iron II. Method using bathophenanthroline Se-iron II (bathophenanthroline). Ann. Biol. Clin. (Paris), 35(3): 275-277.

- Eden, A. and Green, H.H. (1940):** Micro-determination of copper in biological material. *Biochem. J.*, 34(8-9): 1202-1208.
- El Magdoub, A.; El Sayed, I.A.; Samak, M.A.; Hassan, A.A. and El Sawy, M.F. (1993):** Effects of ration and *Fasciola* infection on some blood parameters of ewes. *J. Egypt. Soc. Parasitol.*, 3: 741-749.
- El-Khadrawy, H.H.; El Moghazy, F.M.; Abd El Aziz M.M. and Ahmed, W.M. (2008):** Field Investigation on the Correlation Between Ovarian Activity and Fasciolosis in Buffalo-Cows. *American-Eurasian J. Agri. Environ. Sci.*, 3(4): 539-546.
- El-Merzabani, M.M.; El-Aaser, A.A. and Zakhary, N.I. (1977):** A new method for determination of inorganic phosphorus in serum without deproteinization. *J. Clin. Chem. Clin. Biochem.*, 15(12): 715-718.
- El-Refaei, S.A.; Bassiouny, G.A.; Marie, N.A.M. and Moris, E. (1984):** Concomitant hepatic *Fasciola* and hydatid infections in animals. *J. Egypt. Soc. Parasitol.*, 14(2): 421-427.
- El-Reffai, A.H. (1993):** *Entamoeba bovis*, Liebetanz 1905, recorded from large ruminants in Egypt. *J. Egypt. Soc. Parasitol.*, 23(1): 239-246.
- El-Ridi, R.; Salah, M.; Wagih, A.; William, H.; Tallima, H.; El-Shafie, M.H.; Abdel Khalek, T.; El- Amir, A.; AboAmmou, F.F. and Motawi, H. (2007):** *Fasciola gigantica* excretory-secretory products for immunodiagnosis and prevention of sheep fasciolosis. *Vet. Parasitol.*, 149: 290-293.
- El-Shazly, A.M.; el-Nahas H.A.; Soliman, M.; Sultan, D.M.; Abdel Tawab, A.H. and Morsy, T.A. (2006):** The reflection of control programs of parasitic diseases upon gastrointestinal helminthiasis in Dakahlia Governorate, Egypt. *J. Egypt. Soc. Parasitol.*, 36(2): 467-480.
- Eltohamy, M.M.; Ahmed, W.M. and Abdoon, A.S. (1994):** Ovarian activity in relation to some blood composition. *Proc. IV Wld. Buff. Cong.*, São Paulo, Brazil, pp: 612-614.
- El-Wishy, A.B. (2007):** The postpartum buffalo, I. Endocrinological changes and uterine involution. *Anim. Reprod. Sci.*, 97(3-4): 201-215.

- El-Zarkouny, S.Z.; Cartmill, J.A.; Hensley, B.A. and Stevenson, I.S. (2004):** Pregnancy in dairy cows after synchronized ovulation regimens with or without pre synchronization and progesterone. *J. Dairy Sci.*, 87: 1024-1037.
- Espino, A.M. and Hillyer, G.V. (2003):** Molecular cloning of a member of the *Fasciola hepatica* saposine-like protein family. *J. Parasitol.*, 89: 545-552.
- Esteban, J.G.; Gonzalez, C.; Curtale, F.; Munoz-Antoli, C.; Valero, M.A.; Bargues, M.D.; El-Sayed, M.; El-Wakeel, A.A.W.; Abdel-Wahab, Y.; Montresor, A.; Engels, D.; Savioli, L. and Mas-Coma, S. (2003):** Hyperendemic fascioliasis associated with schistosomiasis in villages in the Nile Delta of Egypt. *Am. J. Trop. Med. Hygiene*, 69: 429-437.
- Ezzat, E.A.; Mousa, A.A.; Eid, R.S.A. and Ashour, A.A. (1994):** Control of fascioliasis. National of General Vet. Org. and Ciba Geigy, Egypt.
- FAO (2005):** Food and Agriculture Organization, Year Book of production, United Nations.
- Farag, H.F. (1998):** Human fascioliasis in some countries of the Eastern Mediterranean Region. *East. Medit. Health J.*, 4(1): 156-160.
- Ferre, I.; López, P.; Gonzalo-Orden, M.; Julian, M.D.; Rojo, F.A. and González-Gallego, J. (1995 a):** The effects of subclinical fascioliasis on hepatic secretory function in sheep. *Parasitol. Res.*, 81: 127-131.
- Ferre, I.; Ortega, L.M. and Rojo-Vazquez, F.A. (1995 b):** Sero-prevalence of *Fasciola hepatica* infection in sheep in Northern Spain. *Parasitol. Res.*, 81(2): 137-142.
- Fleming, M.W. and Fetterer, R.H. (1986):** Peripheral androgen levels in peripuberal rams infected with *Fasciola hepatica*. *Vet. Parasitol.*, 19(3-4): 295-299.
- Forshell, K.P.; Anderson, L. and Pehrson, B. (1991):** The relationships between the fertility of dairy cows and clinical and biochemical measurements with special reference to plasma glucose and milk acetone. *J. Vet. Med.*, 38(A): 608-616.
- Gajewska, A.; Smaga-Kozłowska, K. and Wiśniewski, M. (2005):** Pathological changes of liver in infection of *Fasciola hepatica*. *Wiad Parazytol.*, 51(2): 115-123.

- Ganga, G.; Varshney, J.P.; Sharma, R.L.; Varshney, V.P. and Kalicharan (2007):** Effect of *Fasciola gigantica* infection on adrenal and thyroid glands of riverine buffaloes. Res. Vet. Sci., 82: 61-67.
- Geary, W.; Whiter, J.C.; Downing, E.R.; Lefever, D.G.; Silcox, R.W.; Holland, M.D.; Nett, T.M. and Niswender, G.D. (1998):** Pregnancy rates of postpartum beef cows that were synchronised using Synchromate-B or the ovysynch protocol. J. Anim. Sci., 76: 1523-1527.
- Ghirmire, N.P. and Karki, N.P.S. (1996):** Prevalence of fascioliasis and efficacy of various anthelmintics in buffaloes of Rural Kathamandu vetcon. N. U. A., pp. 43.
- Gindler, M. and King, J.D. (1972):** Rapid colorimetric determination of calcium in biologic fluids with methylthymol blue. Am. J. Clin. Pathol., 58(4): 376-382.
- Gonzalo-Orden, M.; Millán, L.; Álvarez, M.; Sánchez-Campos, S.; Jiménez, R.; González-Gallego, J. and Tuñón, M.J. (2003):** Diagnostic imaging in sheep hepatic fascioliasis: ultrasound, computer tomography and magnetic resonance findings. Parasitol. Res., 90: 359-364.
- Gornal, A.C.; Bardawill, C.J. and David, M.M. (1949):** Determination of serum proteins by means of the biuret reaction. J. Biol. Chem., 177(2): 751-766.
- Hafeez, M.D. (2003):** Helminth parasites of public health importance-Trematodes. J. Parasit. Dis., 27(2): 69-75.
- Hanna, R.E.; Cromie, L.; Taylor, S.M. and Couper, A. (2006):** The effect of a parenteral ivermectin/closantel injection on the growth and reproductive development of early immature *Fasciola hepatica* in cattle. Vet. Parasitol., 142(1-2): 78-90.
- Haseeb, A.N.; el-Shazly A.M.; Arafa, M.A. and Morsy A.T. (2002):** A review on fascioliasis in Egypt. J. Egypt. Soc. Parasitol., 32(1): 317-354.
- Hillyer, G.V.; Galanes, M.S.D.; Buchon, P. and Bjorland, J. (1996):** Herd evaluation by enzyme linked immunosorbent assay for the detection of *Fasciola hepatica* infection in sheep and cattle from Altipano of Bolivia. Vet. Parasitol., 61: 211-220.

- Hutchinson, G.W.; Dawson, K.; Fitzgibbon, C.C. and Martin, P.J. (2009):** Efficacy of an injectable combination anthelmintic (nitroxynil+clorsulon+ivermectin) against early immature *Fasciola hepatica* compared to triclabendazole combination flukicides given orally or topically to cattle. *Vet. Parasitol.*, 162: 278-284.
- Ibrahim, M.A. (2004):** Epidemiological studies on fascioliasis in the Beni-Suef Governorate. *Vet. Med. J. Giza*, 52: 507-515.
- Ishwar, A.K. (1994):** Enzyme activity in Black Bengal goats following estrus synchronization and superovulation. *Small Ruminant Res.*, 13(3): 301-305.
- Ishwar, A.K. and Pandey, J.N. (1994):** Blood metabolite changes in Black Bengal goats following estrus synchronization and superovulation. *Small Ruminant Res.*, 13(3): 251-256.
- Janovick, J.A. and Conn, P.M. (1996):** Progesterone diminishes the sensitivity of gonadotropin releasing hormone stimulated luteinizing hormone (LH) release and protects an LH pool from desensitization: actions opposed by cholera toxin. *Endocrinol.*, 137: 1823-1827.
- Johnson, S.K.; Johnson, A.R.; Keefer, C.L. and Silcox, R.W. (1990):** Blood constituents during the estrous cycle and early pregnancy in dairy cows. *Theriogenol.*, 34: 701-707.
- Kakar, M.N. and Kakarsulemankhel, J.K. (2008):** Prevalence of endo (trematodes) and ecto-parasites in cows and buffaloes of Quetta, Pakistan. *Pakistan. Vet. J.*, 28(1): 34-36.
- Kendal, S.B. (1975):** Some parasites of domestic animals in Aswan Governorate, A.R.E. *Trop. Anim. Health Prod.*, 6(3): 128-130.
- Kerbler, T.L.; Buhr, M.M.; Jordan, L.T.; Leslie, K.E. and Walton, J.S. (1997):** Relationship between maternal plasma progesterone concentration and interferon-tau synthesis by the conceptus in cattle. *Theriogenol.*, 47: 703-14.
- Khalil, M.Z. (1979):** A study of fascioliasis in farm animals and man M.D. *Vet. Parasit.*, Thesis, Alex. Univ., Egypt.
- Kiladze, M.; Chipashvili, I.; Abuladze, D. and Jatchvliani, D. (2000):** Obstruction of common bile duct caused by liver fluke- *Fasciola hepatica*. *Sb. Lek.*, 101: 255-259.

- Kim, L.H.; Suh, G.H.; Nam, N.W.; Kang, R.G. and Kim, L.H. (2005):** Follicular wave emergence, luteal function and synchrony of ovulation following GnRH or estradiol benzoate in a CIDR-treated, lactating Holstein cows. *Theriogenol.*, 63: 260-268.
- Kohram, H.; Mohammadi, G. and Dirandeh, E. (2008):** Alteration of follicular dynamics by GnRH at two different stages of the estrous cycle in Water buffalo. *J. Anim. Vet. Adv.*, 7(8): 996-1001.
- Kumar, N.; Gosh, S. and Gupta, S.C. (2008):** Early detection of *Fasciola gigantica* infection in buffaloes by enzyme-linked immunosorbent assay and dot enzyme-linked immunosorbent assay. *Parasitol. Res.*, 103(1): 141-150.
- Kumar, S. and Sharma, M.G. (1991):** Infertility in rural cows in relation to Fasciolosis. *Ind. J. Anim. Sci.*, 61: 838- 840.
- Lacau-Mengido, I.M.; Mejía, M.E.; Díaz-Torga, G.S.; Gonzalez, I.A.; Formía, N.; Libertun, C. and Becú-Villalobos, D. (2000):** Endocrine studies in ivermectin-treated heifers from birth to puberty. *J. Anim. Sci.*, 78: 817-824.
- Long, S.T.; Yoshida, C. and Nakao, T. (2009):** Plasma progesterone profile in ovariectomized beef cows after intra-vaginal insertion of new, once-used or twice-used CIDR. *Reprod. Domest. Anim.*, 44(1): 80-82.
- Lopez, P.; Gonzalez, P.; Tuñon, M.J. and Gonzalez-Gallego, J. (1994):** The effects of experimental fasciolosis on bilirubin metabolism in the rat. *Exp. Parasitol.*, 78(4): 386-393.
- Lopez-Diaz, M.C.; Carro, M.C.; Cadorniga, C.; Diez-Banos, P. and Mezo, M. (1998):** Puberty and serum concentrations of ovarian steroids during prepuberal period in Friesian heifers artificially infected with *Fasciola hepatica*. *Theriogenol.*, 50(4): 587-593.
- Lotfollahzadeh, S.; Mohri, M.; Ranjbar Bahadori, Sh.; Mokhber Dezfouly, M.R. and Tajik, P. (2008):** The relationship between normocytic, hypochromic anaemia and iron concentration together with hepatic enzyme activities in cattle infected with *Fasciola hepatica*. *J. Helminth.*, 82: 85-88.
- Lotfy, A.; Youssef, H.; Nassar, A. and Abdel-Aziz, N. (1995):** Fascioliasis in slaughtered animals, The incidence and public health importance. *Fleischwirt*, 75(6): 803-804.

- Lotfy, H.S. (2001):** Correlation between *F. gigantica* infection and reproductive models in farm animals before and after treatment. Beni-Suef Vet. Med. J., 11: 705-714.
- Lowry, O.H.; Rosenbrough, N.J.; Farr, A.L. and Randall, R.J. (1951):** Protein measurement with Folin-phenol reagent. J. Biol. Chem., 193: 265-275.
- Loyacano, A.F.; Williams, J.C.; Gurie, J. and DeRosa, A.A. (2002):** Effect of gastrointestinal nematode and liver fluke infections on weight gain and reproductive performance of beef heifers. Vet. Parasitol., 107(3): 227-234.
- Mann, G.E. and Lamming, G.E.(2001):** Relationship between maternal endocrine environment, early embryo development and inhibition of the luteolytic mechanism in cows. Reprod., 121: 175-180.
- Mann, G.E. (2002):** Corpus luteum function and early embryonic death in the bovine. In: Proc. XXII World Buiatric Congress 2002, August, 18-23: 300-306.
- Maqbool, A.; Anjum, A.D.; Ahmad, M. and Raza, A. (1999):** Efficacy of rafoxanide against natural infection of fascioliasis in buffaloes with particular reference to metabolic parameters. Buff. J., 15(3): 353-359.
- Maqbool, A.; Hayat, C.S.; Akhtar, T. and Hashmi, H.A. (2002):** Epidemiology of fasciolosis in buffaloes under different managemental conditions. Vet. Arhiv, 72: 221-228.
- Marcilla, A.; Bargues, M.D. and Mas-Coma, S. (2002):** A PCR-RFLP assay for the distinction between *Fasciola hepatica* and *Fasciola gigantica*. Mol. Cell. Probes, 16(5): 327-333.
- Marques, S.M.T. and Scroferneker, M.L. (2003):** *Fasciola hepatica* infection in cattle and buffaloes in the State of Rio Grande do Sul, Brazil. Parasitol. Latinoam, 58: 169-172.
- Martínez, M.F.; Adams, G.P.; Bergfelt, D.R.; Kastelic, J.P. and Mapletoft, R.J. (1999):** Effect of LH or GnRH on the dominant follicle of the first follicular wave in beef heifers. Anim. Reprod. Sci., 57: 23-33.
- Mas-Coma, S.; Bargues, M.D. and Valero, M.D. (2005):** Fascioliasis and other plant-borne trematode zoonoses. Int. J. Parasitol., 35(11-12): 1255-1278.

- Mehra, U.R.; Verma, A.K.; Dass, R.S.; Sharma, R.L. and Yadav, S.C. (1999):** Effects of *Fasciola gigantica* infection on growth and nutrient utilisation of buffalo calves. Vet. Rec., 145(24): 699-702.
- Mejía, M.E.; Libertun, C.; Díaz-Torga, G.S.; Villafane, P.; Formía, N.; Becú-Villalobos, D. and Lacau-Mengido, I.M. (1999):** Continuous ivermectin treatment from birth to puberty on growth and reproduction in dairy heifers. J. Anim. Sci., 77: 1329-1334.
- Mekroud, A.; Chauvin, A. and Rondelaud, D. (2007):** Variations of biological indicators as highly presumptive markers for fasciolosis in experimentally-infected sheep. Rev. Méd. Vét., 158(8-9): 437-441.
- Mendes, E.A.; Lima, W.S. and de Melo, A.L. (2008):** Development of *F. hepatica* in *Lymnaea columella* infected with miracidia derived from cattle and marmoset infections. J. Helminth., 82: 81-84.
- Mitchell, G.B.; Maris, L. and Bonniwell, M.A. (1998):** Triclabendazole resistant liver flukes in Scottish sheep. Vet. Rec., 143: 399.
- Molina, E.C. and Skerratt, L.F. (2005):** Cellular and humoral responses in liver of cattle and buffaloes infected with a single dose of *Fasciola gigantica*. Vet. Parasitol., 131(1-2): 157-63.
- Molina, E.C.; Gonzaga, E.A. and Lumbao, L.A. (2005 a):** Prevalence of infection with *Fasciola gigantica* and its relationship to carcass and liver weights, and fluke and egg counts in slaughter cattle and buffaloes in Southern Philippines. Trop. Anim. Health Prod., 37(3): 215-221.
- Molina, E.C.; Sinolinding, E.O. and Peralta, A.A. (2005 b):** Clinicopathological indications of resilience of swamp buffaloes to infection with *F. gigantica*. Trop. Anim. Health Prod., 37: 451-455.
- Molina, E.C.; Lozano, S.P. and Barraca, A.P. (2006):** The relationship between haematological indices, serum gamma-glutamyl transferase and glutamate dehydrogenase, visual hepatic damage and worm burden in cattle infected with *Fasciola gigantica*. J. Helminth., 80: 277-279.
- Molina, E.C.; Skerratt, L.F. and Campbell, R. (2008):** Overcoming liver fluke in southeast Asia, Pathology of fasciolosis in large ruminants, 94-98.

- Moll, L.; Gaasenbeek, C.P.; Vellema, P. and Borgsteede, F.H. (2000):** Resistance of *Fasciola hepatica* against triclabendazole in cattle and sheep in The Netherlands. Vet. Parasitol., 91(1-2): 153-158.
- Molloy, J.B.; Anderson, G.R.; Fletcher, T.I.; Landmann, J. and Knight, B.C. (2005):** Evaluation of commercially available enzyme-linked immunosorbent assay for detecting antibody to *Fasciola hepatica* and *Fasciola gigantica* in cattle, sheep and buffaloes in Australia. Aust. Vet. Parasitol., 130: 207-212.
- Munguía-Xóchihua, J.A.; Ibarra-Velarde, F.; Ducoing-Watty, A.; Montenegro-Cristino, N. and Quiroz-Romero, H. (2007):** Prevalence of *Fasciola hepatica* (ELISA and fecal analysis) in ruminants from a semi-desert area in the northwest of Mexico. Parasitol. Res., 101: 127-130.
- Oldham, G. (1983):** Antibodies to *Fasciola hepatica* antigens during experimental infections in cattle measured by ELISA. Vet. Parasitol., 13: 151-158.
- Orellana, P.; Recabarren, S.; Lobos, A.; Islas, A.; Briones, M. and Rubilar, L. (1999):** Effects of winter supplementation and anti-parasite treatment on the productive performance of milk herd in the central-south region of Chile. Preventive Vet. Med., 38: 207-215.
- Overend, D.J. and Bower, F.L. (1995):** Resistance of *Fasciola hepatica* to triclabendazole. Aust. Vet. J., 72: 275-276.
- Pachauri, S.P. (1995):** Clinico-biochemical profiles in fasciolosis of buffaloes. Buff. Bull., 14 (1): 7-10.
- Paczkowski, M.J. (2004):** Effects of experimental fascioliasis on puperty and comparison of mounting activity by Radiotelemetry in pupertal and gestating beef heifers. M. Sci. Thesis, Texas Univ.
- Pal, S. and Dasgupta, C.K. (2006):** Haemto-Biochemical profiles of buffalo in anthelmintics treatment against *Fasciola gigantica* infection. Buff. Bull., 25(2): 40.
- Paul, S.S.; Chawla, D.S. and Lall, D. (2000) :** Serum mineral profile and its relationship with reproductive disorders in Nili Ravi buffaloes. Ind. J. Anim. Nutrition, 17: 324-327.

- Periago, M.V.; Valero, M.A.; El Sayed, M.; Ashrafi, K.; El Wakeel, A.; Mohamed, M.Y.; Desquesnes, M.; Curtale, F. and Mas-Coma, S. (2008):** First phenotypic description of *Fasciola hepatica*/*Fasciola gigantica* intermediate forms from the human endemic area of the Nile Delta, Egypt. *Infect. Genet. Evol.*, 51–58.
- Pfeifer, L.F.M.; Siqueira, L.G.; Mapletoft, R.J.; Kastelic, J.P.; Adams, G.P.; Colazo, M.G. and Singh, J. (2009):** Effects of exogenous progesterone and cloprostenol on ovarian follicular development and first ovulation in prepubertal heifers. *Theriogenol.*, 72: 1054–1064.
- Pfukenyi, D.M.; Mukaratirwa, S.; Willingham, A.L. and Monrad, J. (2006):** Epidemiological studies of *Fasciola gigantica* infections in cattle in the highveld and lowveld communal grazing areas of Zimbabwe. *Onderstepoort J. Vet. Res.*, 73(1): 37–51.
- Phiri, A.M.; Phiri, I.K.; Sikasunge, C.S. and Monrad, J. (2005):** Prevalence of fasciolosis in Zambian cattle observed at selected abattoirs with emphasis on age, sex and origin. *J. Vet. Med. B. Infectious Diseases of Veterinary Public Health*, 52(9): 414–416.
- Prasitirat, P.; Thammasart, S.; Chompoochan, T.; Nithiuthai, S. and Taira, S. (1996):** The dynamics of antibody titres and faecal egg output in cattle and buffalo following infection with 500 and 1,000 *Fasciola gigantica* metacercariae. *Thai. J. Vet. Med.*, 26: 85–89.
- Promdireg, A.; Presicce, G.A.; De Rensis, F.; Singlor, J. and Techakumphu, M. (2008):** Follicular Dynamics Following Estrus Synchronization in Swamp Buffalo Cows (*Bubalus bubalis*). *Thai. J. Vet. Med.*, 38(4): 25–34.
- Pursley, J.R.; Kosorok, M.R. and Wiltbank, M.C. (1997):** Reproductive management of lactating dairy cows using synchronization of ovulation. *J. Dairy Sci.*, 80: 301–306.
- Pursley, J.R.; Mee, M.O. and Wiltbank, M.C. (1995):** Synchronization of ovulation in dairy cows using PGF₂ α and GnRH. *Theriogenol.*, 44: 915–923.
- Raadsma, H.W.; Kingsford, N.M.; Suharyanta; Spithill, T.W. and Piedrafita, D. (2008):** Host responses during experimental infection with *Fasciola gigantica* or *Fasciola hepatica* in Merino sheep: I. Comparative immunological and plasma biochemical changes during early infection. *Vet. Parasitol.*, 154(3–4): 250–261.

- Racioppi, O.; Moriena, R.A.; Alvarez, J.D.; Crudeli, G.A. (2007):** *Fasciola hepatica*: New buffalo (*Bubalus bubalis*) parasitosis in the province of Corrientes, Argentina. Ital. J. Anim. Sci., 6(2): 948-950.
- Radostitis, O.M.; Gay, C.C.; Hinchcliff, K.W. and Constable, P.D. (2007):** Veterinary medicine. 10th Ed. Edinburgh, Saunders Co. 2047-2050.
- Radwan, I.G.H. (1996):** Prevalence of cryptosporidiosis and fascioliasis in farm animals and their case-takers. M. Thesis, Vet. Med., Cairo Univ.
- Randel, R.D.; Lammoglia, M.A.; Lewis, A.W.; Neuendorff, D.A. and Guthrie, M.J. (1996):** Exogenous PGF(2)alpha enhanced GnRH-induced LH release in postpartum cows. Theriogenol., 45: 643-654.
- Rastegarnia, A.; Niasari-Naslaji, A.; Hovareshti, P.; Sarhaddi, F. and Safaei, M. (2004):** The effect of different doses of gonadorelin on ovarian follicle dynamics in river buffalo (*Bubalus bubalis*). Theriogenol., 62(7): 1283-1291.
- Razzaque, W.A.A.; Sahatpure, S.K.; Pawshe, C.H. and Kuralkar, S.V. (2008):** Biometry of ovaries and follicular count in cyclic and non-cyclic Nagpuri buffaloes. Buff. Bull., 27: 150-153.
- Reichel, M.P. (2002):** Performance characteristics of an enzyme-linked immunosorbent assay for the detection of liver fluke (*Fasciola hepatica*) infection in sheep and cattle. Vet. Parasitol., 107: 65-72.
- Reitman, A. and Frankel, S. (1957):** A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. Am. J. Clin. Pathol., 28(1): 56-63.
- Rickard, L.G.; Zimmerman, G.L.; Hoberg, E.P.; Bishop, J.K. and Pettitt, R.J. (1992):** Influence of ivermectin and clorsulon treatment on productivity of a cow-calf herd on the southern Oregon coast. Vet. Parasitol., 41(1-2): 45-55.
- River-Marrero, C.A.; Santiago, N. and Hillyer, G.V. (1988):** Evaluation of immuno-diagnostic antigens in the excretory-secretory products of *Fasciola hepatica*. J. Parasitol., 74: 646-652.
- Roman-Ponce, H.; Caton, D.; Thatcher, W.W. and Lehrer, R. (1983):** Uterine blood flow in relation to endogenous hormones during estrous cycle and early pregnancy. Am. J. Physiol. 245: 843-849.

- Roy, K.S. and Prakash, B.S. (2007):** Seasonal variation and circadian rhythmicity of the prolactin profile during the summer months in repeat breeding Murrah buffalo heifers. *Reprod. Fertil. Develop.*, 19: 569-575.
- Saleh, A.M. (2008):** Circulating oxidative stress status in desert sheep naturally infected with *F. hepatica*. *Vet. Parasitol.*, 154: 262-269.
- Salem, A.A.; Shawkat, M.E.; El- Seify, M.A. and Khateib, A. (1990):** Incidence and seasonal prevalence of fascioliasis in Beni-Suef, Egypt. *Assuit Vet. Med. J.*, 22(44): 62-66.
- Salimi-Bejestani, M.R.; Cripps, P. and Williams, D.J.L. (2008):** Evaluation of an ELISA to assess the intensity of *Fasciola hepatica* infection in cattle. *Vet. Rec.*, 162(4): 109-111.
- Samaha, H. (1989):** The zoonotic importance of fascioliasis in Abbis, Alexandria Governorate. *Assuit Vet. Med. J.*, 21(42): 118-122.
- Sanad, M.M. and Al-Megrin, W.A. (2005):** Fascioliasis among local and imported sheep in Saudi Arabia: Parasitological and serological diagnosis. *J. Egypt. Soc. Parasitol.*, 35: 1121-1134.
- Sangsritavong, S.; Combs, D.K.; Sartori, R.; Armentano, L.E. and Wiltbank, M.C. (2002):** High feed intake increases liver blood flow and metabolism of progesterone and estradiol-17 β in dairy cattle. *J. Dairy Sci.*, 85: 2831-2842.
- Sanyal, P.K. (1996):** Kinetic disposition and critical efficacy of triclabendazole against experimental bovine and bubaline fasciolosis. *J. Vet. Parasitol.*, 10(2): 147-152.
- Satish, K. and Sharma, M.C. (1991):** Level of haemoglobin and certain serum biochemical constituents in rural cows during fertile and non-fertile oestrus. *Ind. Vet. J.*, 69: 361-364.
- Schweizer, G.; Braun, U.; Deplazes, P. and Torgerson, P.R. (2005):** Estimating the financial losses due to bovine fasciolosis in Switzerland. *Vet. Rec.*, 157: 188-193.
- Shah, N.H.S.; Willemse, A.H. and Van de Wiel, D.F., (1990):** Reproductive performance of Nili-Ravi buffaloes after a single injection of GnRH early post partum. *Trop. Anim. Health Prod.*, 22(4): 239-246.
- Shaikh, A.A.; Gill, N.; Khan, M. and Bilqees, F.M. (2007):** Biochemical changes in the livers of bovines naturally infected with *Fasciola gigantica*. *Pakistan J. Biol. Sci.*, 10(16): 2756-2759.

- Sharma, R.K., (2003):** Reproductive problems of buffaloes and their management. Compendium of lectures delivered to the Central Institute of Research on Buffalo, Hisar, India, pp: 119-126.
- Sheikh, G.N.; Qadri, S.G.J.; Willayat, M.M. and Das Gunjan (2006):** Biochemical profile of cattle naturally infected with *Fasciola gigantica* and *Fasciola hepatica*. J. Vet. Parasitol., 20: 1.
- Sherwood, L. (2001):** Human Physiology from Cells to Systems. United States: Brooks/ Cole.
- Shirazi, M.; Rasouli, A. and Keshavarz, M. (2007):** The case report of Closantel poisoning in Buffalo calves in Ahvaz, Iran. Iran. J. Sci., 7(23-24): 12-15.
- Simsek, S.; Risvanli, A.; Utuk, A.E.; Yuksel, M.; Saat, N. and Koroglu, E. (2007):** Evaluation of relationship between repeat breeding and *Fasciola hepatica* and hydatid cyst infections in cows in Elazig district of eastern Turkey. Res. Vet. Sci., 83: 102-104.
- Soulsby, E.J. (1982):** Helminthes, Arthropods and Protozoa of Domesticated Animals. 6th Ed., Bailliere, Tindall and Cassell, London.
- Spithill, T.W.; Smooker, P.M. and Copeman, B. (1999):** *Fasciola gigantica*: epidemiology, control, immunology and molecular biology. In: Fasciolosis (Ed. Dalton, J.P.), CAB Inter. Publishing, Wallingford, UK, pp. 465-520.
- SPSS 11 (2002):** Statistical Package for Social Science, SPSS for windows Release 11.0.0, 12 June, 2002."Standard Version, copyright SPSS Inc., 1989-2002".
- Stevenson, J.S.; Tiffany, S.M. and Lucy, M.C. (2004):** Use of estradiol cypionate as a substitute for GnRH in protocols for synchronizing ovulation in dairy cattle. J. Dairy Sci., 87(10): 3298-3305.
- Subandrio, A.L.; Sheldon, I.M. and Noakes, D.E. (2000):** Peripheral and intrauterine neutrophil function in the cow: The influence of endogenous and exogenous sex steroid hormones. Theriogenol., 53: 1591-1608.
- Trinder, P. (1969):** Determination of blood glucose using an oxidase-peroxidase system with a non-carcinogenic chromogen. J. Clin. Pathol., 22(2): 158-166.

- Uehlinger, H.; Binder, H.; Hauser, B., Rusch, P. and Zerobin, K. (1995):** Comparison by hormone analysis of the intravaginal devices CIDR™ and PRID® in ovariectomised cows. Schweiz. Arch. Tierheilk., 137: 81-86.
- Valeroa, M.A.; Gironèsb, N. ; García-Bodelóna, M.A.; Periagoa, M.V.; Chico-Calero, I.; Khoubbanea, M.; Fresnob, M. and Mas-Comaa, S. (2008):** Anaemia in advanced chronic fasciolosis. Acta Trop., 108: 35-43.
- Van Saun, R.J. and Wustenberg, M. (1997):** Metabolic profiling to evaluate nutritional and disease status. Bov. Practitioner, 1(2): 37-42.
- Vasconcelos, J.L.M.; Sangsritavong, S.; Tsai, S.J. and Wiltbank, M.C. (2003):** Acute reduction in serum progesterone concentrations after feed intake in dairy cows. Theriogenol., 60: 795-807.
- Vaughan, J.L.; Charles, J.A. and Boray, J.C. (1997):** *Fasciola hepatica* infection in farmed emus (*Dromaius novaebollandiae*), Aust. Vet. J., 75: 811-813.
- Vengušt, G.; Klinkon, M.; Bidovec, A. and Vengušt, A. (2003):** *Fasciola hepatica*: effects on blood constituents and liver minerals in fallow deer (*Dama dama*). Vet. Parasitol., 112: 51-61.
- Vera Montenegro, Y.; Ibarra Velarde, F.; Quiroz Romero, H.; Hernández Campos, A. and Castillo, R. (2003):** Field trial on the efficacy of an experimental fasciolicide compared with some commercial compounds in naturally infected cattle. Parasitol. Res., 91: 1-4
- Vignoles, P.; Ménard, A.; Rondelaud, D.; Chauvin, A. and Dreyfuss, G. (2001):** *Fasciola hepatica*: the characteristics of experimental infections in *Lymnaea truncatula* subjected to miracidia differing in their mammalian origin. Parasitol. Res., 87: 945-949.
- Walter, M. and Gerade, H., (1970):** Bilirubin assay. Microchem. J., 15: 231.
- Wamae, L.W.; Hammond, J.A.; Harrison, L.J.S. and Onyango-Abuje, J.A. (1998):** Comparison of production losses caused by chronic *Fasciola gigantica* infection in yearling Friesian and Boran cattle. Trop. Anim. Health Prod., 30: 23-30.
- Weems, C.W.; Weems, Y.S. and Randel, R.D. (2006):** Prostaglandins and reproduction in female farm animals. Vet. J., 171: 206-208.

- Welch, S.; Malone, J. and Geaghan, H. (1987):** Herd evaluation of *Fasciola hepatica* infection in Louisiana cattle by ELISA. Am. J. Vet. Res., 48(3): 345-347.
- Wiedosari, E.; Hayakawa, H. and Copeman, B. (2006):** Host differences in response to trickle infection with *Fasciola gigantica* in buffalo, Ongole and Bali calves. Trop. Anim. Health Prod., 38: 43-53.
- Wolfenson, D.; Luft, O.; Berman, A. and Meidan, R. (1993):** Effect of season, incubation temperature and cell age on progesterone and prostaglandin F₂ α production in bovine luteal cells. Anim. Reprod. Sci., 32: 27-40.
- Wyckoff, J.H. and Bradley, R.E. (1985):** Diagnosis of *Fasciola hepatica* infection in beef calves by plasma enzyme analysis. Am. J. Vet. Res., 46: 1015-1019.
- Yadav, S.C.; Sharma, R.L.; Kalicharan, A.; Mehra, U.R.; Dass, R.S. and Verma, A.K. (1999):** Primary experimental infection of riverine buffaloes with *Fasciola gigantica*. Vet. Parasitol., 82: 285-296.
- Yang, Q.; Mao, W.H.; Ferre, I.; Bayón, J.E.; Mao, X.Z. and González-Gallego, J. (1998):** Plasma aspartate aminotransferase (AST), glutamate dehydrogenase (GLDH) and gamma-glutamyl transpeptidase (GGT) activities in water buffaloes with experimental subclinical fasciolosis. Vet. Parasitol., 78: 129-136.
- Younas, M.; Fuquay, J.W.; Smith, A.E. and Moore, A.B. (1993):** Estrous and endocrine responses of lactating Holsteins to forced ventilation during summer. J. Dairy Sci., 76: 430-436.
- Zulu, V.C.; Nakao, T.; Yamada, K.; Moriyosh, M.; Nakada, K. and Sawamuka, Y. (2000):** Clinical response of inactive ovaries in dairy cattle after PRID treatment. J. Reprod. Develop., 46: 415-22.