List of Histograms

Histogram no.	Histogram Caption	Page no.
1	The 12 species which occur in five or more sites (numbers preceeding	7
1	species names are as in Table 1 above).	,
2	Families represented by two or more species and number of species in	Q
2	each family.	0
3	Number of species recorded for each geological age.	9

List of Text- Figures

Text- Figure no.	Text-Figure Caption	Page no.
1	Map showing sites of collection of fossil wood in Egypt	2
	Map showing location of the study area (two localities) from which the	
2	specimens were collected * Wadi Gabgaba & *-* Cairo-Bahariya Oasis	11
	Desert Road.	
3	Satellite image of Wadi Gabgaba, Egypt (Google Earth, 2009)	13
4	Geological Map showing fossil trees just north to latitude 22° 00` N near	1/1
4	Wadi Gabgaba, at Gebel Korosko and Aswan after Lyons (1894).	14
5	Lithostratigraphic section of Abu-Ballas Formation supplied by Prof. Dr.	17
3	Riffaat Osman, Geology Department, Faculty of Science, Benha University.	17
	Lithostratigraphic section of Lower Member of Sabaya Formation supplied	
6	by Prof. Dr. Riffaat Osman, Geology Department, Faculty of Science,	19
	Benha University.	
	Lithostratigraphic section of Upper Member of Sabaya Formation supplied	
7	by Prof. Dr. Riffaat Osman, Geology Department, Faculty of Science,	20
	Benha University.	
	Generalized lithostratigraphic section exposed at Cairo-Bahariya Oasis	
8	Desert Road supplied by Prof. Dr. Riffaat Osman, Geological Department,	22
	Faculty of Science, Benha University.	
9	Distribution of Agathoxylon in the world: Africa (1-Egypt) & Europe (2-	37
	Hungary).	
	Map showing distribution of gymnosperm fossil woods in Egypt (present work	
	included). 1- Wadi Araba, 2- Gebel Ahmer, 3- Gebel Mokattam, 4- Gebel Qatrani,	
	5- Bahariya Oasis, 6- Gebel Hufhuf, 7- Dakhala Oasis, 8- Kharga Oasis, 9-Esna,	
4.0	10-Road between Esna and Wadi Halfa, 11-Garf Hussein, 12-Sinn-El-Kaddab, 13-	•
10	Korosko, 14- Gilf Kebir, 15-Gebel Kamil, the star* indicates position of Wadi	39
	Gabgaba (the first locality of the present work) based on, Kräusel & Stromer,	14 17 19 20 22 37
	1924, Kräusel, 1939, Boureau, 1950, Dupéron-Laudouenei & Lejal-Nicol,	
	1981, Youssef et al., 2000, Youssef, 2002, Philippe et al., 2003 and Kedves et al., 2004a,b.	
	Distribution of <i>Bombacoxylon</i> in the world: Asia [1-Pakistan]; Africa [2-	
11	Kenya, 3-Somalia, 4-Ethiopia, 5-Egypt, 6-Libya, 7-Tunisia, 8-Algeria, 9-	57
11	Mali & 10-Rio de Oro]; Europe [11-France]; North America [12-USA].	31
	Wight & 10-Nio de Oloj, Europe [11-11dhee], North America [12-USA].	

12	Distribution of <i>Bombacoxylon owenii</i> in Egypt (this work included) [1-Suez, 2-Gebel Geneffa, 3-Gebel Shabraweet, 4-Turra, 5-Gebel Ahmer and Bir El-Fahme, 6-Mokattam, 7-Qattamiya, 8-Wadi Ankebieia, 9-Wadi Sannor, 10-Fayum, 11-Birket Qarun, 12-Gebel El-Khashab (Northern Petrified Forest & Southern Petrified Forest) 13-Gize pyramides & West of Giza Pyramides, 14-Wadi Faregh and Garet Aujan, 15-Der Baramus, 16-Wadi Natrun, 17-Bir Lebuk, 18-Moghara, 19-Cairo-Bahariya Oasis Desert	58
	Road (second locality of the present work), 20-Bahariya], based on Kräusel, 1939; Youssef, 1993; Kamal El-Din, 1996, 2002.	
13	Distribution of <i>Afzelioxylon</i> in Africa [1-Ethiopia, 2-Egypt, 3-Chad & 4-Algeria].	67
14	Distribution of <i>Cynometroxylon</i> in the world: Asia [1-Cambodia, 2-Thailand & 3-India] and Africa [4-Ethiopia, 5-Egypt, 6-Tunisia & 7-Algeria].	77
15	Distribution of <i>Tetrapleuroxylon</i> in the world: Africa [1-Somalia, 2-Sahara Sudanese, 3-Egypt, 4-Tunisia, 5-Algeria & 6-Mali], Europe [7-France], South America [8-Brazil, 9-Peru & 10-Colombia] and North America [11-USA].	88

List of Tables

Table	Table continu	Page
No.	Table caption	No.
1	Petrified wood flora of Egypt: recorded taxa, their ages and localities.	3
2	Size and remarks on specimens of Cairo-Baharyia Oasis Desert Road.	23
3	Names of taxa recorded in the present work, number of their specimens, locality	29
3	they come from and whether they are new or old records to Egypt.	29
4	Wood structure of Agathoxylons lifiyii (present work) compared with specimens	36
4	of A. lifiyii and A. sp. described earlier from Egypt.	30
5	Names of gymnosperm fossil woods reported from Egypt their ages and sites of	38
3	collection (this work included).	
6	Names of gymnosperm fossil wood families in Egypt. Numbers of genera and	40
	species are given for each family.	
7	Anatomical features of the seven species of <i>Bombacoxylon</i> .	53
8	Anatomical features of B. owenii from different ages and different countries of	54
0	the world and present work.	34
	Wood structure of Bombacoxylon owenii from Cairo-Bahariya Oasis Desert	
9	Road (present work) compared with specimens described earlier from other	54
	mainly Tertiary sites in Egypt.	
10	Comparison between B. langstoni from USA and from Cairo-Baharyia Oasis	55
10	Desert Road (present work).	33
11	The distribution of the seven known Bombacoxylon species in different counties	56
11	(present work included).	30
12	Anatomical features of Afzelioxylon species from mainly Tertiary sites from	65
12	Africa & from Cairo-Baharyia Oasis Desert Road (present work).	0.5
13	Difference between Afzelioxylon welkitii from Gebel Ruzza and from Cairo-	66
13	Baharyia Oasis Desert Road.	00
1.4	Anatomical features of eight Cynometroxylon species from mainly Tertiary sites	73
14	& from Cairo-Baharyia Oasis Desert Road (present work).	13
15	Comparison between Cynometroxylon tunesense from Tunisia, Egypt (Cairo	75
13	Petrified Forest) and (Cairo-Baharyia Oasis Desert Road) present work	13

16	The distribution of the eight known <i>Cynometroxylon</i> species in different countries (present work included).	76
17	Comparison between <i>Tetrapleuroxylon acaciae</i> , from Oligocene and Lower Miocene of Egypt and present work.	86
18	Comparison between <i>Tetrapleuroxylon ingaeforme</i> from Brazil (undetermined age) and present work .	86
19	The distribution of the nine known <i>Tetrapleuroxylon</i> species in different countries (present work included).	87
20	Names of Egyptian fossil wood genera and species assigned to the family Leguminosae (this work included).	89
21	Genera of fossil Leguminosae wood in Egypt and their distribution in Africa.	91
22	Summary of characteristics of the six studied dicot species from Cairo-Baharyia Oasis Desert Road.	93

List of Plates

Plate no.	Plate Caption	Page no.
1	Agathoxylon lifiyii Youssef, El-Saadawi, Kedves & Mostafa Fig. (1): A wood log (specimen no. 9) from Wadi Gabgaba. Annual rings are clear on transversely cut face. Fig. (2): Cross section showing 5 unequal growth rings (1-5), resin ducts and diffuse parenchyma with dark contents (arrows), x 35.	32
2	Agathoxylon lifiyii Fig. (3): Cross section magnified to show late wood (white arrow) and resin duct (black arrow), x 40. Fig. (4): Radial section magnified to show biseriate pits on the radial wall of tracheids (white arrow) and cross field (black arrow), x 284.	33
3	Agathoxylon lifiyii Fig. (5): Radial section showing uniseriate pits, x156. Fig. (6): Radial section magnified to show different shapes of biseriate pits on the radial wall of tracheids with round apertures, x 289. Fig. (7): Radial sections magnified to show cross field pits (arrow), x291. Fig. (8): Tangential section showing uniseriate ray (arrow), x 156. Fig. (9): Tangenital section magnified to show spiral wall thickening of the tracheids (arrow), x 292.	34
4	Bombacoxylon owenii (Carr.) Gottwald Fig. (10): A wood log (specimen no.112) from Cairo-Baharyia Oasis Desert Road. Fig. (11): Cross section showing semi-ring-porous wood, solitary and paired vessels and diffuse parenchyma, x 53. Fig. (12): Tangential section showing a simple perforation plate (arrow), x122.	47
5	 Bombacoxylon owenii Fig. (13): Tangential section showing alternate intervessel pitting (arrow), x 196. Fig. (14): Tangential section showing vessel element plugged with tyloses, x 125. Fig. (15): Tangential section showing 1-3 (mostly biseriate) rays, x130. Fig. (16): Radial section showing homocellular rays, composed of procumbent cells (arrow) x98. 	48
6	Bombacoxylon langstoni Wheeler & Lehman Fig. (17): A wood log (specimen no. 140) from Cairo-Baharyia Oasis Desert Road. Fig. (18): Cross section showing diffuse to semi-ring-porous wood, solitary and paired vessels and diffuse parenchyma, x 53. Fig. (19): Cross section showing tyloses and abundant diffuse and diffuse-in-aggregates axial parenchyma (arrows), x 53. Fig. (20): Tangential section showing simple perforation plate (arrow) x 156.	50

	Bombacoxylon langstoni Fig. (21): Tangential section showing crowded alternate intervessel pitting, x 180. Fig. (22): Tangential section showing vessel elements plugged with tyloses,	
7	x 164. Fig. (23): Tangential section showing multiseriate rays, 2-4 seriate, and parenchyma occur in strand, x117. Fig. (24): Radial section showing heterocellular rays, composed of procumbent cells (white arrow), and upright cells (black arrow), x156.	51
	Afzelioxylon welkitii (Lemoigne & Beauchamp) Lemoigne Fig. (25): A wood log (specimen no. 148) from Cairo-Baharyia Oasis Desert Road.	
8	Fig. (26): Cross section showing diffuse-porous, solitary and radial multiples 2-3 of vessels, lozenge-aliforme, confluent parenchyma and banded parenchyma (arrows), x 45. Fig. (27): Tangential section showing vessel elements with horizontal end walls (arrow), x 90. Fig. (28): Tangential section magnified to show alternate intervessel pitting,	62
	x 291.	
9	Afzelioxylon welkitii Fig. (29): Tangential section showing 1-4 (mostly triseriate rays), x20. Fig. (30): Tangential section magnified to show multiseriate ray (arrow), x 180.	63
	Fig. (31): Radial section showing homocellular rays, composed of procumbent cells (arrow), x 98.	
	Cynometroxylon tunesense Delteil-Desneux Fig. (22)	
10	Fig. (32): A wood log (specimen no. 108) from Cairo-Baharyia Oasis Desert Road. Fig. (33): Cross section showing diffuse-porous, solitary vessels and a forked tangential parenchyma band (arrow), x 50.	69
11	Cynometroxylon tunesense Fig. (34): Cross section magnified to show parenchyma bands alternating with fibre bands of nearly same thickness of cell wide, x 156. Fig. (35): Tangential section showing oblique (black arrow) and oblique to	70
	horizontal (white arrow) end walls, x 98. Fig. (36): Tangential section showing alternate intervessel pitting x 291.	
12	Cynometroxylon tunesense Fig. (37): Tangential section showing 1-3 seriate rays (mostly biseriate), x 176. Fig. (38): Radial section showing heterocellular rays composed of	71
	procumbent (white arrow) and upright cells (black arrow), x 156.	
	Tetrapleuroxylon acaciae (Kraüsel) Müller-Stoll & Mädel Fig. (39): A wood log (specimen no. 106) from Cairo-Baharyia Oasis	
13	Desert Road. Fig. (40): Cross section showing diffuse-porous, solitary and radial multiples of vessels, paratracheal lozenge-aliform, confluent and diffuse apotracheal parenchyma, x 63.	79

14	Tetrapleuroxylon acaciae Fig. (42): Tangential section showing rays, x117. Fig. (43): Radial section showing homocellular rays, composed of upright cells (arrow), x 153.	80
15	 Tetrapleuroxylon ingaeforme (Felix) Müller-Stoll & Mädel Fig. (44): A wood log (specimen no. 137) from Cairo-Baharyia Oasis Desert Road. Fig. (45): Cross section showing a growth ring marked by radially flattened late wood fibres (arrows), diffuse-porous, solitary and radial multiples of vessels, x 45. Fig. (46): Cross section magnified to show paratracheal vasicentric (black arrow), lozenged-aliform (red arrow) and diffuse apotrocheal parenchyma, x 96. Fig. (47): Tangential section showing a simple perforation plate (arrow), x 90. 	82
16	Tetrapleuroxylon ingaeforme Fig. (48): Tangential section showing 1-2 seriate rays, x 156. Fig. (49): Radial section showing homocellular rays, composed of procumbent cells (arrow), x 130.	83

Contents

Subject	Page No.
List of Histogram	I
List of Text-Figures	II
List of Tables	IV
List of Plates	VI
Introduction	1
Aim of the work	10
Localities & Materials	11
Methods	24
Collection	24
• Preparation of the samples	24
Description	26
Photography	26
Affinities and comparisons	27
Results	28
Descriptions, comparisons, affinities and discussion	30
I. Araucariaceae	30
1. Agathoxylon lifiyii (old record)	31
II. Bombacaceae	44
2. Bombacoxylon owenii (old record)	46
3. B. langstoni (new record)	49
III. Leguminosae	59
4. Afzelioxylon welkitii (old record)	61
5. Cynometroxylon tunesense (old record)	68
6. Tetrapleuroxylon acaciae (old record)	78
7. T. ingaeforme (new record)	81
Summary	98
References	101
Glossary	112
Arabic summary	1