
-REFERENCES

- Abd El- Rahman, A. M.M; El-Nobi, G. Ahmed and Moustafa, A. M. (2002).** Studies on *Pseudomonas* septicaemia among Tilapia fish in Abbassa fish farm . Zag. Vet. J. 30 (1), 25- 31.
- Abd El Wahab A. M., Hassouna M. M. E., Abd El-Maksoud A. M. S., Abd El Tawwab A. and Abu-Seef A. M. M.; (2007).** Cinnamon as a feed supplemented in Nile tilapia, (*Oreochromis Niloticus*), diets that reared in earthen ponds; Egyptian J. Nutrition and feeds, **10** (2), 881.
- Abd El-Azeem, F. (2006).** Effect of using fenugreek and fennel seeds as natural feed additives on performance of broiler chicks. Egyptian J. Nutrition and Feeds **9** (2): 277 - 297.
- Abd El-Aziz, E.S. (1988).** "Some studies on bacterial agents causing tail and fin rot among fresh water fishes in Egypte". M.Sc. Thesis (Infections diseases) faculty of Vet. Med. Cairo University .
- Abd El-Aziz, E.S. (1994).** "Immunological studies on Aeromonas and Pseudomonas bacterial Infectious in fresh water fishes" Ph.D. thesis (Infectious diseases) Faculty of Vet. Med. Cairo University.
- Abd El-hamid, A.M.; Salem M.F.I and Tolan, A.E. (2005).** Utilization of black seed meal (*Nigella sativa*) in nile tilapia (*Oreochromis niloticus*) Diets. J. Agric. Res. Tanta Univ., **31** (3).
- Abd El-Latif, S.A., A.T. El-Yamany and Eman A.F. Edaly. (2004).** Evaluation of using different levels and sources of medicinal

herbs in growing Japanese quail diets. Egyptian J. Nutrition and feeds 7(1):69-81.

Abd El-Maksoud, A.M.S.; Aboul-Fotoh, G.E.; Allam, S.M. and Abou zied, R.M. (2002). The response of Nile tilapia to animal protein free diets supplemented with some free amino acids and some medicinal plants. Proceedings of the 1th Conference on Aquaculture, organized by Egyptian Aquaculture Society and Suez Canal Univ. pp: 233-260

Abd El-Rahman, A.M.M. (1996). "Studies on bacterial diseases among cultured Tilapias ."M. Sc. Thesis (Fish diseases and Mangement) faculty of Vet. Med. Suez Canal University.

Abd El-Samee, Laila D. (2003). Effect of Dietary black cumin meal and garlic on the performance of muscovy ducklings. J. Agric. Sci. Mansoura Uni., 28(6):4387 – 4398.

Abdel.Zaher A. M. M., Ahmad M. H., Mousallamy A. and Asmaa Samir (2009). Effect of Using Dried Fenugreek Seeds as Natural Feed Additives on Growth Performance Feed Utilization Whole-body Composition and Entropathogenic *Aeromonas Hydrophila*-challenge of Monsex Nile Tilapia *O. Niloticus* (L) Fingerlings. Aus tralian Journal of Basic and Applied Sciences., 3 (2): 1234 – 1245.

Abd-El-Monem, A.I.; Shalaby, S.M.M. and El- Dakar, A.Y. (2002). Response of red tilapia to different level on some medicinal plant by-product: black seed and roquette seed meals. Proc. 1st Sci. Conf Aqua., El-Arish, Dec. 13-15, pp 247-260.

- Aboul-Fotouh, G.E.; Allam, S.M.; Shehata, E. and Abd El-Azeem, S.N. (2000).** Effect of some medicinal plants as feed additives on performance of milk production and composition of lacting buffaloes. *Egypt. J. Nutr. Feeds*, 3: 31-37.
- Aboul-Fotouh, G.E.; Allam, S.M; Shehat E. and Abdel-Azeem S.N. (1999).** Effect of some medicinal plants as feed additives on performance of growing sheep. *Egypt. J. Nutr. Feeds*, 2(2): 79-87.
- AbouZeid, R.M. (1998).** Evaluation of some medicinal plants as a feed additive in diets of Nile tilapia (*Oreochromis niloticus*). MSc Fac. of Agric, El-Fayoum, Cairo Univ.
- Ahen, W.; Popp, W. and Hoffman, R. (1982).** (hogens Ph.D. Thesis (microbiology) Fac. Of Vet. Med. 1982). "Pseudomonas fluorescens as pathogen for tench (*Tinca tinca*)". Bulletin of the European Association of fish pathogenis, 2:56-75.
- Ahmad H. M; Hany I. El-Marakby, Medhat E.A. Seden, Mohsen Abdel-Tawwab and Mohammad E. Abou El-Atta, (2006).** The use of organic selenium (Sel-Plex[®]) in practical diets for *Nile tilapia, Oreochromis niloticus* (L.): Effect on growth performance, feed utilization, whole - body composition and enteropathogenic *Aeromonas hydrophila* - Challenge."ISTA7" international symposium on tilapia Aquaculture. 6-8 September
- Ahmad, M.H.; El Mesallamy, A.M.; Samir, F. and Zahran, Z. (2009).** Effect of different levels of cinnamon (*Cinnamomum zeylanicum*, NEES) on growth performance, feed utilization, whole-body composition and enteropathogenic *Aeromonas hydrophila*-challenge

- of all male Nile tilapia *Oreochromis niloticus* L. Fingerlings.
Egypt. J. Appl. Sci., 24(5A): pp:1-18
- Ahmed, S.M. and Shoreit A.A.M. (2001).** "Bacterial haemorrhagic septicemia in *Oreochromis niloticus* at Aswan fish hatcheries"
Assiut Vet. Med. J. Vol. 45 No. 89, April 2001.
- Ahmed, S.M. and Shoreit A.A.M. (2001).** "Bacterial haemorrhagic septicemia in *Oreochromis niloticus* at Aswan fish hatcheries"
Assiut Vet. Med.
- Ali, M.F.; Saleh, M.S.; Eweedah, N.M. and Mahmoud S.A. (2005).**
Effect of using chamomile (*Mtricaria chamomilla*) flowers as feed additives on performance of growing lambs under desert farming systems. Egyptian. J. Nutrition and Feeds 8 (2): 127 - 137.
- Aligiannis, N., Kalpoutzakis, E., Mitaku, S., & Chinou, I. B. (2001).**
Composition and antimicrobial activity of the essential oils of two Origanum species. Journal of Agriculture Food Chemistry, 49, 4168 4170.
- Aligiannis, N., Kalpoutzakis, E., Mitaku, S., & Chinou, I. B. (2001).**
Composition and antimicrobial activity of the essential oils of two Origanum species. Journal of Agriculture Food Chemistry, 49, 4168 - 4170.
- Allen D.A; Austin, B. and Colwell, R.R. (1983).** Numerical taxonomy of bacterial isolates associated with fresh water fresh water fishery. J.of General Microbiol., 129: 2043 – 2062.

- Alzoreky, N.S. and Nakahara, K. (2003).** "Antibacterial activity of extracts from some edible plants commonly consumed in Asia" International Journal of Food Microbiology 80 (223 – 230).
- Amresh G., Paras Nath Singh and Chandana Venkateswara Rao. (2007).** Toxicological screening of traditional medicine laghupatha (*Cissampelos pareira*) in exeperimental animals. J. of Ethnopharmacology V. 116, Issue P. 454-460.
- Angka, S.L. and Lioc, K.G. (1982).** "Control of bacterial hemorrhagic septicemia in cultured fish "Bull. Penel. Inst. Bogor, 3 (3): 13-14.
- Anna, M.; Marina, K.; Mihkel, J. and Anne, O. (2004).** Identification and characterization of supercritical fluid extracts from herbs. C.R. chimie 7 (628-633).
- AOAC, (1990).** Official Methods of Analyses of the Association of Official Analytical Chemists International. 15th edition, Association of Official Analytical Chemists, Arlington, VA, USA.
- Arafa, R.F. (2000).** "Antibacterial activity of Egyptian limonium pruinsum extracts" 9th Sci cong. Fac Vet. Med. Assuit Univ. Egypt.
- Atwa, A.M. (1997).** Evaluation of the nutritive value of black seed meal (*Nigella sativa*) in diets of Nile tilapia (*Oreochromis niloticus*, Trewavas). MSc, Fac. of Agric, Alexandria Univ.
- Ayfer D. ATES,Ozlem Turgay ERDOGRUL. (2003).** Antimicrobial activities of various medicinal and commercial plant extracts. Turk J. Biol 157-162.
- Ayyad, K.M. (2003).** Studies on some dairy products. Ph.D. Abd El

- Wahab A. M., Hassouna M. M. E., Abd El-Maksoud A. M. S., Abd El Tawwab A. and Abu-Seef A. M. M.; Cinnamon as a feed supplemented in Nile tilapia, (*Oreochromis Niloticus*), diets that reared in earthen ponds; Egyptian J. Nutrition and feeds, **10** (2), 881, (2007).
- Badran, A.F. and Eissa, I.A.M. (1991).** "Studies on bacterial diseases among cultured Freshwater fish (*Oreochromis niloticus*) in relation to incidence of bacterial pathogens at Ismailia Governorate."J. Egypt . Vet .Med. Ass., 51 (4) , 837 – 847.
- Badran, A.F.H.M. (1993).** "Studies on pseudomonas septicemia outbreak among intensively cultured freshwater fishes with special reference to its control" Zag. Vet. J., 21 (4) : 737 – 746 .
- Baker, M.N. and Mostafa, M.A.A. (2006).**The importance of Betafin or chamomile flower as natural feed additives for Nile tilapia fry cultured under cold season conditions. Mansoura Univ. J. Agric. Sci., 31(7): 4145-4153.
- Bauer, J.D. (1982).** Colorimetric Determination of Serum albumin Clinical Laboratory Methods, 4th Ed., 495 - 496, 1121.
- Betts, G. D., Linton, P., & Betteridge, R. J. (1999).** Food spoilage yeasts: Effects of pH, NaCl and temperature on growth. Food Control, 10, 27–33.
- Boyd, C.E. (1990).** Water Quality in Ponds for Aquaculture. Birmingham Publishing Co., Birmingham, Alabama, USA.

- Cheesbrough, M. (1991).** Medical laboratory manual tropical countries. Volume 2: Microbiology. Printed in great Britian at the university press, Cambridge. 479pp.
- Christybapita, D.; Divyagnaneswari, M. and Dinakaran Michael R. (2007).** Oral administration of *Eclipta alba* leaf aqueous extract enhances the non-specific immune responses and disease resistance of *Oreochromis mossambicus*. Fish & Shellfish Immunology Volume 23, Issue 4, Pages 840 - 852.
- Collins, C.H. and Lyne, P.M. (1995).** "Microbiological Method. Weventh Edition. Butter Wroth-Heinemann Ltd.
- Csaba, G.Y.; Prigli, M. Bekesi, L.; Kovacs – Gayer, E.; Bajmocy, E. and Fazekas, B. (1981).** "Septicaemia in silver carp (*Hypophthalmichthys molitrix*, val.) and bighead (*Aristichthys nobilis* Rich.) caused by pseudomonas fluorescence" in : Olah, J.
- Davis, J. (1994).** Inactivation of antibiotics and the dissemination of resistance genes. Science, 264, 375 – 382.
- Deak, T., & Beuchat, L. R. (1996).** Handbook of food spoilage. New York, USA: CRC Press.
- Diab, A.S., M. E. Abo El-Atta and Doaa A. El-Araby. (2006 a).** Efficiency and characterization of an antibacterial substance from (*Origanum vulgare*) against aeromonas hydrophila in some freshwater fishes. Egypy. J. Agric. Res., 84(1B).
- Diab, A.S.; Abdel-Hadi, Y.M.; Ahmad, M.H.; Sakr, S.F. and Abo El-Atta M.E. (2006 b).** Outdoor study on the use of Echinacea (*Echinacea purpurea*), Marjoram (*Origanum Majoranan*) and

- yeast (*Saccharomyces cerevisiae*) as feed additives for *Oreochromis niloticus*. Egypt. J. Agric. Res., 84 (IB).
- Diab, A.S.; El-Nagar, G.O. and Abd-El-Hady, Y.M. (2002).** Evaluation of *Nigella sativa* L. (black seeds; baraka), *Allium sativum* (garlic) and Biogen as feed additives on growth performance and immunostimulants of *O. niloticus* fingerlings. Suez Canal Vet. Med. J.,(SCVMJ), Vol. V (2),pp: 745-753.
- Diab, A.S.; Sakr, S.F. Abd El-Hadi Y.M. and Ahmad, M.H. (2006 c).** Evaluation of time and dose related use of *Allium sativum* (Garlic) and *Nigella sativa* L.E. (Black seeds) mixture as feed additives for *Oreochromis niloticus*. Egypt J. Agric. Res.,84 (IB).
- Dias, M.C.; Spinardi-Barbisan, A.L.T. and Rodrigues, M.A.M. (2006).** Lack of chemopreventive effects of ginger on colon carcinogenesis induced by 1,2 – dimethylhydrazine in rats "Food and Chemical Toxicology".
- Divyagnaneswari, M.; Christyapita, D. and Dinakaran Michael R. (2006).** Enhancement of nonspecific immunity and disease resistance in *Oreochromis mossambicus* by *Solanum trilobatum* leaf fractions. Fish & Shellfish Immunology Vol. 23, Issue 2, P. 249 - 259.
- Doaa, A.El-Araby, (2006):** Antimicrobial Activities of Various Medicinal and Commerical Plant Extracts on Pathogenic Bacteria Isolated from Fish. Library of Faculty of Scinece, Banha University and Library of (CLAR) Central laboratory for Aquaculture Research, Abbassa, Abo Hammad, Sharkia, Egypt.

- Doumas, B.T. (1975):** Colorimetric determination of total proteins. Clin. Chem. 21: 1159 - 1166.
- Dugenci, S.K.; Arda, N. and Candan, A. (2003).** Some medicinal plants as immunostimulant for fish. Journal-of-Ethnopharmacology, 88: 1, 99-106. Journal – of - Herbs, Spices – and - Medicinal - Plants, 6: 4, 29-38.
- Dytham, C. (1999):** Choosing and Using Statistics: A Biologist's Guide. Blackwell Science Ltd., London, UK. p 147.
- Ekanem, A.P.; Wang, M.; Simon, J.E.; Obiekezie, A.I. and Morah, F. (2004).** In vivo and in vitro activities of the seed extract of Piper guineense Schum. and Thonn. against skin and gill monogenean parasites of goldfish (*Carassius auratus auratus*). Phytotherapy-Research; 18 (10): 793-797.
- Ekanem, S.B.; and Okoronkwo T.W. (2003).** Pawpaw seed as fertility control agent on male *Nile tilapia*. NAGA, World fish center Quarterly Vol. 26 No. 2.
- El- Bouhy, Z.M. (1986).** "Studies on diseases affecting the skin in some Nile fishes" Ph.D. Thesis Zag. Universtiy .
- El Dakar, A.Y.;Gaber, D.I.; Hassanien, G.D.I.; Seham, S., Gad and Salah E.Sakr (2003).**Use of Medical and Aromatic Plants in Fish Diets: I. Effect of dried Marjoram leaves on performance of hybrid tilapia *Oreochromis niloticus* x *Oreochromis aureus*, fingerlings. Egypt. J. Aquat. Biol.& Fish., Vol. 7, No 4: 66 – 84.
- El-Ashry, M.A.A.; El-Bordeny N.E.; Khattab H.M. and El-Sayed H.M. (2006).** Effect of diets supplemented with medicinal herbs

- on nutrient digestibility and some blood metabolites of buffalo calves. Egyptian J. Nutrition and Feed 9 (2): 179 - 191.
- El-Attar, A.A. and Moustafa, M. (1996).** "Some studies on tail and fin rot disease among cultured Tilapia Fishes". M.Sc. Thesis (Infections diseases) faculty of Vet. Me. Cairo University .
- El-Bouhy, Z. M. (1995).** "Studies on the fins and rot disease in some commercial culture fish in sharkia province." J. Egypt. Vet. Med. Ass. No. 1, 2:123-140.
- El-Bouhy, Z. M. (1995).** "Studies on the fins and rot disease in some commercial culture fish in sharkia province." J. Egypt. Vet. Med. Ass. No. 1, 2:123-140.
- El-Boushy, M.E. and El-Ashram, A.M.M. (2008).** Effect of dietary bovine lactoferrin on non-specific immune response of Nile tilapia (*Oreochromis niloticus*). Abbassa Int. Aqua. (IB).
- El-Dakar, A.Y. (2004).** Growth response of hybrid tilapia, *Oreochromis niloticus* x *Oreochromis aeneus*, fingerlings to diets supplemented with different levels of caraway seeds. J. Agric. Sci. Mansoura Univ., 29 (11): 6083 - 6094.
- El-Dakar, A.Y.; Hassanien G.D.I.; Gad S.S. and Sakr S.E. (2004).** Use of medical and aromatic plants in fish diets: 2. Effect of dried basil leaves on performance of hybrid tilapia (*Oreochromis niloticus* x *Oreochromis aeneus*) fingerlings. 3rd Intern. Conf. on Anim. Production and Health in Semi-Arid Areas, Suez Canal Univ. J.Egypt.Acad.Soc.Environ.Develop.; (B-Aquaculture) 5(2):43- 67

- El-Dakar, A.Y.; Hassanien, G.D.I.; Gad, S.S. and Sakr, S.E. (2008).**
Use of dried basil leaves as a feeding attractant for hybrid tilapia, *Oreochromis niloticus* x *Oreochromis aureus*, fingerlings. Med. Aquacult. J., 1 (1): 35-44.
- El-Dakar, A.Y.; Shymaa, M.M., Shalaby; S.M., Abead and M.M., Khalafala (2005).** Use of spices as feeding stimulants in diets of marine shrimp, *Penaeus japonicus*. Egypt. J. Nutr Feeds, 8 (1): 1065-1076.
- El-Daly, A.A. and Tolba, M. (2009).** Internal structure of silver crystallites : going beyond nanostructure using novel processing technique. Cryst. Res. Technol. 44, No. 2, 197-202.
- El-Kaiaty, A.M.; Fatma R. Mohamed; Eman M. Abou-Eita and Abeer A.M. Eshra (2005).** Beneficial impact of organic chromium supplementation of broiler diets on growth performance, some blood constituents and immunological status. Egyptian J. Nutrition and Feed, 8 (1) Special Issue: 737 - 759.
- El-Kholy, Kh.F.; El-Azab, A.A.; ABdel Wareth A.A. and Hanan A. Abo State, (2005).** Effect of partial replacement of fish meal by full-fat soybean meal or cottonseed meal supplemented with iron on growth performance and body composition of juvenile nil tilapia (*Oreochromis niloticus* L.). Egyptian. J. Nutrition and Feed, 8 (1) Special Issue: 1157 - 1170.
- El-Mallah, G.M. (2003).** Response of Growing turkey to some medicinal plants as natural growth promoters. J. Agric. Sci. Mansoura Univ., 28 (11): 6621 - 6631.

- El-Saidy, D.M.S.D. (1999).** Effect of different level of dried onion meal supplemented to the diets on growth, feed utilization and body composition of Nile tilapia, *Oreochromis niloticus* fingerlings. Monofiya J. Agri. Res., 24(2): 481-492.
- El-Saidy, D.M.S.D. and Gaber, M.M.A. (1997).** Effect of different levels of dried garlic meal supplemented to the diets on growth performances, survival and body composition of Nile tilapia, *Oreochromis niloticus* fingerlings. Ann. Agri. Sci. Moshtohor, 35(3): 1197-1209
- Enany, M.E., El-sayed, M.E.; Diab, A.S. ;Hassan, S.M and El-Gamal, R.M.(1995).** "Bacterial causes of fin rot in some fresh water fishes."Alex. J. Vet. Sci. ,Vol. 11, No.4, 535-547.
- Faisal, M. and Easa, M. E. (1987).** "Acute septicemia in silver carp (*Hypophthalmichthys molitrix* val.) caused by a capsulated pseudomonas following transport" J. Egypt . Vet . Ass.47 (1/2): (25 – 36) .
- Gaber, A.A.; El-Ayouty, S.A.; Zaki, A.A.; Abou Ammo, F.E. and El-Gohary E.S.I., (1998).** Productive performance of lambs fed diets containing *Nigella sativa* meal. Egypt. J. Nutri. Feeds, 1(2): 97.
- Gatti, R. and Nigrelli, D.A. (1984).** "Haemorrhagic septicaemia in catfish. Pathogenicity of the strain strains isolated and reproducibility of the disease." Obietivie Documenti Veterinari , 5 (6), 61 -63.

- Ghittino, P. (1976).** "International aspects of disease control in aquaculture". Fao Technical conference on aquaculture. Kyoto, Japan. 26 May- 2 june.
- Gittino, P. (1976).** "International aspects of disease control in aquaculture". Fao Technical conference on aquaculture. Kyoto, Japan. 26 May- 2 june.
- goats diets and their effects on milk yield and milk composition. Proc.3th All Africa Conf. Anim. Agric. 11th Conf. Egyptian Soc. Anim. Prod., Alexandria, Egypt, 6-9 November, p 197.
- Goering, H.K. and P.G. Van Soest (1970).** Forage fiber analysis (apparatus, reagent, procedures, and some applications). US Dept. Agric. Handbook, Washington D.C., USA, p. 379.
- Guner, A., Ozhatay, N., Ekim, T., & Baser, K. H. C. (2000).** Flora of Turkey and the East Aegean Islands (Vol. 11 (supplement-II)). Edinburgh: Edinburgh University Press.
- Haenen, O.L. and Davidse . A (2001).** First isolation and pathogen city studies with pseudomonas anguilliseptica from diseased European eel *Anguilla* (L.) in the Netherlands . Aquaculture. 169: 27 – 36 .
- Hanafy M. M.; (1995).** The use of anise, fennel and ginger as a feed additive in the nutrition of broilers, M. Sc. Thesis, Faculty of Agriculture, Alexandria University,
- Hassan G.D.I.; El-Dakar A.Y.; Ghonim S.I. and El-Aiat A.A.O. (2005).** Utilization of some wild plants in north Sinai as energy sources in diets of hybrid tilapia, *Oreochromis nitoticus* x

Oreochromis aureus. Egyptian J. Nutrition and Feed, 8 (1) Special Issue: 975 - 985.

Hassanen, G.D.I.; El-Dakar A.Y.; Ghonim S.L. and El-Aiat A.A.O. (2005). Nutritional evaluation of some field wastes in north sinai as unconventional energy sources in hybrid tilapia, *Oreochromis niloticus* x *Oreochromis aureus*, Dites. Egyptian J. Nutrition and Feed, 8 (1) Special Issue: 963 - 974.

Hayam D. Tonsy; Gomaah, S.A. Agouz H.M. and Abou Zead M.Y. (2005). Effect of partial replacement of yellow corn by olive cake (A non - conventional feed) as dietary energy source on growth performance, liver and kidney enzymes and some blood parameters of Nile tilapia (*Oreochromis niloticus*). Egyptian. J. Nutrition and Feed , 8 (1) Special Issue: 1171 - 1183.

Hsieh, P. C., Mau, J. L. and Huang, S. H. (2001). Antimicrobial effect of various combinations of plant extracts. Food Microbiology, 18, 35– 43.

Ian Bricknell and Roy A. Dalmo, (2005). The use of immunostimulants in fish larval aquaculture. Fish & Shellfish Immunology 19, 457 - 472.

Ibrahim M. R., Abd El-Latif M. S. and El-Yamany A. T.; (1998). Effect of adding some natural growth promoters to broiler chick's diets on growth performance, digestibility and some metabolic functions. J. Agric. Sci., Mansoura Univ, 32 (3), 1029,

- Ietswaart, J. H. (1980).** A taxonomic revision of the genus *Origanum* (Labiatae). Ph.D Thesis. Leiden Botanical Series 4. Leiden University Press, The Hague.
- Iftikhar - Ahmad; Mushtaq - Ahmad; Ashfaq - Ahmad (1994).** Antimicrobial activity of *Dodonaea viscosa* oil. *Fitoterapia*, 65: 2, 167-168 .
- Jens K.S.; Mller, Rodrigo R.; Catharion and Marcos N. Eberlin (2005).** "Electrospray ionization mass spectrometry fingerprinting of essential oils: Spices from the Labiatatae family" *Food Chemistry*.
- Jirawan Oonmetta-areea, Tomoko Suzukib, Piyawan Gasalucka and Griangsak Eumkeb (2005).** "Antimicrobial properties and action of galangal (*Alpinia galangal* Linn.) On *Staphylococcus aureus*" Article in press.
- Josè M. Prieto; Patrizial Lacopini; Pierluigi Cioni and Silvio Chericoni, (2007).** In vitro activity of the essential oils of *Origanum vulgare*, *Satureja montana* and their main constituents in peroxynitrite - induced oxidative processes. *Food Chemistry* Volume 104, Issue 3, Pages 889 - 895.
- Jyrki Nikki, Juhani Pirhonen , Malcolm Joblin and Juha karjalainen (2004).** Compensatory growth in juvenile rainbow trout, *Oncorhynchus mykiss* (Walbaum), hald individually. *Aquaculture* 235 pp 285 - 296.

- Katiyar, S.K.; Agarwal, R. and Mukhtar, H., (1996).** Inhibition of tumor promotion in sencar skin by ethanol extract of *Zingiber officinale* rhizome. *Cancer Research* 56, 1023–1030.
- Khan R.A.; Compbell, J. and Lear, (1981).** "Mortality in captive Atlantic cod, *Gauss morhua* associated with fin rot disease". *J.of wild life diseases*, 17 (4) 521-527.
- Kholif, A.M. (2000).** Medicinal plants seeds supplementation to lactating.
- Konning G.H.; Agyare , C. and Ennison, B. (2004).** "Antimicrobial activity of some medicinal plants from Ghana". *Fitoterapia* 75 (65 – 67).
- Kosar, M.; Dorman, H.J.D. and Hiltunen, R. (2005).** Effect of an acid treatment on the phytochemical and antioxidant characteristics of extacts from selected *Lamiaceae* species. *Food Chemistry* 91, (525-533).
- Krieg, N.R. and Holt, J.G. (1984).** "Bergey's Manual of systemic Bacteriology – Williams and Wilkins, Baltimore, London, Vol. 1."
- Kulkarni, M.G. Street, R.A. and Van Staden, J. (2007).** Germination and seedling growth requirements for propagation of *Dioscorea dregeana* (Kunth) Dur. And Schinz - A tuberous medicinal plant. *South African Journal of Botany* Volume 73, Issue 1, Pages 131 - 137.
- Langner, E., Greifenberg, S. and Gruenwald, J., (1998).** Ginger: history and use. *Advances in Therapy* 15, 25–44.

- Lennette, E. H.; Balows, A.; Hausler, W.J. and Traunt, J. P.(1980).** "Manual of clininca Microbiology". Amer. Soc. Micro., Washington , D.C.
- Lio-Po, G. and Sanvictores, E.G. (1987).** "Studies on the causative organism of niloticus (Linnacus) fry mortalities, 1. primary isolation and pathogenicity experiments". J. Aquacult. Trop. 2: 25-30.
- Loper, J. E.; Henkels, M. D.; Roberts, R. G.; Grove, G. G.; Willett, M. J., and Smith, T. J. (1991).** Evaluation of streptomycin, oxytetracycline, and copper resistance of *Erwinia amylovora* isolated from pear orchards in Washington State. Plant Disease, 75, 287–290.
- Maass N., Bauer J., Paulicks B. R., BÖhmer B. M. and Roth-Maier D. A.(2005).** Efficiency of *Echinacea purpurea* on growth performance and immune status in pig, Journal of animal physiology and nutrition, **89**, 244, (2005).
- Mahady G.B. (2005):** Medicinal plants for the prevention of bacterial : plants as sou proteins .vol.11,n°19,pp. 2405-2427[23 page(s) (article)] (122ref.) ISSN 1381-6128.
- Mariah S.; Alina, A.R.; Chia, K.L.; Babji, A.S.; Nulkirah, M.; Chang-YuShyun (ed.); Mohtar, M, (ed.); Subramaniam, V. (ed.) and Yunos, N.M. (2002).** Antioxidative effects of ginger (*Zingiber officinale*) extract on surimi during frozen storage. Towards-modernisation-of-research-and-technology-in-herbal-industries.-Proceedings-of-the-Seminar-on-Medicinal-and-Aromatic-Plants, 211-215 .

- Marzouk, M.S.; Eissa, I.A.M.; Moustafa, M. and Mervat, S. H. (1989).** Contribution to tail and fin rot disease in catfish (*Clarias lazara*) . Zag . Vet. J. Vol. 17 No. 3P. 244- 257 .
- Megahed, A.A. (2000).** Studies on some Gram negative bacteria of fish M.Sc. Thesis (microbiology) Fac. Of Vet. Med. , Suez Canal University .
- Milos, M., Mastelic, J. and Jerkovic, I. (2000).** Chemical composition and antioxidant effect of glycosidically bound volatile compounds from oregano (*Origanum vulgare* L. ssp. hirtum. Food Chemistry, 71, 79–83.
- Mohamed , A.H.; El-Said, B.E. and El-Seidi, I.A. (2003).** Influence of some medicinal plants supplemental on digestibility, nutritive value, rumen fermentation and some blood biochemical parameters in sheep. Egypt. J. Nutr. Feeds, 6: 139.
- Montasir A.H. and Hassib, M. (1956):** Manual flora of Egypt, Impreimerie Masr. S.A.E.
- Moustafa, B.E.N. (1986).** "Epidemiological on fin rot disease in some Nile fish from Upper Egypt."M.V.Sc., Thesis (fish disease) Assuit University
- Muangsan N. and Senamontee, V. (2009).** Antimicrobial effects of some medicinal plant extracts against bacteria associated with lack disease. ISHS Acta Horticulturae 786: International Workshop on Medicinal and Aromatic Plants.
- Muhl, A. and Liebert, F. (2007).** Growth and parameters of microflora in intestinal and faecal samples of piglets due to application of a

phytogenic feed additive. Journal of animal physiology and animal nutrition Volume 91 Issue 9 - 10 Page 411 - 418.

Neill, C.R.; Nelssen, J.L.; Tokach, M.D. Goodband, R.D.; DeRouchey, J.M.; Groesbeck, S.C.N.; Lawrence, K.R.; Hastad, C.W.; Gottlob, R.O. and Hil, B.M. (2004). Evaluating oregano oil as a growth enhancef in nursery pig diets.

NRC; National Research Council (1994). Nutrient requirements of fish. Committee on Animal Nutrition. Board on Agriculture. National Research Council. National Academy Press. Washington DC, USA.

Okaeme, A.N. (1989). "Bacteria associated with mortality in tilapias (*Heterobronchus bidorsalis*) and *Clarias Lazera* in indoor hatcheries and outdoor ponds". J. Aquacult . Trop. 4 (2): 143 – 146.

Okpokwasili, G.C. and Obaho, (1991). "Relationship between water quality and bacteria associated with the brown patch disease of *Tilapia* fingerlings reades in tropical fresh water, culture ponds ". J.Aquacult . In the tropics 6: 2, 157 – 172.

Osman, A.; Atallah; Mohey, E.I.S. ; Easa Abd-Elmonem A. Aly; Ahmed, S. Diab and Saleh, F. Sakr (1997). Pathological changes associated with the fin –Rot in during Bacterial diseases in fresh – water fish . Alex. J. Vet. Sc. Vol. 13 No. 6:629 – 644 .

Osman, M.F.; Khattab H.M.; Fatama A. Hafez and Mounes H.A.M. (2005). Effect of different nutritional treatments on the productive

performance of *Oreochromis niloticus*. Egyptian. J. Nutrition and Feed, 8 (1) Special Issue: 1033 - 1047.

Oya Keles; Tulay Bakirel; Seyyal A.K. and Abdulkarim Alpmar (2000). "Antibacterial activity of some plants used for medicinal purpose against pathogens of veterinary importance" 9th Sci. Cong. Fac. Vet. Med. Assiut Univ. Egypt

Petrinee, Z.; Naglic, T. Matasin, Z. and Fijan, N. (1985). "Pseudomonas fluorescens septicemia in big head (*Aristichthy nobilis rich*), following handling". Veterinarshi, Arhiv, 55 (6), 277- 284.

Pier paolo Gatta; Kim D. Thompson; Richard Smullen; Andrea Piva; Silvia Testi and Alexandra Adams (2001). Dietary organic chromium supplementation and its effect on the immune response of rainbow trout (*Oncorhynchus mykiss*). Fish & Shellfish Immunology 11, 371 - 382.

Pitt, J.I., & Hocking, A. D. (1997). Fungi and food spoilage (2nd ed.). London, UK: Blackie Academic and Professional, 596 pp.

Post, G. (1987). "Text book of fish health ." T.F.H. publications, Inc., Ltd, 2 nd . Ed. , 182 – 185 .

Post, G. (1983). "Text Book of fish health" T.F.H. publication, Inc., Lid, Wes. Sylrania, New Jersy, USA.

Reitman, S. and Frankel, S. (1957). A colorimetric method for determination of serum glutamic oxaloacetic and glutamic pyruvic transaminases. J. Clin. Path. 28 - 56.

- Rina Chakrabarti and Rao. Y. Vasudeva (2005).** *Achyranthes aspera* stimulates the immunity and enhances the antigen clearance in *Catla catla*. Department of Zoology, University of Delhi, Delhi - 110 007, India.
- Rizk, K; M.H.; N.F. and Easa, M. (1984).** "Progect complete survey on diseases of cultured fish Egyptian ponds. "A progressive report to Scientific Research Academy.
- Sahin, M.; Guluçe, D.; Daferera, A.; Sokmen, M.; Sokmen, M.; Polissiou, G. & Agar and Ozer, H. (2004).** Biological activities of the essential oils and methanol extract of *Origanum vulgare* ssp. *Vulgare* in the Eastern Anatolia region of Turkey . Food Control 15, (549-557).
- Sahu, S. ; Das B.K.; Mishra B.K.; Pradhan J. and Sarangi N. (2007).** Effect of *allium sativum* on the immunity and survival of *Labeo rohita* infected with *Aeromonas hydrophila*. Journal of applied ichthyology volume 23 Issue 1 page 80 - 86.
- Sakr, S. E. (2003).** Studies on the feeding attractants for fish. M.Sc Thesis. Fac. Environ. Agric. Sci., El-Arish, Suez Canal Univ., Egypt. 83:669-675.
- Salama, S.S and Abd El- H . (1999):** Studies on the bacterial diseases of *Oreochromis niloticus* reared in concrete pond. m.V.SC. Thesis (Fish Diseases and Mangagement) Fac. Vet. Med. Suez Canal Univ.
- Schäperclaus, W.; Kulow, H. and Schreckenback, K. (1992).** Fish Disease Vol.I A.A. Balkema / Rotterde .

- Seden, M. E. A.; Fayza E.Abbass and M.H.Ahmed (2009).** Effect of origanum Vulgare as a feed additive on growth performance, feed utilization and whole body composition of Nile Tilapia.(*Oreochromis niloticus*) fingerlings challenged with pathogenic *Aeromonas hydrophila* .J. Agric .Sci. Mansoura University ., 34(3) :1683- 1695.
- Service, R. F. (1995).** Antibiotics that resistance. Science, 270, 724–727
- Shalaby, S.M.M. (2004):** Response of Nile tilapia, *Oreochromis niloticus* fingerlings to diets supplemented with different levels of fenugreek seeds (Hulba). J. Agric. Mansoura Univ., 29: 2231-2242
- Shalaby, S.M.M.; Abd Elmonem, A.L. and El-Dakar, A.Y. (2003):** Enhancement of growth performance, feed and nutrient utilization, of Nile tilapia, *Oreochromis niloticus*, using of licorice roots (Erksous) as a feed attractive. J. Egypt. Acad. Soc. Environ. Develop. (B – Aquaculture), 4 (2): 119 – 142.
- Shariff, M. and Law, A.T. (1980).** An incidence of fish mortality in bakok River . In M.N.H. Hasan , A.L. Mohamed , M.M. Clyde and Tan S.C. eds. (P.153 – 162) . International Symp . On Topley, M. and Wilson, L. (1983): Principles of bacteriology, Virology and immunity Vol. 117th Ed. , Edward Arnold , Great Britain , London .
- Shivanand, D.; Jolad, R.; Clark Lantz; Guan Jie Chen; Robert B.; Bates, Barbara N. and Timmermann (2005).** "Commercially processed dry ginger (*Zingiber officinale*) :Composition and

- effects on LPS- stimulated PGE production" *Phytochemistry* 66 (1614 – 1635).
- Snogerup, S. (1971).** Evolutionary and plant geographical aspects of Chasmophytic communities. In P. H. Davis, P. C. Harper, & I. C. Hedge (Eds.), *Plant life of South–West Asia* (pp. 157–170).
- Soltan, M.A. (2005).** Potential of using raw and processed canola seed meal as an alterantive fish meal protein source in diets for nile tilapia (*Oreochromis niloticus*). *Egyptian. J. Nutrition and Feed.*, 8 (1) Special Issue: 1111 - 1128.
- Suheyla Karatas Dugenci; Nazli Arda and Akin Candan, (2003).** Some medicinal plants as immunostimulant for fish. *Journal of Ethnopharmacology* 88, 99 - 106.
- Sung – Sook Chun; Dhiraj, A.; Vатtem, Yuan; Tong Lin and Kalidas Shetty (2005).** "Phenolic antioxidants from clonal oregano (*Origanum vulgare*) with antimicrobial activity against *Helicobacter pylori* " *Process Biochemistry* 40 (809 – 816).
- Surh, Y. J., (2002).** Anti-tumor promoting potential of selected spice ingredients with antioxidative and anti-inflammatory activities: a short review. *Food and Chemical Toxicology* 40, 1091–1097.
- Surh, Y.J., Lee, E., Lee, J.M., (1998).** Chemopreventive properties of some pungent ingredients present in red pepper and ginger. *Mutation Research* 402, 259–267.
- Tohamy E.Y.; H.M. Atta; M.E. Abou El Atta and D.A. El Arabe. (2006).** Purification and characterization of an antibacterial agent

from (*Origanum vulgare*) against of some pathogenic bacteria isolated from fishes. Egypt. J. Biotechnol. Vol. 22.

Tollba, A.A.H. (2003). Using some Natural additives to improve physiological and productive performance of broiler chicks under high temperature conditions. 1- Thyme (*Thymus vulgaris* L.) or Fennel (*Foeniculum vulgare* Mill). Egypt. Poult. Sci., 23(2): 313-326.

Topley M. and Wilson, L. (1983). Principles of bacteriology. Virology and immunity Vol.11-7th Ed., Edward Arnold. Great Britain, London.

Toranzo, A. E. ; Novoa, B . and Romalde , J. L. (1993). Numez , S.; Devesa, S.; Marino, E.; Silva, R. ; Martinez, E. ; Figueras, A. and Barja, J.L.

Turnbull, J.F. (1993). Bacterial gill diseases and rot . Bacterial diseases of fish , PP. 40 – 57 . Oxford , UK ; Blackwell Sci . publications.

Walker, S.J. (1988). Major spoilage micro-organisms in milk and dairy products. Journal of the Society of Dairy Technology, 41, 91– 92.