

References

Aeschbach A and Mekhail NA: Common nerve blocks in chronic pain management. *Anes Clin North Am.* 2000 Jun; 18(2):429-59.

Ahmed HE, Craig WF, White PF, et al: Percutaneous electrical nerve stimulation (PENS): a complementary therapy for the management of pain secondary to bony metastasis. *Clin J Pain.* Dec 1998;14(4):320-3.

American Cancer Society: Cancer facts & figures. Atlanta. 1995.

American Nurses Association (ANA): Code of Ethics for Nurses with Interpretive Statements. Washington DC: author 2001.

American Pain Society: Principles of analgesic use in the treatment of acute pain and cancer pain, (4 ed.). 1999, Glenview, IL: American Pain Society.

American Pain Society: Guideline for the management of cancer pain in adult and children. 2005.

American Pain Society: Principles of Analgesic Use in the Treatment of Acute and Cancer Pain, 5th ed. Glenview, IL: author. 2005.

Anderson R, Saiers JH, Abram S, et al: Accuracy in equianalgesic dosing: Conversion dilemmas. *J Pain Symptom Manage* 2001; 21: 397-406.

Aram L, Krane EJ, Kozloski LJ, et al: Tunneled epidural catheters for prolonged analgesia in pediatric patients. *Anesth Analg* 2001; 92:1432-1438.

Arbaiza D and Vidal O: Tramadol in the treatment of neuropathic cancer pain: a double-blind, placebo-controlled study. *Clin Drug Investig*, 2007; 27 (1): 75-83.

Aydin G, Tomruk S, Keles I, et al: Transcutaneous electrical nerve stimulation versus baclofen in spasticity: clinical and electrophysiologic comparison. *Am J Phys Med Rehabil.* Aug 2005;84(8):584-92.

Backonja M and Glanzman RL: Gabapentin dosing for neuropathic pain: evidence from randomized, placebo-controlled clinical trials. *Clin Ther* 2003; 25:81-104.

- Backonja MM:** Use of anticonvulsants for treatment of neuropathic pain. *Neurology* 2002; 59(suppl 2):S14-S17.
- Banerjee T. Letter:** Transcutaneous nerve stimulation for pain after spinal injury. *N Engl J Med.* Oct 10 1974;291(15):796.
- Basbaum AI, Fields HL:** Endogenous pain control mechanisms: review and hypothesis. *Ann Neurol.* Nov 1978;4(5):451-62.
- Becker C, Heidersdorf S, Drewlo S, et al.** Efficacy of epidural perineural injections with autologous conditioned serum for lumbar radicular compression: an investigator-initiated, prospective, double-blind, reference-controlled study. *Spine.* Aug 1 2007;32(17):1803-8.
- Behbehani MM:** Sensory receptors and somatosensory system. Chapter 7.in: *Essentials of physiology.* 2nd ed., Eds: Sperlakis N and Banks Ro., Little, Brown and Company, Boston, London, Toronto, New York, PP81-97.
- Benedetti F, Amanzio M, Casadio C, et al:** Control of postoperative pain by transcutaneous electrical nerve stimulation after thoracic operations. *Ann Thorac Surg.* Mar 1997;63(3):773-6.
- Bennett M.** The LANSS Pain Scale: the Leeds assessment of neuropathic symptoms and signs. *Pain.* 2001;92:147-157.
- Berman BM, Swyers JP:** Applying alternative medical approaches to managing chronic pain syndromes: If not now, when?. *APS Bulletin.* 1998;8:4-6.
- Bertalanffy A, Kober A, Bertalanffy P, et al.** Transcutaneous electrical nerve stimulation reduces acute low back pain during emergency transport. *Acad Emerg Med.* Jul 2005;12(7):607-11.
- Bloodworth D, Callvillo O, Smith K, et al.** Chronic pain syndromes: evaluation and treatment. In: Braddom R, ed. *Physical Medicine and Rehabilitation.* 2nd ed. Mosby; 2000:913-33.
- Bogduk N, Aprill C, Derby R.** Epidural steroid injections. In: White AH, Schofferman JA, eds. *Spine Care Operative Treatment.* vol 1. St Louis, Mo: Mosby; 1995:122-32.
- Bonadio WA and Wagner V:** Efficacy of tetracaine-adrenaline-cocaine topical anesthetic without tetracaine for facial laceration repair in children. *Pediatrics* 1990; 86:856-857.

- Bonica JJ and Loeser JD:** History of pain concepts and therapies. In: Loeser JD, Butler SH, Chapman CR, Turk DC, eds. *Bonica's Management of Pain*. 3rd ed. Philadelphia: Lippincott, Williams and Wilkins, 2001:3–16.
- Borjesson M, Eriksson P, Dellborg M, et al.** Transcutaneous electrical nerve stimulation in unstable angina pectoris. *Coron Artery Dis*. Aug-Sep 1997;8(8-9):543-50.
- Bostrom B, Sandh M, Lundberg D and Fridlund B:** Cancer-related pain in palliative care: patients' perceptions of pain management. *Journal of Advanced Nursing* 2004; 45(4):410-9,
- Botwin KP, Gruber RD, Bouchlas CG, et al.** Complications of fluoroscopically guided caudal epidural injections. *Am J Phys Med Rehabil*. Jun 2001;80(6):416-24.
- Breitbart W:** Psychotropic adjuvant analgesics for pain in cancer and AIDS. *Psychooncology*, 1998; 7 (4): 333-45, Jul-Aug.
- Bruera E and Kim HN:** Cancer pain. *JAMA* 2003; 290(18):2476-2479.
- Bowman SJ, Wedderburn L, Whaley A, et al.** Outcome assessment after epidural corticosteroid injection for low back pain and sciatica. *Spine*. Aug 1993;18(10):1345-50.
- Brown FW.** Management of diskogenic pain using epidural and intrathecal steroids. *Clin Orthop*. Nov-Dec 1977;(129):72-8.
- Buenaventura RM, Datta S, Abdi S, et al.** Systematic review of therapeutic lumbar transforaminal epidural steroid injections. *Pain Physician*. Jan-Feb 2009;12(1):233-51.
- Bush K, Hillier S.** A controlled study of caudal epidural injections of triamcinolone plus procaine for the management of intractable sciatica. *Spine*. May 1991;16(5):572-5.
- Byrod G, Rydevik B, Nordborg C, et al.** Early effects of nucleus pulposus application on spinal nerve root morphology and function. *Eur Spine J*. 1998;7(6):445-9.
- Cannon DT, Aprill CN.** Lumbosacral epidural steroid injections. *Arch Phys Med Rehabil*. Mar 2000;81(3 Suppl 1):S87-98; quiz S99-100.
- Campa JA and Payne R:** Pain syndromes due to cancer treatment. In: Patt RB, editor. *Cancer pain*. Philadelphia Lippincott; 1993. p. 41-56.
- Caraceni A, Gorni G, Zecca E, et al.:** More on the use of nonsteroidal anti-inflammatories in the management of cancer pain. *J Pain Symptom Manage* 2001; 21 (2): 89-91.

Caraceni A, Zecca E, Bonezzi C, et al.: Gabapentin for neuropathic cancer pain: a randomized controlled trial from the Gabapentin Cancer Pain Study Group. *J Clin Oncol* 2004; 22 (14): 2909-17.

Carette S, Leclaire R, Marcoux S, et al. Epidural corticosteroid injections for sciatica due to herniated nucleus pulposus. *N Engl J Med*. Jun 5 1997;336(23):1634-40.

Caudill MA. Medical evaluation and management of chronic pain from a primary care perspective. In: Aronoff GM, ed. *Evaluation and Treatment of Chronic Pain*. 3rd ed. Baltimore, Md: Williams & Wilkins; 1998:67-79.

Cetin N, Aytar A, Atalay A, et al.: Comparing hot pack, short-wave diathermy, ultrasound, and TENS on isokinetic strength, pain, and functional status of women with osteoarthritic knees: a single-blind, randomized, controlled trial. *Am J Phys Med Rehabil*. Jun 2008;87(6):443-51.

Chaney MA: Side effects of intrathecal and epidural opioids. *Can. J Anesth*, 1995; 42: 891-898.

Chapman CR and Syrjala KL: Measurement of pain. In: Loeser JD, Butler SH, Chapman CR, Turk DC (Eds). *Bonica's Management of Pain*. Philadelphia: Lippincott, Williams and Wilkins, 2001, pp 310–328.

Charlton E: Ethical guidelines for pain research in humans. Committee on Ethical Issues of the International Association for the Study of Pain; *Pain*, 1995; 63: 277–278.

Cheing GL, Hui-Chan CW: Transcutaneous electrical nerve stimulation: nonparallel antinociceptive effects on chronic clinical pain and acute experimental pain. *Arch Phys Med Rehabil*. Mar 1999;80(3):305-12.

Chen L, Tang J, White PF, et al: The effect of location of transcutaneous electrical nerve stimulation on postoperative opioid analgesic requirement: acupoint versus nonacupoint stimulation. *Anesth Analg*. Nov 1998;87(5):1129-34.

Cherny NI, Coyle N and Foley KM: Suffering in the advanced cancer patient: a definition and taxonomy. *J Palliat Care*, 1994; 10: 57–70.

Cherny N, Ripamonti C, Pereira J, et al.: Strategies to manage the adverse effects of oral morphine: an evidence-based report. *J Clin Oncol* 2001; 19 (9): 2542-54.

- Cipriano G Jr, de Camargo Carvalho AC, Bernardelli GF, et al:** Short-term transcutaneous electrical nerve stimulation after cardiac surgery: effect on pain, pulmonary function and electrical muscle activity. *Interact Cardiovasc Thorac Surg*. Aug 2008;7(4):539-43.
- Clark MR, Treisman GJ.** Perspectives on pain and depression. *Adv Psychosom Med*. 2004;25:1-27.
- Cl Von Baeyer and Cl Hicks:** Echelle de la douleur chez les enfants: en faveur d'un systeme commun, pain research and management. Summer 2000; 5, number 2: 157-160.
- Cleeland CS, Mendoza TR, Wang XS, Chou C, and Harle MT:** Assessing symptom distress in cancer patients: the M.D. Anderson Symptom Inventory. *Cancer* 2000; 89:1634–46.
- Cleeland CS and Syrjala KL:** How to assess cancer pain. In: Turk DC, Melzack R (Eds). *Handbook of Pain Assessment*, 2nd ed. New York: Guilford Press, 2001, pp 362–390.
- Clement-Jones V, McLoughlin L, Tomlin S, et al:** Increased beta-endorphin but not met-enkephalin levels in human cerebrospinal fluid after acupuncture for recurrent pain. *Lancet*. Nov 1 1980;2(8201):946-9.
- Collins VJ:** Principles of Anesthesiology. Vol 2, 3rd ed., ed Lee and Febiger, Philadelphia (1993).PP 1517-1630 and 1317-1345.
- Conn A, Buenaventura RM, Datta S, et al.** Systematic review of caudal epidural injections in the management of chronic low back pain. *Pain Physician*. Jan-Feb 2009;12(1):109-35.
- Cottingham B, Phillips PD, Davies GK, et al:** The effect of subcutaneous nerve stimulation (SCNS) on pain associated with osteoarthritis of the hip. *Pain*. Jul 1985;22(3):243-8.
- Cooper MG, Keneally JP and Kinchington D:** Continuous brachial plexus neural blockade in a child with intractable cancer pain. *J Pain Symptom Manage*, 1994; 9:277-281.
- Craig KD, Prkachin KM and Grunau RE:** The facial expression of pain. In: Turk DC and Melzack R (Eds). *Handbook of Pain Assessment*, 2nd ed. New York: Guilford Press, 2001, pp 153– 169. Dworkin SF, Whitney CW. Relying on objective and subjective measures of chronic pain. In: Turk DC, Melzack R (Eds). *Handbook of Pain Assessment*, 2nd ed. New York: Guilford Press, 2001, pp 429–446.

Cross, SA. Pathophysiology of Pain. This article on pain mechanisms is still by far the best summary we have encountered!. *Proc*1994 (69) 375-383

D'Alberto A. Auricular acupuncture in the treatment of cocaine/crack abuse: a review of the efficacy, the use of the National Acupuncture Detoxification Association protocol, and the selection of sham points. *J Altern Complement Med.* Dec 2004;10(6):985-1000.

Danziger N, Rozenberg S, Bourgeois P, et al. Depressive effects of segmental and heterotopic application of transcutaneous electrical nerve stimulation and piezo-electric current on lower limb nociceptive flexion reflex in human subjects. *Arch Phys Med Rehabil.* Feb 1998;79(2):191-200.

Derby R, Date ES, Lee C, et al. Size and aggregation of corticosteroids used for epidural injections. *ISIS Sci Newsletter.* 2006;5(4):30-7.

Dersh J, Polatin PB, Gatchel RJ. Chronic pain and psychopathology: research findings and theoretical considerations. *Psychosom Med.* 2002;64:773-786.

Deyo RA, Walsh NE, Martin DC, et al. A controlled trial of transcutaneous electrical nerve stimulation (TENS) and exercise for chronic low back pain. *N Engl J Med.* Jun 7 1990;322(23):1627-34.

Dickenson AH, Chapman V and Green GM: The pharmacology of excitatory and inhibitory amino acid-mediated events in the transmission and modulation of pain in the spinal cord. *Gen Pharmacol* 1997; 28:633-638.

Dickenson AH, Matthews EA and Suzuki R: Central nervous system mechanisms of pain in peripheral neuropathy. In: Hansson PT, Fields HL, Hill RG, Marchettini P (Eds). *Neuropathic Pain: Pathophysiology and Treatment, Progress in Pain Research and Management, Vol. 21.* Seattle: IASP Press, 2001, pp 85–106.

Dickenson AH, Suzuki R and Matthews EA, et al: Balancing excitations and inhibitions in spinal circuits. In: L. Villanueva, A. H. Dickenson and H. Ollat (Eds.). *The Pain System in Normal and Pathological States: A Primer for Clinicians, Progress in Pain Research and Management, Vol. 31.* Seattle: IASP Press, 2004, pp 79–106.

Dickenson, AH: Gate control theory of pain stands the test of time. *Br. J. Anaesth:* 2002; 88(6): 755–757.

- Doi K, Nikai T, Sakura S, et al:** Intercostal nerve block with 5% tetracaine for chronic pain syndromes. *J Clin Anes*, 2002; 14:39-41.
- Dominguez E.** The practice of cervical transforaminal injection of steroids deserves more scrutiny. *Pain Physician*. Jul 2005;8(3):323-4.
- Doralina L. Charles B and Kenneth J:** Practical Guidelines in Oncology, V.1, 2007; 18.
- Downie PA:** The place of physiotherapy in hospice care principles and practice. New York: Springer, 1983 chapter 12.
- Doyle D, Hanks GWC and Cherny NI:** Oxford textbook of palliative medicine, 3rd ed. New York: Oxford University Press, 2004.
- Doyle E, Robinson D and Morton NS:** Comparison of patient-controlled analgesia with and without a background infusion after lower abdominal surgery in children. *Br J Anaesth*, 1993; 71: 670–3.
- Dray A, Urban L and Dickenson A:** Pharmacology of chronic pain. *Trends in Pharmacology sciences*, 1994; 15: 19.
- Dray A:** Peripheral mediators of pain. In: Dickenson AH, Besson JM (Eds). *The Pharmacology of Pain*, Handbook of Experimental Pharmacology, Vol. 130. Springer-Verlag, 1997, pp 21–42.
- Dunteman ED:** Levetiracetam as an adjunctive analgesic in neoplastic plexopathies: case series and commentary. *J Pain Palliat Care Pharmacother*.2005; 19(1):35 -43.
- Dworkin RH, O'Connor AB, Backonja M, et al.:** Pharmacologic management of neuropathic pain: evidence-based recommendations. *Pain*, 2007; 132 (3): 237-51.
- Eisenberg E, Berkey CS, Carr DB, Mosteller F and Chalmers TC:** Efficacy and safety of nonsteroidal anti-inflammatory drugs for cancer pain: a meta-analysis. *J Clin Oncol*, 1994; 12: 2756-65.
- Eisenberg E, River Y Shifrin A and Krivoy N:** Antiepileptic drugs in the treatment of neuropathic pain. *Drugs*, 2007; 67:1265 -1289.
- Ellershaw JE and Dickman A:** Clonazepam as an adjuvant analgesic in patients with cancer-related neuropathic pain. *J Pain Symptom Manage* 2003; 26 (6): 1073-4.

- Ellison N, Loprinzi CL, Kugler J et al:** Phase III placebo-controlled trial of capsaicin cream in the management of surgical neuropathic pain in cancer patients. *J Clin Oncol*, 1997; 15:2974–2980.
- Epstein JB, Epstein JD, Epstein MS, Oien H and Truelove EL:** Management of pain in cancer patients with oral mucositis: follow-up of multiple doses of Doxepin oral rises. *J Pain Symptom Manage*.2007; 33(2):111 -114.
- Eriksson MB, Sjolund BH, Sundbarg G.** Pain relief from peripheral conditioning stimulation in patients with chronic facial pain. *J Neurosurg*. Jul 1984;61(1):149-55.
- Esmail Z, Montgomery C, Courtrn C, Hamilton D and Kestle J:** Efficacy and complications of morphine infusions in postoperative pediatric patients. *Paediatr Anaesth*: 1999; 9:321-327.
- Etches RC:** Patient-controlled analgesia.*Surg Clin North Am* 1999; 79: 297–312
- Fagade OO, Oginni FO, Obilade TO.** Comparative study of the therapeutic effect of a systemic analgesic and transcutaneous electrical nerve stimulation (TENS) on post-IMF trismus and pain in Nigerian patients. *Niger Postgrad Med J*. Jun 2005;12(2):97-101.
- Fahnenstich H, Steffan J, Kau N and Bartmann P:** Fentanyl-induced chest wall rigidity and laryngospasm in preterm and term infants. *Crit Care Med* 2000; 28:836-839.
- Faccenda, Kathryne A, Finucane and Brendan T:** Complications of Regional Anaesthesia: Incidence and Prevention. *Drug Safety*.2001; 24(6):413-442.
- Fainsinger RL and Bruera E:** Is this opioid analgesic tolerance? *J Pain Symptom Manage* 1995; 10 (7): 573-7.
- Fernandez-Del-Olmo M, Alvarez-Sauco M, Koch G, et al:** How repeatable are the physiological effects of TENS?. *Clin Neurophysiol*. Aug 2008;119(8):1834-9.
- Fields HL, Basbaum AI and Heinricher MM:** Central nervous system mechanisms of pain modulation. In: McMahon, Koltzenburg M (Eds). *Textbook of Pain*, Edinburgh: Churchill Livingstone, 2005.
- Fields HL:** Can opiates relieve neuropathic pain? *Pain*, 1988; 35: 365-368.
- Foley KM:** Pain syndrome in patients with cancer. In **Portenoy RK**, Kanner RM, eds *Pain Management: Theory and Practice* Philadelphia: FA Davis: 1996; 191-215.

Frampton V. Transcutaneous electrical nerve stimulation and chronic pain. In: Wells PE, Frampton V, Borosher D, eds. *Pain Management in Physical Therapy*. Norwalk, CT: Appleton & Lange; 1988.

Frick U, Kharraz A, Hauser S, et al. Comparison perception of singular transcranial magnetic stimuli by subjectively electrosensitive subjects and general population controls. *Bioelectromagnetics*. May 2005;26(4):287-98.

Gadsby G, Flowerdew M. Nerve stimulation for low back pain--a review. *Nurs Stand*. Jul 16 1997;11(43):32-3.

Gagnon B, Almahrezi A and Schreier G: Methadone in the treatment of neuropathic pain. *Pain Res Manag* 2003; 8 (3): 149-154.

Gaillet JC, Biraud JC, Bessou M, et al. Modifications of baropodograms after transcutaneous electric stimulation of the abductor hallucis muscle in humans standing erect. *Clin Biomech (Bristol, Avon)*. Dec 2004;19(10):1066-9.

Gammaitoni AR and Davis MW: Pharmacokinetics and tolerability of lidocaine patch 5% with extended dosing. *Ann Pharmacother*; 2002; 36: 236-240.

Ganong WF: Review of Medical physiology. 15th ed, Appleton and Lange, Norwalk, California. PP 124-135.

George RM and Ahmedzai SH: The management of neuropathic pain in cancer: clinical guidelines for the use of adjuvant analgesics. *Indian J Cancer* 2000; 37 (1): 4-9.

Gilles FJ, Shankle W and Dooling EC: Myelinated tracts: growth patterns. In: Gilles FH, Leviton A, Dooling EC, eds. *The developing human brain: growth and epidemiologic neuropathology*. Boston: John Wright, 1983: 117-83.

Ghoname EA, Craig WF, White PF. Percutaneous electrical nerve stimulation (PENS): an alternative to opioid analgesics for treating headache pain after electroconvulsive therapy. *Headache*. 1999;39:502-5.

Ghoname EA, Craig WF, White PF, et al. Percutaneous electrical nerve stimulation for low back pain: a randomized crossover study. *JAMA*. Mar 3 1999;281(9):818-23.

Gisela M and Johannes B: Pocket compendium of peripheral nerve blocks, 2006; 3.

Goldberg G R and Morrison R S: Pain Management in Hospitalized Cancer Patients: A Systematic Review. *JCO*, 2007; 25: 1792-1801.

Graves KD: Social cognitive theory and cancer patients' quality of life: A meta-analysis of psychosocial intervention components. *Health Psychology* 2003; 22:210-219.

Grant DJ, Bishop-Miller J, Winchester DM, et al. A randomized comparative trial of acupuncture versus transcutaneous electrical nerve stimulation for chronic back pain in the elderly. *Pain*. Jul 1999;82(1):9-13.

Goodwin J, Bajwa ZH. Evaluating the patient with chronic pain. In: Warfield CA, Bajwa ZH, eds. *Principles and Practice of Pain Medicine*. 2nd ed. New York, NY: McGraw-Hill; 2004:61-68.

Gyory AN, Caine DC. Electric pain control (EPC) of a painful forearm amputation stump. *Med J Aust*. Jul 30 1977;2(5):156-8.

Guay DR: Adjunctive agents in the management of chronic pain. *Pharmacotherapy*, 2001; 21 (9): 1070-81.

Gutstein A: Opioid analgesics. In: Goodman LS, Hardman JG, Limbird LE, et al.: *Goodman & Gilman's. The Pharmacological Basis of Therapeutics*; 10th ed. New York, NY: McGraw-Hill, 2001, pp 596-7.

Guyton A and Hall J: *Text book of Medical Physiology*. 11th ed., W.b. Sanders Company. Philadelphia, Montrial, London, Toronto, Tokyo (2005) pp 520-531.

Hadjistavropoulos and Craig K D: A theoretical framework for understanding self-report and observational measures of pain: a communications model. *Behaviour Research and Therapy*. Volume 40, Issue 5, May 2002, Pages 551-570.

Hall and Anand: *Physiology of Pain and Stress in the Newborn Neoreviews*. 2005; 6: 61.

Hamill, RJ & Rowlingson, JC: *Handbook of Critical Care Pain Management*. McGraw-Hill. 1994. (ISBN 0-07-025814-7).

Hamza MA, White PF, Ahmed HE, et al: Effect of the frequency of transcutaneous electrical nerve stimulation on the postoperative opioid analgesic requirement and recovery profile. *Anesthesiology*. Nov 1999;91(5):1232-8.

Hanks GW, Conno F, Cherny N, et al.: Morphine and alternative opioids in cancer pain: the EAPC recommendations. *Br J Cancer*: 2001; 84 (5): 587-93.

Hanks GW, Nugent M, Higgs CM, et al.: Oral transmucosal fentanyl citrate in the management of breakthrough pain in cancer: an open, multicentre, dose-titration and long-term use study. *Palliat Med*, 2004; 18 (8): 698-704.

Hansson Pand Lundeborg T. In: Wall P, Melzack R (editors). *Textbook of pain*. 4th edition. Edinburgh: Churchill Livingstone, 1999; Pp.1341-6.

Haughney A: Nausea and vomiting in end-stage cancer. *American Journal of Nursing*: 2004; 104(11):41–48.

Hu WQ, Zhang Z, Zhou P, et al. [Clinical observation on effect of qiangji recipe combined with physical therapy in treating ankylosing spondylitis]. *Zhongguo Zhong Xi Yi Jie He Za Zhi*. Jan 2005;25(1):64-6.

Hunt SP and Mantyh PW: Molecular basis of pain control. *Nat Rev*, 2001; 2:83–91.

Jackson K, Ashby M and Martin P.: "Burst" ketamine for refractory cancer pain: an open-label audit of 39 patients. *J Pain Symptom Manage* 2001; 22 (4): 834-42.

Jackson KC and Stanford BL: *Opioid Use in Clinical Practice*.2003.

Jarvik JG, Deyo RA. Diagnostic evaluation of tow back pain with emphasis on imaging. *Ann Intern Med*. 2002;137:586-597.

Jayme MK, Concha ME, Sarino JA, Uy HG: Effectivity of the electronic dental anesthesia in controlling pain caused by local anesthetic injections. *J Philipp Dent Assoc*. Dec-1999 Feb 1998;50(3):39-52.

Joint Commission on Accreditation of Healthcare Organizations (JACHO): *Pain Assessment and Management*. Oakbrook Terrace, IL: author 2000.

Johnson MI, Ashton CH, Thompson JW: An in-depth study of long-term users of transcutaneous electrical nerve stimulation (TENS). Implications for clinical use of TENS. *Pain*. Mar 1991;44(3):221-9.

Jost T: Medicare and Medicaid financing of pain management. *The Journal of Pain*: 2000; 1(3),

183–94.

Karl HW, Tyler DC and Krane EJ: Respiratory depression after low-dose caudal morphine. *Can J Anaesth* 1996; 43: 1065–7.

Katon W. The impact of major depression on chronic medical illness. *Gen Hosp Psychiatry*. 1996;18:215-219.

Keefe FJ and Williams DA: Assessment of pain behaviors. In: Turk DC, Melzack R (Eds). *Handbook of Pain Assessment*, 2nd ed. New York: Guilford Press, 2001, pp 257–274.

Keefe FJ, Rumble ME, Scipio CD, Giordano LA, Perri LM. Psychological aspects of persistent pain: current state of the science. *J Pain*. 2004;5:195-211.

Keskinbora K, Pekel AF and Aydinli I: Gabapentin and an opioid combination versus opioid alone for the management of neuropathic cancer pain: a randomized open trial. *J Pain Symptom Manage* 2007; 34 (2): 183-9.

Khojainova N, Santiago-Palma J, Kornick C et al: Neuroleptics in the management of cancer pain. *J Pain Symptom Manage* 2002; 23:346–350.

Kumar RJ, Menon KV and Ranjith TC: Use of epidural analgesia for pain management after major spinal surgery. *J Orthop Surg (Hong Kong)*. 2003 Jun; 11(1):67-72.

Law PP, Cheing GL: Optimal stimulation frequency of transcutaneous electrical nerve stimulation on people with knee osteoarthritis. *J Rehabil Med*. Sep 2004;36(5):220-5.

Lebovits AH. The psychological assessment of patients with chronic pain. *Curr Rev Pain*. 2000;4:122-126.

Lee BH, Scharff L and Sethna NF: Physical therapy and cognitive-behavioral treatment for complex regional pain syndromes. *J Pediatr* 2002; 141:135-140.

Lesko SM and Mitchell AA: The safety of acetaminophen and ibuprofen among children younger than two years old. *Pediatrics*: 1999; 104:952-952.

Levine J, Fields H and Basbaum A: Peptide and the primary afferent nociceptor. *J. of Neuroscience*, 1993; 13: 2273-2286.

Lichtor JL, Sevarino FB, Joshi GP, Busch MA and Nordbrock E, Ginsberg B: The relative potency of oral transmucosal fentanyl citrate compared with intravenous morphine in the

treatment of moderate to severe postoperative pain. *Anesth Analg*: 1999; 89:732-738.

Lossignol DA, Obiols-Portis M and Body JJ: Successful use of ketamine for intractable cancer pain. *Support Care Cancer* 2005; 13 (3): 188-93.

Lubenow T, Ivankovich A and McCarthy R: Management of acute postoperative pain. Chapter 45: Clinical Anesthesia. 3rd ed., Eds: Barash G, Gullen B and Stoelting R. Lippincott-Raven Publishers, Philadelphia, New York. 1997; PP 1305-1334.

Lucas LK and Lipman AG: Recent advances in pharmacotherapy for cancer pain management. *Cancer Pract*. 2002; 10 Suppl 1:S14-S20.

Lussier D, Huskey AG and Portenoy RK: Adjuvant analgesics in cancer pain management. *Oncologist* 2004; 9 (5): 571-91.

Lussier D and Portenoy RK: Adjuvant analgesics in pain management. In: Doyle D, Hanks G, Cherny N, et al., eds. *Oxford Textbook of Palliative Medicine*, Third Edition. Oxford, England: Oxford University Press, 2003:349-377.

Macintyre PE and Ready LB: acute Pain Management – a Practical Guide, 2nd edn. London: WB Saunders, 2001.

Maddox RR Williams CK and Fields M: "Respiratory monitoring in patient-controlled analgesia," *AM J Health-Sys Pharm* 2004; 2628-9.

Maizels M and McCarberg B: Antidepressants and antiepileptic drugs for chronic cancer pain. *Am Fam Physician*.2005; 71:483 -490.

Malan Jr: Opioid pharmacology in anesthesia pain management. *ASA Refresher Lectures* 2000:422.

Manfredi PL, Gonzales GR, Sady R et al: Neuropathic pain in patients with cancer. *J Palliat Care* 2003; 19: 115-118.

Mansfield MD. et al. BJA. 1996 (76) 358-361. Influence of dose and timing of administration of morphine on postoperative pain and analgesic requirements.

Martin JH: Neuroanatomy text and atlas. 2nd ed., Appleton and Lange, Asimon and Schuster Company 1996; PP 125-140.

Martinez-Lavin M, Lopez S, Medina M, Nava A. Use of the leeds assessment of neuropathic symptoms and signs questionnaire in patients with fibromyalgia. *Semin Arthritis Rheum.* 2003;32:407-411.

Masand PS and Gupta S: Selective serotonin-reuptake inhibitors: an update. *Harvard Rev Psychiatry*, 1999; 7:69-84.

McDonnell FJ, Sloan JW and Hamann SR: Advances in cancer pain management. *Curr Pain Headache Rep* 2001; 5 (3): 265-71.

McNeely JK and Trentadue NC: Comparison of patient-controlled analgesia with and without nighttime morphine infusion following lower extremity surgery in children. *J Pain Symptom Manage* 1997; 13:268-273.

McNicol E, Strassels S, Goudas L, et al.: Nonsteroidal alone or combined with opioids, for cancer pain: *Cochran Database Syst Rev J Clin Oncol* 2005; Jan (1).

McNicol E, Strassels S, Goudas L, et al.: Nonsteroidal anti-inflammatory drugs, alone or combined with opioids, for cancer pain: a systematic review. *J Clin Oncol* 2004; 22 (10): 1975-92.

McQueen FM, Stewart N, Crabbe, J, et al. Magnetic resonance imaging of the wrist in early rheumatoid arthritis reveals a high prevalence of erosions at four months after symptom onset. *Ann Rheum Dis.* 1998;57:350-356.

Meizack K and Wau P D: Pain mechanism: a new theory. *Current contents*, 1982; 7: 23.

Mercadante S, Arcuri E, Fusco F, et al.: Randomized double-blind, double-dummy crossover clinical trial of oral tramadol versus rectal tramadol administration in opioid-naive cancer patients with pain. *Support Care Cancer*, 2005; 13 (9): 702-7.

Mercadante S, Villari P, Ferrera P, et al.: Opioid plasma concentrations during a switch from transdermal fentanyl to methadone. *J Palliat Med* 2007; 10 (2): 338-44.

Miaskowski C, Cleary J, Burney R, Coyne P, et al: American Pain Society Clinical Practice Guideline Series, No. 3: Guide for the management of cancer pain in adults and children. Glenview, IL: American Pain Society; 2005.

Monitto CL, Greenberg RS, Kost-Byerly S, et al: The safety and efficacy of parent-/nurse-

controlled analgesia in patients less than six years of age. *Anesth Analg* 2000; 91:573-579.

Morgan G and Mikail M: Clinical Anesthesiology, 2nd ed., Appleton and Lange, Asimon and Schuster Company, 1996; PP 130-145,193-200 and 274-315.

Murina F, Bianco V, Radici G, et al. Transcutaneous electrical nerve stimulation to treat vestibulodynia: a randomised controlled trial. *BJOG*. Aug 2008;115(9):1165-70.

Nakamura Y and Chapman CR: Measuring pain: an introspective look at introspection. *Conscious Cogn* 2002; 11:582–592.

Otsuka M and Yoshioka K: Neurotransmitter functions of mammalian tackykinins. *Physiological Reviews*, 1993; 73: 229-303.

Paice JA and Fine BR: Pain at the end of life. In *Textbook of Palliative Medicine*, B. Ferrell and N. Coyle (eds.) 2001; (pp. 76–90). Oxford: Oxford University Press.

Panke JT: Difficulties in managing pain at the end of life. *American Journal of Nursing*: 2002; 102 (7):26–33

Pereira J, Lawlor P, Vigano A, et al.: Equianalgesic dose ratios for opioids. a critical review and proposals for long-term dosing. *J Pain Symptom Manage* 2001; 22 (2): 672-87.

Peters K, Carrico D, Burks F: Validation of a sham for percutaneous tibial nerve stimulation (PTNS). *Neurourol Urodyn*. Jul 31 2008.

Pike PM. Transcutaneous electrical stimulation:. Its use in the management of postoperative pain. *Anaesthesia*. Feb 1978;33(2):165-71.

Poitras S, Brosseau : Evidence-informed management of chronic low back pain with transcutaneous electrical nerve stimulation, interferential current, electrical muscle stimulation, ultrasound, and thermotherapy. *Spine J*. Jan-Feb 2008;8(1):226-33.

Portenoy RK: Pain syndromes in patients with cancer and HIV/AIDS. In: Portenoy RK, ed. *Contemporary diagnosis and management of pain in oncologic and AIDS patients*. Newtown, PA: Handbooks in Healthcare, 1998: 44–70.

Portenoy R: Management of cancer pain. *Lancet*, 1999;353:1695–1700.

Portenoy RK and Frager G: Pain management: pharmacological approaches. In: von Gunten CF, ed.: *Palliative Care and Rehabilitation of Cancer Patients*. Boston, Mass: Kluwer Academic Publishers, 1999, pp 1-29.

- Potter J, Higginson IJ, Scadding JW, Quigley C.** Identifying neuropathic pain in patients with head and neck cancer: use of the Leeds Assessment of Neuropathic Symptoms and Signs Scale. *J R Soc Med.* 2003;96:379-383.
- Price D, Mao J and Frenk H:** The N-methyl-D-aspartate antagonist dextromethorphan selectiviely reduces temporal summation of secand pain. *Pain*, 1994; 59: 165-174.
- Radecki SE, Brunton SA.** Randomized clinical trial of a diagnostic instrument for pain complaints, *Fam Med.* 1999;31:713-721.
- Ragawski MA:** Therapeutic potential of excitatory aminoacid antagonist; Channel blocks and 2, 3-benzodiazepines. *Trends in Pharmacology Science*, 1993; 14: 325-331.
- Reddy S and Patt RB:** The benzodiazepines as adjuvant analgesics. *J Pain Symptom Manage* 1994; 9:510-514.
- Rievi T, Wadhwa S and Bijlani V:** Development of spinal substrate for nociception. *Pain*^{Suppl} 1987; 4:195.
- Richardson, BP.** *Ann NY Acad Sci* 1990 (600) 511-519. Serotonin and Nociception.
- Riew KD, Yin Y, Gilula L, et al.** The effect of nerve-root injections on the need for operative treatment of lumbar radicular pain. A prospective, randomized, controlled, double-blind study. *J Bone Joint Surg Am.* Nov 2000;82-A(11):1589-93.
- Riley J, Ross JR, Rutter D, et al.:** No pain relief from morphine? Individual variation in sensitivity to morphine and the need to switch to an alternative opioid in cancer patients. *Support Care Cancer* 2006; 14 (1): 56-64.
- Ripamonti C and Bianchi M:** The use of methadone for cancer pain. *Hematol Oncol Clin North Am* 16 (3): 543-55, 2002.
- Robb KA, Bennett MI, Johnson MI, et al:** Transcutaneous electric nerve stimulation (TENS) for cancer pain in adults. *Cochrane Database Syst Rev.* Jul 16 2008;CD006276.
- Robinson SL, Rowbotham DJ and Mushambi M:** Electronic and disposable patient-controlled analgesia systems. A comparison of the Graseby and Baxter systems after major gynaecological surgery. *Anaesthesia* 1992; 47: 161–3.
- Rudolph H, Cade JF, Morley PT, Packer JS and Lee B:** Smart technology improves patient-controlled analgesia: a preliminary report. *Anesth Analg* 1999; 89: 1226–32
- Roscoe JA, Matteson SE, Morrow GR, et a.:** Acustimulation wrist bands are not effective for the

control of chemotherapy-induced nausea in women with breast cancer. *J Pain Symptom Manage*. Apr 2005;29(4):376-84.

Rowbotham DJ: Recent advances in the non-pharmacological management of postoperative nausea and vomiting. *Br J Anaesth*. Jul 2005;95(1):77-81.

Saper JR, Lake AE 3rd, Cantrell DT et al: Chronic daily headache prophylaxis with tizanidine: a double-blind, placebo-controlled, multicenter outcome study. *Headache* 2002; 42:470–482.

Schaible H, Hope P and Lang C: Calcitonin gene related peptide causes intrasoinal spreading of substance P released by peripheral stimulation. *European Journal of Neuroscience*, 1992; 4: 750-757.

Sharma SK, Gajraj NM, Sidawi JE, et al: EMLA cream effectively reduces the pain of spinal needle insertion. *Reg Anesth*, 1996; 21:561-564.

Sharp R, Olawson A and Jones R: Postoperative pain relief. Chapter 26: In: *Clinical Anesthesia*. Eds: Aitkenhead A and Jones R, Churchill Livingstone, New York, Edinburgh, London, Madried, Tokyo 1996; PP637-665.

Shipton, EA: Pain, Acute & Chronic.1993;17.

Shipton EA, Tait B : "Flagging the pain: preventing the burden of chronic pain by identifying and treating risk factors in acute pain". *European journal of anaesthesiology* **22** (6) 2004; 405–12. doi:10.1017/S0265021505000694. PMID 15991501.

Sicard A. Les injections médicamenteuses extra-durales par voie sacro-coccygienne. *CR Soc Biol Paris*. 1901;53:369.

Simone DA: Peripheral Mechanisms of Pain Perception. In: Abram E, ed. *The atlas of anesthesia, Pain management*, Philadelphia, PA: Churchill Livingston, 1998:1.10-21.

Sluka and Walsh: Sluka KA, Walsh DM: Transcutaneous electrical nerve stimulation: basic science mechanisms and clinical effectiveness. *Journal of Pain* 2003; 4(3):109-121.

Smith GA, Strausbaugh SD, Harbeck-Weber C, Cohen DM and Shields BJ: New non-cocaine-containing topical anesthetics compared with tetracaine-adrenaline-cocaine during repair of lacerations. *Pediatrics*: 1997; 100:825-830.

Solomon RA, Viernstein MC, Long DM: Reduction of postoperative pain and narcotic use by transcutaneous electrical nerve stimulation. *Surgery*. Feb 1980;87(2):142-6.

- Stockler M, Vardy J, Pillai A, et al.:** Acetaminophen (paracetamol) improves pain and well-being in people with advanced cancer already receiving a strong opioid regimen: a randomized, double-blind, placebo-controlled cross-over trial. *J Clin Oncol* 2004 22 (16): 3389-94.
- Stoelting RK and Miller RD:** Basic of anesthesia 3rd Ed's. Churchill Livingstone. New York. Edinburgh, London, Madrid and Tokyo 1994; 443-453.
- Swanwick M, Haworth M and Lennard RF:** The prevalence of episodic pain in cancer: a survey of hospice patients on admission. *Palliat Med* 2001; 15:9–18.
- Swarm RA and Cousins MJ:** Anesthetic techniques for pain control. IN: Doyle D, Hanks GWS, MacDonald N, eds. *Oxford textbook of palliative medicine*, 2nd ed. New York: Oxford University Press, 1998; 390.
- Sweeder J:** Educating clinicians on effective pain management. *The Pain Clinic*: 2002; 4(1):11–19.
- Sze FK, Chung TK, Wong E, et al:** Pain in Chinese cancer patients under palliative care. *Palliat Med*, 1998; 12:271–277.
- Taylor P, Hallett M, Flaherty .:** Treatment of osteoarthritis of the knee with transcutaneous electrical nerve stimulation. *Pain*. Oct 1981;11(2):233-40.
- Thompson W:** Placebos: A review of the placebo response. *American Journal of Gastroenterology*: 2000; 95(7):1637.
- Tremont-Lukats IW, Megeff C and Backonja MM:** Anticonvulsants for neuropathic pain syndromes: mechanisms of action and place in therapy. *Drugs* 2000; 60:1029-1052.
- Tucker K:** Deceptive placebo administration. *American Journal of Nursing*: 2001; 101(8):55.
- Turk DC and Gatchel RG:** Psychological approach to pain management .New York: The Guilford press.2002; 102-105.
- Tugay N, Akbayrak T, Demirtürk F, et al:** Effectiveness of transcutaneous electrical nerve stimulation and interferential current in primary dysmenorrhea. *Pain Med*. May-Jun 2007;8(4):295-300.
- Turk DC and Melzack R:** Handbook of Pain Assessment, 2nded. New York: Guilford Press, 2001, pp 295–314.

- Vetter T:** Pediatric patient-controlled analgesia with morphine versus meperidine. *J Pain Symptom Manage* 1992; 7:204-208.
- Vickers A, Wilson P and Kleijnen J:** Acupuncture. *Quality and Safety in Health Care* 2002; 11:92-97
- Wall PD and Melzack R:** (Eds). Textbook of Pain, 4th ed. Edinburgh: Churchill Livingstone, 1999.
- Wallace MS.** Pharmacologic treatment of neuropathic pain. *Curr Pain Headache Rep.* 2001;5:138-150.
- Waston CBN:** antipressent drugs as adjuvant therapy *J pain symptom manage:* 1994; 9:392.
- Weinstein SM.** Physical examination. In: Ashburn MA, Rice LJ, eds. *The Management of Pain.* Philadelphia, Pa: Churchill Livingstone; 1998:17-25.
- Weisman SJ, Bernstein B and Schechter NL:** Consequences of inadequate analgesia during painful procedures in children: *Arch Pediatr Adolesc Med* 1998; 152:147-149.
- Woolf CJ:** Pain: moving from symptom control toward mechanism-specific pharmacologic management. *Ann Intern Med* 2004; 140: 441-451.
- Wright J:** The double-edged sword of COX-2 selective NSAIDs. *Can Med Assoc J* 2002; 167:1131-1136.
- Yaksh TL and Malmberg AB:** Central pharmacology of nociceptive transmission. In: Wall PD, Melzack R, eds. *Textbook of Pain.* 3rd ed. Edinburgh:Churchill Livingstone; 1994.
- Zempsky WT and Schechter NL:** What's new in the management of pain in children? *Pediatr Rev:* 2003; 24:337-347.
- Zhukovsky DS, Gorowski E, Hausdorff J, Napolitano B and Lesser M:** Unmet analgesic needs in cancer patients. *J Pain Symptom Manage* 1995; 10:113-119.