

References



References

- Abbas, A.M.A. (1967): Comparative study of methods used of the isolation of *Toxoplasma gondii*. Bull. WHO, 36: 344. Quoted from Mahgoub, A.M.A. (2001): Characterization of *Toxoplasma gondii* antigen and its use in serodiagnosis of toxoplasmosis. M.Sc. Thesis in Parasitology Faculty of Medicine Cairo University.
- Abdel-Aziz ,F.; Habib, M.and Mohamed, M.K. (2000): Hepatitis C virus infection in a community in the Nile Delta. Hepatology., 32:111-115.
- Abdel-Wahab, M.F.; Zakaria ,S. and Kamal, M. (2004): High seroprevalence of hepatitis C infection among risk groups in Egypt. Am. J. Tropical Medicine and Hygiene, 5: 63-67.
- Aboul- Enein, E.; Abdel Wahab, F.; El-Basestar, S. and Abdel-Aal, A.M. (1983): Toxoplasmosis among people at risk of exposure to infection using IHA test. J. Egypt. Soc. Parasitol., 13: 435-440.
- Aboul- Magd, L.A. ; Arafa, M.S. and El-Ridi, A.M.S. (1987): *Toxoplasma* infection in Zagazig, Egypt: an analytical study. J. Egypt. Soc. Parasitol., 17: 519-524.
- Abraham, B.; Tamby, I. Reynes, J. and Bastien, P. (2000): Polymerase Chain Reaction on sputum for the diagnosis of pulmonary toxoplasmosis in AIDS patients. J. Egypt. Soc. Parasitol. , 19: 20 – 25.
- Abramowicz, M.(2004): Drugs for parasitic infections .J.Egypt.Soc.Parasitol.,46:1-12.

- Akiawa, M.D.; Komata, Y.; Asai, T. and Osamu, M.(1977). Transmission and scanning electron microscopy of host cell entry by *Toxoplasma gondii*. Am. J. Pathol., 87: 285 – 96.
- Alagirl (2009): Relationship between toxoplasmosis and Hepatitis C virus. The American Journal of Gastroenterology., (21) 2: 271-275.
- Al-Amari,O.M.(1994):Prevalence of antibodies to *T.gondii* among blood donors in Abha,Asir Region,Southwestern Saudi Arabia.J.Egypt.Pub.Hlth.Assoc.,69(2):77.
- Alexander, J.H.; Scharton-Krsten, TM.; Yap, C.; Roberts, C.W.and Liew, F. A. (1997): Mechanisms of innate resistance to *Toxoplasma gondii* infection philos Trans R. Soc. Land. B. Biol. Sci .,(12) 3: 1355- 1359.
- Al-Harthi,S.A.;Jamjoom,M.B.and hazi,H.O.(2006):Seroprevalence of *Toxoplasma gondii* among pregnant women in Makkch,Saudi Arabia Umm Al-Qura University.J.Sci.Med.,Engne.,18(2):217-227.
- Ali, S.M.; Allam, S.R.; Negm, A.Y. and EL-Zawawy, L.A. (2003): Vaccination against congenital toxoplasmosis. J. Egypt. Soc. Parasitol., 33 : 863 – 874.
- Allam, S.R.; Ibrahim, I.R. and El-Kerdany, I.D. (1999): Effect of salinity and microwave cooking on the infectivity of encysted forms of *Toxoplasma gondii* and *Trichinella spiralis*. Egypt. J. Med. Sci., 20 (1): 11 – 25.
- Al-Qurashi, A.R.; Ghandour, A.M.; Obeid, O.E.; Al-Mulhim, A. and Makki, S.M.(2001): Seroepidemiological study of *Toxoplasma*

- gondii infection in the human population in the Eastern region. Saudi-Med. J., 22 (1): 13 – 18.
- Al-Ramadi, B.K.; Meissler, J.J.r.; Hunag, D. and Eisentein, T.K. (1992): Immunosuppression induced by nitric oxide and its inhibition by interleukin-4. Eur. J. Immunol., 22 : 2245 – 2254.
 - Alter ,H. (1992): New Kit on the block: evolution of antibody to the hepatitis C virus. Hepatology; 15: 350-353.
 - Alter ,H.J. (1991): Hepatitis C virus in current prospective study. Intern. Med., 115:644-649.
 - Alter ,H.J. (1995): To C or not to C: Journal of the Royal Australian college., 85: 1681-1695.
 - Alter ,M.J.; Hadler,S.C.and Judson, F.N (1990): Risk factors for acute non A, non B hepatitis in the United States and association with hepatitis C virus infection. Journal of the Royal Australian college., 64:2231-2235.
 - Alter ,M.J.;Hutin, Y.J. and Armistrong ,G.L. (2000): Hepatitis C epidemiology. Son Digo academic press, 189-193.
 - Alter,M.J. (1997): Epidemiology of hepatitis C. Journal of Gastroentrol- Hepatol., 26:625, 655.
 - Alter,M.J.(2007): Epidemiology of hepatitis C virus infection. World J Gastroentrol.,13(17):2436-2441.
 - Alvarado – Esquivel. (2006): Laboratory diagnosis of *Toxoplasma gondii* infection and toxoplasmosis. J. infect. Dis .,(16) 7: 573-598.
 - Alvarado,C.;Suarez,M.F.;Rodríguez,A.and Torres,L.(2007) Seroepidemiology of infection *Toxoplasma gondii* in healthy blood donors of Durango,Mexico.B.M.C.Infect.Dis.,7:75-80.

- Ambroise, T.P.; Garin, J.P. and Kien, T. (1971): Our experience of immunofluorescence in the serological diagnosis of toxoplasmosis. *J. infect. Dis.* ,(12) 7: 506-5540.
- Amendoeira, M.R.R.; Sobral, C.A.Q.; Teva, A.; Lima, J.N. and Klein, C.H. (2003): Inquerito serologico para a infeccao por *Toxoplasma gondii* em ameindios isolados, Mato Grosso. *Rev Soc Bras Med Trop* 36: 671-676.
- Anderson, S.E. and Remington, J. S. (1975): Current concepts in diagnosis : The diagnosis of toxoplasmosis. 5th *Med. J.*, 68 (11): 1433 – 1443
- Araujo, F.G. (1994): "Immunization against *Toxoplasma gondii*". *Parasitol. Today*, 10: 358 – 360.
- Araujo, F.G.; Handman, E. and Remington, J.S. (1980): "Use of monoclonal antibodies to detect antigens of *Toxoplasma gondii* in serum and other body fluids. *Infect. Immunol.*, 30 : 42.
- Araya.; Eric.; Luisa ,Garcia.; Jase ,Maria .and Ricardo. (1997): Mangement guide for hepatits C virus. *Journal of the Royal Australian college.*,12(6):210-214.
- Armstrong,G.L.; Wasly, A.and Simard, E.P. (2008): The prevalence of hepatits C virus infection in the united states. *An intern med.*, 44:705-714.
- Arther, R.R.; Hassan ,N.F.and Abdallah, M.Y. (1997): Hepatits C antibody prevalence in blood donors in different governorates in Egypt. *Transactions of the Royal Society of Trop. Med. Hyg.*, 19: 271-344.

- Asai, T. ;Mizuno, F.; Kojima, S.; Takeuchi, T.; Kobayashi, A. and Suzuki, Y. (1992): High correlation in antibody titre between Sabin-Feldman dye test and enzyme-linked immunosorbent assay detecting immunoglobulin G antibodies to the nucleoside triphosphate hydrolase to *Toxoplasma gondii*. J. Clin. Microbiol., 30 (5): 1291 – 1299.
- Ashrafunnessa,K.;Islam,N.N.and Huq,T.(2003):Seroprevalence of toxoplasmosis antibodies among antenatal population in Bangladish.J.Obstet.Gynecol.Res.,24(2):115-119.
- Aspinall, T.V.; Joynson, D.H.; Guy, E.; Hyde, J.E. and Sims, P.F. (2002): The molecular basis of sulfonamide resistance in toxoplasmosis and implication for clinical management of toxoplasmosis. J. Infect. Dis., (11): 1637-1643.
- Balfour, A.H.; Bridges, J.B. and Hardford, J.P. (1980): An evaluation of the *Toxoplasma* IHA test for the detection of antibodies to *Toxoplasma gondii* in human serum. J. Clin. Pathol., 33 : 64.
- Barbosa, I.R.; de Carvalho, C.M. and de Andrade-Neto, V.F. (2008): Toxoplasmosis screening and risk factors amongst pregnant females in natal, northeastern Brazil. Trans. R. Soc. Trop. Med. Hyg. 10 (4): 377-409.
- Bassiounni,S.O.(1992):Parasitological and serological studies of *Toxoplasma* infection in abortion and stillbirth .M.D.Thesis, Faculty of Medicine, Ain shams University.
- Beazley, D.m. and Egerman, R.S. (1998): Toxoplasmosis. Seminars in Perinatology.Trans. R. Soc. Trop. Med. Hyg, 22: 332 – 338 .

- Bellowbuono.; Rojas, M.; HakYangkim.; Yong Bum. and SooTae - Kim. (1995): Effect of combination therapy in patients with chronic hepatitis C, Journal of Gastroenterol- Hepatol., (4) 9: 476-480
- Bennett.; Nalpas, B.; Vallet-Pichard. and Fontaine. (1997): "Management and efficacy of antiviral treatment in patients with hepatitis C virus, Journal of Antiviral therapy., (6) 10: 769-776.
- Bessiere, M.H.; Roques, C.; Berrebi, A.; Barre, V.; Cazaux, M. and Seguela, J.P. (1992): IgA antibody response during acquired and congenital toxoplasmosis. J. Clin. Pathol., 45: 605 – 608.
- Bisceglie, A.M.; Goodman, Z.D. and Ishak, K.G. (1991): Long term clinical and histopathological follow up of chronic post transfusion hepatitis. Journal of Gastroenterol- Hepatology., 18(6): 894- 969.
- Black, M.W. and Boothroyd, J.C. (2000): Life cycle of *Toxoplasma gondii*. Microbiol. Mol. Biol. Rev; 64 (3): 607-632.
- Blanco, J.C.; Angela, S.O.; Maero, E.; Psyzenny, V; Serpente, P. and Garberi, J.C. (1992): Cloning of repetitive DNA sequences from *Toxoplasma gondii* and their usefulness of parasitic detection. Am. J. Tropical Medicine and Hygiene., 46(3): 350-357.
- Bobic, B.; Jevermovic, I.; Marikovic, J.; Sibalic, D. and Djurkovic, O. (1998): Risk Factors for *Toxoplasma* infection in a reproductive age female population in the area of Belgrade, Yugoslavia. Eur. J. Epidemiol. 14 (6): 605 – 610.
- Bobic, D.; Nikolic, A.; Djurkovic-Djakovic, O. (2007): Identification of risk factors for infection with *Toxoplasma gondii* in Serbia as a basis of a program for prevention of congenital toxoplasmosis. Srp. Arch Celok Le. Mar.-Apr; 131 (3-4) : 162 – 167.

- Bosch-Driessen, L.E.; Berendschot, T.T.; Ongkosuwito, J.V. and Rothova, A. (2002): Ocular toxoplasmosis: clinical features and prognosis of 154 patients. *Ophthalmology*, 19: 869-878.
- Bulow, R. and Boothyroid, J.C. (1991): Protection of mice from fatal *Toxoplasma gondii* infection by immunization with P 30 antigen in liposomes. *J. Immunol.*, 147 : 3496 – 3500.
- Busch ,M.P.; Glynn, S.A.and Stramer, S.I. (2005): New Strategy for estimating risk of transfusion transmitted viral infections based on rates of detection of recently infected donors. *Transfusion. J. Immunol.*, 45:254-264.
- Buskila ,D. (2000): Hepatitis C associated arthritis. *Curropin Rheumatol.*, 1: 95.
- Cai, G.; Radzarowsk, T.; Villegas, E.N.; Kastelein, R. and munter, C.A., (2000): Identification of STAT T4-dependent and independent mechanisms of resistance to *Toxoplasma gondii* *J. Immunol.*, 165 (5) : 2619 – 2627.
- Camargo, M.E.; Ferreira, A.W.; Mineo, J.R.; Takiguti, C.K. and Nakahara, O.S. (1978): Immunoglobulin G and immunoglobulin M enzyme-linked immunosorbent assays and defined toxoplasmosis serological patterns. *Infect. Immunol.*, 21 : 55 – 58.
- Camma and Carxi.(2000): Sustained response to combination therapy in patient with chronic hepatitis C., *Journal Gastroenterol Hepatol.*, (5) 15: 567-610.
- Capron, A.; Bout, D. and Dugimont, J.C. (1975): A port de methodes immunoenzymologiques utilisant des antigenes purifies au diagnostic specifique et automatise des infections purasitories.

- Annales de la Societe Beleya de Medicine Tropicale, 55 : 443 – 454.
- Carithers ,R.L. (2000): Extrahepatic manifestations of hepatitis C. update on viral hepatitis. American Association for the liver disease. 153-160.
 - Cesbron-Delauw, M.F.; Guy, B. and Torpier, G. (2003): Molecular characterization of a 23 –KDa major antigen secreted by *Toxoplasma gondii* Proc. Natl. Acad. Sci. USA, 86 : 7537 – 7541.
 - Chalupsky, J. (1981): Application of India ink immunoreaction for the diagnosis of toxoplasmosis. Folia parasitol. Praha, 28 : 131.
 - Chan, T.T.; Liao, T.C.and Yaung ,K.C. (1994): Intrafamilial transmission of hepatitis C virus .Journal Gastroenterol Hepatol., 35(3): 343-352.
 - Chevaliez, S.; Pawlotsky, J.M. (2007): Hepatitis C virus diagnosis and management. J Gastroenterol.,13: 2461-2466.
 - Coelho,R.;Kobayashi,M.and Carvalho,L.B.Jr.(2003):Prevalence of IgG antibodies specific to *Toxoplasma gondii* among blood donors in Rrcife,Northeast Brazil.Rev.Inst.Med.Trop.Sao Paulo,45:229-231.
 - Combe, C.L.; Curiel, T.J.; Moretto, M.M. and Khan, I.A. (2005): NK cells help to induce CD8(+)-T- cell immunity against *Toxoplasma gondii* in the absence of CD4 (+) T cells. Infect. Immunol. , 73 (8): 4913 – 4934.
 - Comparcola,D.;Nobili,V.and Sartorelli,M.R.(2005):Childhood hepatitis C virus infection.J Gastroentrol Hepatol.,20(3):1948-1952.

- Contini, C.o.; Seraceni, S.; Misurati, E.; Grilli, A.; Romani, R. and Cultrera, R. (2001): A multiplex PCR assay for molecular recognition of *T. gondii* stage- specific genes. *J. Eukaryot. Microbiol.*, 1: 191-192.
- Corde ,L.N.; Chosidow ,O.and Frances ,C. (2000): Cutaneous disorder associated hepatitis C virus infection. *Ann Intern Med.*, 151-201.
- Couvereur, G.; Sadak, A.; Fortier, B. and Dubremetz, J.F. (1988): Surface antigens of *Toxoplasma gondii*. *Parasitology*, 9: 1-100.
- Cox, F.E. G. and Waklin, D. (1998): Immunology and immunopathology of human parasitic infection. Quoted from Topley Wilson's Microbiology and microbial infection 5(9) 5: 57 – 84.
- Crucerescu, E. (1998): Epidemiological data on toxoplasmosis: The aspects of congenital toxoplasmosis. *Bacteriologica, Virologica, Parasitologica, Epidemiologica.*, 43 : 147 – 155.
- Darcy,F.;Deslee,D.;Santoro,F.and Charif,H.(1988):Introduction of protective antibody dependant response against toxoplasmosis by in vitro excreted/secreted antigens from tachyzoites of *Toxoplasma gondii*. *Parasite immunology*,10:553-567.
- David, R.G. (2005): Diagnostic tests for hepatitis C. *Hepatology* 3(27): 43-47.
- Davis, G.L. (1997): Early virologic response to treatment with peg-interferon alfa-2b plus ribavirin in patients with chronic hepatitis C. *Journal of Gastroenterol- Hepatology.*, 38:645-652.

- Dawoud,H.A.;Ageely,H.M.;Elshake,A.H.and Heiba,A.A.(2009):Latex agglutination and indirect immunofluorescence tests in the diagnosis of *Toxoplasma gondii* in Saudi Arabia.J.Egypt.Soc.Parasitol.39(1):1-9.
- Delic, D. (2006): Efficacy and safety of peginterferon alfa- 2a and ribavirin in patients with chronic hepatitis C virus infection. Journal of Gastroenterol- Hepatology., 59 (9-15): 415-422.
- Denkers, E.Y. and Gazzinelli, R.T. (1998): Regulation and function of T-cell mediated immunity during *Toxoplasma gondii* infection. Clin. Microbiol. Rev., 11 : 559 – 568
- Derouin ,F. (2001): Anti-toxoplasmosis drugs. Curr. Opin. Investing. Drugs., 2(10): 1368-1374.
- Derouin, F.; Lsullebe, G. and Ballet, J.J. (1987): Sequential determination of IgG subclasses and IgA specific antibodies in primary and reactivation toxoplasmosis. Biomed. Pharmacother., 41 (8): 429 – 462.
- Des jarlis ,D.C.; Diaz, T.and Perlis, T. (2003): Variability in the incidence of human immunodeficiency virus, hepatitis B virus and hepatitis C virus infection among young injecting drug users in New York city. Am. J. Epidemiol., 157:467-471.
- Desmonts, G. and Remington, J.S. (1980): Direct agglutination test for diagnosis of *Toxoplasma* infection method for increasing sensitivity and specificity. J. Clin. Microbiol., 11: 562 – 568.
- Deufic, S.; Buffat, L.and Poynard ,T. (1999): Modeling the hepatitis C virus epidemic in France .Journal of Gastroenterol- Hepatology., 29: 1596-1601.

- Diab, A.K. (2000): Immunological relationship between viral hepatitis and *Toxoplasma gondii* infection in man and animals. M.SC. Thesis in immunology and microbiology Faculty of Medicine Assiut University.
- Dibisceglie ,A.M. (2003): Hepatits C and HCC. Hepathology; 26: 345
- Dlugonska, M. (2000): Immunity in *Toxoplasma gondii* infection. J. parasitol., 54 (1): 53 – 65.
- Domingue, Z.A.; Bruguera ,M.and Vidal, J. (2001): Community based seroepidemiological survey of HCVinfection in Catalonia, Spain, J Med Viral.,65:688-693.
- Druwe, P.M.; Michelsen, .P. and Raman, A.M. (1994): Hepatits C and nephrology. Nephrol. Dial. Transplant, 9:230-237.
- Dubey, J.P. (1996): Infectivity and pathogenicity of *Toxoplasma gondii* oocysts for cats. J. Parasitol., 82: 957-961.
- Dubey, J.P. (1998): Toxoplasmosis. In Topley and Wilson's Microbiology and Microbial Infections, edited by Cox, F.E.G.; Kreier, J.P and Wakelin, D.; Oxford University Press, Inc.; New York.,5(9): 303 – 318
- Dubey, J.P. (2002): Tachyzoite-induced life cycle of *Toxoplasma gondii* in cats. J. parasitol., 88 (4): 713-717.
- Dubey, J.P. (2005): Toxoplasmosis. in Topley and Wilson's: Microbiology and microbial infections,edited by Cox,F.E.G.; Wakelin, Gillespie, S.H and Despommie, D.D) ., Oxford University Press, Inc.; New York. 6(10): 422-442.

- Dubey, J.P.; Kotula, A.W.; Sharar, A.; Andrews, C.D. and Lindsay, D.S. (1990): Effect of high temperature on infectivity of *Toxoplasma gondii* tissue cysts in pork. J. Parasitol., 76 : 201 – 204.
- Dunn, D.; Wallon, M.; Peyron. F.; Petersen, E.; Pechham, C. and Gilbert, R. (1999): Mother-to-child transmission of toxoplasmosis: risk estimates for clinical counseling. Lancet, 353: 1829-1833.
- Dussol,B.;Berthezene, P.and Brunet. (1995): Hepatitis C virus infection among chronic dialysis patients in the Saone of France: A collaborative study. Am. J. Kidney. Dis., 25:399-404.
- El –Gaafary, M.M.; Rekacewicz, C.and Michelsen, P.P.(2005): Surveillance of acute hepatitis C in Cairo, Egypt. J .Med .Virol,76:520–5.
- El.Sayed,M.N.;Gomatos,P.J.andRodier,G.R.(1996):Seroprevalence Survey of Egyptian Worker for hepatitis B virus, HCV, HIV, Treponema Pallidum infection. Am. J. Tropical Medicine and Hygiene.,55(2): 179-184.
- EL-Fakahany, A.F.; Abdel-Maboud, A.I., EL-Garhy, M.F.and Eraky, M.A. (2002): Comparative study between ELISA IgG, IgM and PCR in diagnosing and studying toxoplasmosis in Qalyobia Governorate, Egypt. J. Egypt Soc. Parasitol. , 32 (2) : 475 – 486.
- El-Gohary, A.; Hassan, A.and Nomman ,Z.(1995): High prevalence of hepatitis C virus among urban and rural population group in Egypt. Acta. Trop, 59: 155-161.
- El-Raziky,M.S.;El-Hawary,M.andEl-Koofy,N.(2007):Hepatitis C virus infection in Egyptian children :single center experience.J Viral Hepatol.,11(5):471-477.

- El-Ridi, A.M.S.; Nada, S.M.M.; Abdel Fattah, M.M.; Habeeb, Y.S.M. and Awad, M.B. (1991): Prevalence of congenital and acquired uveitis as evidenced by two serological methods. J. Egypt. Soc. Parasitol., 21 (2): 357 – 371.
- El-Sahly, A.M.; Nour El-Din, M.A.B. and Abdel-Rahmen .M.(1991): prevalence of antibodies to hepatitis C virus in Egyptian patients with schistosomiasis. Egypt. Soc. of Hepatology., 9:17-24.
- El-Shennawy, S.F.A.; EL-Gozamy, B.M.R.M.; Mahmoud, M.S.E. and Thabet, H.S. (2000): Characterization of tachyzoites and bradyzoite antigen of *Toxoplasma gondii* recognized by immunoglobulins G, M and A in experimental model of reactivated toxoplasmosis. J. Egypt. Soc. Parasitol., 31 (2): 349 – 34.
- Engvall, E. and Perlmann, P. (1971): Enzyme linked immunosorbent assay (ELISA). Quantitative assay of immunoglobulin G. J. Immunochemistry., 8 : 871 – 874.
- Esche, C.; Stellato, C. and Beck, L.A. (2005): Chemokines: Key players in innate and adaptive immunity. J. Invest. Dermatol. Oct; 125 (4): 615 – 28.
- Esteban, J.I.; Cordoba, J. and Saulead, S. (1998): Hepatitis C: Molecular Biology, pathogenesis, epidemiology, clinical feature and prevention in: Boyer JL, Okner RK (eds). Progress in liver disease. Philadelphia, W. B. Saunders., 10: 253-282.
- Etheredge, G.D. and Frenkel, J.K. (1995): Human *Toxoplasma* infection in Kuna and Embera children in the Bayano and San Blas. Eastern Panama. Am. J. Tropical Medicine and Hygiene., 35(5): 448-457.

- Everson, W.V.; Ware, M.W.; Dubey, J.P. and Lindquist, H.D. (2002): Isolation of purified oocyst walls and sporocysts from *Toxoplasma gondii*. J. Eukaryot. Microbiol., 49(4): 344-349.
- Eyster ,M.W.; Alter, H.J.and Aldort, L.M. (1991): Heterosexual Co-transmission of hepatitis C virus and human HIV. Ann. Int. Med., 115:764-768.
- Fahovich ,G.; Giustina, G.and Degos ,L. (1997): Morbidity and mortality in compensated cirrhosis type C. Aretrospective follow up study of 384 Pat. Gastroenterology.,35 (12) 112: 463.
- Ferguson, D.J.; Parmley, S.F. and Tomavo, S. (2002): Evidence for nuclear localization of two stage specific isoenzymes in *Toxoplasma gondii* correlates with active parasite replication. Int. J. Parasitol. ,32(4): 1399-1410.
- Ferreira, M.S. and Borges, A.S. (2002): Some aspects of protozoan infections in immunocompromised patients- a review. Mem. Inst. Oswaldo. Cruz., 97 (4): 443-457.
- Ferrer, A.; Hoebeke, J. and Bout, D. (1999): Purification and characterization of two amylases from *Toxoplasma gondii*. Exp. Parasitol., 9 : 64 – 72.
- Figueroa, M.S.; Bou, G.; Marti-Belda, P.; Lpez-Velez R. and Guerreo, A. (2000): Diagnostic value of Polymerase Chain Reaction in blood and aqueous humor in immunocompetent patients with ocular toxoplasmosis. Int. J. Parasitol., 20 (6): 614 – 619.
- Foster, B.G. and McCulloch, W.F. (1968): Studies of active and passive immunity in animals inoculated with *Toxoplasma gondii*. Canad. J. Microbiol. 14 : 103- 110. Quoted from Mahgoub, A.M.A. (2001): Characterization of *Toxoplasma gondii* antigen

and its use is serodiagnosis of toxoplasmosis. M.Sc. Thesis in Parasitology Faculty of Medicine Cairo University.

- Foulon, W.; Naessens, A. and Ho-Yen, D. (2000): Prevention of congenital toxoplasmosis. J. Perinat. Med., 28 (5): 337-345.
- Foulon, W.; Villena, I.; Stray-Pedersen, B.; Decoster, A. and Lappalainen, M. (1999): Treatment of toxoplasmosis during pregnancy. Am J. Obstet. Gynecol. 180 : 410 – 415.
- Frenkel, J.K. (1981): Congenital toxoplasmosis: prevention or palliation. AM. J. Obstet. Gynecol., 141: 359 – 361.
- Frenkel, J.K. (1982): Common questions on toxoplasmosis, veterinary, medical and public health considerations. VMSAC., 77 : 1188 – 1196.
- Frenkel, J.K. (1985): Toxoplasmosis. Pediatric Clinicals of North America, 32 : 917 – 932.
- Frenkel, J.K.; Hassanein, K.M.; Hassanein, R.; Brown, E.; Thulliez, P. and Quintero-Nunez, R. (1995): Transmission of *Toxoplasma gondii* in panama city, a five-year prospective cohort study of children, cats, rodents, birds, and soil. Am. J. Trop. Med. Hyg., 53 (5): 458-468.
- Garcia, J.L.; Navarro, I.T.; Ogawa, L.; de Oliveira, R.C. and Kobilka, E. (1999): Seroprevalence, epidemiology and ocular evaluation of human toxoplasmosis in the rural zone Jauguapita Brazil. Rev. Panam. Salud Publica., 6 (3): 157-163.
- Garcia, L.S. (2007): Tissue protozoa: *Toxoplasma gondii*. In: Diagnostic Medical parasitology. American Society for Microbiology., 4: 130-141.

- Garcia, L.S. and Bruckner, D.A. (1997): Tissue protozoa. In Diagnostic medical parasitology. 3rd edition, ASM Press, Washington, D.C.
- Gazzinelli, S.; Hieny, T.A.; Wynn, S.F.; Wolf, S.F. and Sher, A. (1993): Interleukin-12 is required for the T-lymphocyte – independent induction of interferon by an intracellular parasite and induces resistance in T-cell-deficient hosts. Proc. Natl. Acad. Sci. USA; 90 : 6115 – 6119.
- Gillbert, R.E.; Cool, A. and Dunn, D. (2000): Sources of *Toxoplasma* infection in pregnant women: a European multicentre case control study. Br. Med. J. 312, 142-147.
- Gola, J.; Czuba, B.; Mazurek, U.; Kaminski, K. and Wilczok, T. (2000): Usefulness of quantitative assessment of *Toxoplasma gondii* genome using PCR in amniotic fluid, maternal and newborn blood in selected complications in pregnancy. Ginekol-Pol., 71(9):954-958.
- Goldman, N. (1957): Staining of *Toxoplasma gondii* with fluorescein labeled antibody. II. A new serological test for antibodies to *Toxoplasma* based upon inhibition of specific staining. J. Exp. Med., 105 : 549. Quoted from Mahgoub, A.M.A. (2001): Characterization of *Toxoplasma gondii* antigen and its use in serodiagnosis of toxoplasmosis. M.Sc. Thesis in Parasitology Faculty of Medicine Cairo University.
- Graille, M.; Stura, E.A.; Bossus, M.; Muller, B.H.; Letourneur, O.; Battail-Poirot, N.; Sibai, G.; Gauthier, M.; Rolland, D.; Le Du, M.H. and Ducancel, F. (2005) Crystal structure of the complex between the monomeric form of *Toxoplasma gondii* surface

- antigen 1 (SAG1) and a monoclonal antibody that mimics the human immune response. J. Mol. Biol. Nov25; 354 (2): 447-458.
- Gretch ,D.; Dela Rosa ,D. Corey ,L. and Carithers ,R. (2006): Assessment of hepatitis C Viremia using molecular amplification technologies. Viral Hepatitis Rev., 2: 85-96.
 - Gross, U.; Keksel., O.; and Darde, M.L. (1997): Value of detecting immunoglobulin E antibodies for the serological diagnosis of *Toxoplasma gondii* infection. Clin. Diag. Lab. Immunol., 4 (3): 247 – 251.
 - Guay, J.M.; Dubois, D.; Morency, M.J.; Gagnon, S.; Mercier, J. and Levesque, R.C. (1993): Detection of the pathogenic parasite *Toxoplasma gondii* by specific amplification of ribosomal sequences using comultiplex polymerase chain reaction. J. Clin. Microbiol., 31 (2): 203 – 207.
 - Haber ,M.M.; West ,A.B.; Haber, A.D. and Reuben, A. (1996): Relationship of aminotransferases to liver histological status in chronic hepatitis C. Am J Gastroenterol., 90: 1250-1257.
 - Habib,M.;Mohamed,M.K.and Abdel-Aziz,F.(2004) Hepatitis C virus infection in a community in the Nile Delta:risk factors for seropositivity.Hepatology.33(6):248-253.
 - Hagan, H.; Snyder, N.and Haugh ,E. (2002): Case reporting of acute hepatitis B and C among injection drug users. J. Urban Health., 97:579-585.
 - Hammouda, N.A.; Gl-Gebaly, W.M. and Sadaka, H.M. (1993): Seroprevalence of toxoplasmosis in complicated pregnancies. J. Egypt. Soc. Parasitol., 26 (1): 177-189.

- Hashem ,El-Serag.; Bhupinder, J.; Peter ,R. and Linda, R. (2008): Association between hepatitis C infection and other infectious diseases: a case for targeted screening. *The American Journal of Gastroenterology.*, 25(3): 167-174.
- Hauri, A.M.; Armstrong, G.L. and Hutin, Y.J. (2004): The global burden of disease attributed to contaminated injection given in health care setting. *Int. J. STD. AIDS*, 15:7-16.
- Hedman, K.; Lappalainen, M.; Seppala, I. and Makela, O. (1989): Recent primary *Toxoplasma* infection indicated by a low avidity of specific IgG. *J. Infect. Dis.*, 159 : 736 – 740.
- Hewitt, J.; Coates, A R. M.; Mitchison, D.A. and Ivanyi, J. (1982): The use of murine monoclonal antibodies without purification of antigen in the serodiagnosis of tuberculosis, *J. Immunol. Methods.*, 55: 205- 211.
- Hill, D.E. and Dubey; J.P. (2002): *Toxoplasma gondii*: transmission, diagnosis and prevention. *Clin. Microbiol. Infect.* , 8(10): 634-640.
- Hoballah, A. and Sabry, H. (1996): Hepatitis C virus infection. *Egyptian Military Medical Journal.*,41(2): 14-16.
- Holliman, R.E. (1996): *Toxoplasma gondii*. In *Manson's Tropical Diseases*, 20th ed., edited by Cook, G.C.; W.B. Saunders Company Ltd, P. 1246 – 1254.
- Howard ,J. and Worman,M.D. (2002): The hepatitis C source book, disease of the liver, national institutes of health consensus statement on management of hepatitis C, A consensus statement on the management of hepatitis C prepared by a non advocate, non-federal panel of experts, from the U.S. centers for disease control

- and prevention. The koop institute at Dartmouth college, hepatitis foundation international.,P.688-753.
- Hughes, H.P.A. and Van Knapen, F. (1982): Characterization of secretory antigen from *Toxoplasma gondii*. And its role circulating antigen production. Int. J. parasitol.,12: 433-437.
 - Huskinson, J.; Stepick-Biek, P. and Remington, J.S. (1989): Detection of antigens in urine during acute toxoplasmosis. J.Clin. Microbiol., 27 (5): 1099-1101.
 - Hussein, A.H.; Ali, A.E.; Saleh, M.H.; Nagaty, I.M. and Rezk, A.Y. (2001): Prevalence of *Toxoplasma* infection in Qualyobia Governorate, Egypt. J. Egypt. Soc. Parasitol., 31 (2): 355 – 363.
 - Hutchison, W.M and Dunachie, J.F. (1970): Coccidian-like nature of *Toxoplasma gondii*. Br Med. J.I. 142-146.
 - Hyder,S.N.;Hussain,W.;Aslam,M.and Maqbool,S.(2005):Seroprevalence of anti- HCV in asymptomatic children.Pak Ped J .25(4):89-93.
 - Irshad,M.and Peter,S.(2006):Spectrum of viral hepatitis in thalasemic children receiving multiple blood transfusion .India J Gastroenterol.21(3):183-184.
 - Isaac Renton, J.; Bowie, W.R.; King, A.; Irwin, G.S.; Ong, C.S.; Fung, C.P.; Shokeir, M.O. and Dubey, J.P. (1998): Detection of *Toxoplasma gondii* oocysts in drinking water. Appl. Environ. Microbiol., 64 (6): 2278 – 2340.
 - Isabelle Dimier Poisson, Fleur Aline, Marie-Noelle Mevelec, Celine Beauvillain, Dominique Buzoni-Gatel, and Daniel Bout (2003): Protective mucosal Th2 immune response against

- Toxoplasma gondii* by murine mesenteric lymph node dendritic Cells. Infect. and Immunity., 71(9): 5254 – 5265.
- Jacobs, L. (1953): The biology of *Toxoplasma*. Am. J. Trop. Med. Hyg., 2 : 365 – 389. Quoted from Mahgoub, A.M.A. (2001): Characterization of *Toxoplasma gondii* antigen and its use in serodiagnosis of toxoplasmosis. M.Sc. Thesis in Parasitology Faculty of Medicine Cairo University.
 - Jacobs, L. and Lunde, M.N. (1967): Differences in *Toxoplasma* dye test and haemagglutination antibodies shown by antigen fractionation. Am.J. Trop. Med. Hyg., 16 : 26 – 30. Quoted from Mahgoub, A.M.A. (2001): Characterization of *Toxoplasma gondii* antigen and its use in serodiagnosis of toxoplasmosis. M.Sc. Thesis in Parasitology Faculty of Medicine Cairo University
 - Jacobson, J.M.; Hafner, R.; Remington, J.; Farthing, C.; Holden-Wiltse, J.; Bosler, E.M.; Harris, C.; Jayaweera, D.T.; Roque, C. and Luft, B.J. (2001): Dose- escalation, phase I/II study of azithromycin and pyrimethamine for the treatment of toxoplasmic encephalitis in AIDS., 15 (5): 583-589.
 - Jaeckel, E.; cornberg, M. and Wedemeyer, H. (2005): Treatment of acute hepatitis C with interferon Alpha 2b. New England Journal of Medicine., 45(3): 1452-1457.
 - James, G.J.; Sintghenko, V.G.; Dickeson, D.J. and Gilbert, G.L. (1996): Comparison of cell culture, mouse inoculation and PCR for detection of *Toxoplasma gondii*. Effects of storage condition on sensitivity. J. Clin. Microbiol., 34 (6): 1572 – 1575.

- Jara,P.;Resti,M.and Hierro,L.(2003):Chronic hepatitis C virus infection in childhood:Clinical patterns and evolution in 224 white cases .Clinical infectious diseases. 36(5):275-280.
- Jeannel, D.; Costagliola, D.; Niel, G.; Hubert, B. and Danis, M. (1990): What is known about the prevention of congenital toxoplasmosis. Lancet., 336: 359 – 361.
- Johnson, A.M. and Dubey, J. P. (1994): The diagnosis of toxoplasmosis. and Food borne diseases,. J. Clin. Microbiol., 4 (6): 572 – 595.
- Johnson, A.M. and Illana, S. (1991): Cloning of *Toxoplasma gondii* gene fragments encoding diagnostic antigens.J. Clin. Microbiol., 99 : 127- 132.
- Johnson, A.M.; Haynes, W.D.; Leppard, P. J.; DeDonald, P.J. and Neoh, S.H. (1983): Ultrastructural and biochemical studies on the immunohistochemistry of *Toxoplasma gondii* antigens using monoclonal antibodies. Histochemistry, 77 : 209.
- Johnson, R.N.; Sargon, R.; Woollard, G. and Davidson, J. (2005): An evaluation of the abbott IMx sirolimus assay in relation to a high-performance liquid chromatography-ultraviolet method. Ann. Clin. Biochem: 42(Pt5). 394-397.
- Joiner, K.A. Beckers, C.J.M.; Bermudes, D.; Ossoria, P.N.; Schwab, J.C. and Dubremetz, J.F. (1994): Structure and function of the parasitophorous vacuole membrane surrounding *Toxoplasma gondii*. Ann. N.Y. Acad. Sci., 730 – 1- 5.
- Jones, J.L.; Lopez, A.; Wilson, M.; Schulkin, J. and Gibbs, R. (2001): Congenital toxoplasmosis: a review. Obstet. Gynecol. Surv., 56(5): 296-305.

- Joss, A.W.L.; Skinner, L.J.; Moir, I.L.; Chatterton, J.M.W.; Williams, H. and Ho-Yen, D.O. (1989): Biotin-labelled antigen screening test for *Toxoplasma* IgM antibody. J. Clin. Pathol., 42 : 206 – 209.
- Kabil , S.M.;El-Zayadi, A.and Kamel, R. (1990): The true situation of liver disease in Egypt with the beginning of declension of virus B hepatits J. Trop. Med, 1(1): 1-10.
- Kabil, S.M.; Abdel-Hamid, M.and Constantine, N. (1993): HCV RNA in sera and saliva of Egyption patients. J. Trop.med, 2(4):1-2.
- Kabil, S.M.; Salwa ,M.; Youssef .and Hussien, A.T. (1995): About HCV in Egypt, J. H. i. D, 3(4): 1-3.
- Kabil, S.M.; Salwa, M.; Youssef and Atta, M.M.E.. (1996):High prevalence of hepatits C virus RNA in saliva of patients with chronic HCV infection.J.H.G.I.D,4(2):1-6.
- Kamel, M.A.; Ghaffar, Y.A.and Wasef ,M.A.. (1992): High HCV prevelance in Egyptian blood donors. Lancet, 15 (340): 427.
- Kamitsaukasa, H. Harda, H. and Yakura ,M. (1989): Intrafamilia transmission of HCV. Lancet, 2(8669):987.
- Kasper, L.H. and Boothroyd, J.C. (1993): *Toxoplasma* gondii and toxoplasmosis. In immunology and Molecular biology of Parasitic infections, 3rd ed., edited by Warren, K.S., Blackwell scientific Publications, P. 269 – 301.
- Kasper, L.H. and Buzoni-Gatel, D. (1998): Some opportunistic parastic ,infections in AIDS; Candidiasis, Pneumocytosis, Cryptosporidiosis, Toxoplasmosis. Parasitology Tody, 14 (14): 150 – 156

- Katz, M.; Despommier, D.D. and Gwadz, R.W. (1989): *Toxoplasma gondii*. In parasitic Diseases, 2nd edition, Springer-Verlag, New York. P. 146-152.
- Kelen, A.E.; Ayllon-Leindl, L. and Labzoffsky, N.A. (1962): Immunofluorescent antibody method in sero-diagnosis of toxoplasmosis. Canad. J. Microbiol., 8: 545 – 554. Quoted from Mahgoub; A.M.A. (2001): Characterization of *Toxoplasma gondii* antigen and its use in serodiagnosis of toxoplasmosis. M.Sc. Thesis in Parasitology Faculty of Medicine Cairo University.
- Khalifa, A.S.; Mitchell, B.S. and Watt, D.M. (1993): Prevalence of hepatitis C viral antibody in transfused and non transfused Egyptian children. Am. J. Trop. Med. Hyg, 49 (3): 316-321.
- Khalil, H.M.; Azab, M.E.; Safar, E.; EL-Shennawy, S.F. and EL-Din, M.H.N. (1990): Primary and secondary immune responses in experimental *Toxoplasma* infection. N. Egypt. J. Med., 4 (2) : 17 – 19.
- Kiyosawa, K.; Sodeyama, T. and Tanaka, E. (1994): Hepatitis C virus infection in health care workers. Viral hepatitis and liver disease Tokyo: Springer-Verlag, 479-482.
- Klaren, V.N. and Kijlstra, A. (2002): Toxoplasmosis, an overview with emphasis on ocular involvement. Ocul Immunol Inflamm., 10 (1): 1-26.
- Kleinman, S.; Alter, H.; Bush, M.; Holland, P.; Tegtmeir, G.; Nelles, M. and Lee, S. (1992): Increased detection of hepatitis C virus infected blood-donors by a multiple- antigen HCV enzyme immunoassay. Transfusion; 32: 805-813.

- Konishi, E. (2006): Naturally occurring immunoglobulin M antibodies to *Toxoplasma gondii* in Japanese populations. *Parasitology*, 102 : 157 – 162.
- Kopecky, D.; Hayde, M.; Prusa, A.R. and Adlassnig, K.P. (2001): Knowledge- based interpretation of toxoplasmosis serology test results including fuzzy temporal concepts. *J. Infect. Dis.*, 10: 484-488.
- Koskiniemi, M.; Lappalainen, M. and Hedman, K. (1989): Toxoplasmosis needs evaluation. *AJDC*, 143 : 727 – 72.
- Koskiniemi, M.; Lappalainen, M.; Koskela, P.; Hedman, K.; Ammala, P.; Hiilesmaa, V. and Teramo, K. (1992): The program for antenatal screening of toxoplasmosis in Finland: a prospective cohort study. *Scand. J. Infect. Dis. Suppl.*, 84: 70-74.
- Kovacs, J.A. and Masur, H. (2002): Prophylaxis against opportunistic infections in patients with human Immunodeficiency virus infection. *N. Engl. J. Med.* 342: 1416-1429.
- Krahenbuhl, J.L. and Remington, J.S. (1982): "Immunology of *Toxoplasma* and Toxoplasmosis". In "Immunology of parasitic infections". (eds. Cohen S. and Warren K.S.), Blackwell scientific publications
- Kwantes, W. (1987): Toxoplasmosis. In *Oxford textbook of medicine*, edited by Weatherall, D.J.; Ledingham J.G.G. and Warrell, D.A., 2nd ed., Oxford University press, Oxford, 5 : 506 – 509
- Laura Del Rio, Barbara, A.; Butcher, Soumaya Bennouna, Sara Hieny, Alan Sher and Eric, Denkers. (2004): *Toxoplasma gondii* triggers myeloid differentiation factor 88-dependent IL-12 and

- chemokine ligand 2 (monocyte chemoattractant protein 1) responses using distinct parasite molecules and host receptors. *J. Immunol.*, 172: 6957 – 6960.
- Law, M.G.; Dore, G.J. and Bath, N. (2001): Modelling hepatitis C virus incidence, prevalence and long term sequelae in Australia. *Int. J. Epidemiol.*, 32:717-724.
 - Lee, Y.F.; Chen, S.J.; Chung, Y.M.; Liu, J.H. and Wong, W.W. (2000): Diffuse toxoplasmic retinochoroiditis as the initial manifestation of acquired immunodeficiency syndrome. *J. Formos. Med. Associac.*, 99 (3): 219 – 223.
 - Leightly, J.C. (1990): Strategies for control of toxoplasmosis. *J. Egypt. Soc. Parasitol.*, 190 : 281 – 286.
 - Leriche, M.A. and Dubremetz, J.F. (1991): Characterization of the protein contents of rhoptries and dense granules of *Toxoplasma gondii* tachyzoites by subcellular fractionation and monoclonal antibodies. *Mol Biochem. Parasitol.*, 45 : 24 – 260
 - Liesenfeld, O. and Remington, J.S. (2000): "Toxoplasmosis" In Cecil textbook of medicine (eds. L. Goldman, J.C. Bennett), W.B. Saunders company; p: 963-967.
 - Liesenfeld, O.; Montoya, J.G.; Kinney, S.; Press, C. and Remington, J.S. (2001): Effect of testing for IgG avidity in the diagnosis of *Toxoplasma gondii* infection in pregnant women: experience in a US reference laboratory. *J. Infect. Dis.*, 183: 1248-1253.
 - Lin, T. M Chin-See, M.W.; Halbert, S.P. and Joseph, J.M. (1986): An enzyme immunoassay for immunoglobulin M-antibodies to

Toxoplasma gondii which is not affected by rheumatoid factor or immunoglobulin G antibodies. J. Clin. Microbiol., 23 : 77 – 82.

- Luft, R.L.; Naot, Y.; Araujo, F.G.; Stinson, E.B. and Remington, J.S. (1983): Primary and reactivated *Toxoplasma* infection in patients with cardiac transplants clinical spectrum and problems in diagnosis in a defined population. Ann: Intern. Med., 99 (1): 27-31.
- Major ,M.E. and Feinstone, S.M. (1997): The molecular virology of hepatitis C. Hepatology,25:1527-1537.
- Manson-Bahr, P.E.C. and Bell, D.R. (2005): Toxoplasmosis. In: Manson's tropical diseases, 19th ed., Bailliere Tindall, London, P. 207-212.
- Many, A. and Koren, G. (2006): Toxoplasmosis during pregnancy. Can. Fam. Physician., 52: 29:32.
- Marazuela ,M.; Garcia ,B.L.and Gonzalez. (1996): Thyroid autoimmune disorder in patients with chronic Viral hepatitis during interferon therapy. Clin Endo., 44: 635.
- Mariuz, P.; Bosler, E.M. and Luft, B.J. (1994): Toxoplasmosis in individuals with AIDS. Infect. Dis. Clin. North Am., 8(2): 365-381.
- Marshall, A.J. and Denkers, E.Y. (1998): *Toxoplasma gondii* triggers granulocyte-dependent cytokine-mediated lethal shock in D-galactosamine sensitized mice. Infect. Immun., 66 : 1325 – 1333
- Matsumori ,A. (2000): Hepatitis C virus and cardiomyopathy. Ann Intern Med; 25: 249.
- McCabe, R.E. (2001): Anti-*Toxoplasma* chemotherapy. In: Joynson, D.H.M. and wrephilt. T.G. (eds). Toxoplasmosis. A

- comprehensive clinical guide. Cambridge: Cambridge University press. 319-359.
- Mchutchison ,J.G.; Gordon ,S.C.and Schiff ,E.R. (1998): Interferon alfa-2b alone or in combination with ribavirin as initial treatment for chronic hepatitis C. Hepatitis Interventional therapy Group. N Engl J Med; 339: 1485-1492.
 - McLeod, R. and Remington, J.S. (1985): "Toxoplasmosis" In "Harrison's Principles of Internal Medicine" (eds. R.G. Petersdorf, R.D. Adams, E. Braunwald, K.J. Isselbacher, J.B. Martin, and J.D. Wilson,), 10th ed., McGraw-Hill International Book Company.
 - Medhat ,A.; Shehata, M.and Magder ,L.S. (2002): Hepatitis C in community in upper Egypt and risk factors for infection. Am. J. Trop.Med. Hyg, 66:633-638.
 - Meeka, B.; Goolb.; Gilisb, H. and Peeka, R. (2007): Dissecting the IgM antibody response during the acute and latent phase of toxoplasmosis. Diagnostic microbiology and infectious disease. 41: 131-137.
 - Mehta ,S.H.; Brancatift and Sulkowski.(2000): Prevalence of type2 Diabetes mellitus among personse with HCVin united state. N Engl J Med; 133: 592.
 - Mele, A.; Paterson, P.J.; Prentice, H.G.; Leoni, P. and Kibbler, C.C. (2002): Toxoplasmosis in bone marrow transplantation: a report of two cases and systematic review of the literature. Bone Marrow Transpl. 29, 691-698.
 - Mohammed, M.k.; Abdel-Hamid, M.and Mikhail, N.N. (2009): Intrafamilial transmission of hepatitis C in Egypt. Hepatology, 42:683-687.

- Mohan,P.;Colvin,C.and Glymph,C.(2009):Clinical spectrum and histopathological features of chronic hepatitis C infection in children .30(5):168-174.
- Montoya, J.C. (2006): Laboratory diagnosis of *Toxoplasma gondii* infection and toxoplasmosis. J. infect. Dis. ,8(3): 73-82.
- Montoya, J.G and Remington, J.S. (2000): *Toxoplasma gondii*. In Mandell GL, Bennett JE, Dolin R (eds), Principles and practice of infectious disease, 5th edn. Churchill Livingstone: Philadelphia, PA; 2858-2888.
- Montoya, J.G. and Liesenfeld, O. (2004): Toxoplasmosis. Lancet.; 36 (94): 1965-76.
- Morishima ,C. and David, R.G. (2005): Virological tests for hepatitis C. Update on viral hepatitis sec. 4 page 127-133. American Association for the study of liver diseases.
- Moscatelli, G.; Altcheh, J.; Biancardi, M.; Lapena, A.; Ballering, G.and Freilij, H. (2006): Acute toxoplasmosis: clinical and laboratory data in eleven patients. Ann. Pediatr. (Barc), 65 (6): 551-555.
- Mousa,M.M.(2004):Astudy on toxoplasmosis and pregnancy.M.D.Thesis,Faculty of Medicine,Cairo University.
- Murray, J.M.; Law, M.G.and Gao ,Z. (2003): The impact of behavioural changes on the prevalence of human immunodeficiency virus and hepatitis C among injecting drug users. Int. J. Epidemiol, 32: 708-714.

- Nabias, R.; Ngouamizokou, A.; Migot-Nabias, F.; Mbou-Moutsimbi, R.A. and Soukate, L.J. (1998): Serological investigation of toxoplasmosis in patients of M.I.P. Center of Franceville (Gabon). Bull. Soc. Pathol. Exot., 91 (4): 314 – 420.
- Naot, Y. and Remington, J.S. (1980): An enzyme-linked immunosorbent assay for detection of IgM antibodies to *Toxoplasma gondii*: use for diagnosis of acute acquired toxoplasmosis. J. Infect. Dis., 142 : 757 – 766.
- Naot, Y.; Guptill, D.R. and Remington, J.S. (2004): Duration of IgM antibodies of *Toxoplasma gondii* after acute acquired toxoplasmosis. J. Infect. Dis., 145 (5): 770.
- National Institutes of Health. (1997):Panal statement: Management of hepatitis C Hepatology, Vol. 26, No. 3, suppl. 1:25-105.
- Nielsen, H.V.; Lauemoller, S.L.; Christiansen, L.; Buus, S.; Fomsgaard, A. and Petersen, E. (1999): Complete protection against lethal *Toxoplasma gondii* infection in mice immunized with a plasmid encoding the SAG1 gene. Infect. Immunol., 67 (12) 6358 – 6363.
- Nigro, G.; Piazze-J.; Paesano, R.; Mango-T.; Provivedi, S.; Capunao, Q. and Pollastrini, L. (1999): Low levels of natural killer cells in pregnant woman transmitting *Toxoplasma gondii*. Prenat. Diagn., 19 (5): 401 – 404.
- Noureldin, M.S.A. Shaltout, A.A.; Hamshary, E.M. and Ali, M.E. (1999): Opportunistic intestinal protozoal infections in immunocompromised children. J. Egypt. Soc. Parasitol., 229 (3): 951 – 961.

- Nowakowska, D.; Stray- Pederson, B.; Spiewak, E.; Sobala, W.; Malafiej, E. and Wilezynski, J. (2006): prevalence and estimated incidence of *Toxoplasma* infection among pregnant women in Poland, ClinMicrobiol Infec; 12 (9), P: 913-920.
- Obtata, K. (1996): Phylum Apicomplexa: Gregarines, Coccidia. In: Oberts. L.S., Janovy J.Jr. eds. Foundations Parasitology, 5th ed. Brown Publishers. P: 113-135.
- Ogasawara ,S.; Kage, M. and Kosal, K.I. (1993): HCVRNA in saliva and breast milk of hepatitis C cancer mother. Lancet, 341:561-563.
- Ohato, H.; Terzawa, S. and Sasaki, N. (2004): Transmission of hepatitis C virus from mothers to infants. N. Engl, J. Med., 330:747-750.
- Okayama ,A.; Stuver, S.O.and Tabor ,E. (2002): Incident hepatitis C virus infection in a community-based population in Japan, J. Hepatol 55:44-49.
- Parija, S. (2004): Textbook of Medical Parasitology. Protozoology and helminthology .Bull. Soc. Pathol. Exot., 172-181.
- Partanen, P.; Turunen, H.J.; Paasivuo, R.T.A. and Leinikki, P.O. (2005): Immunoblot analysis of *Toxoplasma gondii* antigens by human immunoglobulin G,M. and A Antibodies at different stages of infection.
- Pereira, B.J.; Milford ,E.L. and Kirkman ,R.L. (1992): Prevalence of hepatitis C virus RNA in organ donors positive for hepatitis C antibody and in the recipients of their organs. N. Engle. J. Med., 327:910-915.

- Perz, J.F. and Alter, M.J. (2006): The coming wave of HCV-related liver disease. *J. Hepatology*, 44:441-443.
- Perz, J.F.; Armstrong, G.L.; and Farrington, L.A. (2006): The concentration of hepatitis B virus and hepatitis C virus infections to cirrhosis and primary liver cancer world wide. *J. hepatol*, 45:529-538.
- Petersen, E. and Liesenfeld, O. (2007): Clinical Disease and Diagnostics. In *Toxoplasma gondii* The model Apicomplex: Perspectives and Methods *Bull. Soc. Pathol. Exot.*, (4) : 81-101.
- Peyron, F.; Wallon, M.; Liou, C. and Garner, P. (2000): Treatment for toxoplasmosis in pregnancy. *Cochrane-Database-Syst. Rev.*, (2): 1670 – 1684.
- Pfefferkor, E.R. (1990): Cell biology of *Toxoplasma gondii*. In *Modern Parasite Biology: Cellular, immunological and molecular aspects*, edited by Wyler, D.J., Freeman and Company, New York: 26-50.
- Philipperi, M. and Bastani, B. (2001): Kidney disease in patients with chronic Hepatitis C. *Gastroenterol*; 3: 79.
- Pinlaor, S.; Ieamviteevanich, K.; Pinlaor, P.; Maleewong, W. and Pipitgool, V. (2000): Seroprevalence of specific total IgG and IgM antibodies to *Toxoplasma gondii* in blood donors from Loei Province, Northeast Thailand. *Southeast Asian J. Trop. Publ. Hlth.*, 31:123-127.
- Pons, J.C.; Sigrand, C.; Grangeot-Keros, L.; Frydman, R. and Thulliez, P. (1995): Congenital toxoplasmosis: mother-to-fetus transmission of pre-pregnancy infection. *Press Med.*, 24: 179-182.

- Poynard ,T.; Yuen, M.F.and Ratziu ,V. (2003): Viral hepatitis C. Lancet, 362:2095-100.
- Poynard, T.; Deuffic, S. and Valleron, A.J. (1999): Correlation between hepatitis C virus prevalence and hepatocellular carcinoma mortality in Europe J. Viral hepatitis, 6: 411-443.
- Prati ,D.; Capelli, C.and Silvani ,C.. (1997): The incidence and risk factors of community acquired hepatitis C in a cohort of Italian blood donors. Hepatology; 25:702-704.
- Punda Polic, V.; Tonkic, M. and Capkun, V. (2000): prevalence of antibodies to *Toxoplasma gondii* in the female population of the country of split dalmatia in Croatia, Eur J Epidemiol, P: 875-881.
- Puro, V.; Petrosillo, N. and Ippolito, G. (1995): Risk of hepatitis C seroconversion after occupational exposure in health care workers: Italian study group on occupational risk of HIV and other Blood born infections. Am. J. Infect. Control, 23:273-277.
- Quer ,J. and Esteban,J.I. (1997): Epidemiology and prevention of hepatitis C virus infection hepatitis C 1997: Essay, Diagnosis and Therapy. Abbott Diagnostics Division, 26-38.
- Quinny;El-Gassani and Divizia, M. (1995): Seroprevalence of HCVinfection in Alexandria, Egypt. Zentral, Blatt für, Bacteriologie, 283
- Raizman, R.E. and Neva, F.A. (1975): Detection of circulating antigen in acute experimental infections with *Toxoplasma gondii*. J. Infect. Dis., 132 – 44.
- Remington, B.L and Desmonts, C. (1983): Toxoplasmosis infectious diseases of the fetus and newborn infant. ,13:223-225

- Remington, J.S. and Cavanaugh, E.N. (1965): Isolation of the encysted form of *Toxoplasma gondii* from human skeletal muscle and brain. New Engl. J. Med., 273 : 1308. Quoted from Mahgoub, A.M.A. (2001): Characterization of *Toxoplasma gondii* antigen and its use in serodiagnosis of toxoplasmosis. M.Sc. Thesis in Parasitology Faculty of Medicine Cairo University.
- Remington, J.S. and Desmonts, G. (2005): Toxoplasmosis infectious diseases of the fetus and newborn infant. New Engl. J. Med., 27 : 134-150.
- Remington, J.S. and McLeod, R. (1998): Toxoplasmosis. In Gorbach, S.I.; Bartlett, J.G. and Blacklow, N.R. (ed.) "Infectious diseases" 2nd ed., Pub. W.B. Saunders Co., Philadelphia: 1620-1640.
- Remington, J.S.; Araujo, F.G. and Desmonts, G. (2003): Recognition of different *Toxoplasma* antigens by IgM and IgG antibodies in mothers and their congenitally infected new-born. J. Infect. Dis., 15 : 1020 – 1024.
- Remington, J.S.; Thulliez P. and Montoya, J.G. (2004): Recent developments for diagnosis of toxoplasmosis. J. Clin. Microbiol. 42(3): 941-945.
- Ridel, R.R.; Auriault, C.; Darcy, F.; Pierce, R.J. and Capron, A. (1988): Protective role of IgE in immuno-compromised rat toxoplasmosis. J. Immunol., 141: 978 – 983.
- Ritter, S.E. and Vermund, S.H. (1985): Congenital toxoplasmosis. J. Obstet. Gynecol. Neonatal. Nurs., 14 : 435 – 439.
- Roberts, E.A. and Yeung, L. (2002): Maternal infant transmission of hepatitis C virus infection. Hepatology, 636:1065-1135.

- Rodriguez, J.C.; Martinez, M.M.; Martinez, A.r. and Royo, G. (1997): Evaluation of different techniques in the diagnosis of *Toxoplasma* encephalitis. J. Med. Microbiol., 46 : 597 – 601.
- Romand, S.; Wallon, M.; Franck, J. Thulliez, P. Peyron, F. and Dumon, H. (2001): Prenatal Diagnosis using polymerase Chain Reaction on amniotic fluid for congenital toxoplasmosis . Obstet. Gynecol., 97 (2): 296 – 300.
- Rorman, E.; Zamir, C.S.; Rilkis, I. and Ben-David, H. (2006): Congenital toxoplasmosis: prenatal aspects of *Toxoplasma gondii* infection. Report toxicol 21:458-72.
- Roth ,D.; Zucker ,K.and Cirocco, R. (1996): A prospective study of hepatitis C virus infection in renal allograft recipient. Transplantation, 61(6): 886-9.
- Rumpo,C.(2005) Hepatitis C current approaches in pediatrics.Pediatr Transplant .9(20:662-665.
- Ruskin, J. and Remington, J.S. (1976): Toxoplasmosis in the compromised host. Ann. Inter. Med., 54 : 193 – 199.
- Saadeh, S.; Cammell ,G. and Carey, W.D.(2001): The role of liver biopsy in chronic hepatitis C. Hepatology.33:196-200.
- Sabin, A.B. and Olitsky, P.K. (1948): *Toxoplasma*, an obligate intracellular parasitism. Science, 85 : 336. Quoted from Mahgoub, A.M.A. (2001): Characterization of *Toxoplasma gondii* antigen and its use in serodiagnosis of toxoplasmosis. M.Sc. Thesis in Parasitology Faculty of Medicine Cairo University.
- Safar, E.H.; Azab, M.E. and Osman, Z.M. (1984): "Diagnosis of ocular and glandular toxoplasmosis using the Indian ink

- immunoreaction". Trans. Roy. Soc. Trop. Med. Hyg., 78: 169 – 172.
- Sanad,M.M.and Al-Ghabban,A.J.(2007):Serological survey on toxoplasmosis among slaughtered sheep and goat in Tabook,Saudi Arabia.J.Egypt.Soc.Parasitol.,37(10;329-340.
 - Sanchez, M.R.; Gordo, B.R.; Amador, A.E. and Berriol, A. (1994): Prevalence of toxoplasmosis in pregnant women in Sao Paulo. Rev. Inst. Med. Trop. Sao, Paub., 36 (5): 445 – 450.
 - Sarwat,M.A.;Ahmed,A.B.;Zamzami,O.M.andMorsy,T.A.(1993): *Toxoplasma gondii* in Saudi blood donors:Serologic study using 3 tests.J.Egypt.Soc.Parasitol.,23,3:751-757.
 - Schmidt, G.D. and Roberts, L.S. (1981): Foundation of parasitology (2nd ed.) Mosby C.V.Co., Trnto, 128 – 139.
 - Schmidt, G.D. and Roberts, L.S. (1987): Phylum Apicomplexa: Gregaines. Coccidea and related organisms. Foundation Parasitology, 4th ed. St. Lois: Mirro/Mosby collage. P: 123- 129.
 - Schoeman, M.N.; Liddle, C.and Bilous, M. (1996): Lack of correlation between biochemical and morphological activity and effects of immuno suppressive therapy on disease progressive. J Med; 20:56.
 - Schwartzman, J.D. (2001): Toxoplasmosis. In: Principles and Practice of Clinical parasitology, edited by Gillespie, S. and Pearson, R.D. John Wiley and Sond Ltd. England. ,3(2): 113-138.
 - Seff, L.B. and Hofnagle, J.H. (2008): National institutes of health consensus development conference: management of Hepatits C: hepatology; 36: 51-52.

- Sensini, A. (2006): *Toxoplasma gondii* infection in pregnancy: opportunities and pitfalls of serological diagnosis. Clin. Microbiol. Infect., 12 (6): 504-512.
- Shepard, C.W.; Finelli, L. and Fiore, A.E. (2005): Epidemiology of hepatitis B and hepatitis C virus infection in United States children. Pediatr Infect. Dis. J.; 24:755-760.
- Sher, A.L.; OsWald, O.; Hieny, S. and Gazzinelli, R.T. (1993): *Toxoplasma gondii* induces a T-independent INF- γ response in NK cells which requires both adherent accessory cells and TNF- α . J. Immunol., 150 : 3982 – 3989.
- Sherlock, S. and Dooley, J. (1997): Diseases of the liver and biliary system. Pediatr. Infect. Dis. J., 24:755-760.
- Sibley, L.D.; Adams, L.B. and Krahenbuhl, J.L. (1993): Macrophage interactions in toxoplasmosis. Res. Immunol., 144: 38-40.
- Simmonds, P. (1995): Variability of hepatitis C virus. Hepatology; 21: 570-582.
- Simpore, J.; Savadogo, A.; Ilboudo, D.; Nadambega, M.C.; Esposito, M.; Yara, J.; Pignatelli, S.; Pietra, V. and Musumeci, S. (2006): *Toxoplasma gondii*, HCV and HBV seroprevalence and co-infection among HIV-positive and-negative pregnant women in Burkina Faso. J Med Virol., 6(4): 730-732.
- Skidmore, S.J.; Collingham, K.E. and Drake, M. (1994): Sexual transmission of HCV. J. Med. Virol, 42:247-248.
- Skinner, L.J.; Chatterton, J.M.; Joss, A.W.; Moir, I.L. and Ho-Yen, D.O. (1989): The use of an IgM immunosorbent agglutination

- assay to diagnose congenital toxoplasmosis. J. Med. Microbiol. 28: 125 – 128.
- Spalding, S.M.; Amendoeira, M.R.; Klein, C.H. and Ribeiro, L.C. (2005): Serological screening and toxoplasmosis exposure factors among pregnant women in South of Brazil. Rev. Soc. Bras. Med. Trop., 38(2): 173-177.
 - Speer, C.A. and Dubey, J.P. (2005): Ultrastructural differentiation of *Toxoplasma gondii* schizonts (types B to E) and gamonts in the intestines of cats fed bradyzoites
 - Speer, C.A.; Clark, S. and Dubey, J.P. (1998): Ultrastructure of the oocysts, sporocysts, and sporozoites of *Toxoplasma gondii*. J. Parasitol., 84 (3): 505 – 512.
 - Speer, E.A.; Dubey, J.P.; Bixt, J.A. and Prokop, K. (1997): Time lapse video microscopy and ultrastructure of penetrating sporozoites, type 1 and 2 parasitophorous vacuoles, and the transformation of sporozoites to tachyzoites of the VEG strain of *Toxoplasma gondii*. J. Parasitol., 83: 565 – 574.
 - Splendore, A. (1990): Search about new protozo. J. Inf. Dis., 2: 462-467.
 - Stepick-Biek, P.; Thulliez, P.; Araujo, F.G. and Remington, J.S. (1990): IgA antibodies for diagnosis of acute congenital and acquired toxoplasmosis. J. Inf. Dis., 162 : 270 – 273.
 - Stroffolini, A.; Yamasaki, K. and Yamanishi, C. (1998): RNA carriers: a 10-year follow-up study. J. Gastroenterol Hepatol; 16: 536–40.

- Sulzer, A.J. and Hall, E.C. (1967): Indirect fluorescent antibody test for parasitic diseases, statistical study of variation in the indirect fluorescent antibody test for toxoplasmosis. Am. J. Epidemiol., 86: 401 -- 407. Quoted from Mahgoub, A.M.A. (2001): Characterization of *Toxoplasma* gondii antigen and its use in serodiagnosis of toxoplasmosis. M.Sc. Thesis in Parasitology Faculty of Medicine Cairo University.
- Sundar,P.;Mahadevan,A.;Jayshree,R.andShanker,S.(2007): *Toxoplasma* seroprevalence in healthy voluntary blood donors from urban Karnataka Indian J.Med.Res.,126:50-55.
- Suzuki, Y.(2002): Host resistance in the brain against *Toxoplasma* gondii. J. Infect. Dis. 1 : 58 – 65.
- Suzuki, Y.; Conley, F.K. and Remington, J.S. (1989): Importance of endogenous INF- γ for the prevention of toxoplasmic encephalitis in mice. J. Immunol.,143 : 2045 – 2050.
- Tenter, A.M.; Heckeroth, A.R. and Weiss, L.M. (2000): *Toxoplasma* gondii: from animals to humans. Int. J. parasitol., 30 (12 – 13): 1217 – 58.
- Thrope, L.E.; Quellet, L.T.and Hershow, R. (2002): Risk of hepatitis C virus infection among young adult injection drug users who share injection equipments. Am. J. Epidemiol, 155:645-653.
- Thulliez, P. (1992): Screening programme for congenital toxoplasmosis in france. Scand. J. Infect. Dis., 84: 43-45.
- Tong, M.J.; El Farra, N.S.and Reikes ,A.R. (1995): Clinical outcomes after transfusion associated hepatitis G. N Engl J Med; 332:1463.

- Tsai, H.C.; Lee, S.S.; Lin, H.H.; Lin, W.R.; Chen, Y.S.; Huaug, C.K.; Liu, Y.C. and Chen, E.R. (2002): Treatment of *Toxoplasma* brain abscess with clindamycin and sulfadiazine in an AIDS patient with concurrent atypical pneumocystis carinii pneumonia. J. Formos Med. Assoc., 101(9): 646-649.
- Van derpoel C.L, Cwrypers H.T, and Resinb HW (1994): Hepatitis C virus Six Years on. Lancet; 244: 1475-1479.
- Van Knapen, F. and Panggabean, S.O. (1977): Detection of circulating antigen during acute infections with *Toxoplasma gondii* by enzyme-linked immunosorbent assay. J. Clin. Microbiol., 6 : 545.
- Verhofstede, C.; Van Gelded, P. and Rabaey, M. (1988): The infection-stage – related IgG response to *Toxoplasma gondii* studied by Immunoblotting. Parasitol. Res., 8: 516 – 520.
- Wallon, M.; Gandilhon, F.; Peyron, F. and Mojon, M. (1994): Toxoplasmosis in pregnancy. Lancet, 344 : 451.
- Wassim, H.; Shaker, W. and Sherif ,A. (1996): Prevalance of hepatitis C virus in Egypt. Hepatology, 25(1): 167.
- Wastling, J.M.; Nicoll, S. and Buxton, D. (1993): Comparison of two gene amplification methods for the detection of *Toxoplasma gondii* in experimentally infected sheep. J. Med. Microbiol. 38: 360 – 365.
- Weiss, L. and Dubey, J. (2009): Toxoplasmosis: A history of clinical observations. Int. J. Parasitol., 39(8): 895-901.
- Wilber, J. C. (1993): Development and use of laboratory test for hepatitis C infection. A review. J. clin. Immunoassay., 12 : 16- 204.

- Wilson, M., Ware, D.A. and Juranek, D. (1990): "Serologic aspects of toxoplasmosis". JAVMA, 196 : 227 – 281.
- Woldemichael, T.; Fontanet, A.L.; Sahl, T.; Gilis, H.; messele, De-wit, T.F.; Yeneneh, H.; Coutinho, R.A. and Van Gool, T. (1998): Evaluation of the eiken latex test for anti-*Toxoplasma* antibodies and seroprevalence to *Toxoplasma* infection among factory workers in addis Ababa, Ethiopia. Trans. R. Soc. Trop. Med. Hyg., 92 (4): 401 – 403.
- Wong, A.; Tan, K. H.; Tee, C.S. and Yeo, G. S. (2007): seroprevalence of Cytomogalovirus, *Toxoplasma* and Parvovirus in pregnancy. Singapore Med. J., 41(4): 151-155.
- Wong, S.Y. and Remington, J.S. (1993): Biology of *Toxoplasma gondii*. Editorial review. AIDS, 7 (3): 299 – 316
- Wong, J.B. ; sansonno, D.; Gasparott. and Ferracciolo, G. (1997): "Efficacy and safety of peginterferon alfa- 2a and ribavirin in patients with chronic hepatitis C virus infection". Med Pregl., 59 (10) :415 – 460.
- World Health Organization (1993): The central of schistosomiasis. 2nd report of WHO expert committee. WHO Tech. Rep. Ser., 30:70-71.
- Wreghitt, T.G.; Gray, J.J. and Allain, J.P. (1992): Transmission of hepatitis C virus by organ transplantation in the united kingdom. J. Hepatol; 20:768-722.
- Yazar, S.; Esar, B. and Yay, M. (2006): Prevalence of anti-*Toxoplasma gondii* antibodies in Turkish blood donors. Ethiop. Med. J., 44:257-261.

- Yoo, M.; Pinar. ; Roberto ,Jiron. ; Luftig. and Mercedes, Somarria.(1994): "Gene therapy for hepatitis C virus", Journal of medical virology, volume (77), Issue (2), pages: 209-215 .
- Yosry,A.;Esmat,G.and El-Serafy,M. (2008):Outcome of liver transplantation for Egyptian patients with hepatitis C related cirrhosis.4095);481-485.
- Zamorano,C.G.;Contreras,M.;Villalobos,S.andSalinas,P.(1999):Seroprevalence of human toxoplasmosis in Osorno,Region X,Chile.Bol.Chil.Parasitol.,54:33-36.
- Zarger,H.A.;Wani,A.I.Masoodi,S.R.and Bashir,M.I.(1999):Seroprevalence of toxoplasmosis in women with recurrent abortion/neonatal deaths .Indian Journal of Parasitology,42(4):483-486.
- Zawawy , L.A.; El-Temsahi, M.M.; El-Azzouni, M.Z.; El-Gebaly W.M.; Sharaky,O.A. and Awadalla, H.N. (2000): Diagnosis of toxoplasmosis in immuno-suppressor mice using the PCR and the ELISA techniques. J. Egypt. Soc. Parasitol., 30 (1): P. 27 – 42.
- Zhang M, Sun XD, Mark SD, et al. (2005): Hepatitis C virus infection, Linxian and China. Emerg. Infect. D.S, 11:17-21.
- Zuber, P. and Jacquier, P. (1995): Epidemiology of toxoplasmosis worldwide status. Schweiz Med. Wochenschr., 65 (suppl.):19-22.