

- **Abaci a, Oguzhan a, Kahraman S, et al.** Effect of diabetes mellitus on formation of collateral vessels. *Circulation* 1999; 99:2239.1999.
- **Abd El Kader MR, Tantawy WH, Shaker AF, et al.** Non invasive assessment of coronary arteries using 64-slice MSCT. *Thesis for partial fulfillment of MD cardiology, Ain shams university, 2009.* P120-122
- **Achenbach S, Ropers D, Holle J, et al.** In-plane coronary arterial motion velocity: measurement with electron-beam CT. *Radiology* 2000; 216:457–463
- **Achenbach S, Ulzheimer S, Baum U, et al.** Noninvasive coronary angiography by retrospectively ECG-gated multislice spiral CT. *Circulation* 2000; 102:2823–2828.
- **Achenbach S, Moshage W, Ropers D, et al.** Curved multiplanar reconstructions for the evaluation of contrast-enhanced electron beam CT of the coronary arteries. *AJR Am J Roentgenol* 1998; 170: 895–899.
- **Achenbach S.** Detection of coronary stenoses by multidetector computed tomography: It's all about resolution. *J Am Coll Cardiol* 2004; 43: 840-841
- **Adler AI, Stratton IM, Neil HA, et al.** Association of systolic blood pressure with macrovascular and microvascular complications of type 2 diabetes (UKPDS 36): prospective observational study. *BMJ* 2000; 321:412.

- **Alkadhi H, Scheffel H, Desbiolles L, et al.** Dual-source computed tomography coronary angiography: influence of obesity, calcium load, and heart rate on diagnostic accuracy. *Eur Heart J* 2008; 29:766–776
- **Anand DV, Lim E, Lahiri A, Bax JJ.** The role of non-invasive imaging in the risk stratification of asymptomatic diabetic subjects. *Eur Heart J* 2006; 27:905.
- **Anoop KP and Rajgopal K.** Estimation of missing data using windowed linear prediction in laterally truncated projections in cone-beam CT. *Conf Proc IEEE Eng Med Biol Soc* 2007; 2903–2906,
- **Bernard J, Gersh MB, ChB, David M, et al.** Prevalence of and risk factors for coronary heart disease in diabetes mellitus. *www.uptodate.com*, 2009, P1-4
- **Blake DR, Meigs JB, Muller DC, et al.** Impaired glucose tolerance, but not impaired fasting glucose, is associated with increased levels of coronary heart disease risk factors: results from the Baltimore Longitudinal Study on Aging. *Diabetes* 2004; 53:2095.
- **Bland JM and Altman DG:** Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986; 1:307–310,

- **Budoff MJ, Achenbach S, Blumenthal RS, et al.** Assessment of coronary artery disease by cardiac computed tomography: a scientific statement from the American Heart Association Committee on Cardiovascular Imaging and Intervention, Council on Cardiovascular Radiology and Intervention, and Committee on Cardiac Imaging, Council on Clinical Cardiology. *Circulation* 2006; 114:1761–91
- **Budoff MJ, Dowe D, Jollis JG, et al.** Diagnostic performance of 64-multidetector row coronary computed tomographic angiography for evaluation of coronary artery stenosis in individuals without known coronary artery disease: results from the prospective multicenter ACCURACY (Assessment by Coronary Computed Tomographic Angiography of Individuals Undergoing Invasive Coronary Angiography) trial. *J Am Coll Cardiol* 2008; 52:1724.
- **Cademartiri F, Nieman K, Lugt A, et al.** Intravenous contrast material administration at 16-detector row helical CT coronary angiography: test bolus versus bolus-tracking technique. *Radiology* 2004;233:817–23
- **Cerqueira MD, Weissman NJ, Dilsizian V, et al.** Standardized myocardial segmentation and nomenclature for tomographic imaging of the heart: a statement for healthcare professionals from the Cardiac Imaging Committee of the Council on Clinical Cardiology of the American Heart Association. *Circulation* 2002; 105:539–42.

- **Cesmeli E, Edic M, Iatrou M, et al.** A novel reconstruction algorithm for multiphasic cardiac imaging using multislice CT. *Proc SPIE Int Symp Med Imag* 2001; 4320:645-54.
- **Chobanian AV, Bakris GL, Black HR, et al.** The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: *The JNC 7 Report*. *JAMA* 2003; 289:2560.
- **Coutinho M, Gerstein HC, Wang Y, et al.** The relationship between glucose and incident cardiovascular events. A meta regression analysis of published data from 20 studies of 95,783 individuals followed for 12.4 years. *Diabetes Care* 1999; 22:233.
- **Crawford CR and King KF.** Computed tomography scanning with simultaneous patient translation. *Med Phys* 1990; 17:967-82
- **Di Carli MF, Bianco-Battles D, Landa ME, et al.** Effects of autonomic neuropathy on coronary blood flow in patients with diabetes mellitus. *Circulation* 1999; 41:1387.
- **Dirksen MS, Bax JJ, de Roos A, et al.** Usefulness of dynamic multislice computed tomography of left ventricular function in unstable angina pectoris and comparison with echocardiography. *Am J Cardiol* 2002; 90:1157– 1160.
- **Dirksen MS, Lamb HJ, Doornbos J, et al.** Coronary magnetic resonance angiography: technical developments and clinical applications. *J Cardiovasc Magn Reson* 2003; 10: 219-27.

- **Earls JP, Berman EL, Urban BA, et al.** Prospectively gated transverse coronary CT angiography versus retrospectively gated helical technique: improved image quality and reduced radiation dose. *Radiology* 2008; 246:742.
- **Ehara M, Kawai M, Surmely JF, et al.** Diagnostic accuracy of coronary in-stent restenosis using 64-slice computed tomography: comparison with invasive coronary angiography. *J Am Coll Cardiol* 2007; 49:951.
- **Faerman I, Faccio E, Milei J, et al.** Autonomic neuropathy and painless myocardial ischemia in diabetic patients: histologic evidence of their relationship. *Diabetes* 1977; 26:1147.
- **Fleiss JL:** Statistical Methods for Rates and Proportions. 2nd ed. New York, Wiley,1981
- **Flohr T, McCollough CH, Bruder H et al.** First performance evaluation of a dual source CT (DSCT) system. *Eur Radiol* 2006; 16:256-268
- **Flohr T and Ohnesorge B.** Heart rate adaptive optimization of spatial and temporal resolution for electrocardiogram gated multislice spiral CT of the heart. *J Comput Assist Tomogr* 2001; 25:907–923

- **Flohr T, Stierstorfer K, Bruder H, et al.** New technical developments in multislice CT, part 1: Approaching isotropic resolution with sub-millimeter 16-slice scanning. *Röfo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr* 2002; 174:839-845
- **Flohr T, Stierstorfer K, Raupach R, et al.** Performance evaluation of a 64-slice CT system with z-flying focal spot. *Röfo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr* 2004; 176:1803-1810
- **Flohr T, Schoepf U, Kuettner A, et al.** Advances in cardiac imaging with 16-section CT systems. *Acad Radiol* 2003; 10(4):386–401.
- **Fuchs T, Krause J, Schaller S, et al.** Spiral interpolation algorithms for multislice spiral CT–Part 2: Measurement and evaluation of slice sensitivity profiles and noise at a clinical multislice system. *IEEE Trans Med Imag* 2000; 19:835-847
- **Gallagher MJ and Raff GL.** Use of multislice CT for the evaluation of emergency room patients with chest pain: the so-called "triple rule-out". *Catheter Cardiovasc Interv* 2008; 71:92-99
- **Gerber TC, Krause J, Schaller S, et al.** Current results and new developments of coronary angiography with use of contrast-enhanced computed tomography of the heart. *Mayo Clin Proc*, 2002; 77: 55–71

- **Giesler T, Baum U, Ropers D, et al.** Noninvasive visualization of coronary arteries using contrast-enhanced multidetector CT: influence of heart rate on image quality and stenosis detection. *AJR Am J Roentgenol* 2002; 179:911–916.
- **Granger C, Califf R, Young S, et al.** Outcome of patients with diabetes mellitus and acute myocardial infarction treated with thrombolytic agents. *J Am Coll Cardiol* 1993; 21:920.
- **Grundy SM, Benjamin IJ, Burke GL, et al.** Diabetes and Cardiovascular Disease: A Statement for Healthcare Professionals From the American Heart Association. *Circulation* 1999; 100:1134
- **Haberl R, Becker A, Leber A, et al.** Correlation of coronary calcification and angiographically documented stenoses in patients with suspected coronary artery disease: results of 1,764 patients. *J Am Coll Cardiol* 2001; 37:451–457.
- **Haffner SM, Lehto S, Ronnema T, et al.** Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N Engl J Med* 1998; 339:229.
- **Halliburton SS, Petersilka M, Schvartzman PR, et al.** Evaluation of left ventricular dysfunction using multiphasic reconstructions of coronary multi-slice computed tomography data in patients with chronic ischemic heart disease: validation against cine magnetic resonance imaging. *Int J Cardiovasc Imaging* 2003; 19:73–83.

- **Hammoud T, Tanguay JF, Bourassa MG et al.** Management of coronary artery disease: therapeutic options in patients with diabetes. *J Am Coll Cardiol* 2000; 36:355.
- **Hamon M, Biondi-Zoccai GL, Malagutti, et al.** Diagnostic performance of multislice spiral computed tomography of coronary arteries as compared with conventional invasive coronary angiography. *J Am Coll Cardiol* 2006; 48: 1896-1910.
- **Hausleiter J, Meyer T, Hermann F, et al.** Estimated radiation dose associated with cardiac CT angiography. *JAMA* 2009; 301:500.
- **Hendel RC, Patel MR, Kramer CM, et al.** Appropriateness criteria for cardiac computed tomography and cardiac magnetic resonance imaging: a report of the American College of Cardiology Foundation Quality Strategic Directions Committee Appropriateness Criteria Working Group, American College of Radiology, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, American Society of Nuclear Cardiology, North American Society for Cardiac Imaging, Society for Cardiovascular Angiography and Interventions, and Society of Interventional Radiology. *J Am Coll Cardiol* 2006; 48:1475.

- **Henneman, MM, Schuijf, JD, Pundziute, G, et al.** Noninvasive evaluation with multislice computed tomography in suspected acute coronary syndrome: plaque morphology on multislice computed tomography versus coronary calcium score. *J Am Coll Cardiol* 2008; 52:216.
- **Herzog C, Abolmaali N, Balzer JO, et al.** Heart-rate-adapted image reconstruction in multidetector-row cardiac CT: influence of physiology and technical prerequisite on image quality. *Eur Radiol* 2002; 12:2670–78.
- **Herzog C, Nguyen SA, Savino G et al.** Does two-segment image reconstruction at 64-section CT coronary angiography improve image quality and diagnostic accuracy? *Radiology* 2007; 244:121–129
- **Hoffmann MH, Shi H, Manzke R, et al.** Noninvasive coronary angiography with 16-detector row CT: effect of heart rate. *Radiology* 2005; 234:86–97
- **Hong C, Becker CR, Huber A, et al.** ECG gated reconstructed multi-detector row CT coronary angiography: effect of varying trigger delay on image quality. *Radiology* 2001;220:712–717
- **Hsieh J, Londt J, Vass M, et al.** Step-and-shoot data acquisition and reconstruction for cardiac x-ray computed tomography. *Med Phys* 2006; 33:4236–4248

- **Hu G, Jousilahti P, Qiao Q, et al.** The gender-specific impact of diabetes and myocardial infarction at baseline and during follow-up on mortality from all causes and coronary heart disease. *J Am Coll Cardiol* 2005; 45:1413.
- **Husmann L, Leschka S, Desbiolles L, et al.** Coronary artery motion and cardiac phases: dependency on heart rate—implications for CT image reconstruction. *Radiology* 2007; 245:567–576
- **Huxley R, Barzi F, Woodward M, et al.** Excess risk of fatal coronary heart disease associated with diabetes in men and women: meta-analysis of 37 prospective cohort studies. *BMJ* 2006; 332:73
- **Jakobs TF, Becker CR, Ohnesorge B, et al.** Multislice helical CT of the heart with retrospective ECG gating: reduction of radiation exposure by ECG controlled tube current modulation. *Eur Radiol* 2002;12:1081–86.
- **Jeroen JB, Schuijf JD, Don Poldermans, et al.,** Screening for coronary heart disease in patients with diabetes mellitus, 2008 www.uptodate.com, P2-5.
- **Juergens KU, Grude M, Maintz D, et al.** Multi-detector row CT of left ventricular function with dedicated analysis software versus MR imaging: initial experience. *Radiology* 2004; 230:403–410.

- **Juutilainen A, Lehto S, Ronnema T, et al.** Type 2 diabetes as a "coronary heart disease equivalent": an 18-year prospective population-based study in Finnish subjects. *Diabetes Care* 2005; 28:2901.
- **Kannel W and McGee D.** Diabetes and cardiovascular risk factors: The Framingham Study. *Circulation* 1979; 59:8.
- **Kannel, WB.** Lipids, diabetes, and coronary heart disease: insights from the Framingham Study. *Am Heart J* 1985; 110:1100
- **Kawano H, Motoyama T, Hirashima O, et al.** Hyperglycemia rapidly suppresses flow-mediated endothelium-dependent vasodilatation of brachial artery. *J Am Coll Cardiol* 1999; 34:146.
- **Khaw KT, Wareham N, Bingham S, et al.** Association of hemoglobin A1c with cardiovascular disease and mortality in adults: the European prospective investigation into cancer in Norfolk. *Ann Intern Med* 2004; 141:413.
- **Knez A, Petersilka M, Schvartzman PR et al.** Usefulness of multislice spiral computed tomography angiography for determination of coronary artery stenoses. *Am J Cardiol*, 2001; 88: 1191.
- **Kosar F, Barzi F, Woodward M, et al.** Effect of ectasia size or the ectasia ratio on the thrombosis in myocardial infarction frame count in patients with isolated coronary artery ectasia. *Heart Vessels*, 2005, 20, 199–202.

-
- **Krolewski AS, Kosinski EJ, Warram JH, et al.** Magnitude and determinants of coronary artery disease in juvenile-onset, insulin-dependent diabetes mellitus. *Am J Cardiol* 1987; 59 :750.
 - **Kuettner A, Lehto S, Ronnema T, et al.** Diagnostic accuracy of multidetector computed tomography coronary angiography in patients with angiographically proven coronary artery disease. *J Am Coll Cardiol*, 2004; 43: 831–9.
 - **Kuettner A, Wareham N, Bingham S, et al.** Diagnostic accuracy of noninvasive coronary imaging using 16-detector slice spiral computed tomography with 188 ms temporal resolution. *J Am Coll Cardiol*, 2005, 45, 123–7.
 - **Langer A, Freeman MR, Josse R, et al.** Metaiodobenzylguanidine imaging in diabetes mellitus: Assessment of cardiac sympathetic denervation and its relation to autonomic dysfunction and silent myocardial ischemia. *J Am Coll Cardiol* 1995; 25:610.
 - **Lawler LP, Pannu HK, and Fishman EK.** MDCT evaluation of the coronary arteries: how we do it-data acquisition, postprocessing, display, and interpretation. *Am J Roentgenol*, 2005; 184:1402–12.
 - **Leber AW, Knez A, von Ziegler F, et al** Quantification of obstructive and nonobstructive coronary lesions by 64-slice computed tomography. A comparative study with quantitative coronary angiography and intravascular ultrasound. *J Am Coll Cardiol* 2005;46:147–154

- **Lee CD, Folsom AR, Pankow JS, et al.** Cardiovascular events in diabetic and nondiabetic adults with or without history of myocardial infarction. *Circulation* 2004; 109:855.
- **Lembcke A, Rogalla P, Mews J, et al.** Imaging of the coronary arteries by means of multislice helical CT: optimization of image quality with multisegmental reconstruction and variable gantry rotation time. *Fortschr Roentgenstr* 2003; 175:780–785.
- **Leschka S, Alkadhi H, Plass A, et al.** Accuracy of MSCT coronary angiography with 64-slice technology: first experience. *Eur Heart J* 2005; 26:1482-1487
- **Leschka S, Scheffel H, Desbiolles L, et al.** Image quality and reconstruction intervals of dual-source CT coronary angiography: recommendations for ECG pulsing windowing. *Invest Radiol* 2007;42:543–549
- **Leschka S, Wildermuth S, Boehm T, et al.** Noninvasive coronary angiography with 64-section CT: effect of average heart rate and heart rate variability on image quality. *Radiology* 2006; 241:378–385
- **Levitan EB, Song Y, Ford ES, et al.** Is nondiabetic hyperglycemia a risk factor for cardiovascular disease? A meta-analysis of prospective studies. *Arch Intern Med* 2004; 164:2147.
- **The DCCT Research Group.** Lipid and lipoprotein levels in patients with IDDM diabetes control and complication. Trial experience. *Diabetes Care* 1992; 15:886.

- **Lu B, Mao SS, Zhuang N, et al.** Coronary artery motion during the cardiac cycle and optimal ECG triggering for coronary artery imaging. *Invest Radiol* 2001;36 :250–256.
- **Lu W, Resnick HE, Jablonski KA, et al.** Non-HDL cholesterol as a predictor of cardiovascular disease in type 2 diabetes: The Strong Heart Study. *Diabetes Care* 2003; 26:16.
- **Mahnken AH, Spuentrup E, Niethammer M, et al.** Quantitative and qualitative assessment of left ventricular volume with ECG-gated multislice spiral CT: value of different image reconstruction algorithms in comparison to MRI. *Acta Radiol* 2003; 44: 604–611.
- **Manhnken AH, Katoh M, Bruners P et al.** Acute myocardial infarction:assessment of left ventricular function with 16-detector row spiral CT versus MR imaging study in pigs, *Radiology* 2005;236:112-117
- **Mao S, Budoff MJ, Oudiz RJ, et al.** A simple single slice method for measurement of left and right ventricular enlargement by electron beam tomography. *Int J Card Imaging* 2000;16 :383–390.
- **Mao SS, Bakhsheshi H, Lu B, et al.** Effect of electrocardiogram triggering on reproducibility of coronary artery calcium scoring. *Radiology* 2001; 220(3):707–711.
- **Martuscelli E, Romagnoli A, D’Eliseo A, et al.** Accuracy of thin slice computed tomography in the detection of coronary stenoses. *Eur Heart J* 2004; 25: 1043–1048.

- **Mather KJ, Verma S and Anderson TJ.** Improved endothelial function with metformin in type 2 diabetes mellitus. *J Am Coll Cardiol* 2001; 37:1344.
- **Matt D, Scheffel H, Leschka S, et al.** Dual-source CT coronary angiography: image quality, mean heart rate, and heart rate variability. *AJR Am J Roentgenol* 2007; 189:567–573.
- **Mautner SL, Lin F and Roberts WC.** Composition of atherosclerotic plaques in the epicardial coronary arteries in juvenile (type I) diabetes mellitus. *Am J Cardiol* 1992; 70: 1264.
- **McGill, JB, Schneider DJ, Arfken CL, et al.** Factors responsible for impaired fibrinolysis in obese subjects and NIDDM patients. *Diabetes* 1994; 43:104.
- **McNitt-Gray MF.** AAPM/RSNA Physics Tutorial for Residents: Topics in CT. Radiation dose in CT. *Radiographics* 2002; 22:1541–1553
- **Meijboom WB, Meijs MF, Schuijf JD, et al.** Diagnostic accuracy of 64-slice computed tomography coronary angiography: a prospective, multicenter, multivendor study. *J Am Coll Cardiol* 2008; 52:2135.
- **Miller JM, Rochitte CE, Dewey M, et al.** Diagnostic Performance of Coronary Angiography by 64-Row CT. *N Engl J Med* 2008; 359:2324.

- **Miura H, Wachtel RE, Loberiza FR, et al.** Diabetes mellitus impairs vasodilation to hypoxia in human coronary arterioles: reduced activity of ATP-sensitive potassium channels. *Circ Res* 2003; 92:151.
- **Mogensen CE, Hansen KW, Pedersen MM, et al.** Renal factors influencing blood pressure threshold and choice of treatment for hypertension in IDDM. *Diabetes Care* 1991; 14 Suppl 4:13.
- **Mollet NR, Cademartiri F, van Mieghem CA, et al.** High-resolution spiral computed tomography coronary angiography in patients referred for diagnostic conventional coronary angiography. *Circulation* 2005; 112:2318–2323
- **Moreno PR, Murcia AM, Palacios IF, et al.** Coronary composition and macrophage infiltration in atherectomy specimens from patients with diabetes mellitus. *Circulation* 2000; 102:2180.
- **Mori S, Endo M, Obata T, et al.** Properties of the prototype 256-row (cone beam) CT scanner. *Eur Radiol* 2006; 16:2100-2108
- **MRC/BHF Heart Protection Study** of cholesterol lowering with simvastatin in 20 536 high-risk individuals. *Lancet* 2002; 360:7.
- **Muhlenbruch G, Seyfarth T, Soo CS, et al.** Diagnostic value of 64-slice multi-detector row cardiac CTA in symptomatic patients. *Eur Radiol* 2007; 17: 603-9.
- **Napel S, Marks MP, Ruben GD, et al.** CT angiography with spiral CT and maximum intensity projection. *Radiology* 1992; 195:607–610.

- **Niakan E, Harati Y, Rolak L, et al.** Silent myocardial infarction and diabetic cardiovascular autonomic neuropathy. *Arch Intern Med* 1986; 146:2229.
- **Nieman K, Cademartiri F, Lemos PA, et al.** Reliable noninvasive coronary angiography with fast submillimeter multislice spiral computed tomography. *Circulation* 2002; 106:2051–2054.
- **Nieman K, Rensing BJ, van Geuns RJM, et al.** Non-invasive coronary angiography with multislice spiral computed tomography: impact of heart rate. *Heart* 2002; 88:470–474.
- **Nieman K, Rensing BJ, van Geuns RJM et al.** Coronary angiography with multi-slice computed tomography. *Lancet*, 2001, 357, 599–603.
- **Nikolaou K, Knez A, Rist C, et al.** Accuracy of 64-MSCT in the diagnosis of ischemic heart disease. *Am J Roentgenol* 2006; 187: 111-117.
- **Norhammar A, Tenerz A, Nilsson G, et al.** Glucose metabolism in patients with acute myocardial infarction, and no previous diagnosis of diabetes mellitus: A prospective study. *Lancet* 2002; 359:2140.
- **O'Brien T, Nguyen TD and Zimmerman BR.** Hyperlipidemia and diabetes mellitus. *Mayo Clin Proc* 1998; 73:969.

- **Okmen E.** Left main coronary artery aneurysm associated with extensive coronary arterial calcification: case report and review. *Int J Cardiovasc Imaging*, 2004, 20, 231–5.
- **Ong TK, Chin SP, Liew CK, et al.** Accuracy of 64-row multidetector computed tomography in detecting coronary artery disease in 134 symptomatic patients: influence of calcification. *Am Heart J* 2006; 151:1323, e1321–1326
- **Pavone P, Fioranelli M, Dowe DA, et al.** Basic Techniques in the Acquisition of Cardiac Images with CT. CT Evaluation of Coronary Artery Disease. *Springer-Verlag Italia* 2009.
- **Pajunen P, Taskinen MR, Nieminen MS, et al.** Angiographic severity and extent of coronary artery disease in patients with type 1 diabetes mellitus. *Am J Cardiol* 2000; 86:1080.
- **Perez A, Wagner AM, Carreras G, et al.** Prevalence and phenotypic distribution of dyslipidemia in type 1 diabetes mellitus: effect of glycemic control. *Arch Intern Med* 2000; 160:2756.
- **Poirier P, Giles TD, Bray GA, et al.** Obesity and cardiovascular disease: Pathophysiology, evaluation, and effect of weight loss: An update of the 1997 American Heart Association Scientific Statement on Obesity and Heart Disease from the Obesity Committee of the Council on Nutrition, Physical Activity, and Metabolism. *Circulation* 2006; 113:898.

- **Poll LW, Cohnen M, Brachten S, et al.** Dose reduction in multi-slice CT of the heart by use of ECG-controlled tube current modulation (“ECG pulsing”): phantom measurements. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr* 2002; 174(12):1500–1505.
- **Quinones MJ, Hernandez-Pampaloni M, Schelbert H, et al.** Coronary vasomotor abnormalities in insulin-resistant individuals. *Ann Intern Med* 2004; 140:700.
- **Raff GL, Gallagher MJ, Kayali F, et al.** Diagnostic accuracy of non invasive coronary angiography using 64-slice spiral computed tomography. *J Am Coll Cardiol* 2005; 46: 552-7.
- **Ranjadayalan K, Umachandran V, Ambepityla G, et al.** Prolonged anginal perceptual threshold in diabetes: Effects on exercise capacity and myocardial ischemia. *J Am Coll Cardiol* 1990; 16:1120.
- **Ravipati G, Aronow WS, Ahn C, et al.** Association of hemoglobin A(1c) level with the severity of coronary artery disease in patients with diabetes mellitus. *Am J Cardiol* 2006; 97:968.
- **Rispler S, Keidar Z, Ghersin E, et al.** Integrated single-photon emission computed tomography and computed tomography coronary angiography for the assessment of hemodynamically significant coronary artery lesions. *J Am Coll Cardiol* 2007; 49:1059.

- **Robertson WB and Strong JP.** Atherosclerosis in persons with hypertension and diabetes mellitus. *Lab Invest* 1968; 18:538.
- **Ropers D, Baum U, Pohle K, et al.:** Detection of coronary artery stenoses with thin-slice multi-detector row spiral computed tomography and multiplanar reconstruction. *Circulation* 2003 107:664–666.
- **Ropers U, Ropers D, Pflederer T, et al.** Influence of heart rate on the diagnostic accuracy of dualsource computed tomography coronary angiography. *J Am Coll Cardio* 2007; 50:2393–2398
- **Sacks FM, Tonkin AM, Craven T, et al.** Coronary heart disease in patients with low LDL-cholesterol: benefit of pravastatin in diabetics and enhanced role for HDL-cholesterol and triglycerides as risk factors. *Circulation* 2002; 105:1424.
- **Saito I, Folsom AR, Brancati FL, et al.** Nontraditional risk factors for coronary heart disease incidence among persons with diabetes: the Atherosclerosis Risk in Communities (ARIC) Study. *Ann Intern Med* 2000; 133:81.
- **Scheffel H, Alkadhi H, Leschka S, et al.** Low-Dose CT Coronary Angiography in the Step-and-Shoot Mode: Diagnostic Performance. *Eur Heart J* 2008; 29:766–776
- **Schiller NB, Acquatella H, Ports TA et al.** Left ventricular volume from paired biplane two-dimensional echocardiography. *Circulation* 1979; 60: 547–555.

- **Schinkel AF, Elhendy A, Van Domburg RT et al.** Prognostic value of dobutamine-atropine stress myocardial perfusion imaging in patients with diabetes. *Diabetes Care* 2002; 25: 1637–1643.
- **Schlosser T.** Assessment of left ventricular parameters using 16-MDCT and new software for endocardial and epicardial border delineation. *Am J Roentgenol* AJR, 2005, 184, 765–73.
- **Schroeder S, Kopp AF, Kuettner A, et al.** Influence of heart rate on vessel visibility in noninvasive coronary angiography using new multislice computed tomography: experience in 94 patients. *Clin Imaging* 2002; 26:106–111.
- **Schuijf JD, Bax JJ, Jukema JW et al.** Non-invasive angiography and assessment of left ventricular function using multislice computed tomography in patients with type 2 diabetes. *Diabetes care* 2004; 27: 2905-2910.
- **Schuijf, JD., Bax JJ, Jukema JW et al.** Noninvasive coronary imaging and assessment of left ventricular function using 16-slice computed tomography. *Am J Cardiol*, 2005, 95, 571–574.
- **Scognamiglio R, Negut C, Ramondo A, et al.** Detection of coronary artery disease in asymptomatic patients with type 2 diabetes mellitus. *J Am Coll Cardiol* 2006; 47:65.
- **Selvin E, Marinopoulos S, Berkenblit G, et al.** Meta-analysis: glycosylated hemoglobin and cardiovascular disease in diabetes mellitus. *Ann Intern Med* 2004; 141:421.

- **Shechter M, Merz CN, Paul-Labrador MJ, et al.** Blood glucose and platelet-dependent thrombosis in patients with coronary artery disease. *J Am Coll Cardiol* 2000; 35:300.
- **Sheth T, Amlani S, Ellins ML et al.** Computed tomographic coronary angiographic assessment of high-risk coronary anatomy in patients with suspected coronary artery disease and intermediate pretest probability. *Am Heart J* 2008; 155: 918-23.
- **Shlipak MG, Elmouchi DA, Herrington DM, et al.** The incidence of unrecognized myocardial infarction in women with coronary heart disease. *Ann Intern Med* 2001; 134:1043.
- **Singer DE, Nathan DM, Anderson KM, et al.** Association of HbA1c with prevalent cardiovascular disease in the original cohort of the Framingham Heart Study. *Diabetes* 1992; 41:202.
- **Small M, Lowe GD, MacCuish AC, et al.** Thrombin and plasmin activity in diabetes mellitus and their association with glycaemic control. *Q J Med* 1987; 65:1025.
- **Sozzi FB, Elhendy A, Roelandt JR, et al.** Prognostic value of dobutamine stress echocardiography in patients with diabetes. *Diabetes Care* 2003; 26:1074– 1078.
- **Stamler J, Vaccaro O, Neaton JD, Wentworth D.** Diabetes, other risk factors, and 12-yr cardiovascular mortality for men screened in the Multiple Risk Factor Intervention Trial. *Diabetes Care* 1993; 16:434.

- **Stein PD, Yaekoub AY, Matta F et al.** 64-slice CT for diagnosis of coronary artery disease: a systematic review. *Am J Med* 2008; 121:715.
- **Stevens MJ, Raffel DM, Allman KC, et al.** Cardiac sympathetic dysinnervation in diabetes: Implications for enhanced cardiovascular risk. *Circulation* 1998; 98:96
- **Sun Z, Lin CH, Davidson R, et al.** Diagnostic value of 64-slice CT angiography in coronary artery disease: A systematic review. *Eur J Radiol* 2008; 67: 78-84.
- **Taegtmeyer H, McNulty P, Young ME.** Adaptation and maladaptation of the heart in diabetes: Part I: general concepts. *Circulation* 2002; 105:1727.
- **Third report of the National Cholesterol Education Program (NCEP).** Expert Panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III): Final report. US Department of Health and Human Services; Public Health Service; National Institutes of Health; National Heart, Lung, and Blood Institute. *Circulation* 2002; 106:3143.
- **Tops LF, Krishnàn SC, Schuijf JD, et al.** Non coronary Applications of Cardiac Multi-detector Row Computed Tomography. *JACC: Cardiovascular imaging* 2008; 1: 94-106.

- **Vanhoenacker PK, Heijenbrok-Kal MH, Van Heste R, et al.** Diagnostic performance of multidetector CT angiography for assessment of coronary artery disease: meta-analysis. *Radiology* 2007; 244:419.
- **Vogl TJ.** Techniques for the detection of coronary atherosclerosis: multi-detector row CT coronary angiography. *Radiology* 2002; 223: 212–20.
- **Wachtell K, Ibsen H, Olsen MH, et al.** Albuminuria and cardiovascular risk in hypertensive patients with left ventricular hypertrophy: the LIFE study. *Ann Intern Med* 2003; 139:901.
- **Waller BF, Palumbo PJ, Lie JT, et al.** Status of the coronary arteries at necropsy in diabetes mellitus with onset after age 30 years. Analysis of 229 diabetic patients with and without clinical evidence of coronary disease and comparison to 183 control subjects. *Am J Med* 1980; 69:498.
- **Winocour P.** Platelet abnormalities in diabetes mellitus. *Diabetes* 1992; 41:26.
- **Wintersperger BJ, Nikolaou K, von Ziegler F et al.** Image quality, motion artifacts, and reconstruction timing of 64-slice coronary computed tomography angiography with 0.33-second rotation speed. *Invest Radiol* 2006; 41:436–442.

- **Yamamuro M, Heijenbrok-Kal MH, Van Heste R, et al.** Cardiac functional analysis with Multi-Detector Row CT and segmental reconstruction algorithm: comparison with echocardiography, SPECT, and MR imaging. *Radiology*, 2005, 234, 381–90.
- **Yokoyama I Momomura S-I, Ohtake T, et al.** Reduced myocardial flow reserve in non-insulin-dependent diabetes mellitus. *J Am Coll Cardiol* 1997; 30:1472.
- **Yusuf S, Hawken S, Ounpuu S, et al.** Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004; 364:937.
- **Zhang J, Ren S, Sun D, Shen GX.** Influence of glycation on LDL-induced generation of fibrinolytic regulators in vascular endothelial cells. *Arterioscler Thromb Vasc Biol* 1998; 18:1140.