

References

- Abraham DJ, Eckes B, Rajkumar V and Krieg T (2007):** New developments in fibroblast and myofibroblast biology: implications for fibrosis and scleroderma. *Curr. Rheumatol. Rep.*, 9: 136–143.
- Adam M and Hudgins L (2003):** The importance of minor anomalies in the evaluation of the newborn. *NeoReviews*, 4(S4): 99–104.
- Adam R (2008):** Skin care of the diaper area. *Pediatr. Dermatol.*, 25(S4): 427-433.
- Akin F, Spraker M, Aly R, Leyden J, Raynor W and Landin W (2001):** Effects of breathable disposable diapers: reduced prevalence of *Candida* and common diaper dermatitis. *Pediatr. Dermatol.*, 18(S4): 282-290.
- Akinbi HT, Narendran V, Pass AK, Markart P and Hoath SB (2004):** Host defense proteins in vernix caseosa and amniotic fluid. *Am. J. Obstet. Gynecol.*, 191: 2060-2096.
- Akiyama M, et al. (2003):** The clinical spectrum of nonbullous congenital ichthyosiform erythroderma and lamellar ichthyosis. *Clin. Exp. Dermatol.*, 28: 235.
- Al-Dahiyat (2006):** Neonatal Skin lesions in Jordan, Study of Consecutive 500 neonates at King Hussein Medical Center .*Calicut Medical Journal*, 4(S4): 1.

- Aldana-Valenzuela C, Morales-Marquec M, Castellanos-Martnez J, et al. (2005):** Congenital candidiasis: a rare and unpredictable disease. *J. Perinatol.*, 25: 680-682.
- Al-Waili NS (2005):** Clinical and mycological benefits of topical application of honey, olive oil and beeswax in diaper dermatitis. *Clin. Microbiol. Infect.*, 11(S2): 160-163.
- Amagai M, Matsuyoshi N, Wang Z, Andl C and Stanley J (2000):** Toxins in bullous impetigo and staphylococcal scalded-skin syndrome targets desmoglein 1. *Nature Med.*, 6: 1275-1277.
- Amitai DB et al (2000):** Cutis marmorata telangiectatica congenita: Clinical findings in 85 patients. *Pediatr. Dermatol.*, 17: 100.
- Argoubi H, Fitchner C, Richard O, Lavocat MP, Cambazard F and Stéphan JL(2007):** Pustular miliaria rubra and systemic type 1b pseudohypoaldosteronism in a newborn. *Ann. Dermatol. Venereol.*, Mar; 134: 253-256.
- Arya V, Molinaro MJ, Majewski SS, et al. (2003):** Pediatric scabies. *Cutis.*, 71: 193 -196.
- Asboth D, Kassay E, Noll J and Szalai Z (2000):** Neonatal lupus erythematosus: deep and ulcerating form. *Borgyogaszati Venerol. Szemle*, 76: 263-265.
- Ashton GH, Sorelli P, Mellerio JE, Keane FM, Eady RA and McGrath JA (2001):** Alpha 6 beta 4 integrin abnormalities in junctional epidermolysis bullosa with pyloric atresia. *Br. J. Dermatol.*, 144: 408–414.

- Aslan D, Gürsel T and Kaya Z (2004):** Supernumerary nipples in children with hematologic disorders. *Pediatr. Hematol. Oncol.*, Jul-Aug; 21(S5): 461-463.
- Atay E, Akin M, Yuzkoller E, et al. (2003):** Neonatal purpura fulminans associated with early-onset gram-negative enterobacter septicemia: a case report. *Int. Pediatr.*, 18: 162-163.
- Atherton D (2005):** Maintaining healthy skin in infancy using prevention of irritant napkin dermatitis as a model. *Community Pract.*, 78(S7): 255-257.
- Atherton D (2001):** The aetiology and management of irritant diaper dermatitis. *J. Eur. Acad. Dermatol. Venereol.*, 15(S1): 1-4.
- Aumailley M and Gayraud B (1998):** Structure and biological activity of the extracellular matrix. *J. Mol. Med.*, 76: 253–265.
- Aumailley M, El Khal A, Knoss N and Tunggal L (2003):** Laminin 5 processing and its integration into the ECM. *Matrix Biol.*, 22: 49–54.
- Aumailley M, Has C, Tunggal L and Bruckner-Tuderman L (2006):** Molecular basis of inherited skin-blistering disorders, and therapeutic implications. *Expert Rev. Mol. Med.*, 8: 1–21.
- Avalos-Diaz E, Olague-Marchan M, LopezSwiderski AL, Herrera-Esparza R and Diaz L (2000):** Trans-placental passage of maternal pemphigus foliaceus autoantibodies induces neonatal pemphigus. *J. Am. Acad. Dermatol.*, Dec; 43: 1130-1134.

- Aviram-Goldring A, Goldman B, Netanelov-Shapira I, et al. (2000):** Deletion patterns of the STS gene and flanking sequences in Israeli X-linked ichthyosis patients and carriers: analysis by polymerase chain reaction and fluorescence in situ hybridization techniques. *Int. J. Dermatol.*, Mar; 39(S3): 182-187.
- Bello YM, Falabella AF and Schachner LA (2003):** Management of epidermolysis bullosa in infants and children. *Clin. Dermatol.*, 21: 278-282.
- Berg C, Geipel A, Kohl M, Krokowski M, Baschat AA, Germer U, et al. (2003):** Prenatal sonographic features of Harlequin ichthyosis. *Arch. Gynecol. Obstet.*, Apr; 268(S1): 48-51.
- Berk DR and Bayliss SJ (2008):** Milia: a review and classification. *J. Am. Acad. Dermatol.*, Dec; 59(S6): 1050-1063.
- Bernier V, Weill FX, Hirigouyen V, Eleau C, Feyler A, Labreze C, et al. (2002):** Skin colonization by *Malassezia* species in neonates. A prospective study and relationship with neonatal cephalic pustulosis. *Arch. Dermatol.*, 138: 215-218.
- Bieber T (2008):** Atopic dermatitis. *N. Engl. J. Med.*, 358: 1483–1494.
- Bigorre M, Van Kien AK and Valette H (2009):** Beta-blocking agent for treatment of infantile hemangioma. *Plast. Reconstr. Surg.*, Jun; 123(S6): 195-196.
- Bix G and Iozzo RV (2005):** Matrix revolutions: “tails” of basement membrane components with angiostatic functions. *Trends Cell Biol.*, 15: 52–60.

- Bjarnason B and Flosadottir E (1999):** Childhood, neonatal and stillbirth pemphigus vulgaris. *Int. J. Dermatol.*, 38: 680-688.
- Boiko S (2000):** Making rash decisions in the diaper area. *Pediatr. Ann.*, 29: 50–56.
- Bongain A, Benoit B, Ejnes L, Lambert JC and Gillet JY (2002):** Harlequin fetus: three-dimensional sonographic findings and new diagnostic approach. *Ultrasound Obstet. Gynecol.*, Jul; 20(S1): 82-85.
- Boon L and Vikkula M (2008):** Vascular malformation. In: Wolf K., Goldsmith L., Katz S., Gilchrest B., Paller A. and Leffell D.(editors): *Fitzpatrick`s dermatology in general medicine*. Seventh edition, McGraw Hill Medica; Ch 137: p 1657-1660.
- Boulais N and Misery L (2007):** Merkel cells. *J. Am. Acad. Dermatol.*, 57: 147–165.
- Boulais N and Misery L (2008):** The epidermis: a sensory tissue. *Eur. J. Dermatol.*, 18(S2): 119-127.
- Braverman IM (2000):** The cutaneous microcirculation. *J. Investig. Dermatol. Symp. Proc.*, 5: 3–9.
- Bruckner AL (2008):** Epidermolysis bullosa. In: Eichenfield L, Frieden IL, Esterly NB, (editors): *Neonatal dermatology*. 2nd ed, London: Saunders Elsevier; p: 159-173.
- Bruckner AL and Frieden IJ (2003):** Hemangiomas of infancy. *J. Am. Acad. Dermatol.*, 48: 477-493.

- Bruckner-Tuderman L (2002):** Epidermolysis bullosa. In: Royce PM and Steinmann B (editors): Connective tissue and its heritable disorders. Molecular, genetic and medical aspects. 2nd editions, Wiley-Liss, New-York, p: 689–725.
- Bruckner-Tuderman L, Hopfner B, Hammami-Hauasli N (1999):** Biology of anchoring fibrils: lessons from dystrophic epidermolysis bullosa. *Matrix Biol.*, 18: 43–54.
- Buckely DA, Munn SE and Higgins EM (2001).** Neonatal eosinophilic pustular folliculitis. *Clin. Exp. Dermatol.*, 26: 251-255.
- Burns DA, Breathnach SM, Cox N and Griffiths CE, et al. (2004):** Rook's Textbook of Dermatology. Seventh edition, Wiley-Blackwell.
- Buyon JP and Clancy RM. (2005):** Neonatal lupus: basic research and clinical perspectives. *Rheum. Dis. Clin. N. Am.*, 31: 299–331.
- Campbell JM and Banta-Wright SA (2000):** Neonatal Skin Disorders: A Review of Selected Dermatologic Abnormalities. *J. Perinat. Neonat. Nurs.*, 14(S1): 63–83.
- Campo-Voegeli C, Muniz F, Mascaro JM, et al. (2002):** Neonatal pemphigus vulgaris with extensive muco-cutaneous lesions from a mother with oral pemphigus vulgaris. *Br. J. Dermatol.*, Oct; 147(S4): 801-805.
- Candi E, Schmidt R and Melino G (2005):** The cornified envelope: a model of cell death in the skin. *Nat. Rev. Mol. Cell Biol.*, 6: 328–340.

- Carpo BG, Grevelink JM and Grevelink SV (1999):** Laser treatment of pigmented lesions in children. *Seminars in Cutaneous Medicine and Surgery*, 18: 233-243.
- Cartlidge P (2000):** The epidermal barrier. *Semin. Neonatol.*, 5(4): 273-280.
- Chang CJ, Yu JS and Nelson JS (2008):** Confocal microscopy study of neurovascular distribution in facial port wine stains (capillary malformation). *J. Formos. Med. Assoc.*, Jul; 107(S7): 559-566.
- Chang MW and Orlow SJ (2008):** From birth to old age. In: Wolf K., Goldsmith L. , Katz S., Gilchrest B., Paller A., Leffell D.(editors): *Fitzpatrick's dermatology in general medicine*. Seventh edition, McGraw Hill Medica; Ch 19: p 936l.
- Chaudhari P and Marinkovich MP (2007):** What's new in blistering disorders? *Curr. Allergy Asthma Rep.*, 7: 255–263.
- Cimaz R, Biggioggero M, Catelli L, Muratori S and Cambiaghi S (2002).** Ultraviolet light exposure is not a requirement for the development of cutaneous neonatal lupus. *Lupus*, 11(S4): 257-260.
- Concannon P, Gisoldi E, Phillips S and Grossman R (2001):** Diaper dermatitis: a therapeutic dilemma. Results of a double-blind placebo controlled trial of miconazole nitrate 0.25%. *Pediatr. Dermatol.*, 18(S2): 149-155.
- Conde DM, Kashimoto E, Torresan RZ and Alvarenga M (2006):** Pseudomamma on the foot: an unusual presentation of

supernumerary breast tissue. *Dermatol Online J.*, May 30; 12(S4): 7.

Conlon JD and Drolet BA. (2004): Skin lesions in the neonate. *Pediatr. Clin. North. Am.*, 51: 863-888.

Cotsarelis G (2006): Epithelial stem cells: a folliculocentric view. *J. Invest. Dermatol.* , 126: 1459–1468.

Cribier B and Grosshans E (2002): Histologie de la peau normale et lésions histopathologiques élémentaires. In: *Encycl Méd Chir (Dermatologie)*, Éditions Techniques (Paris), 85-98 A10; 16 p.

Darmstadt GL, Dinulos JG and Miller Z. (2000): Congenital cutaneous candidiasis: clinical presentation, pathogenesis, and management guidelines. *Pediatrics*, 105: 438-444.

De Raeve LE and Roseeuw DI (2002): Curettage of giant congenital melanocytic nevi in neonates: a decade later. *Arch. Dermatol.*, Jul; 138(S7): 943-947.

De Wet P, Rode H, Van A, et al. (1999): Perianal candidiasis a comparative study with mupirocin and nystatin. *Int. J. Dermatol.* , 38: 618 –622.

Ehretsmann C, Schaefer P and Adam R (2001): Cutaneous tolerance of baby wipes by infants with atopic dermatitis, and comparison of the mildness of baby wipe and water in infant skin. *J. Eur. Acad. Dermatol. Venereol.*, 15(S1): 16-21.

- Eichenfield LF and Bogen ML (2007):** Absorption and efficacy of miconazole nitrate 0.25% ointment in infants with diaper dermatitis. *J. Drugs Dermatol.* , 6(S5): 522-526.
- Eichenfield LF and Hardaway CA (1999):** Neonatal dermatology. *Curr. Opin. Pediatr.* , 11: 471–474.
- Eichenfield LF, Frieden IJ and Esterly NB (2008):** Neonatal Dermatology. Second edition, Philadelphia, Pa: Saunders Elsevier; 142.
- Eichenfield LF, Lucky AW, Boguniewicz M, et al. (2002):** Safety and Efficacy of pimecrolimus (ASM 981) cream 1% in the treatment of mild and moderate atopic dermatitis in children and adolescents. *J. Am. Acad. Dermatol.*, 46(S4): 495.
- Eigenmann PA (2001):** Clinical features and diagnostic criteria of atopic dermatitis in relation to age. *Pediatr. Allergy Immunol.* , 14 (S12): 69-74.
- El Helali N, Carbonne A, Naas T, et al. (2005):** Nosocomial outbreak of staphylococcal scalded skin syndrome in neonates: epidemiological investigation and control. *J. Hosp. Infect.*, 61: 130-138.
- Elias PM (2007):** The skin barrier as an innate immune element. *Semin. Immunopathol.* , 29(S1): 3-14.
- Elias PM, Crumrine D, Rassner U, et al. (2004):** Basis for abnormal desquamation and permeability barrier dysfunction in RXLI. *J. Invest. Dermatol.* , Feb; 122(S2): 314-319.

- Elias PM, Schmuth M, Uchida Y, et al. (2002):** Basis for the permeability barrier abnormality in lamellar ichthyosis. *Exp. Dermatol.*, 11: 248-256.
- Elish D and Silverberg NB (2006):** Neonatal lupus erythematosus. *Cutis.*, Feb; 77(S2): 82-86.
- Enright AM and Prober CG (2002):** Neonatal herpes infection: diagnosis, treatment and prevention. *Semin. Neonatol.*, 7: 283.
- Erasala GN, Merlay I and Romain C (2007):** Evolution of disposable diapers and reduction of diaper dermatitis. *Arch. Pediatr.*, 14(S5): 495-500.
- Evangelista F, Dasher DA, Diaz LA, Prisayanh PS and Li N (2008):** E-cadherin is an additional immunological target for pemphigus autoantibodies. *J. Invest. Dermatol.*, 128: 1710–1718.
- Fabbri P, Amato LK, Coronella G, et al. (2003):** Cutaneous lupus erythematosus: diagnosis and management. *Am. J. Clin. Dermatol.*, 4: 449 .
- Feingold KR, Schmuth M and Elias PM (2007):** The regulation of permeability barrier homeostasis. *J. Invest. Dermatol.*, 127: 1574–1576.
- Fine JD, Eady RA, Bauer EA, et al. (2000):** Revised classification system for inherited epidermolysis bullosa: report of the Second International Consensus Meeting on diagnosis and

classification of epidermolysis bullosa. *Am. Acad. Dermatol.*, 42: 1051- 1066.

Fischer J, Faure A, Bouadjar B, et al. (2000): Two new loci for autosomal recessive ichthyosis on chromosomes 3p21 and 19p 12-q 12 and evidence for further genetic heterogeneity. *Am. J. Hum. Genet.*, 66: 904-913.

Fluhr JW, Kao J, Jain M, Ahn SK, Feingold KR and Elias PM (2001): Generation of free fatty acids from phospholipids regulates stratum corneum acidification and integrity. *J. Invest. Dermatol.*, 117(S1): 44-51.

Folkman J (2003): Fundamental concepts of the angiogenic process. *Curr. Mol. Med.*, 3: 643-651.

Fonder MA, Mamelak AJ, Kazin RA and Cohen BA (2007): Port-wine-stain-associated dermatitis: implications for cutaneous vascular laser therapy. *Pediatr. Dermatol.*, Jul-Aug; 24(4): 376-379.

Friedlander SF and Bradley JS (2008): Viral infections. In: Eichenfield L, Frieden IL and Esterly NB (editors): *Neonatal Dermatology*. The second edition, London: Saunders Elsevier; p. 195.

Fuchs E (2007): Scratching the surface of skin development. *Nature*, 445: 834–842.

Fuchs E (2008): Skin stem cells: rising to the surface. *J. Cell Biol.*, 180: 273–284.

- Galli SJ and Tsai M (2008):** Mast cells: versatile regulators of inflammation, tissue remodeling, host defense and homeostasis. *J. Dermatol. Sci.*, 49: 7–19.
- Gallup E and Plott T (2005):** A multicenter, open-label study to assess the safety and efficacy of ciclopirox topical suspension 0.77% in the treatment of diaper dermatitis due to *Candida albicans*. *J. Drugs Dermatol.*, 4(S1): 29-34.
- Gardella C, Handsfield HH and Whitley R (2008):** Neonatal herpes, the forgotten perinatal infection. *Sex. Transm. Dis.*, 35: 22-24.
- Ghadially R, Halkier-Sorensen L and Elias PM (1992):** Effects of petrolatum on stratum corneum structure and function. *J. Am. Acad. Dermatol.*, 26: 387–396.
- Gilliam AE, Pauporte M and Frieden IJ (2008):** Vesiculobullous and erosive diseases in the newborn. In: Bologna JL, Jorizzo JL, Rapini RP (editors): *Dermatology*. 2nd edition, New York, NY: Elsevier; p. 475-478.
- Green KJ and Simpson CL (2007):** Desmosomes: new perspectives on a classic. *J. Invest. Dermatol.*, 127: 2499–2515.
- Greve B, Hammes S and Raulin C (2001):** The effect of cold air cooling on 585 nm pulse dye laser treatment of port-wine stains. *Dermatol. Surg.*, 27(S7): 633.
- Griffiths WAD, Judge MR and Leigh IM (1998):** Disorders of keratinization. In: Champion RH, et al. (editors): *Rook's textbook of dermatology*. The sixth edition, Malden, MA: Blackwell; p. 1483–1588.

- Grossl NA (2000):** Supernumerary breast tissue: historical perspectives and clinical features. *South. Med. J.*, 93: 29.
- Gu LH and Coulombe PA (2007):** Keratin function in skin epithelia: a broadening palette with surprising shades. *Curr. Opin. Cell Biol.*, 19: 13–23.
- Gupta AK and Skinner AR (2004):** Management of diaper dermatitis. *Int. J. Dermatol.*, 43(S11): 830-834.
- Gupta R, Warren T and Wald A (2007):** Genital herpes. *Lancet*, 370: 2127-2137.
- Haass NK, Smalley KS, Li L and Herlyn M (2005):** Adhesion, migration and communication in melanocytes and melanoma. *Pigment Cell Res.*, 18: 150–159.
- Hairston BR, Davis MD, Pittelkow MR, et al. (2006):** Livedoid vasculopathy: further evidence for procoagulant pathogenesis. *Arch. Dermatol.*, 142: 1413-1418.
- Hamm H (2000):** Developmental abnormalities. In: John H, Arnold O and Neil P (editors): *Pediatric dermatology*. 2nd edition, Blackwell science, p.8: 91.
- Handsfield HH, Waldo AB, Brown ZA, et al. (2005):** Neonatal herpes should be a reportable disease. *Sex. Transm. Dis.*, 32: 521-525.
- Hanson M, Lupski J, Hicks J and Metry D (2003):** Association of dermal melanocytosis with lysosomal storage disease. *Arch. Dermatol.*, 139: 916-920.

- Happle R (2008):** What is a capillary malformation? *J. Am. Acad. Dermatol.*, Dec; 59(S6): 1077-1079.
- Hariharan D, Manno CS, and Seri I (2000):** Neonatal lupus erythematosus with microvascular hemolysis. *Journal of Pediatric Hematology and Oncology*, 22: 351-354.
- Has C, Chang YR, Volz A, Hoeping D, Kohlhase J and Bruckner-Tuderman L (2006):** Novel keratin 14 mutations in patients with severe recessive epidermolysis bullosa simplex. *J. Invest. Dermatol.*, 126: 1912–1914.
- Hashimoto K, Gee S and Tanaka K (2000):** Lamellar ichthyosis: response to etretinate with transglutaminase1 recovery. *Am. J. Dermatopathol.*, 22: 277–280.
- Hernandez-Machin B, Reyes CS, Vargas MM, et al. (2007):** Picture of the month: congenital cutaneous candidiasis. *Arch. Pediatr. Adolesc. Med.*, 161: 907-908.
- Hernandez-Martin A, Gonzalez-Sarmiento R, De Unamuno P (1999):** X-linked ichthyosis: an update. *Br. J. Dermatol.*, 141: 617–627.
- Heukelbach J and Feldmeier H (2006):** Scabies. *Lancet*, 367: 1767-1774.
- Hickey P, Piantanida E, Lentz-Kapua S, et al. (2003):** Neulaxova syndrome: a case report. *Tripler Army Med Center Hawaii. Pediatr. Dermatol.*, 20: 25–27.

- Hidano A, Purwoko R and Jitsukawa K (1996):** Statistical survey of skin changes in Japanese neonates. *Pediatr. Dermatol.*, 3: 140-144.
- Hidano A., Parwoko R and Jitsukawa K. (1986):** Changes in Japanese neonates. *Paediatr. Dermatol.*, 3: 140 – 144.
- Hoath SB (2004):** Physiologic development of the skin. In: Polin RA, Fox WW and Abman SH (editors): *Fetal and Neonatal Physiology*. Philadelphia, Pa: Elsevier.
- Hoath SB (2005):** The skin and neonatal nursing from the biology of vernix to neonatal adaptation. Program and abstracts of the 21st Annual Conference of the National Association of Neonatal Nurses; September 28-October 1; Anaheim, California.
- Hoath SB, Narendran V and Visscher M (2001):** Role and biology of vernix. *Neonatal Infant Nurs. Rev.*, 1: 53–58.
- Hoeger PH and Enzmann CC (2002):** Skin physiology of the neonate and young infant: a prospective study of functional skin parameters during early infancy. *Pediatr. Dermatol.*, 19(S3): 256-262.
- Hoffman D and Ratner D (2006):** Diagnosis and management of a changing congenital melanocytic nevus. *Skinmed*, Sep-Oct; 5(S5): 242-5.
- Hohenester E and Engel J (2002):** Domain structure and organisation in extracellular matrix proteins. *Matrix Biol.*, 21: 115–128.

- Honeycutt KA, Koster MI and Roop DR (2004):** Genes involved in stem cell fate decisions and commitment to differentiation play a role in skin disease. *J. Investig. Dermatol. Symp. Proc.*, 9: 261–268.
- Huxley-Jones J, Robertson DL and Boot-Handford RP (2007):** On the origins of the extracellular matrix in vertebrates. *Matrix Biol.*, 26: 2–11.
- Hyttiainen M, Penttinen C and Keski-Oja J (2004):** Latent TGF beta binding proteins: extracellular matrix association and roles in TGF-beta activation. *Crit. Rev. Clin. Lab. Sci.*, 41: 233–264.
- James W, Berger T and Elston D (2006):** Atopic Dermatitis, Eczema and Noninfectious Immunodeficiency Disorder. In Andrews' (editor): *Andrews' Clinical dermatology*. The tenth edition, Elsevier.
- Jameson J and Havran WL (2007):** Skin gamma delta T-cell functions in homeostasis and wound healing. *Immunol. Rev.*, 215: 114–122.
- Jamora C, DasGupta R, Kocieniewski P and Fuchs E (2003):** Links between signal transduction, transcription and adhesion in epithelial bud development. *Nature*, 422: 317–322.

- Jobard F, Bouadjar B, Caux F, Hadj-Rabia S, Has C, Matsuda F, Weissenbach J, Lathrop M, Prud'homme JF and Fischer J (2003):** Identification of mutations in a new gene encoding aFERM family protein with a pleckstrin homology domain in Kindler syndrome. *Hum. Mol. Genet.*, 12: 925–935.
- Jones P and Simons BD (2008):** Epidermal homeostasis: do committed progenitors work while stem cells sleep? *Nat, Rev, Mol, Cell Biol.*, 9: 82–88.
- Jonkman MF, Pasmooij AM, Pasmans SG, van den Berg MP, Ter Horst HJ, Timmer A and Pas HH (2005):** Loss of desmoplakin tail causes lethal acantholytic epidermolysis bullosa. *Am. J, Hum. Genet.*, 77: 653–660
- Kadler KE, Baldock C, Bella J and Boot-Handford RP (2007):** Collagens at a glance. *J. Cell Sci.*, 120: 1955–1958.
- Kahana M, Feldman M, Abudi Z and Yurman S (1995):** The incidence of birthmarks in Israeli neonates. *Int. J. Dermatol.*, 34: 704-706.
- Kalayciyan A, Engin B, Serdaroglu S, Mat C, Aydemir EH and Kotogyan A (2002):** *Br. J. Dermatol.*, Aug; 147(S2): 396-397.
- Kaplan DH, Jenison MC, Saeland S, Shlomchik WD and Shlomchik MJ (2005):** Epidermal langerhans cell-deficient mice develop enhanced contact hypersensitivity. *Immunity*, 23: 611–620.

- Karvonen SL, Vaaljalaht P, Marenk M, et al. (1992):** Birthmarks in 4346 Finnish newborns. *Acta. Dermal. Venereol.*, 72: 55 – 57.
- Kaur N, Bhakoo ON, et al. (1989):** Survey of cutaneous lesions in Indian newborns. *Paediatr. Dermatol.*, 6: 39 – 42.
- Kaur P (2006):** Interfollicular epidermal stem cells: identification, challenges, potential. *J. Invest. Dermatol.*, 126: 1450–1458.
- Kelly KM, Nanda VS and Nelson JS (2002):** Treatment of port-wine stain birthmarks using the 1.5-msec pulsed dye laser at high fluences in conjunction with cryogen spray cooling. *Dermatol. Surg.*, 28(4): 309.
- Kellsell DP, et al. (2005):** Mutations in ABCA12 underlie the severe congenital skin disease harlequin ichthyosis. *Am. J. Hum. Genet.*, 76: 794.
- Kemp DJ, Walton SF, Harumal P and Currie BJ. (2002):** The scourge of scabies. *Biologist*, 49:19- 24.
- Kern JS, Kohlhasse J, Bruckner-Tuderman L and Has C (2006):** Expanding the COL7A1 mutation database: novel and recurrent mutations and unusual genotype-phenotype constellations in 41 patients with dystrophic epidermolysis bullosa. *J. Invest. Dermatol.*, 126: 1006–1012.
- Kimberlin DW, Lin CY, Jacobs RF, et al. (2001):** Safety and efficacy of high dose intravenous acyclovir in the management of neonatal herpes simplex virus infections. *Pediatrics.*, 108: 230-238.

- Kissenpfennig A and Malissen B (2006):** Langerhans cells– revisiting the paradigm using genetically engineered mice. *Trends Immunol.*, 27: 132–139.
- Kjaer KW, Hansen L, Schwabe GC, Marques-de-Faria AP, Eiberg H, Mundlos S, Tommerup N and Rosenberg T (2005):** Distinct CDH3 mutations cause ectodermal dysplasia, ectrodactyly, macular dystrophy (EEM syndrome). *J. Med. Genet.*, 42: 292–298.
- Kohelet D and Arbel E (2000):** A prospective search for urinary tract abnormalities in infants with isolated preauricular tags. *Pediatrics*, 105(S5).
- Kono T, Groff WF and Sakurai H (2006):** Treatment of port wine stains with the pulse dye laser. *Ann. Plast. Surg.*, Apr; 56(S4): 460-463.
- Koster MI, Huntzinger KA and Roop DR (2002):** Epidermal differentiation: transgenic/knockout mouse models reveal genes involved in stem cell fate decisions and commitment to differentiation. *J. Investig. Dermatol. Symp. Proc.*, 7: 41–45.
- Kulkarni ML and Rajeev S (1996):** Normal variants of skin in neonates. *Indian J. Dermatol. Venereol. Leprol.*, 62: 83-86.
- Kuller JM (2001):** Skin Breakdown Risk Factors, Prevention, and Treatment. *Clinical practice*, 1: 35–42.

- Ladhani S and Joannu C (2000):** Difficulties in diagnosis and management of the staphylococcal scalded-skin syndrome. *Pediatric Infec. Dis. J.*, 19: 819–821.
- Ladhani S., Robbie S., Chapple DS., et al. (2003):** Isolating staphylococcal scalded-skin syndrome is not clinically useful. *Pediatr Infect Dis*: 22: 284–286.
- Lai-Cheong JE, Arita K and McGrath JA (2007):** Genetic diseases of junctions. *J. Invest. Dermatol.*, 127: 2713–2725.
- Lane EB and McLean WH (2004):** Keratins and skin disorders. *J. Pathol.*, 204: 355-366.
- Lanigan SW (2001):** Treatment of vascular naevi in children. *Hosp. Med.*, 62(S3): 144.
- Laufer MR and Goldstein DP (2005):** The breast: examination and lesions. In: Sydor A. (editor): *Pediatric & Adolescent Gynecology* (5th edition), Philadelphia, Lippincott Williams & Wilkins, p 729-759.
- Laughter D, Istvan JA, Tofte SJ, et al. (2000):** The prevalence of atopic dermatitis in Oregon school children. *J. Am. Acad. Dermatol.*, 43: 649-655.
- LaVoo EJ and Paller AS (1994):** Common skin problems during the first year of life. *Pediatric Clinics of North America*, 41: 1105–1119.
- Leaute-Labreze C, Dumas de la Roque E, Hubiche T, Boralevi F, Thambo JB and Taieb A(2008):** Propranolol for severe

hemangiomas of infancy. *N. Engl. J. Med.*, Jun 12; 358(S24): 2649-2651.

Lee LA (2004): Neonatal lupus erythematosus: clinical findings and pathogenesis. *J. Investig. Dermatol. Symp. Proc.*, 9: 52–56.

Lee LA (2004): Neonatal lupus: clinical features and management. *Pediatr. Drugs.*, 6: 71-78.

Lee LA (2009): The clinical spectrum of neonatal lupus. *Arch. Dermatol. Res.*, Jan; 301(s1): 107-110.

Lee LA, Sokol RJ, and Buyon JP (2002): Hepatobiliary disease in neonatal lupus: Prevalence and clinical characteristics in cases enrolled in a national registry. *Pediatrics.* 109: 1–5.

Leena Bruckner-Tuderman and Cristina Has (2010): Inherited bullous disease. In Thomas Krieg, David R. Bickers and Yoshiki Miyachi (Editors): *In Therapy of Skin Diseases.* Springer-Verlag Berlin Heidelberg .Ch. 10: p. 549-559.

Leighty L, Li N, Diaz LA and Liu Z (2007): Experimental models for the autoimmune and inflammatory blistering disease, Bullous pemphigoid. *Arch. Dermatol. Res.*, 299: 417–422.

Leung DY (2000): Atopic dermatitis: new insights and opportunities for therapeutic intervention. *J. Allergy Clin. Immunol.*, 105(S5): 860.

- Lippens S, Denecker G, Ovaere P, Vandenabeele P and Declercq W (2005):** Death penalty for keratinocytes: apoptosis versus cornification. *Cell Death Differ.*, 12(S2): 1497–1508.
- Litjens SH, De Pereda JM and Sonnenberg A (2006):** Current insights into the formation and breakdown of hemidesmosomes. *Trends Cell. Biol.*, 16: 376–383.
- Liu C, Feng J, Qu R, Zhou H, Ma H, Niu X, et al. (2005):** Epidemiologic study of the predisposing factors in erythema toxicum neonatorum. *Dermatology*; 210: 269-272.
- Loden M, Buraczewska I and Edlund F (2003):** The irritation potential and reservoir effect of mild soaps. *Contact Dermatitis*, 49(2): 91-96.
- Long WB, Edlich RF, Winters KL, et al. (2005):** Cold injuries .J. Long term effect *Med. Implants.*, 15: 67.
- Lorenz S, Maier C, Segerer H, Landthaler M and Hohenleutner U (2000):** (Skin changes in newborn infants in the first 5 days of life). *Hautarzt.*, Jun; 51(6): 396-400.
- Loser K and Beissert S (2007):** Dendritic cells and T cells in the regulation of cutaneous immunity. *Adv. Dermatol.*, 23: 307–333.
- Maari C and Frieden IJ (2004):** Klippel-Trénaunay syndrome: the importance of "geographic stains" in identifying lymphatic disease and risk of complications. *J. Am. Acad. Dermatol.*, Sep; 51(3): 391-398.

- Mac-Moune Lai F, To KF, Choi PC, et al. (2001):** Kaposiform hemangioendothelioma: five patients with cutaneous lesion and long follow-up. *Mod. Pathol.*, 14: 1087- 1092.
- Madison KC (2003):** Barrier function of the skin: “la raison d’etre” of the epidermis. *J. Invest. Dermatol.*, 121: 231–241.
- Maghoob AA (1996):** Large congenital melanocytic nevi and the risk for developing malignant melanoma: A prospective study .*Arch. Dermatol.*, 132: 170.
- Magin TM, Vijayaraj P and Leube RE (2007):** Structural and regulatory functions of keratins. *Exp. Cell Res.*, 313: 2021–2032.
- Mallory S B and Colven R (2000):** Disorders of subcutaneous tissue in the newborn .In: John H, Arnold O and Neil P (editors): *Pediatric dermatology* .2nd edition, Blackwell science; p.5: 64.
- Mallory SB (1991):** Neonatal Skin Disorders. *Pediatric Clinics of North America.*, 38: 745–761.
- Mancini AJ (2006):** Vascular disorders of infancy and childhood. *Hurwitz Clinical Pediatric Dermatology*. 3rd edition, London: Elsevier; p. 322-331.
- Marchini G, Hultenby K, Nelson A, et al. (2007):** Increased expression of HMGB-1 in the skin lesions of erythema toxicum. *Pediatr. Dermatol.*, Sep-Oct; 24(S5): 474-482.

- Marchini G, Lindow S, Brismar H, et al. (2002):** The newborn infant is protected by an innate antimicrobial barrier: peptide antibiotics are present in the skin and vernix caseosa. *Br. J. Dermatol.*, Dec; 147(S6): 1127-1134.
- Marchini G, Nelson A, Edner J, Lonne-Rahm S, Stavreus-Evers A and Hultenby K (2005):** Erythema toxicum neonatorum is an innate immune response to commensal microbes penetrated into the skin of the newborn infant. *Pediatr. Res.*, Sep; 58(S3): 613-616.
- Marchini G, Stabi B, Kankes K, Lonne-Rahm S, Ostergaard M and Nielsen S (2003):** AQP1 and AQP3, psoriasin, and nitric oxide synthases 1-3 are inflammatory mediators in erythema toxicum neonatorum. *Pediatr. Dermatol.*, Sep-Oct; 20(S5): 377-384.
- Marchini G, Ulfgren AK, Lore K, Stabi B, Berggren V and Lonne-Rahm S (2001):** Erythema toxicum neonatorum: an immunohistochemical analysis. *Pediatr. Dermatol.*, May-Jun; 18(S3): 177-187.
- Margulis A, Adler N and Bauer BS (2009):** Congenital melanocytic nevi of the eyelids and periorbital region. *Plast. Reconstr. Surg.*, Oct; 124(4): 1273-1283.
- Martin JM and Green M. (2006):** Group A streptococcus. *Semin. Pediatr. Infect. Dis.*, 17: 140-148.

- Matter K and Balda MS (2007):** Epithelial tight junctions, gene expression and nucleo-junctional interplay. *J. Cell. Sci.*, 120: 1505–1511.
- Mc Atee-Smith J, Herbert AA, Rapini RP, et al. (1994):** Skin lesions of the spinal axis and spinal dysraphism. Fifteen cases and a review of the literature. *Arch. Pediatr. Adolesc. Med.*, 148: 740-748.
- McCarthy JS, Kemp DJ, Walton SF and Currie BJ (2004):** Scabies: more than just an irritation. *Postgrad. Med. J.*, 80: 382 -387.
- McGill DJ and Mackay IR (2007):** Capillary vascular malformation response to increased ambient temperature is dependent upon anatomical location. *Ann. Plast. Surg.*, Feb; 58(S2): 193-199.
- McGrath JA and Mellerio JE (2006):** Epidermolysis bullosa. *Br. J. Hosp. Med.*, 67: 188-191.
- McLaughlin MR, O'Connor NR and Ham P (2008):** Newborn skin: part II. Birthmarks. *Am. Fam. Physician.*, 77: 56-60.
- McMillan JR, Akiyama M and Shimizu H (2003):** Epidermal basement membrane zone components: ultrastructural distribution and molecular interactions. *J. Dermatol. Sci.*, 31: 169–177.
- Medina YN, Rapaport R (2009):** Evolving diagnosis of McCune-Albright syndrome. Atypical presentation and follow up. *J. Pediatr. Endocrinol Metab.*, Apr; 22(S4): 373-377.

- Mengesha YM and Bennett ML (2002):** Pustular skin disorders: diagnosis and treatment. *Am. J. Clin. Dermatol.*, 3(S6): 389-400.
- Mese G, Richard G and White TW (2007):** Gap junctions: basic structure and function. *J. Invest. Dermatol.*, 127: 2516–2524.
- Moll I, Roessler M, Brandner JM, Eispert AC, Houdek P and Moll R (2005):** Human Merkel cells—aspects of cell biology, distribution and functions. *Eur. J. Cell Biol.*, 84: 259–271.
- Montanez E, Piwko-Czuchra A, Bauer M, Li S, Yurchenco P and Fassler R (2007):** Analysis of integrin functions in peri-implantation embryos, hematopoietic system and skin. *Methods Enzymol.*, 426: 239–289.
- Moosavi Z and Hosseini T (2006):** One-year survey of cutaneous lesions in 1000 consecutive Iranian newborns. *Pediatr. Dermatol.*, 23: 61-63.
- Morelli JG (2001):** Vascular neoplasms. In: Fitzpatrick JE (Editor): *Dermatology Secrets*. The second edition, Philadelphia, Hanley and Belfus, Inc, p 303–306.
- Muhle C, Jiang QJ, Charlesworth A, Bruckner-Tuderman L, Meneguzzi G and Schneider H (2005):** Novel and recurrent mutations in the laminin-5 genes causing lethal junctional epidermolysis bullosa: molecular basis and clinical course of Herlitz disease. *Hum. Genet.*, 116: 33–42.
- Muller EJ, Williamson L, Kolly C and Suter MM (2008):** Outside-in signaling through integrins and cadherins: a central

mechanism to control epidermal growth and differentiation?
J. Invest. Dermatol., 128: 501–516.

Nanda A, Kaur S, Bhakoo ON and Dhall K (1998): Survey of cutaneous lesions in Indian newborns. *Pediatr. Dermatol.*, 6: 39-42.

Nanda S, Reddy BS, Ramji S, et al. (2002): Analytical study of pustular eruptions in neonates. *Pediatr. Dermatol.*, 19: 210- 215.

Nanno M, Shiohara T, Yamamoto H, Kawakami K and Ishikawa H (2007): Gamma delta T cells: firefighters or fire boosters in the front lines of inflammatory responses. *Immunol. Rev.*, 215: 103–113.

Narendran V, Wickett RR, Pickens WL and Hoath SB (2000): Interaction between pulmonary surfactant and vernix: a potential mechanism for induction of amniotic fluid turbidity. *Pediatr. Res.*, 48: 120-124.

Neiman AR, Lee LA, Weston WL and Buyon J.P (2000): Cutaneous manifestations of neonatal lupus without heart block: Characteristics of mothers and children enrolled in a national registry. *The Journal of Pediatrics*, 137: 674-680.

Nelson A, Ulfgren AK, Edner J, Stabi B, Brismar H and Hultenby K (2007): Urticaria Neonatorum: accumulation of tryptase-expressing mast cells in the skin lesions of newborns with Erythema Toxicum. *Pediatr. Allergy Immunol.*, Dec; 18(S8): 652-658.

- Nickoloff BJ, Xin H, Nestle FO and Qin JZ (2007):** The cytokine and chemokine network in psoriasis. *Clin Dermatol*; 25: 568–573.
- Nield LS and Kamat D (2007):** Prevention, diagnosis, and management of diaper dermatitis. *Clin. Pediatr.*, 46(S6): 480-486.
- Niessen CM (2007):** Tight junctions/adherens junctions: basic structure and function. *J. Invest. Dermatol.*, 127: 2525–2532.
- Niessen CM and Gottardi CJ (2008):** Molecular components of the adherens junction. *Biochim. Biophys. Acta.*, 1778: 562–571.
- Nishifuji K, Sugai M and Amagai M. (2008):** Staphylococcal exfoliative toxins: “molecular scissors” of bacteria that attack the cutaneous defense barrier in mammals. *J. Dermatol. Sci.*, 49: 21-31.
- Nolan J and Sinclair R (2001):** Review of management of purpura fulminans and two case reports. *Br J Anaesth*; 86: 581-586.
- Nonato LB, Lund CH, Kalia YN and Guy RH (2000):** Transepidermal water loss in 24 and 25 weeks gestational age infants. *Acta. Paediatr.*, 89(6): 747-748.
- Nordlund JJ (2007):** The melanocyte and the epidermal melanin unit: an expanded concept. *Dermatol. Clin.*, 25: 271–281.
- Norlen L (2006):** Stratum corneum keratin structure, function and formation – a comprehensive review. *Int. J. Cosmet. Sci.*, 28: 397–425.

- North PE, Waner M, Mizeracki A, et al. (2000):** GLUT1: a newly discovered immunohistochemical marker for juvenile hemangiomas. *Hum. Pathol.*, 31: 11- 22.
- Ochiai T, Suzuki Y, Kato T, et al. (2007):** Natural history of extensive Mongolian spots in mucopolysaccharidosis type II (Hunter syndrome): a survey among 52 Japanese patients. *J. Eur. Acad. Dermatol. Venereol.*, 21: 1082-1085.
- O'Connor NR, McLaughlin MR and Ham P (2008).** Newborn skin: Part I. Common rashes. *Am. Fam. Physician.*, Jan 1; 77(S1): 47-52.
- Odio M, Streicher-Scott J and Hansen RC (2001):** Disposable baby wipes: efficacy and skin mildness. *Dermatol. Nurs.*, 13(S2): 107-121
- Oliver G and Detmar M (2002):** The rediscovery of the lymphatic system: old and new insights into the development and biological function of the lymphatic vasculature. *Genes Dev.*, 16: 773–783.
- Orkin MI and Maibach HI (2001):** Scabies and pediculosis. In Freedberg IM et al. (Editor): *Fitzpatrick's Dermatology in general medicine*. The fifth edition, New York, McGraw-Hill, Health Professions Division: 2677–2695.
- Osburn K, Schosser RH and Everett MA (1987):** Congenital pigmented and vascular lesions in newborn infants. *J. Am. Acad. Dermatol.*, 16: 788-792.

Oztas MO and Gurer MA (2007): Dermoscopic features of accessory nipples. *Int. J. Dermatol.*, Oct; 46(S10): 1067-1068.

Paller A, Mancini AJ and Hurwitz S (2006): Hurwitz Clinical Pediatric Dermatology: A Textbook of Skin Disorders of Childhood and Adolescence. The third edition, Philadelphia, Pa.: Elsevier Saunders: 737-746.

Paller AS (2001): Use of nonsteroidal topical immunomodulators for the treatment of atopic dermatitis in the pediatric population. *J. Pediatr.*, 138(S2): 163.

Paller AS, McAlister RO, Dayle JJ, Jackson A, et al. (2002): Perceptions of physicians and pediatric patients about atopic dermatitis, its impact, and its treatment. *Clin. Pediatr.*, 41(S5): 323.

Paris L, Tonutti L, Vannini C and Bazzoni G (2008): Structural organization of the tight junctions. *Biochim. Biophys. Acta.*, 1778: 646–659.

Parlowsky T, Welzel J, Amagai M, Zillikens D, Wygold T (2003): Neonatal pemphigus vulgaris: IgG4 autoantibodies to desmoglein 3 induce skin blisters in newborns. *J. Am. Acad. Dermatol.*, Apr; 48 (S4): 623-625.

- Pasmooij AM, Pas HH, Jansen GH, Lemmink HH and Jonkman MF (2007):** Localized and generalized forms of blistering in junctional epidermolysis bullosa due to COL17A1 mutations in the Netherlands. *Br. J. Dermatol.*, 156: 861–870.
- Penate Y, Lujan D, Rodriguez J, et al (2005):** (Neonatal lupus erythematosus: 4 cases and clinical review). *Actas. Dermosifiliogr*, Dec; 96(S10): 690-696.
- Pickens W, Zhou Y, Wickett R, Visscher M and Hoath SB (2000):** Antioxidant defense mechanisms in vernix caseosa: potential role of endogenous vitamin E. *Pediatr. Res.*, 47(S5): 425A.
- Pickens WL, Warner RR, Boissy YL, Boissy RE and Hoath SB (2000):** Characterization of vernix caseosa: water content, morphology, and elemental analysis. *J. Invest. Dermatol.*, 115(S5): 875-881.
- Pickering LK (2003):** Candidiasis. In *Red Book: Report on the Committee on Infectious Disease*. Elk Grove Village: American Academy of Pediatrics, p. 229-231
- Plensdorf S and Martinez J (2009):** Common pigmentation disorders. *Am. Fam. Physician.*, Jan 15; 79(S2): 109-16.
- Powell J. (2007):** Skin physiology. *Acute care*, 193-196.
- Prendiville JS. (2000):** Scabies and lice. In: Harper J, Oranje A and Prose N (editors): *Textbook of pediatric dermatology*. The seven edition, Oxford Blackwell Science; p. 555- 569.

- Prigent F, Vige P and Martinemt C. (1999):** Coetaneous lesions during the first week of life in 306 consecutive newborns infants. *Ann. Dermatol. Venereol.*, 118: 697 – 699.
- Rahm S (2001):** Erythema toxicum neonatorum. *Pediatr. Dermatol.*, 18: 177-187.
- Raposo G and Marks MS (2007):** Melanosomes–dark organelles enlighten endosomal membrane transport. *Nat. Rev. Mol. Cell Biol.*, 8: 786–797.
- Requena L, Sanchez Yus E and Kutzner H (2008) :** Panniculitis Disorders of subcutaneous tissue. In: Wolf K., Goldsmith L., Katz S., Gilchrest B., Paller A. and Leffell D.(editors): *Fitzpatrick`s dermatology in general medicine. The seventh edition, McGraw Hill Medica; section 10, Ch 68: p 576.*
- Richard G (2005):** Connexin disorders of the skin. *Clin. Dermatol.*, 23: 23–32.
- Rinn JL, Bondre C, Gladstone HB, Brown PO and Chang HY (2006):** Anatomic demarcation by positional variation in fibroblast gene expression programs. *PLoS Genet.*, 2: 119.
- Ripmeester P and Dunn S (2002):** Against all odds: breastfeeding a baby with harlequin ichthyosis. *J. Obstet. Gynecol. Neonatal. Nurs.*, Sep-Oct; 31(S5): 521-525.
- Ripple F, SchrienerV and Schwantiz HJ (2002):** The acidic medium of the horny layer. *Am. J. Clin. Dermatol.*, 3: 261-272

- Rivers JK, Frederiksen PC and Dibdin C (1990):** A prevalence survey of dermatoses in the Australian neonate. *J. Am. Acad. Dermatol.*, 23: 77-81.
- Runeman B (2008):** Skin interaction with absorbent hygiene products. *Clin. Dermatol.*, 26(1): 45-51.
- Rutter KJ and Judge MR (2009):** Profuse congenital milia in a family. *Pediatr. Dermatol.*, Jan-Feb; 26(S1): 62-64.
- Sachdeva M, Kaur S, Nagpal M and Dewan SP (2002):** Cutaneous lesions in new born. *Indian J. Dermatol. Venereol. Leprol.*, 68: 334-337.
- Sandhu K and Kanwar AJ. (2004):** Generalized bullous impetigo in a neonate. *Pediatr. Dermatol.*, 21: 667-669.
- Sandilands A, Smith FJ, Irvine AD and McLean WH (2007):** Filaggrin's fuller figure: a glimpse into the genetic architecture of atopic dermatitis. *J. Invest. Dermatol.*, 127: 1282–1284.
- Scheinfeld NS, Silverberg NB, Weinberg JM and Nozad V (2004):** The preauricular sinus: a review of its clinical presentation, treatment, and associations. *Pediatr. Dermatol.*, 21(s3): 191–196.
- Schlotawa L, Steinfeld R, von Figura K, Dierks T and Gartner J (2008):** Molecular analysis of SUMF1 mutations: stability and residual activity of mutant formylglycine-generating enzyme determine disease severity in multiple sulfatase deficiency. *Hum. Mutat.*, Jan; 29(S1): 205.

- Schmid-Wendtner MH and Korting HC (2006):** The pH of the skin surface and its impact on the barrier function. *Skin Pharmacol. Physiol.*, 19(S6): 296-302.
- Schmuth M, Gruber R, Elias PM and Williams ML (2007):** Ichthyosis update: towards a function-driven model of pathogenesis of the disorders of cornification and the role of corneocyte proteins in these disorders. *Adv. Dermatol.*, 23: 231–256.
- Schroder JM, Reich K, Kabashima K, Liu FT, Romani N, Metz M, Kerstan A, Lee PH, Loser K, Schon MP, Maurer M, Stoitzner P, Beissert S, Tokura Y and Gallo RL (2006):** Who is really in control of skin immunity under physiological circumstances - lymphocytes, dendritic cells or keratinocytes? *Exp. Dermatol.*, 15: 913–929.
- Schunck M, Neumann C and Proksch E (2005):** Artificial barrier repair in wounds by semi-occlusive foils reduced wound contraction and enhanced cell migration and reepithelization in mouse skin. *J. Invest. Dermatol.*, 125(S5): 1063-1071.
- Sen K and Roy A (2006):** Management of neonatal purpura fulminans with severe protein C deficiency. *Indian Pediatr.*, 43: 542-545.
- Sengler C, Lau S, Wahn U, et al. (2002):** Interactions between genes and environmental factors in asthma and atopy: new developments. *Respir. Res.*, 3(S1): 7.

- Sethuraman G and Mancini AJ (2008):** Neonatal Skin Disorders and the Emergency Medicine Physician. In: Clinical pediatric emergency medicine; 9: 200-209.
- Shahab RK and Loo DS (2003):** Bullous scabies. J. Am. Acad. Dermatol., 49: 346- 350.
- Shieh S, Frang YV, Becker JL, Holm A, Beutner EH and Helm TN (2004):** Pemphigus, pregnancy and plasma-pheresis. Cutis., May; 73(S5): 327-329.
- Shin HT (2005):** Diaper dermatitis that does not quit. Dermatol. Ther. 18(2): 124-135.
- Shwayder T (2004):** Disorders of keratinization, diagnosis, and management. Am. J. Clin. Dermatol., 5: 17–29.
- Shwayder T and Akland T (2005):** Neonatal skin barrier: structure, function, and disorders. Dermatologic Therapy, 18: 87–103.
- Siegel DH, Ashton GH, Penagos HG, Lee JV, Feiler HS, Wilhelmsen KC, South AP, Smith FJ, Prescott AR, Wessagowit V, Oyama N, Akiyama M, Al Aboud D, Al Aboud K, Al Githami A, Al Hawsawi K, Al Ismaily A, Al Suwaid R, Atherton DJ, Caputo R, Fine JD, Frieden IJ, Fuchs E, Haber RM, Harada T, Kitajima Y, Mallory SB, Ogawa H, Sahin S, Shimizu H, Suga Y, Tadini G, Tsuchiya K, Wiebe CB, Wojnarowska F, Zaghloul AB, Hamada T, Mallipeddi R, Eady RA, McLean WH, McGrath JA and Epstein EH (2003):** Loss of kindlin-1, a human homolog of the *Caenorhabditis elegans* actin-extracellular-matrix linker

protein UNC-112, causes Kindler syndrome. *Am. J. Hum. Genet.*, 73: 174–187.

Siegfried EC, Keenan WJ and Al-Jureidini S (2008): More on propranolol for hemangiomas of infancy. *N. Engl. J. Med.*, Dec 25; 359(S26): 2846-2847.

Simpson C. (2003): The management of staphylococcal scalded skin syndrome. *Nursing Times*, 99: 59–61.

Singalavanija S and Limpongsanurak W (2003): Immuno-bullous disease in Thai children. *J. Med. Assoc. Thai.*, Aug; 86(S3): 681-688.

Singh S, Bhura M, Maheshwari A, Kumar A, Singh CP and Pandey SS (2001): Successful treatment of harlequin ichthyosis with acitretin. *Int. J. Dermatol.*, Jul; 40(S7): 472-473.

Sixt M, Bauer M, Lammermann T and Fassler R (2006): Beta1 integrins: zip codes and signaling relay for blood cells. *Curr. Opin. Cell Biol.*, 18: 482–490.

Sladden MJ and Sladden JA (2008): A novel treatment for accessory nipples. *Br. J. Dermatol.*, Feb; 158(S2): 405.

Smolinski KN, Shah SS, Honig PJ, et al. (2005): Neonatal cutaneous fungal infections. *Curr. Opin. Pediatr.*, 17: 486-493.

Sonnenberg A and Liem RK (2007): Plakins in development and disease. *Exp. Cell Res.*, 313: 2189–2203.

Sorrell JM and Caplan AI (2004): Fibroblast heterogeneity: more than skin deep. *J. Cell Sci.*, 117: 667–675.

- Spergel JM and Paller AS. (2003):** Atopic dermatitis and the atopic march. *J. Allergy Clin. Immunol.*, 112(S6): 118-127.
- Sprecher E, Bergman R, Richard G, Lurie R, Shalev S, Petronius D, Shalata A, Anbinder Y, Leibu R, Perlman I, Cohen N and Szargel R (2001):** Hypotrichosis with juvenile macular dystrophy is caused by a mutation in CDH3, encoding P-cadherin. *Nat. Genet.*, 29: 134–136.
- Stanley JR and Amagai M. (2006):** Pemphigus, bullous impetigo, and the staphylococcal scalded-skin syndrome. *N. Engl. J. Med.*, 355: 1800-1810.
- Stratigos AJ, Dover JS and Arndt KA (2000):** Laser treatment of pigmented lesions- 2000: how far have we gone? *Arch. Dermatol.*, 136(S7): 915-921.
- Suresh S, Vijayalakshmi R, Indrani S, and Lata M. (2004):** Short foot length: a diagnostic pointer for harlequin ichthyosis. *J. Ultrasound Med.*, Dec; 23(S12): 1653-1657.
- Taïeb A and Boralevi F (2007):** Hypermelanoses of the newborn and of the infant. *Dermatol. Clin.*, 25: 327-336.
- Tan T, Constantinides H and Mitchell TE (2005):** The preauricular sinus: a review of its aetiology, clinical presentation and management. *Int. J. Pediatr. Otorhinolaryngol.*, 69(S11): 1469–1474.
- Tekin M, Bodurtha JN and Riccardi VM (2001):** Cafe au lait spots: the pediatrician's perspective. *Pediatr. Rev.*, Mar; 22(S3): 82-90.

- Thacher TD, Fischer PR, Pettifor JM and Darmstadt GL (2004):** Nutritional rickets in ichthyosis and response to calcipotriene. *Pediatrics*, Jul; 114(S1): 119-123.
- Thami GP, Kaur S and Kanwar AJ (2002):** Surgical Pearl: Enucleation of milia with a disposable hypodermic needle. *J. Am. Acad. Dermatol.*, Oct; 47(S4): 602-603.
- Travers J and Mousdicas N (2008):** Gram –positive bacteria associated with toxin production . In: Wolf K., Goldsmith L., Katz S., Gilchrest B., Paller A., Leffell D.(editors): *Fitzpatrick`s dermatology in general medicine*. The seventh edition, by: McGraw Hill Medica; 178 : p1714 .
- Tschachler E (2007):** Psoriasis: the epidermal component. *Clin. Dermatol.*, 25: 589–595.
- Tzu J and Marinkovich MP (2008):** Bridging structure with function: structural, regulatory, and developmental role of laminins. *Int. J. Biochem. Cell Biol.*, 40: 199–214.
- Uitto J and Richard G. (2004):** Progress in epidermolysis bullosa: genetic classification and clinical implications. *Am. J. Med. Genet.*, 131: 61-74.
- Uitto J, Richard G and McGrath JA (2007):** Diseases of epidermal keratins and their linker proteins. *Exp. Cell Res.*, 313: 1995–2009.
- Van Gysel D, Lijnen RL, Moekti SS, et al. (2002):** Collodion baby: a follow-up study of 17 cases. *J. Eur. Acad. Dermatol. Venereol.*, 16: 472-473.

- Varki R, Sadowski S, Pfendner E and Uitto J (2006):** Epidermolysis bullosa. I. Molecular genetics of the junctional and hemidesmosomal variants. *J. Med. Genet.*, 43: 641–652.
- Verbov J (2000):** Common skin conditions in newborn. *Semin. Neonatol.*, 5: 303-310.
- Visscher M (2003):** Environmental Interactions. In: Hoath SB and Maibach HI (editors): *Neonatal Skin Structure and Function*. The second Edition, Marcel Dekker, NY, USA.
- Visscher M (2005):** Infant skin research. Presented at: The 2nd Annual Joint Meeting of the International Society of Bioengineering and the Skin and the International Society for Skin Imaging. Philadelphia, PA, USA, 29 September-1 October.
- Visscher M and Hoath SB (2006):** Diaper dermatitis. In Chew A (editor): *Irritant Dermatitis*. Springer, Berlin, Germany, 37-51.
- Visscher M, Hoath SB, Conroy E and Wickett RR (2001):** Effect of semipermeable membranes on skin barrier repair following tape stripping. *Arch. Dermatol. Res.*, 293(S10): 491-499.
- Visscher M, Taylor T, White T et al. (2007):** Wipes versus current standard for diaper skin care. Presented at: The National Association of Neonatal Nurses 23rd Annual. Conference. San Diego, CA, USA, 26-29 September.
- Visscher M, Taylor T, White T et al. (2008):** Wipes versus Current Standard for Diaper Skin Care. Presented at: The 9th

European Society for Pediatric Dermatology. Athens, Greece, 15-17 May.

Visscher MO, Chatterjee R, Munson KA, Pickens WL and Hoath SB (2000): Changes in diapered and non diapered infant skin over the first month of life. *Pediatr. Dermatol.*, 17(S1): 45-51.

Visscher MO, Narendran V, Pickens WL et al. (2005): Vernix caseosa in neonatal adaptation. *J. Perinatol.*, 25(S7): 440-446.

Visscher SB (2005): Diaper Dermatitis. In Chew A (editor): *Irritant Dermatitis*. Springer, San Francisco, CA, USA: 37-51.

Vural E, Ramakrishnan J, Cetin N, Buckmiller L, Suen JY and Fan CY (2008): The expression of vascular endothelial growth factor and its receptors in port-wine stains. *Otolaryngol. Head Neck Surg.*, Oct; 139(S4): 560-564.

Wagner A. (1997): Distinguishing vesicular and pustular disorders in the neonate. *Curr. Opin. Pediatr.*, 9: 396-405.

Walker VP, Akinbi HT, Meinzen-Derr J, Narendran V, Visscher M and Hoath SB (2008): Host defense proteins on the surface of neonatal skin: implications for innate immunity. *J. Pediatr.*, 152(S6): 777-781.

Wallach D (2003): Diagnosis of common benign neonatal dermatoses. *Clin. Dermatol.*, 21: 264-268.

- Ward DB, Fleischer AB Jr, Feldman SR and Krowchuk DP (2000):** Characterization of diaper dermatitis in the United States. Arch. Pediatr. Adolesc. Med., 154(S9): 943-946.
- Watt FM, Lo Celso C and Silva-Vargas V (2006):** Epidermal stem cells: an update. Curr. Opin. Genet. Dev., 16: 518–524.
- Weedon D (2002):** Skin Pathology. The second edition, London, Churchill Livingstone.
- Weston W, Lane A and Morelli JG (2002):** Structure and function of the skin. In: Color textbook of pediatric dermatology. The third edition, St. Louis: Williams & Wilkins: 1–8.
- White SJ, McLean WH (2005):** Kindler surprise: mutations in a novel actin-associated protein cause Kindler syndrome. J. Dermatol. Sci., 38: 169–175.
- Willman JH, Golitz LE and Fitzpatrick JE (2003):** Clear cells of Toker in accessory nipples. J. Cutan. Pathol., Apr; 30(4): 256-260.
- Wolf R, Wolf D, Tuzun B and Tuzun Y (2000):** Diaper dermatitis. Clin. Dermatol., 18(S6): 657-660.
- Woodley DT, Remington J and Chen M (2007):** Autoimmunity to type VII collagen: epidermolysis bullosa acquisita. Clin. Rev. Allergy Immunol., 33: 78–84.
- World Health Organization (2003):** Pregnancy, Childbirth, Postpartum and Newborn Care.

- Wu H, Brandling-Bennett HA and Harris TJ. (2004):** Noninfectious vesiculobullous and vesiculopustular diseases. In Elder D, Elenitsasis R, Jaworsky C, and Johnson Jr B (editors): *Lever's Histopathology*. 10th. Philadelphia, Pa: Lippincott-Raven.
- Wu H, Wang ZH, Yan A et al. (2000):** Protection against pemphigus foliaceus by Desmoglein 3 in neonates. *N. Engl. J. Med.*, July; 343(S1): 31-35.
- Yamasaki O, Yamaguchi T, Sugai M, et al. (2005):** Clinical manifestations of staphylococcal scalded-skin syndrome depend on serotypes of exfoliative toxins. *J. Clin. Microbiol.*, 43: 1890-1893.
- Yosipovitch G, Maayan-Metzger A, Merlob P and Sirota L (2000):** Skin barrier properties in different body areas in neonates. *Pediatrics*, 106: 105-108.
- Yung-Weng W and Ying-Ju C (2004):** A preliminary study of bottom care effects on premature infants' heart rate and oxygen saturation. *J. Nurs. Res.*, 12(S2): 161-168.
- Zeeuwen PL (2004):** Epidermal differentiation: the role of proteases and their inhibitors. *Eur. J. Cell Biol.*, 83: 761–773.
- Zhu HY, Li HB, Wu LQ, et al (2008):** (Multiplex quantitative PCR detection for female carrier in an X-linked ichthyosis family). *Zhonghua Yi Xue Za Zhi*. Dec16; 88(S46): 3246-3249.

Zvulunov A and Metzker A (2002): Hemangiomas and vascular malformations: unapproved treatments. *Clin. Dermatol.*, 20: 660- 667.

Zweers MC, Davidson JM, Pozzi A, Hallinger R, Janz K, Quondamatteo F, Leutgeb B, Krieg T and Eckes B (2007): Integrin alpha2 beta1 is required for regulation of murine wound angiogenesis but is dispensable for reepithelialization. *J. Invest. Dermatol.*, 127: 467–478.