

REFERENCES

Abdel-Khalek, M.; Sheir, K.Z.; Mokhtar, A.A.; Eraky, I.; Kenawy, M. Bazeed, M. (2004):

Prediction of success rate after extracorporeal shock-wave lithotripsy of renal stones- a multivariate analysis model. Scand J Urol Nephrol., 38: 161 - 167.

Aldret-Neilson, L.; Bjerring, P. and Nielson, J. 1999):

Regional variations in analgesic efficacy of EMLA cream. Acta Dermatol Scand., 70: 314–8.

Avalos, H. R.; Sanchez, L. M. and Lechuga, A.B. (2003):

The analgesic effect of petroleum jelly as a means of contact in ESWL. J. Mexican colleague. Urol. 18 (4):141-144.

Basar, H.; Yilmaz, E.; Ozcan, S. et al (2003):

Four analgesic techniques for shockwave lithotripsy: Eutectic mixture of local anaesthetic is a good alternative. J Endourol., 17:36.

Basar, M.M.; Samli, M.M.; Erbil, M.; Ozergin, O.; Basar, R. and Atan,A. (2004):

Early effects of extracorporeal shock-wave lithotripsy exposure on testicular sperm morphology. Scand J Urol Nephrol., 38: 38 - 41.

Becker, A.J.; Stiff, C.G.; Truss, M.C.; Oelke, M.; Machtens, S. and Jonas, U. (1999):

Petroleum jelly is an ideal contact medium for pain reduction and successful treatment with extra corporeal shock wave lithotripsy, J. Urol., 162: 18-22.

Bierkens, A.F.; Hendrikx, A.J.; Lemmens, W.A. and. Debruyne, F.M. (1991):

Extracorporeal shock-wave lithotripsy for large renal calculi: the role of ureteral stents. A randomized trial. J Urol., 145: 699 – 702.

Bierkens, A.F.; Maes, R.M.; Hindriks, A.J.M.; Erdos, A.f.; Vries, J.D.; and Debruyne, F.M.(1991):

The use of local anaesthesia in extracorporeal shock wave lithotripsy: eutectic mixture of local anaesthetics. J. Urol., 146:287- 289.

Brinkmann, O.A.; Griebel, A.; Kuwertz-Broking, E.; Bulla, M. and Hertle, L. (2001):

Extracorporeal shock wave lithotripsy in children. Efficacy, complications and long-term follow-up. Eur Urol., 39: 591- 597.

Chaussy, C.; Brendel, W. and Schmiedt E (1980): Extracorporeally induced destruction of kidney stones by shock waves. Lancet 316: 1265- 1275.

Chaussy, C.; Schmiedt, E.; Jocham, D.; Brendel, W.; Forssmann, B. and Walther, V. (1982):

First clinical experience with extracorporeally induced destruction of kidney stones by shock waves. J Urol., 127: 417.

Chaussy, C.; Schuller, J.; Schmiedt, E.; Brandl, H.; Jocham, D. and Liedl, B. (1984):

Extracorporeal shock-wave lithotripsy (ESWL) for treatment of urolithiasis. Urol., 23: 59 - 66.

Chen, R.N. and Streem, S.B. (1996):

Extracorporeal shock wave lithotripsy for lower pole calculi: long term radiographic and clinical outcome. J Urol., 156: 1572- 1575.

Cleveland, R.O.; Bailey, M.R.; Hartennaum, B., et al (2000):

Design and characterization of a research electrohydraulic lithotripter patterned after the Dornier HM3. Rev Scientific Inst., 71: 2514-2525.

Coffman, J.D. (1993):

Cutaneous changes in peripheral vascular disease. In: Dermatology in General Medicine, 4th ed. Edited by T.B. Feitzpatrick. New Yourk: Mcgraw-Hill, pp. 2077-2102.

Cohen, T.D. and Preminger, G.M. (1997):

Management of calyceal calculi. Urol Clin North Am., 24: 81-96.

Delevecchio, F. C. and Preminger, G. M. (2000):

Management of residual stones. Urol Clin North Am., 27: 347-354.

Demir, E.; Kilciler, M.; Bedir, S.; Erten, K. and Ozgok Y. (2007):

Comparing Two Local Anaesthesia Techniques for Extracorporeal Shock Wave Lithotripsy: Urol., 69: 625-628.

Elbahnasy, A.M.; Shalhav, A.L.; Hoenig, D.M., et al. (1998):

Lower caliceal stone clearance after shock wave lithotripsy or ureteroscopy. The impact of lower pole radiographic anatomy. J Urol., 159: 676-682.

Freinkel, R.K. (1993):

Cutaneous changes in peripheral vascular disease. In: Dermatology in General Medicine, 4th ed. Edited by T.B. Feitzpatrick. New Yourk: Mcgraw-Hill, pp. 2113-2131.

Ganapathy, S.; Razvi, H.; Moote, C.; Parkin, J.; Yee, I.; Gverzdys, S.; Dain, S. and Denstedt, J.D. (1996):

Eutectic mixture of local anaesthetics is not effective for extracorporeal shock wave lithotripsy. Can J Anaesth 43: 1030.

Gofrit, O.N.; Pode, D. Meretyk, S.; Katz, G.; Shapiro, A.; Golijanin, D .et al, (2001):

Is the pediatric ureter as efficient as the adult ureter in transporting fragments following extracorporeal shock wave lithotripsy for renal calculi larger than 10 mm. J Urol., 166: 1862 – 1864.

Grasso, M. (2008):

Extracorporeal Shockwave Lithotripsy. [http:// emedicine. medscape. com/](http://emedicine.medscape.com/) updated article. 2008.

Greenstein, A. and Matzkin, H. (1999):

Does the rate of extracorporeal shock wave delivery affect stone fragmentation?. Urology 54 (1999) (430 - 432).

Gupta, N. and Kumar A. (2008):

Analgesia for pain control during extracorporeal shock wave lithotripsy: Current status. Indian J Urol., 24: 155-8.

Heidenreich, A.; Bonfig, R.; Wilbert, D.M. and Engelmann, U.H. (1995):

Painless ESWL by cutaneous application of vaseline. Scand J Urol Nephrol., 29: 155–60.

Jain, A. and Shah, TK. (2007):

Effect of Air Bubbles in the Coupling Medium on Efficacy of Extracorporeal Shock Wave Lithotripsy. J Europ Urol., 51: 1680-1687.

Jenkis, A. D. (1987):

Patient selection and education. In: Riehle, R. A. and Newman, R. C. (eds), Principles of extra corporeal shockwave lithotripsy. Churchill Livingstone, New York, Chapt.5, p.67.1987.

Jocham, D.; Hager, H. and Weber, W. (1988):

Technical, clinical, and other aspects. In Kamper-Jorgensen, F., Challah, S., Folmer Andersen, T. Technology Assessment and New Kidney Stone Treatment Methods. Oxford University Press, Oxford, U.K., p. 25.

Kumar, A.; Gupta, N.P.; Hemal, A.K. and Wadhwa, P. (2007):

Comparison of three analgesic protocols for pain control during shock wave lithotripsy using dornier delta compact lithotripter: A prospective randomized clinical trial. J Endourol., 21:578–82.

Lichtenberg, m.H.; Miskowlak, J.; Mogensen, P. and Andersen, J.T. (1992):

Local anaesthesia for extra corporeal shock wave lithotripsy: a study comparing eutectic mixture of local anesthesia cream and lidocaine infiltration. J. Urol., 147:96.

Lingeman, J.E.; Coury, T.A.; Newman, D.M.; Kahnoski, R.J.; Mertz, J.H.; Mosbaugh, P.G. et al. (1987):

Comparison of results and morbidity of percutaneous nephrolithotomy and extracorporeal shock wave lithotripsy. J Urol., 138: 485 - 490.

Lingeman, J.E.; Matlaga, B.R. and Evan, A.O. (2007):

Surgical Management of Upper Urinary Tract Calculi. In: Kavoussi, L.R., Novick, A.C., Partin, A.W., Peters, C.A. editors: Campbell's Urology, 9th edit, Vol. 4, section XI. Ch.44, W.B. Saunders, PP. 89-128.

Lingeman, J.E.; Siegel, Y.I.; Steele, B., et al. (1994):

Management of lower pole nephrolithiasis: a critical analysis. J Urol., 151: 663-667.

Longo, J.A. and Netto Junior, N.R. (1995):

Extracorporeal shock wave lithotripsy in children. Urol., 46: 550-552.

Madbouly, K.; Sheir, K.Z.; Elsobky, E.; Eraky, I. and Kenawy, M. (2002):

Risk factors for the formation of steinstrasse after extracorporeal shock wave lithotripsy: a statistical model. J Urol., 167: 12349-12442.

McCullough, D. L. (1989):

Extracorporeal shock wave lithotripsy and residual stone fragments in the lower calices (letter to the editor). J Urol, 141: 140.

Menon, M. and Koul, H. (1992):

Clinical review 32: Calcium oxalate nephrolithiasis. J Clin Endocrinol Metab; 74:703-707.

Monk, T.G.; Boure, B.; White, P.F.; Meretyk, S. and Clayman, R.V. (1991):

Comparison of intravenous sedative-analgesic techniques for outpatient immersion lithotripsy. Anesth Analg., 72: 616-21.

Monk, T.G.; Ding, Y.; White, P.F.; Albala, D.M. and Clayman, R.V. (1994):

Effect of topical eutectic mixture of local anaesthetics on pain response and analgesic requirement during lithotripsy procedures. Anaesth Analg., 79: 506-11.

Moody, J.A.; Evans, A.P. and. Lingeman, J.E. (2001):

Extracorporeal shock-wave lithotripsy. In Weiss, R.M. George, N.J.R., O'Reilly, P.H. (Eds). Comprehensive Urology, Mosby International Limited, p. 623 – 636.

Muslumanoglu, A.Y.; Tefekli, A.; Sarilar, O.; Binbay, M.; Altunrende, F. and Ozkuvanci, U. (2003):

Extracorporeal shock wave lithotripsy as first line treatment alternative for urinary tract stones in children: a large scale retrospective analysis. J Urol., 170: 2405 - 2408.

Ohmori, K.; Matsuda, T.; Horii, T. and Yoshida, O. (1994):

Effects of shock waves on the mouse foetus. J Urol., 151: 255 – 258.

Okeke, Z.; Lam, J.S.; and Gupta, M. (2004):

Use of ureteral access sheath to facilitate removal of large stone burden during extracorporeal shock wave lithotripsy. *Urol.*, 63: 574-576.

Orsola, A.; Diaz, I.; Caffaratti, J.; Izquierdo, F.; Alberola, J. and Garat, J.M. (1999):

Staghorn calculi in children: treatment with monotherapy extracorporeal shock wave lithotripsy. *J Urol.*, 162: 1229 - 1233.

Osman, M.M.; Alfano, Y.; Kamp, S.; Haecker, A. ; Alken, P.; Michel, T. and Knoll, M.S.(2005):

5-year-follow-up of patients with clinically insignificant residual fragments after extracorporeal shockwave lithotripsy. *Eur Urol* 47: 860 – 864.

Ozcan, S.; Yilmaz, E.; Buyukkocak, U.; Basar, H. and Apan, A. (2002):

Comparison of three analgesics for extracorporeal shock wave lithotripsy. *Scand J Urol Nephrol.*, 36: 281.

Pettersson, B.; Tiselius, H.G.; Andersson, A. et al (1989): Evaluation of extracorporeal shock wave lithotripsy without anaesthesia using a Dornier HM3 lithotripter without technical modifications. *J Urol.*, 142: 1189-1192.

Picramenos, D.; Deliveliotis, C.; Alexopoulou, K.; Makrichoritis, C.; Kostakopoulos, A. and Dimopoulos, C. (1996):

Extracorporeal shock wave lithotripsy for renal stones in children. *Int Urol.*, 56: 86-89.

Preminger, G.M.; Kettelhut, M.C.; Elkins, S.L.; Seger, J. and Fetner, C.D. (1989):

Ureteral stenting during extracorporeal shock wave lithotripsy: Help or hindrance. *J Urol.*, 142:L 32-36.

Rassweiler, J.J.; Tailly, G.G. and Chaussy, C. (2005):

Progress in lithotripter technology. Eau Update Series, 3:17.

Reichelt, O.; Zermann, D.H.; Wunderlich, H.; Janitzky, V. and Schubert, J. (1999):

Effective analgesia for extracorporeal shock wave lithotripsy: transcutaneous electrical nerve stimulation. Urol., 54: 433.

Sampaio F., and Aragão, A. (1992):

Inferior pole collecting system anatomy. Its probable role in extracorporeal shock wave lithotripsy. J Urol., 147: 322-324.

Sampaio, F. (2001):

Renal collecting system anatomy: its possible role in the effectiveness of renal stone treatment. Urol., 11: 359-366.

Skolarikos, A.; Alivizatos, G. and Rosette J. (2006):

Extracorporeal Shock Wave Lithotripsy 25 Years Later: Complications and Their Prevention. Europ Urol., 50: 873-1128.

Stoller, M.L. (2000):

Extracorporeal shock wave lithotripsy. In: Tanago, E.A., McAninch, J.W. (eds), Smith's general urology, 15th ed. Mc Graw-Hill, San Francisco, p.321.

Stoller, M.L. (2008):

Urinary stone disease. In: Tanago EA, McAninch JW (eds), Smith's general urology, 17th ed. Mc Graw-Hill, San Francisco, 16: p.264.

Streem, S.B.; Yost, A. and Mascha, E. (1996):

Clinical implications of clinically insignificant stone fragments after extracorporeal shock wave lithotripsy. J Urol., 155: 1186.

Sun, B.Y.; Lee, Y.H.; Jiaan, B.P.; Chen, K.K.; Chang, L.S. and Chen, K.T. (1996):

Recurrence rate and risk factors for urinary calculi after extracorporeal shock wave lithotripsy. J Urol., 156: 903 - 905.

Tiselius, H.G. (1989):

Treatment of large staghorn stones and ureteral stones without anaesthesia. Scand J Urol Nephrol Suppl., 122: 25-28.

Tritrakarn, T.; Lertakyamanee, J.; Koompong, P.; Soontrapa, S.; Somprakit, P.; Tantiwong, A. and Jittapapai, S. (2000):

Both EMLA and placebo cream reduced pain during extracorporeal piezoelectric shock wave lithotripsy with the Piezolith 2300. Anesthesiology 92: 1049.

Tritrakarn, T.; Lertakyamanee, J.; Koompong, P.; Soontrapa, S.; Somprakit, S.; Tantiwong A. et al. (2000):

Both EMLA and placebo cream reduced pain during extracorporeal piezoelectric shock wave lithotripsy with the piezolith 2300. Anesthesiol., 92:1049-54.

Vickers, M.D.; Morgan, M. and Sencer, P.S. (1991):

Drugs in anaesthetic practice. In: Systemic analgesics. Butterworth-Heinmann: Oxford., p. 162-8.

Weber, A.; Koehrmann, K.U.; Denig, N.; Michel M.S. and Alken P. (1998):

What are the parameters for predictive selection of patients requiring anaesthesia for extracorporeal shock wave lithotripsy? Eur Urol., 34:85-92.

Wirth, M.P.; Theiss, M. and Frohmuller, H.G. (1992):

Primary extracorporeal shockwave lithotripsy of staghorn renal calculi. Int Urol., 48: 71-75.

Xavier, B.; Caffaratti, J.; Orsola, A.; Garat, J.M. and Vicente, G.J. (1996):

Topical anaesthesia with the EMLA cream: Application in paediatric urology. Actas Esp. Urol., 20:883-5.

Yilmaz, E.; Batislam, E.; Basar, M.M.; Tuglu, D.; Ozcan, S. and Basar H. (2005):

Effectiveness of eutectic mixture of local anaesthetic cream and occlusive dressing with low dosage of fentanyl for pain control during shock wave lithotripsy. J Endourol., 19:589-94.

Zhong, P. and Preminger, G.M. (1994):

Mechanisms of differing stone fragility in extracorporeal shock wave lithotripsy. J Endourol., 8: 263 - 268.

Zhou, Y.; Cocks, F.H.; Preminger, G.M. and Zhong, P. (2004):

The effect of treatment strategy on stone comminution efficiency in shock wave lithotripsy. J Urol., 172: 349 - 354.