



REFERENCES

ACOG: American College of Obstetricians and Gynecologists
:Practice Bulletin No. 80: premature rupture of membranes. Clinical management guidelines for obstetrician gynecologists. *Obstet Gynecol.* 2007; 109(4): 1007- 19.

ACOG: American College of Obstetricians and Gynecologists.
(ACOG): Premature rupture of membranes. Clinical management guidelines for obstetricians & gynecologists ACOG practice bulletin no 1 Int Gynecol Obstet 1998,63:75-84.

ACOG: American College of Obstetricians and Gynecologists.
Antenatal corticosteroid therapy for fetal maturation. Committee Opinion No. 402: *Obstet Gynecol.* 2008; 111(3):805-7.

ACOG: American College of Obstetricians and Gynecologists.
Premature rupture of membranes. Clinical management guidelines for obstetricians & gynecologists ACOG practice bulletin no 1 Int Gynecol Obstet 1998,63:75-84.

Alexander JM, Mercer BM, Miodovnik M, Thuranu GR, Goldenberg RL, Das AF: The impact of digital cervical examination on expectantly managed preterm rupture of membranes. *Am J Obstet Gynecol* 2000; 183:1003-1007.



- Althuisius SM, Dekker GA, Hummel P, Bekedam DJ, Van Geijn HP:** Final results of the cervical incompetence prevention Randomized Cerclage Trial (CIPRACT): therapeutic cerclage with bed rest versus bed rest alone. *Am Obstet Gynecol* 2001;185(5):1106-1112.
- Alvarez J, Bardequez A, Iffy L, Apuzzio J:** The Journal of Maternal-Fetal & Neonatal Medicine 2007; 1476-4954, 20 (12): 853 – 857.
- Aly H, Alhabashi G, Hammad TA, Owusu-Ansah S, Bathgate S, Mohamed M.** ABO phenotype and other risk factors associated with chorioamnionitis. *J Pediatr.* Jul 2008; 153(1):16-8.
- Anai, T, Tanaka Y, Hirota Y, Miyakawa I:** Vaginal fluid hCG levels for detecting premature rupture of membranes. *Obstet Gynecol* 1997; 89:261-4.
- Anderson BL, Simhan HN, Simons KM, Wiesenfeld HC.** Untreated asymptomatic group B streptococcal bacteriuria early in pregnancy and chorioamnionitis at delivery. *Am J Obstet Gynecol.* Jun 2007;196(6):524.e1-5.
- Arias F, Arias F, Rodriguez L, Rayne SC, Kraus FT.** Maternal placental vasculopathy and infection: two distinct subgroups among patients with preterm labor and preterm



ruptured membranes. Am J Obstet Gynecol 1993; 168:589–591.

Askling J, Erlandsson G, Kaijser M, Akre O, Ekbom A :Sickness in pregnancy and sex of child. Lancet 1999; **354** (9195): 2053.

Ayala AR, Serrano J, de la Luna-Olsen E, Leis MT, Bustos H, López-García R: Chorionic gonadotropin (beta-HCG) in the amniotic fluid in normal pregnant women. Arch Invest Med (Mex). 1989;20(3):223-7.

Baptisi Arthur JR: Chemical tests for the determination of ruptured membranes. Am J Obstet Gynecol,1938;35:688.

Beesley R, Trevino N, Smyth M, Gould C et al.: Vaginal fluid human chorionic gonadotropin for detection of rupture of membranes.Reprod Med.2008;53(11):823-6.

Bendon RW, Faye-Petersen O, Pavlova Z, Qureshi F, Mercer B, Miodovnik M. Fetal membrane histology in preterm premature rupture of membranes: comparison to controls, and between and antibiotic and placebo treatment. Pediatr Dev Pathol 1999; 2: 552-8.

Bentiz WE, Gould JB, Brazin MI: Atntimicrobial prevention of early onset group B streptococcal sepsis estimates of risk



reduction based on a critical literature review.
Pediatrics.2000;160-377.

Bernstein PS, Stern R, Lin N, Furgiuele J, Krmen A, Comerford-Freda M: B-Human chorionic gonadotropin in cervicovaginal secretions as apredictor or preterm delivery. Am J Obstet & Gynecol 1998; 79 : 870.

Bogdanovic G, Gric G: Stimulation of labor with premature rupture of membranes. Med Arch,2004;58(5):295-7.

Brosens J, Gordon MBH: The cytological diagnosis of ruptured membranes using Nile Blue Sulphate Stain. B J Obstet Gynecol.2005;72(3):342-46.

Buhimschi CS, Buhimschi IA, Norwitz ER: Sonographic myometrial thickness predict the latency interval of women with preterm premature rupture pf membranes and oligohydramnios. Am J Obstet Gynecol 2005;193:762-770.

Buhimschi CS, Sfakianaki AK, Hamar BG. Allow vaginal pool amniotic fluid glucose measurement is a predictive but not a sensitive marker for rupture of membranes, Am J Obstet Gynecol 2006;194(2);309-16.

Burton BK, Prins GS, Verp MS. A prospective trial of prenatal screening of down syndrome by means of maternal serum



alpha-fetoprotein, human chorionic gonadotropin and unconjugated estriol. *Am Obstet Gynecol*:1993;169:526-530.

Buyukbayrak EE, Turan C, Unal O, Dansuk R, Cengijoglu B.

Diagnostic power of the vaginal washing fluid prolactin assay as an alternative method for the diagnosis of premature rupture of membranes. *J Mat. Fetal Neonatal M*, 2004; 15:120-125.

Canavan TP, Simhan HN, Caritis S: An evidence-based approach

to the evaluation and treatment of premature rupture of membranes: part II. *Obstet Gynecol Survey* 2004; 59(9):678-689.

Carranza LS, Negrete LM, Quinzanos FC et al., Usefulness of

the qualitative detection of hCG in cervicovaginal lavage for the diagnosis of premature rupture of membranes. *Gunecol Obstet Mex*.2009;77(3):142-6.

Casey BM, Cox SM: Chorioamnionitis and endometritis .*Infect*

Dis Clin North Am 1997;11(1):203-222.

Caughey AB, Robinson JN, Norwitz ER: Contemporary

diagnosis and management of preterm premature rupture of membranes. *Rev Obstet Gynecol*,2008;1(1):11-22.

Chaiworapongsa T, Romero R, Yoshimatsu J et al.,: Soluble

adhesion molecule profile in normal pregnancy and pre-eclampsia. *J Mat. Fetal. Neonatal M*,2002;12:19-27.



Cooper AL, Vermillion ST, Soper DE: Qualitative human chorionioc gonadotropin testing of cervicovaginal washings for the detection of preterm premature rupture of membranes Am J Obstet Gynecol, 2004; 191, 593-7.

Cunningham FG, Kenneth JL, Steven LB, John C, Hauth, Larry C, Gistrop III, Katharine DW: Premature rupture of fetal membranes. Williams Obstetrics, Mcgrowhill professional publishing, 22nd edition, 2005:850-880.

Darj E, Lyrenas: S. Insulin-like growth factor binding protein-1: A quick way to detect PROM. Acta Obstet Gynecol 1998; 77(3):295-9.

De Meeus JB, Sima Ole B, Bascou V, Magnin G: Biological diagnosis or premature rupture of membranes: respective values of diamine oxidase activity compared to vaginal fluid pH(Amnicator). J Gynecol Obstet Biol reprod 1997;26(7): 730-33.

Doust RH, Mobarez Am: Collagenase activity in prevotella bivirus isolated from patients with premature rupture of membranes. National Research Center of Medical Science of I.R.IRAN 2004;18(1):61-66.

Duff P: Premature rupture of the membranes term patients. Semin Perinatol 1996;20(5):401-7.



Edwards RK. Chorioamnionitis and labor . Clin Obstet Gynecol North Am, 2005; 32:287-296.

Emma L. Stuart, Gareth S. Evans, Yu Shan Lin, and Hilary J. Powers: Reduced Collagen and Ascorbic Acid Concentrations and Increased Proteolytic Susceptibility with Prelabor Fetal Membrane Rupture in Women. Biology of Reproduction 2005; 72, 230–235 .

Eriksen NL, Parisi VM, Daoust S, Flamm B, Garite TJ, Cox SM. Fetal fibronectin: a method for detecting the presence of amniotic fluid. Obstet Gynecol 1992;80:451-4.

Esim E, Turan C, Unal O, Dansuk R, Cengliglu B: Diagnosis of premature rupture of membranes by identification of β -hCG in vaginal washing fluid. Eur J Obstet Gynecol Reprod Biol 2003;107(1):37-40.

Faro S: Post partum endometritis. Clin Perinatol. 2005; 32(3):803-814.

Flenady V, King J: Antibiotics for prelabor rupture of membranes at or near term. Cochrane Database of Systematic Reviews 2002;(3):435-40.



Fortunato SJ, Menon R, Rombarid J: Stromelysins in placental membranes and amniotic fluid with PROM. *Obstet Gynecol* 1999 ;94(3):435-40.

Fortunato SJ, Menon R, Swan KF: Inflammatory cytokines (interleukin1,6 and8 and tumor necrosis factor- α) release from cultured human fetal membranes in response to endotoxic lipopolysaccharide mirrors amniotic fluid concentration. *Am J Obstet Gynecol* 1996;174(1):855-62.

Gaucherand P, Guibaud S, Awada A, Rudigoz RC. : Comparative study of three amniotic fluid markers in premature rupture of membranes: fetal fibronectin, alpha-fetoprotein, diamino-oxidase. *Acta Obstet & Gynecol Scand* 1995; 74(2) : 118-21.

Ghidini A, Salafia CM, Kirn V. Biophysical profile in predicting acute ascending infection in preterm rupture of membranes before 32 weeks. *Obstet Gynecol.* 2000; 96 (2):201-6.

Gibbs RS, Blanco JD. Premature rupture of the membranes. *Obstet Gynecol* 1982;60(6):671-9.

Guaschino S, De Seta F, Piccoli M, Maso G, Alberico S: Aetiology of preterm labour: bacterial vaginosis *Br J Obstet Gynaecol*,2006; 113: s3: 46-51.



Hadley CB, Main DM, Gabbe SG: Risk factors for preterm premature rupture of the fetal membranes. *Am J Perinatol* 1990;374-379.

Hagberg H, Mallard C, Jacobsson B. Role of cytokines in preterm labour and brain injury. *Br.J Obstet Gynecol.* 2005;112 Suppl 1:16-8.

Halit P, Edgar S, Singer DB: Diagnosis and Discussion: Turquoise Discoloration of the Umbilical Cord and Membranes After Intra-amniotic Injection of Indigo Carmine Dye for Premature Rupture of Membranes. *Arch Pediatr Adolesc Med.* 1998;152:199-200.

Harding JE, Pang J, Knight DB, Liggins GC: DO antenatal corticosteroids help in the setting of preterm rupture of membranes? *Am J Obstet Gynecol* 2001; 184: 131-139.

Hassan S, Romero R, Hendler I. A sonographic short cervix as the only clinical manifestation of intra-amniotic infection. *J Perinat Med.* 2006;34(1):13-9.

Hitti J, Hillier SL, Agnew KJ, Krohn MA, Reisner DP, Eschenbach DA. Vaginal indicators of amniotic fluid infection in preterm labor. *Obstet Gynecol.* 2001 ; 97 (2):211-9.



Holt PG: Immune and inflammatory function in cigarette smokers. Thorax 1987;42(4):241-9.

Humphries SE: The stromelysin-1 (Matrix Metaloprotein -3) gene and risk of coronary artery disease: a candidate gene that has won the election? Tromb Haemost.Jul 2003; 90 (1):132-9.

Kafali H and Oksuzler C: Vaginal fluid urea and creatinine in diagnosis of premature rupture of membranes.Arch. Gynecol.Obstet.2007;275:157-160.

Kariman NS, Jafari E, Moghadam HRA et al: comparing the diagnostic power of qualitative and quantitative measurements β -hCG in cervicovaginal washing-fluid for the diagnosis of PROM. Journal of reprod and infert.2007;8(1):30-37.

Kayisli U, Selam B, Guzeloglu-Kayisli O, Demir R, Arici A : "Human chorionic gonadotropin contributes to maternal immunotolerance and endometrial apoptosis by regulating Fas-Fas ligand system"2003. *J. Immunol.* **171** (5): 2305–13.

Kendal-wright CE: Stretching, mechanotransduction and proinflammatory cytokines in the fetal membranes.Reprod Science 2007;12(8):35-41.



Kenyon A, Boulvain M, Neilson J: Antibiotics for preterm premature rupture of membranes. Cochrane Database syst Rev.(4):CD001058;2001.

Kilholma P, Gronross M, Erkkola R: The role of calcium, copper, iron and zinc in preterm delivery and premature rupture of fetal membranes. Obstet Gynecol invest 1994;17:194.

Kim D. Detection of premature rupture of membranes. Clinical Obstetrics and Gynecology.1991;34(4):715-722.

Kim YH, Park YW, Kwon HS, Kwon JY et al,: Vaginal fluid beta-human chorionic gonadotropin level in the diagnosis of premature rupture of membranes. Acta Obstet Gynecol Scand.2005;84(8):802-5.

Kletzky O, Rossman F, Bertolli SI, Platt LD et al.,: Dynamics of human chorionic gonadotropin, prolactin and growth hormone in serum and amniotic fluid throughout normal human pregnancy. Am J Obstet Gynecol.1985;151:878-84.

Kovavisarach E, Sermask P: Risk factors related to PROM in term pregnant women. Aust N Z J Obstet Gynecol 2000 ;40(1):30-2

Kumar d, Fung W, Moore RM, Pandey V, Fox J, Stetzer B, Mansour JM, Murcer BM, Redline RW, Moore JJ:



Alpha lipoid acid inhibit tumor necrosis factor induced remodeling and weakening of human fetal membrane Biology of reproduction 2006;74:29-34.

Larsen JW, Sever JL. Group B Streptococcus and pregnancy: a review. Am J Obstet Gynecol. 2008;198(4):440-8.

Leeson SC, Maresh MJA, Martindale EA. Detection of fetal fibronectin as a predictor of preterm delivery in high risk asymptomatic pregnancies. Br J Obstet Gynaecol 1996;103 48-53.

Lewis DF, Barrilleaux PS, Wang Y, Adair CD, Baier J, Kruger T: Detection of interleukin-6 in maternal plasma predict neonatal and infectious complications in preterm premature rupture of membranes. Am J Perinatol 2001;18:387-391.

Lewis DF, Major CA, Towers CV, Asrat T, Harding JA, Garite TJ: Effects of digital vaginal examinations on latency period in preterm premature rupture of membranes. Obstet Gynecol 1992;80:630-634.

Li Hy, Chang TS: Vaginal fluid creatinine, human chorionic gonadotropin and alpha-fetoprotein levels for detecting premature rupture of membranes. Zhonghua Yi Xue Za Zhi (Taipei).2000;63(9):686-90.



Lockwood CJ, Wein R, Chien D, Ghidini A, Alvarez M, Berkowitz RL. Fetal membrane rupture is associated with the presence of insulin-like growth factor-binding protein-1 in vaginal secretions. *Am J Obstet Gynecol* 1994;171:146-50.

Major CA, Vaciana M, Lewis DF, Morgan AM: Preterm premature rupture of membranes and abruptio placentae: Is there an association between these pregnancy complications? *Am J Obstet Gynecol* 1995;172:672-8.]

Malak TM, Bell SC. Structural characteristics of term human fetal membranes—a novel zone of extreme morphological alteration within the rupture site. *Br J Obstet Gynaecol* 1994; 101:375–386.

Malik N, Gittens L, Gonzalez D, Bardequez A, Ganesh V: Clinical amnionitis and endometritis in patients with premature rupture of membranes: endocervical prostaglandin E2 gel versus oxytocin for induction of labor. *Obstet Gynecol* 1996; 88(4pt 1):540-3.

Marowitz A, Jordan R. Midwifery management of prelabor rupture of membranes at term. *J Midwifery Womens Health*. 2007 May-Jun;52(3):199-206.



McGregor H, French J, Parker R: Prevention of premature birth by screening and treatment of common genital tract infection: results of a prospective controlled trial. *Am J Obstet Gynecol* 1995;157-167.

McParland PC, Taylor DJ: Preterm prelabor rupture of the membranes. In: *Recent advances in Obstetrics and Gynecology*: Royal Society of Medicine Press Ltd 2005; 23(3):27-38.

Medina TM, Hill DA: preterm premature rupture of membranes: Diagnosis and management. *American Family Physician*. 2006;73:659-64.

Menon R, Fortunato SJ: Fetal membranes inflammatory cytokines: A switching mechanism between preterm premature rupture of the membranes and preterm labor pathways. *J Perinate med* 2004;32(5):391-399.

Mercer BM, Arheart KL: Antimicrobial therapy in expectant management of preterm premature rupture of the membranes. *Lancet* 1995; 346:1271-1279.

Mercer BM, Goldberg RL, Moawad AH: The preterm prediction study: effect of gestational age and cause of preterm birth on subsequent obstetric outcome. *Am J Obstet Gynecol* 2000;183:738-745. (LevelII-2).



- Mercer BM, Miodovnik M, Thuranu GR, Goldenberg RL et al.,:** Antibiotic therapy for reduction of infant morbidity after preterm premature rupture of the membranes. A randomized controlled trial. JAMA 1997;278:989-995.
- Mercer BM.** Preterm premature rupture of the membranes. Obstet Gynecol 2003;101:178-93.
- Michels KB, Xue F, Colditz GA, Willett WC:** "Induced and spontaneous abortion and incidence of breast cancer among young women: a prospective cohort study". Arch. Intern. Med 2007; **167** (8): 814–20.
- Miller WC, Ford CA, Morris M:** Prevalence of chlamydial and gonococcal infections among young adults in the united states. JAMA 2004;291:2229.
- Mingione MJ, Pressman E K, Woods JR:** Prevention of PPRM: Current and future strategies. J Maternal-Fetal and Neonatal Medicine 2006;19(12):783-789.
- Modena AB, Kaihura C and Fieni S:** Prelabor rupture of the membranes: recent evidence. Acta biomed Ateneo Parmense 2004;75 Suppl 2:9-12.
- Moore RM, Mansour JM, Redline RW:** The physiology of fetal membrane rupture: insight gained from the determination



of physical properties. *Placenta* 2006;27(11-12):1037-1051.

Murtha AP, Sinclair T, Hauser ER, Swamy G, Herbert WNR, Heine RP: Maternal serum cytokines in preterm premature rupture of membranes. *Obstet Gynecol* 2007;109:121-7.

Naeye RL, Peters EC: Causes and Consequences of premature rupture of membranes . *Am J Obstet Gynecol* 1997;21:98.

Obiekwe BC, Chard T. Human chorionic gonadotropin levels in maternal blood in late pregnancy: relation to birth-weight, sex and condition of the infant at birth. *Br J Obstet Gynaecol.* 1982;89:543-6.

Osmanagaoglu MA, Unal S, Bozkaya H: Chorioamnionitis risk and neonatal outcome in preterm premature rupture of membranes. *Arch Gynecol Obstet* 2004;269:254-258.

Ozturk M, Brown N, Milunsky, Wands J: Physiological Studies of Human Chorionic Gonadotropin and Free Subunits in the Amniotic Fluid Compartment Compared to Those in Maternal Serum. *Journal of Clinical Endocrinology & Metabolism* 1988; 67(6) 1117-1121.



- Park JS, Yoon BH, Romero R, Moon JB, Young S:** The relationship between oligohydramnios and the onset of preterm premature rupture of membranes .Am J Obstet Gynecol 2001;1804:459-64.
- Parry S, Straus JF.** Mechanisms of disease: premature rupture of the fetal membranes. N Engl J Med 1998; 338:663–670.
- Pfeffer F, Casanueva E, Kamar J, Guerra A, Perichart O, Vadillo-Ortega F:** Modulation of 72-kilodalton type IV collagenase (matrix metalloproteinase-2) by ascorbic acid in culture human amnion derived cells. Biol Reprod 1998;59:326-329.
- Plessinger MA, Woods JR, Miller RK.** Pretreatment of human amnion- chorion with vitamins C and E prevents hypochlorous acid-induced damage. Am J Obstet Gynecol 2000; 183:979–985.
- Polzin WJ, Brandy K:** the etiology of premature rupture of membranes.Clin Obstet Gynecol 1998;41(4):801-6.
- Pritchard JA, Brekken AL:** Clinical and laboratory studies on severe abruptioplacentae. Am J Obstet Gynecol 1967;97:681.



Pritchard JA, Cunningham FG, Pritchard SA: On reducing the frequency of severe abruption placentae. Am J Obstet Gynecol 1991;165:1345.

Raatikainen K, Heiskanen N, Heinonen S. Transition from overweight to obesity worsens pregnancy outcome in a BMI-dependent manner. Obesity. Silver Spring Md. 2006;14(1):165-71.

Raboni S, Kaihura CT, Fieni S: Amnioscopy: Is it actual?: Acta Bio Medica Atened Paramense. 2004;75(1)59-61.

Randelović G, Kocić B, Miliković-selimović B, Maldenović-Antić S, Stojanović P, Stefanović M: High-density cervical ureaplasma urealyticum colonization in pregnant women as a risk factor for premature rupture of membranes. Vojnosanit Pregl.2006;63(8):737-741.

Richard A. McPherson, Matthew R. Pincus: Chapter.9 :Henry's Clinical Diagnosis and Management by Laboratory Methods. Hardcover, 21st edition, 2006:219-44.

Robert W, Naef W, John R, Elian R, Rick W, John C. premature rupture of membranes at 34-37 weeks' gestation: aggressive versus conservative management. Am J Obstet Gynecol 1998;178: 126-30.



- Robson MS, Turner MJ, Stronge JM, O Herlihy C:** Is amniotic fluid quantitation of value in the diagnosis and conservative management of prelabor rupture of membranes at term? *Br J Obstet Gynaecol* 1990;97:324-28.
- Rochelson BL, Rodke G, White R:** A rapid colorimetric AFP monoclonal antibody test for the diagnosis of preterm rupture of the membranes. *Obstet Gynecol* 1987; 69:163.
- Roman AS, Pernoll ML:** Late pregnancy complications in: *Current Diagnosis and Treatment. Obstet Gynecol*, 1999; 15: 292-294.
- Romero R, Chaiworapingsa T, Espinoza J:** Fetal plasma (MMP9) Matrix Metalloproteinase 9 concentrations are elevated in preterm premature rupture of membranes. *Am J Obstet Gynecol* 2002;187:125.
- Sagol S, Sagol O, Ozkal S, Asena U:** Role of apoptosis and bax protein expression in premature rupture of fetal membranes. *J Reprod Med*. 2002;47:809–815.
- Savitz DA, Blackmore CA, Throp JM.** Epidemiologic characteristics of preterm delivery: etiologic heterogeneity. *Am J Obstet Gynecol* 1991; 164:467-71.



Schoitz H: the evaporation test for detecting rupture of fetal membranes. *Acta Obstet Gynecol Scand* 1987;66:245-6.

Seaward PG, Hannah ME, Myhr TL. International multicenter term PROM study: evaluation of predictors of neonatal infection in infants born to patients with premature rupture of membranes at term. *Premature Rupture of the Membranes. Am J Obstet Gynecol.* 1998;179(3 Pt 1):635-9.

Shahin M, Raslan H: Comparative Study of Three Amniotic Fluid Markers in Premature Rupture of Membranes: Prolactin, Beta Subunit of Human Chorionic Gonadotropin, and Alpha-Fetoprotein. *Gynecol Obstet Invest.* 2007; 63 (4): 195-199.

Shalev J. Diagnosis of ruptured fetal membranes in pregnancy with the use of urograffin. *Journal of Obstetrics & Gynaecology* 1999;19(2):150-151.

Shimoya K, Moriyama A, Matsuzaki N et al. Human placental cells show enhanced production of interleukin (IL)-8 in response to lipopolysaccharide (LPS), IL-1 and tumor necrosis factor (TNF- α), but not to IL-6. *Mol. Hum. Reprod.*,1999; 5: 885–891.



Simhan HN, Caritis SN, Krohn MA, Hillier SL: The vaginal inflammatory milieu and the risk of early premature rupture of membranes. Am J Obstet Gynecol 2005;192:213-218.

Snyder M, Crawford P, Jamieson B, Neher JO. Clinical inquiries. What treatment approach to intrapartum maternal fever has the best fetal outcomes?. J Fam Pract. 2007;56(5):401-2.

Song GA, Han MS: Effect of antenatal corticosteroid and antibiotics in pregnancies complicated by premature rupture of membranes between 24 and 28 weeks of gestation. J Korean Med Sci 2005 :20(1):88-92.

Speroff L, Marc A, Fritz A: Chapter 8.(Endocrinology of pregnancy). Clinical Gynecologic endocrinology and infertility. Lippincot Williams&Wilkins. 7th edition 2005;259-319.

Steer Ph, Flint C: ABC of labour care: preterm labour and premature rupture of membranes. BMJ 1999;318(7190):1059-1062.

Steier JA, Myking OL, Bergsjø PB. Human chorionic gonadotropin in maternal plasma after induced abortion,



spontaneous abortion, and removed ectopic pregnancy. Obstet Gynaecol 1984;64:391-4.

Steier JA, Myking OL, Ulstein M. Human chorionic gonadotropin in cord blood and peripheral blood in singleton and twin pregnancies at delivery. Acta Obstet Gynaecol scand 1989;68:689-92.

Stuart EL, Evans GS, Lin YS, Powers HJ. Reduced collagen and ascorbic acid concentration and increased proteolytic susceptibility with prelabor fetal membrane rupture in women. Biol Reprod 2005;72:230-5.

Vogel I, Poul Thorsen, Allison Curry, Puk Sandager, Niels Uldbjerg: Biomarkers for the prediction of preterm delivery. Acta Obstet Gynecol 2005;84:516-525.

Waddell, Smith R: "FertilityPlus.org". Home Pregnancy Test hCG Levels and FAQ. <http://www.fertilityplus.org/faq/hpt.html>. Retrieved on 2006-06-17.

Wagner MV, Chin VP, Peters CJ, Drexler B, Newman LA: A comparison of early and delayed induction of labor with spontaneous rupture of membranes at term. Obstet Gynecol 1989;74:93-7.



Wang J: Bacterial vaginosis. Prim Care Update. Ob Gyns,2000;1:7(5):181-185.

Wiberg-Itzel E, Cnattingius S, Nordström L: Lactate determination in vaginal fluid: a new method in the diagnosis of prelabor rupture of membranes. Br J Obstet Gynaecol,2005; 112(6):754-8.

Wilkes PT, Galan H: Persistent pulmonary hypertension in a premature new born after 16 hours.Am J Obstet Gynecol.2004;191:304-308.

Yamada H, Kishida T, Negishi H, Sagawa T Fujimoto S. Silent premature rupture of membranes detected and monitored serially by an AFP kit. J Obstet Gynecol Res 1998;24:103-108.

Yoon BH, Jun JK, Romero R. Amniotic fluid inflammatory cytokines (interleukin-6, interleukin-1beta, and tumor necrosis factor-alpha), neonatal brain white matter lesions, and cerebral palsy. Am J Obstet Gynecol. 1997;177(1):19-26.

Yoon BH, kim YH, Romero R, Kim JC, Park KH: Association of oligohydramnios in women with preterm premature rupture of membranes with inflammatory response in



fetal, amniotic and maternal compartments. *Am J Obstet Gynecol* 1999;181(4).

Yost NP, Owen J, Berghella V. Effect of coitus on recurrent preterm birth. *Obstet Gynecol.* 2006;107(4):793-7.

Zaga-Clavellina V, Lopez GC, Estrada-Gutierrez G: Incubation on human chorioamniotic membranes with candida albicans induces differential synthesis and secretion of interleukin 1,interleukin6, prostaglandin E and type IV collagenase.*Myocases* 2006;49(1):6-13.