

## REFERENCES

**Arran TL** : AChR phosphorylation and indirect inhibition of AChR function in seronegative MG. *Neurology* **2003**; 59:1682.

**Auroy Y, Benhamou D, Barges L, et al** : Major complications of regional anesthesia in France: The SOS regional anesthesia hotline service. *Anesthesiology* **2002**; 97: 1274

**Atkinson P, Vijayaragavan K, Chahine M**: Differential modulation of Nav 1.7 and Nav 1.8 peripheral nerve sodium channels by the local anesthetic lidocaine. *Br J Pharmacol* **2006**; 142:576-584.

**Beer D, Thomas M L**. Caudal additives in children – solutions or problems? *Br J Anaesth* **2003**; 90(4): 487.

**Brown BL, Katz J, Benirschke K**: Functional anatomy of the sciatic nerve implications for anesthesia .*Anesthesiology* **2003**; 66: 743.

**Basbaum AI** : Spinal mechanisms of acute and persistent pain. *Reg Anesth Pain Med* **1999**; 24:59.

**Besson JM** : The neurobiology of pain. *Lancet* 2006; 353:1610. *Lancet* **2006**; 353:1610.

**Beck GP:** Anterior approach to sciatic nerve Anesthesiology 2001; 24: 222

**Benzon HT, Kim C, Benzon HP, et al :** Correlation between evoked motor response of the sciatic nerve and sensory blockade. Anesthesiology **2004**; 87 :547.

**Bruelle P, Chan V, Greger J, et al:** Lower extremity peripheral nerve blocks: Essentials of our current understanding. Reg Anesth Pain Med **2006**; 30:4.

**Bertini L ,di Benedetto P, Casati A,.** Continuous subgluteus sciatic nerve block after orthopedic foot and ankle surgery: comparison of two infusion techniques. Reg Anesth Pain Med **2002**;27:168-172.

**Bruelle P, Cuvillon P, Ripart J, et al.** Sciatic nerve block: parasacral approach. Reg Anesth Pain Med **2006**;23 :7882

**Casati A, di Benedetto P, Bertini L .** Continuous subgluteus sciatic nerve block after orthopedic foot and ankle surgery: comparison of two infusion techniques. Reg Anesth Pain Med **2002**;27: 168-172.

**Casati A, Beck GP,Ronie VM:** subgluteal approach to sciatic nerve. Anesthesiology **2007**; 24:222.

**Carlo D, Franco MD, :** A modified mid-femoral approach to the sciatic nerve block: A correlation between evoked motor response and sensory block. Anesth Analg **2007**; 105:528.

**Catterall W** : From ionic currents to molecular mechanisms. The structure and function of voltage-gated sodium channels. *Neuron* 2000; 26:13-25.

**Corning JL, Michael WE, Edward NH et al**: Major complications of regional anesthesia, *Anesthesiology* 2006; 97:1274.

**Chan VW, Nova H, Abbas S, et al**: Ultrasound examination and localization of the sciatic nerve: A volunteer study. *Anesthesiology* 2008; 104:309-314.

**Cederholm KR, Kendig JJ, Cohen EN**: Frequency-dependent conduction block :The role of nerve impulse pattern in local anesthetic potency. *Anesthesiology* 2002; 48: 111-117.

**Clarkson CW, Narchi P, Messiah A, et al**: Serious complications related to regional anesthesia. *Anesthesiology* 2007; 87:479.

**Chelly JE, Delauney L**. A new anterior approach to the sciatic nerve block. *Anesthesiology* 2001 ;91: 1655-1660.

**Chelly JE, Greger J, Casati A, et al**. Continuous lateral popliteal blocks for acute postoperative pain management after major ankle and foot surgery. *Foot Ankle Int* 2008; 23:749-752.

**Condouris GA, Goebel RH, Brady T**: Computer simulation of local anesthetic effects using a mathematical model of myelinated nerve. *J Pharmacol Expiriment* 2008; 196:737-745.

**Di Benedetto P, Bertini L, Casati A, et al :** a comparative study between neuroaxial&Sciatic lock in foot surgery. *Anesth Analg* **2007**;93:1040-1044.

**Di Benedetto P, Casati A, Bertini L, et al.** postoperative analgesia with continuous sciatic nerve block after foot surgery: a prospective, randomized comparison between the popliteal and subgluteal approaches. *Anesth Analg* **2002**;94:996-1000.

**Engleson S:** The influence of acid-base changes on central nervous system toxicity of local anaesthetic agents. I. An experimental study. *Acta Anaesthesiol Scand* **2004**; 18:79-87 .

**Eisenach S, Kehlet H:** Chronic pain as an outcome of surgery. A review of predictive factors. *Anesthesiology* **2001**; 93: 1123.

**Fornage BD:**Complications of Sonographically guided nerve block **2006**; 164:1022-1023.

**Feldman DS,Cavino BS:** Evaluation of clinical tolerance of local anaesthetic agents. *Br J Anaesth* **2002**; 47:328-331 .

**Fanelli G, Casati A, Garancini P, Torri G :** Nerve simulator and multiple injection technique for upper and lower limb blocked: Failure rate, patient acceptance, and neurologic complications. Study Group on Regional Anesthesia. *Anesth* **2000**; 88:847.

**Fink BR, Cairns AM:** Differential slowing and block of conduction by lidocaine in individual afferent myelinated and unmyelinated axons. *Anesthesiology* **2001**; 60:111-120.

**Fertleman C, Maker M, Parker K, et al:** SCN9 A mutation in paroxysmal extreme pain disorder: Allelic variants underlie distinct channel defects and phenotypes. *Neuron* **2006**; 52:767-774.

**Gaertner E, Lascurain P, Venet C, et al:** Continuous parasacral sciatic block: A radiographic study. *Anesth Analg* **2004**; 98:831.

**Gaertner E, Petit S, Cuby C, et al.** Parasacral catheter placement for lower limb surgery. *Reg Anesth* **2001** ;26:71 .

**Galindo A, Benavides O, De Munos SO, et al:** Comparison of anesthetic solutions used in lumbar and caudal peridural anesthesia. *Anesth Analg* **2002**; 57: 175-179.

**Gissen AJ, Covino BG, Gregus J:** Differential sensitivities of mammalian nerve fibers to local anesthetic agents. *Anesthesiology* **2007**; 53:467-474.

**Gray AT:** Ultrasound-guided regional anesthesia: current state of the art. *Anesthesiology* **2006**; 104:368-373.

**Goldstein A:** The effect of acoustic velocity on phantom measurements. *Ultrasound Med Biol* **2000**; 26:1133-1143.

**Gokin AP, Philip B, Strichartz GR:** Preferential block of small myelinated sensory and motor fibers by lidocaine: In vivo electrophysiology in the rat sciatic nerve. *Anesthesiology* **2001**; 95: 1441-

1454.

**Gotta J, Jayamanne A, Vaughan CW, et al:** JLO-conotoxin MrVIB selectively blocks Nav1.8 sensory neuron specific sodium channels and chronic pain behavior without motor deficits. Proc Natl Acad Sci U S A **2006**; 103: 17030-17035.

**Gullberg D:** Cell biology: The molecules that make muscle. Nature **2003**; 424:138.

**Hyvariment RT ,Zelter KH:** Successful use of a 20% lipid emulsion to resuscitate patient after a presumed bupivacaine-related cardiac arrest Anesthesiology **2006**; 105:217-218.

**Hadzic A, Vloka JD.** Peripheral nerve stimulator for unassisted nerve blockade. Anesthesiology**2007** ;84: 15281529.

**Hadzic A, Vloka JD.** A comparison of the posterior versus lateral approaches to the block of the sciatic nerve in the popliteal fossa. Anesthesiology**2007**;88:1480-1486.

**Hyvariment and Zeiter:** The site of action and active form of local anesthetics. 1. Theory and pH experiments with tertiary compounds. J Pharmacol Exp Ther **2005**; 171 :32-44.

**Hefner LV, Parks JA, Goldstein A:** Transducer beam pattern test object. J Clin Ultrasound **2005**; 8:5-10.

**Huczko EL, Gray AT, , Schafhalter-Zoppoth I:** Lateral popliteal nerve block with ultrasound guidance. Reg Anesth Pain Med **2007**; 29:507-509.

**Havner PL, Jackson JO, Waxman SG, et al:** ANA 1.7 channel mutation associated with hereditary erythromelalgia contributes to neuronal hyperexcitability and displays reduced lidocaine sensitivity. J Physiol **2001**; 581:1019-1031.

**Harold Ellis:** clinical anatomy, 3rd edition, BlackWell publishing; **2002**:272-278

**Huczko EL, Gray AT, , Schafhalter-Zoppoth I:** Lateral popliteal nerve block with ultrasound guidance. Reg Anesth Pain Med **2007**; 29:507-509.

**Iskandar H, Benard A, Ruel-Raymond J et al .** The analgesic effect of interscalene block using clonidine as an analgesic for shoulder arthroscopy. Anesth Analg **2003**; 96(1): 260–2.

**Jonsson M, Dabrowski M, Gurley DA, et al:** Activation and inhibition of human muscular and neuronal nicotinic acetylcholine receptors by succinylcholine. Anesthesiology **2006**; 104:203

**Jonsson GE, McCormick DJ, De Aizpurua HJ, et al:**  $\alpha$ -Bungarotoxin binds to human  $\alpha$ 1 acetylcholine receptor  $\alpha$ -subunit peptide 2006; 54: 154-189

**Julius D, Basbaum AI:** Molecular mechanisms of nociception. Nature **2003**; 413:203.

**Kehlet H, Holte K:** Effect of postoperative analgesia on surgical outcome. Br J Anaesth **2005**; 87:62

**Kitain E, Vloka JD, Hadzic A.** Anatomic considerations for sciatic nerve block in the popliteal fossa through the lateral approach. Reg Anesth 2006 ;21 :414-418.

**Klein SM, Greengrass RA, Steele SM, et al:** A comparison of 0.5% bupivacaine, 0.5% ropivacaine, and 0.750/0 ropivacaine for interscalene brachial plexus block. Anesth Analg **2007**; 87:1316-1319.

**Lanz E, Crawford RD, Scurlock JE:** Severe hypoxia and acidosis following local anesthetic – induced convulsions. Anesthesiology 2004; 53: 259-260.

**Lanie K, Strichartz GR, Raymond SA:** On the mechanisms of potentiation of local anesthetics by bicarbonate buffer: Drug structure-activity studies on isolated peripheral nerve. Anesth Analg **2008**; 76:131-143.

**Lukas RJ, Bencherif M:** Heterogeneity and regulation of nicotinic acetylcholine receptors. Int Rev Neurobiol **2003**; 34:45.

**Mansour NY.** Reevaluating the sciatic nerve block: another landmark for consideration. Reg Anesth **2001**; 18:322-323.

**McCammon:** Effect of added epinephrine on spinal anesthesia with lidocaine. Anesth Analg **2002**; 60:417-420

**Mansour NY:** combination of sciatic nerve block with Ither nerve blocks in lower limb surgery ,Anesthesiology :**2005**; 8 7: 547-552

**Missias AC, Chu GC, Klocke BJ, et al:** Maturation of the acetylcholine receptor in skeletal muscle: Regulation of the AChR  $\gamma$ -to- $\epsilon$  switch. Dev Biol **2004**; 179:223.

**Morishima SD:** The pH-dependent rate of action of local anesthetics on the node of Ranvier. J Gen Physiol **2000**; 69:475-496.

**Moore YI, Mather LE:** Pharmacology of local anaesthetic agents. Pharmacokinetics of local anaesthetic agents. Br J Anaesth **2002**; 47:213-224

**Morton RJ, :** Successful resuscitation of a patient with ropivacaine-induced asystole after axillary plexus block using lipid infusion. Anaesthesia **2003**; 61:800-801.

**Mc Intyre :** Kinetics of Anesthetic Binding to Sodium Channels: Role of pKa. Boston, Massachusetts Institute of Technology, **2005**;34: 123-311

**McCarthy MP, Stroud RM:** Conformational states of the nicotinic acetylcholine receptor from *Torpedo californica* induced by the binding of agonist, antagonists, and local anesthetics. Equilibrium measurements using tritium-hydrogen exchange. Biochemistry **2006**; 28:40.

**McCammon JJ:** An SCN9A channelopathy causes congenital inability to

experience pain. Nature **2006**; 444:894-898.

**Morgan JR, Mikhail : Clinical anesthesiology, 3<sup>rd</sup> edition, Appleton& 2006: 16:269-227**

**Near JM, Brudie WA, Berde CB:** Convulsions associated with pediatric regional anesthesia. Anesth Analg 2003; 75:164-166.

**Nolsoe C, Nielsen L, Torp-Pedersen S, Holm HH:** Major complications and deaths due to interventional ultrasonography: A review of 8000 cases. J Clin Ultrasound **2007**;18:179-184.

**Ojj DF and Ojj SA :** Effects of concentration of local anaesthetic drugs in extradural block. Br J Anaesth **2006**; 52:1033-1037.

**Pigg H, Garancini P, Torri G:** Nerve stimulator and multiple injection technique for upper and lower limb blockade: Failure rate, patient acceptance, and neurologic complications. Study Group on Regional Anesthesia. Anesth Analg **2004**; 88:847.

**Pianezza A, Gilbert ML, Minville V, et al:** A modified mid-femoral approach to the sciatic nerve block: A correlation between evoked motor response and sensory block. Anesth Analg **2007**; 105:528.

**Paqueron X, Bouaziz B, Macalou D, et al:** The lateral approach to the sciatic nerve at the popliteal fossa: One or two injections? Anesth Analg **2007**; 89:1221.

**Pedersen SE, Cohen JB:** d-Tubocurarine binding sites are located at alpha-gamma and alpha-delta subunit interfaces of the nicotinic acetylcholine receptor. *Proc Natl Acad Sci USA* **2008**; 87:2785 .

**PennockR Brody T, Larnner J, et al :** Physiology and biochemistry of the peripheral nervous system. **2004**; 27:122-125.

**Pautrell GD :** The systemic absorption of local analgesic drugs. *Br J Exp Anaesth* **2005**; 37:396.

**Pither C, Raj P, Ford D:** The use of peripheral nerve stilmulators for regional anesthesia: a review of experimental characteristics, technique, and clinical applications. *Reg Anesth***2007**; 10:49-58.

**Proc HT, Lacazette E, Courtet M, et al :** Synapses form in skeletal muscles lacking neuregulin receptors. *Science* **2005**; 308:1920.

**Pautrell GD:** Lidocaine constricts or dilates rat arterioles in a dose-dependent manner. *Anesthesiology* **2005**; 62:141-144.

**Raymond SA:** Lack of size-related differential sensitivity to equilibrium conduction block among mammalian myelinated axons exposed to Benoxinate*Anesth Analg* **2006**; 66:948-953 .

**Reynolds SA:** Pharmacokinetics and efficacy of ropivacaine for continuous epidural infusion in neonates and infants. *Pediatr Anesth* **2001**; 15:739-749.

**Rongstad K, Mann RA, Prieskom D, et al.** Popliteal sciatic nerve block for postoperative analgesia. *Foot Ankle Int* **1996**;17:378-382.

**Rorie DK, Byer DE, Nelson DO, et al.** Assessment of block of the sciatic nerve in the popliteal fossa. *Anesth Analg* **2007**;59:3 71-3 76.

**Raines DE:** Anesthesia and nonanesthetic volatile compounds have dissimilar activities on nicotinic acetylcholine receptor desensitization kinetics. *Anesthesiology* **1996**; 84:663.

**Ritchie JM, Rogart RB:** Density of sodium channels in mammalian myelinated nerve fibers and nature of the axonal membrane under the myelin sheath. *Proc Natl Acad Sci USA* **2006**; 74:2

**Ritchie B, Greengard P :** The effect of the nerve sheath on the action of local anesthetics. *J Pharmacol Exp Ther* **2002**; 150:160.

**Sia S, Sarro F, Lepri A et al.** The effect of exogenous epinephrine on the incidence of hypotensive/bradycardic events during shoulder surgery in the sitting position during interscalene block. *Anesth Analg* **2003**; 97(2): 583–8.

**Serenire, principi RT :** Interrelationships of hepatic blood flow and blood levels of lidocaine in man. *Circulation* **2003**; 43:205-211.

**Sandu F, Capan G:** A prospective, randomized comparison between ultrasound and nerve stimulation guidance for sciatic nerve block *Anesthesiology* **2007**; 106:992.

**Singelyn F, Aye F, Gouverneur JM:** Continuous popliteal sciatic nerve block: an original technique to provide postoperative analgesia after foot surgery. *Anesth Analg* **2004**;84:383-386.

**Sinha A, Chan VW:** Ultrasound imaging for popliteal sciatic nerve block. *Reg Anesth Pain Med* **2004**; 29: 130-134.

**Schafhalter-Zoppoth I, Younger SJ, Collins AB, Gray A T:** The "seesaw" sign: improved sonographic identification of the sciatic nerve. *Anesthesiology* **2004**; 101 :141

**Sinsky ER and Patel QS:** Micromolar lidocaine selectively blocks propagating ectopic impulses at a distance from their site of origin. *Pain* **2002**; 99:333-340.

**Taylor AC, Fink BR, Karin DS:** A comparison of 0.5% bupivacaine, 0.5% ropivacaine, and 0.75% ropivacaine for interscalene brachial plexus block. *Anesth Analg* 1998; 87:1316-1319.

**Takaman T: Local anesthetics:** Hydrophilic and hydrophobic pathways for the drug-receptor reaction. *J Gen Physiol* **1975**; 69:497-515.

**Tansey MG, Chu GC, Merlie JP:** ARIA/HRG regulates AChR epsilon subunit gene expression at the neuromuscular synapse via activation of phosphatidylinositol 3-kinase and Ras/MAPK pathway. *J Cell Biol* **2008**; 134:46

**Tsui BC, Finucane BT:** The importance of ultrasound landmarks: a "traceback" approach using the popliteal blood vessels for identification of the sciatic nerve. *Reg Anesth Pain Med* **2006**; 31 :481-482

.

**Van oss AD:** Fundamental properties of local anesthetics. I. The dependence of Carticaine's ionization and octanol:buffer partitioning on solvent and temperature. *Anesth Analg* **2004**; 66:159-165.

**Van Eistraete AC, Poey C, Lebrun T, et al:** role of sciatic nerve block in vascular surgery:2008:16:354-367 Valade N, Ripart J, Nouvellon E, et al: Does sciatic parasacral injection spread to the obturator nerve? An anatomic study. *Anesth Analg* **2008**; 106:664.

**Vloka JD, Hadzic A, April E, Thys DM:** The division of the sciatic nerve in the popliteal fossa: Anatomical implications for popliteal nerve blockade. *Anesth Analg* **2001**; 92:215.

**Wildsmith WE:** Structural elements which determine local anesthetic activity. In: Schwartz GR, ed. *Handbook of Experimental Pharmacology: Local Anesthetics*, Berlin: Springer-Verlag; **2003** :53-89.

**WeberY, Jain P:** Serious complications related to regional anesthesia. *Anesthesiology* **2005**; 87:479.

**Wild smith J, Armitage H:** Continuous parasacral sciatic block: A radiographic study. *Anesth Analg* 2003; 98:831.

**Winnie AP :**sciatic block as only method in hip surgery: **2006**;54: 861-892.

**Wilson GT, :** Pharmacology of local anaesthetic agents. Pharmacokinetics of local anaesthetic agents. *Br J Anaesth* **2001**; 47:213-224.

**Wilson DJ:** Comparison of anesthetic solutions used in lumbar and caudal peridural anesthesia. *Anesth Analg* **2001**; 57:175-179.

**Winnie AP, Arthur GR, Strichartz GR:** Fundamental properties of local anesthetics. I. The dependence of lidocaine's ionization and octanol:buffer partitioning on solvent and temperature. *Anesth Analg* **2003**; 66:159-165. 43.

**Yamaoka K, Vogel SM, Seyama I:** Na<sup>+</sup> channel pharmacology and molecular mechanisms of gating. *Curr Pharm Des* **2006**; 12:429.

**Zetlaoui PJ, Bouaziz H:** Lateral approach to the sciatic nerve in the popliteal fossa. *Anesth ,Analg* **2002**; 87:79.







