

References

- **Adamcová M (2003):** Troponins in children and neonates. *Acta Paediatrica*; 92(12):1373–5.
- **Adams J, Schectmann K B and Landty S (1994):** Comptible detection of acute myocardial infarction. *Clin chem*; 40: 1291-95.
- **Adams JE, Bodor GS, Davila-Roman VG. (1993):** Cardiac troponin I. a marker with high specificity for cardiac injury (see comments). *Circulation*; 88: 101-106.
- **Adcock LM and Papile Lu-Ann (2008):** Perinatal asphyxia. In cloherty J P, Eichenwald EC and Stark AR(eds). *Manual of Neonatal Care*, 6th ed. Philadelphia. Wolters Kluwer. P 518-28.
- **Adcock LM and Papile Lu-Ann (2008):** Perinatal asphyxia. In cloherty J P, Eichenwald EC and Stark AR(eds). *Manual of Neonatal Care*, 6th ed. Philadelphia. Wolters Kluwer. P 518-28.
- **Aigrain EJ and Anderson BJ (2006):**Pain control: Non-steroidal anti-inflammatory agents. *Semin Fetal Neonatal Med*; 11(4): 251-59.
- **Alshuler G (1996):** Role of placenta in perinatal pathology (revisted). *Pediatric Path Lab Med*; 16:207.
- **American Academy of Pediatrics And American College Of obstetricians, Quotedfrom (2002) :** Guidelines for Perinatal Care.
- **Anteman E.M, Gradizenc And Sacks D (1995):** Evaluation of rapid assay or detection of Ctn 1. *J Am Chem*, 273-1279.

- **Anthony M, Costello L, and Manadhar S (2000):** Risk factors for neonatal encephalopathy in Kathmandu, Nepal, A developing country unmatched case-control study. *BMJ*; 320: 1229-1250.
- **Apgar V (1953):** A proposal for a new method of evaluation of the newborn infant. *Curr. Res. Anesth. Analg*; 32 (4): 260-67.
- **Araujo K, da Silva J, Sanudo A, Kopelman B(2004):**plasma concentration of cardiac troponin I in newborn infants.*Clin Chem* :50:1717-1718
- **Arisoy AE(2004).:-** Cord blood cardiac Troponin I as an early predictor of short term outcome in perinatal hypoxia. *Biol Neonate*;86:131–137
- **Arvidson J and Hagberg B (1990):** Delayed onset dyskinetic cerebral palsy (a late effect of perinatal asphyxia). *Acta Pediatr Scand*;79: 1121.
- **Aurora s and Snyder Y E (2004):** Perinatal asphyxia. *Manual of Neonatal Care*. Cloherty j. P. (eds); 4th edition, Little Brown and Lippincott-Raven Co.27: 536-555.
- **Ballard RA, Hansen TN and Corbet A (2005):** Respiratory failure in the term infant in: Taeusch HW, Ballard RA, Gleason CA (eds). In *Avery's disease of newborn*. 8th edition, pp.705-722.
- **Barkovich AJ, Westmark K, Partridge C (1995).** Perinatal asphyxia: MR findings in the first 10 days. *AJNR Am J Neuroradiol*;16:427–438.
- **Barkovich and Truwit (2008):** "Brain damage from perinatal asphyxia correlation of MR findings with gestational age -- Barkovich and Truwit

- 11 (6): 1087 -- American Journal of Neuroradiology"Retrieved on 2008-03-27.
- **Basu P, Som S, Choudhuri N and Das H(2008):** Correlation between apgar score and urinary uric acid to creatinine ratio in perinatal asphyxia *indian journal of clinical biochemistry*, 2008 / 23 (4) 361-364 .
 - **Bear MF, Connors BW and Paradiso MA (2001):** Mitochondria, free radicals and neurodegeneration. **In:** Bear MF, Connors B W, Paradiso MA (eds). *Neuroscience Exploring The Brain*. 2nd ed. Philadelphia. Lippincott Williams & Wilkins. P 36.
 - **Berger R and Garnier Y (1999):** Pathophysiology of perinatal brain damage. *Brain Res Rev Aug*; 30(2): 107-34[Medline]
 - **Biagioni E (2001):** Combined use of electroencephalogram and magnetic resonance imaging in full-term neonates with acute encephalopathy. *Pediatrics* ; 107:461.
 - **Blennow M, Sayman K and Lives P et al (2001):** Brain specific protein in cerebrospinal fluid of severely asphyxiated newborn infants. *Acta Pediatr*; 90: 1171-5.
 - **Carter BS, Haverkamp AD and Nerenstein GB (1993):** The definition of acute perinatal asphyxia. **In:** *Clinics in Perinatology (perinatal asphyxia)*. Shankaran S. (eds), 4th edition, W. B. Saunders Co., Philadelphia; 20(2): 287.
 - **Carteux P, Cohen H and Check J et al (2003):** Evalution and development of potentially better practices for the prevention of brain

hemorrhage and ischemic brain injury in very low birth weight infants. Pediatrics: 111(4, Pt2): 489 – 96.

- **Casey BM, McIntire DD and Leveno KJ (2001):** The continuing value of the Apgar score for the assessment of newborn infants. NEJM: 344(7): 467-71.
- **Clancy RR and Legido A (1991):** Postnatal epilepsy after EEG-confirmed neonatal seizures. Epilepsia Jan-Feb; 32(1): 69-76[Medline].
- **Courtesy A.C.Van Huffless (2006):** Development and disorders of organ systems – The Central Nervous System - Hypoxic Ischemic Encephalopathy. From: Fanaroff and Martin's (2006): Neonatal-Perinatal Medicine, 8th ed. Mosby, An Imprint of Elsevier. Hummel T, Shreiner J, Polizzano F, Nikelsky J, Owens KO. USA.
- **Cowan F (2000):** Outcome after intrapartum asphyxia in term infants. Semin Neonatol ; 5:127.
- **Cranmer H and Shannon M (2007):** Pediatric hypoglycemia. e-Medicin, Emergency Medicine Pediatrics. Availble at <http://www.emedicine.com/emerg/topic384.htm>.
- **Dambinova S (2007):** Diagnostic challenges in traumatic brain injury. IVD Technology-, 4:41-8.
- **Davies G A (2000):** Antenatal fetal assessment. J Soc Obstet Gynaecol Can; 22(6): 456-62.

- **De Menezes M S and Shaw D W (2006):** Hypoxic – Ischemic Brain Injury in the Newborn. E-Medicine. Pediatrics. Neurology. Available at <http://www.emedicine.com/neuro/topic696.htm>.
- **Deulofeut M and Sola A (2005):** Injury to the developing brain in term and preterm infants, mechanisms of injury and potential for prevention and repair. J Arab Neonatal Forum; 2:34-47.
- **Dragunow M, Sirimanne E, and Lawlor PA (1992):** A role for calcitonin gene related peptide during recovery from hypoxic ischemic brain injury in the infant rat. Brain Res;14:267-272.
- **El Metwally D (2006):** Hypoxic ischemic encephalopathy, still haunting! Neuropediatrics: SI (37).
- **Fanaroff A and Martin RJ (2001):** Asphyxiated thoracic dystrophy In: Neonatal perinatal medicine 3rd edition, Chuchill-livingstone, London, p 465-466.
- **Ferriero DM (2004):** Neonatal brain injury. N Engl J. Med; 351 (19): 1985-95.
- **Finster M and Wood M (2005):** The Apgar score has survived the test of time. Anesthesiology: 102 (4): 855-57.
- **Florio P, Luisi S and Moataza B et al (2007):** High urinary concentrations of activin A in asphxiated full-term newborns with moderate of severe hypoxic ischemic encephalopathy. Clin Chem J; 10:1373.

- **Franck LS, Quinn D and Zahr L (2000):** Effect of less frequent bathing of preterm infants on skin flora and pathogen colonization. *J Obstet Gynecol Neonatal Nurs*; 29(6):584-89.
- **Gaze DC, Collinson PO,(2006):** Interpretation of cardiac troponin
- **Golianu B, Krane E and Seybold J et al (2007):** Non-pharmacological techniques for pain management in neonates. *Sem Perinat*; 31(5): 318-22.
- **Gomella T, Cunningham M and Eyal F (2003):** Perinatal asphyxia. In: *Neonatology (Management, procedures, on call problems, diseases and drugs)*. Gomella (Ed), Appleton and Lang, California, p : 516.
- **Granelli SL and McGrath JM (2004):** Neonatal seizures: diagnosis. Pharmacologic interventions, and outcomes. *J perinat Neonatal Nurs*; 18(3):275-87.
- **Grant A (1989):** Monitoring the fetus during labour. In: *Effective care in pregnancy and childhood*. Chalmers A, Enkin M and Keirse MJ(eds), vol(2), Oxford University press, Oxford, p: 848.
- **Gray PH, Tudehope DI, Masel JP (1993):** Perinatal hypoxic-ischaemic brain injury: Prediction of outcome. *Dev Med Child Neurol* 35:965-973.
- **Griessen G (1992):** Effect of cerebral blood flow and cerebrovascular autoregulation on distribution and type of cerebral palsy. *Brain patho*;2: 223
- **Grojean-Gibondi F (1988):** Cerebral Doppler in neonates by J. Messer. *Etude de la velocimetrie arterielle intracerebrale chez le nouveau-ne*

normal, a terme, a J1 et J3. Thesis, Louis Pasteur University, Strasbourg, p: 109.

- **Gunn AJ and Bennet L (2001):** Is temperature important in delivery room resuscitation? *Semin Neonatol*: 6(3):241-9.
- **Halliday H (1998):** Pulmonary disorders and apnea. In: Forfar and Arneil's textbook of pediatrics; 5th edition; by Compbell A. and Neil McIntoch (eds.) Churchill livingstone; P: 198-221.
- **Hill A and Volpe J (1992):** Hypoxic ischemic encephalopathy and intracranial hemorrhage. In: Textbook of Neonatology. Robertson N (ed), 2nd edition, Churchill Livingstone, London, pp: 1061.
- **Hill A, and Volpe J (1989):** Hypoxic ischemic encephalopathy of the newborn. In: Pediatric Neurology, Principles and Practice. Swaiman KF (ed), CV Mosby, St. Louis; 1: 373.
- **Holmes GL and Lombroso CT (1993):** Prognostic value of background pattern in the neonatal EEG. *J Clin Neurophysiol* ; 10:323.
- **Ismail I, Awad H and El-Badawi N (1997):** Increased serum creatine kinase isoenzyme and urinary alpha-microglobulin in perinatal asphyxia. *Ain Shams Medical Journal*; 47:31.
- **Johnston MV (2004):** The biochemical neurotoxic cascade in hypoxic-ischemic injury. In: Donn SM, et al (ed.) Birth Asphyxia and the Brain: Basic Science and Clinical Implications, Armonk, NY: Futura; :51.

- **Johnston MV, Trescher WH and Ishida A, et al (2001):** Neurobiology of hypoxic-ischemic injury in the developing brain. *Pediatr Res*; 49(6):735-741.
- **Kanik E, et al (2009):** Clinics of Pediatrics, Tepecik Training and Research Hospital, Yenisehir, Izmir, Turkey *The Journal of Maternal-Fetal and Neonatal Medicine*, March 2009; 22(3): 239–242
- **Kattwinkel J ,Short J and Shaven L et al (2006):** **Resuscitation** involving meconium. **In:** Kattwinkel J and Short J(eds). *Textbook of neonatal resuscitation*, 5th Ed, Am Acad Ped ;(2):3-10.
- **Katus H.A., Remppis S. and Neumann F.J (1991):** Diagnostic deficiency of troponin T measurements in acute myocardial damage. *Circulation*, 83: 902-912.
- **Khreisat WH and Habahbeh Z (2005):** Risk factors of birth asphyxia: A study at Prince Ali Ben Al-Hussein Hospital, Jordan, *Pak J Med Sci*; 21 (1): 30 —32.
- **Kilpatrick SJ (2004):** Fetal heart rate monitoring. *Internet J Obstetric Anesth*; 13(2): 135.
- **Kliegman RM (1996):** The fetus and neonatal infant (hypoxia-ischemia). **In:** *Nelson Textbook of Pediatrics*. Nelson WE, Behrman RE, Kliegman RM and Arvin (eds); 15th ed; WB Saunders Co; Philadelphia; 11: 469-71.

- **Kleigman, Behrman, Jensen and Stanton (2008):** The Cardiovascular System. Echocardiography. In Nelson textbook of pediatrics, 18th edition, Saunders Elsevier; Philadelphia; library of Congress Cataloging, p: 1869.
- **Klinger G, Beyene, J and Shah, P et al (2005):** Do hyperoxaemia and hypocapnia add to the risk of brain injury after intrapartum asphyxia? Arch. Dis. Child. Fet. Neon. Ed; 90(1): 49-52.
- **Knusel B, Michel P and Schwaber J (1990):** Selective and non-selective stimulation of central cholinergic dopaminergic development in vitro by nerve growth factor and basic fibroblast growth factor. J Neuroscience; 10:558.
- **Levene MI (1998):** Birth asphyxia and Hypoxic ischemic encephalopathy. In: Forfar textbook of pediatrics. Campbell A.G.M. and McIntoch Neil (eds); 5th edition, Churchill Livingstone, New York, 5: 123-131.
- **Levene MI, Evans DJ, Mason S, Brown J (1999):** An international network for evaluating neuroprotective therapy after severe birth asphyxia. Semin Perinatol; 23:226-233.
- **Liu J, Cao HY, He CY (1998):** Hemodynamics of multiple organs in asphyxiated neonates. Zhonghua Er Ke Za Zhi; 36:69-73.
- **Lives P, Talvic R, Talvic T (1998):** Changes in Doppler ultrasonography in asphyxiated term infants with HIE. Acta Paediatr; 87:680-4.
- **Lokesh L, Kumar P and Murki S (2004):** A randomized controlled trial of sodium bicarbonate in neonatal resuscitation-effect on

immediate outcome. Resuscitation ; 60(2):219-23.

- **Lou H, Henricksen L and Bruhn B (1990):** Focal cerebral dysfunction in developmental learning disabilities. Lancet;335:8.
- **Mair G., Dient L.F. and Puschendorf B (1992):** Cardiac specific troponin T in the diagnosis of myocardial injury. Clin Lab Med; 29: 27-35.
- **Majnemer A and Mazer B (1998):** Neurologic evaluation of the newborn infant: definition and psychometric properties. Dev Med Child Neurol Oct; 40(10): 708-15[Medline].
- **Maschmann J, Heinemann Mk and Ziemer G et al (2000):** valuation of protein s-100 serum concentrations in heralthy newborns and seven newborns with perinatal acidosis. Acta Pediatr; 89-553-55.
- **McAuliffe F., K. Mears, S. Fleming, H. Grimes, and J. Morrison, (2004),** “Fetal cardiac troponin I in relation to intrapartum events and umbilical artery pH,” American Journal of Perinatology, vol. 21, no. 3, pp. 147–152.
- **McGuire W, McEwan P and Fowlie P (2004):** Care in the early newborn period. BMJ; 329:1087-89.
- **Mcintosh N and Stenson B (2003):** The newborn. In: Mcintosh N, Helms PJ and Smyth RL (eds). Forfar and Ameil’s Textbook of pediatrics. 6th ed. Edinburgh. Churchill Livingstone, P 177-391.
measurements in neonates—the devil is in the details. Biology of the
- **Meinerts T. and Hamm C.W. (1998):** Rapid testing for cardiac

troponins in patients with acute chest pain in the emergency room. European Heart Journal; 19:973-974.

- **Menache CC (2002):** Prognostic value of neonatal discontinuous EEG. *Pediatr Neurol* ; 27:93.
- **Miller SP, Latal B and Clark H et al (2004):** Clinical signs [redict 30 month neurodevelopmental outcome after neonatal encephalopathy. *Am J Obstetrics Gynecology*; 190:93-9.
- **Nagdyman N, komen W and Ko-H-k et al, (2001):** Early biochemical indicators of hypoxic-Ischemic encephalopathy after birth asphyxia. *Pediatric Res*: 49:502-6.
- **Nagdyman N, Komen W, Ko HK, Muller C, Obladen M (2001):** Early biochemical indicators of hypoxic-ischemic encephalopathy after birth asphyxia. *Pediatric Research.*; 49(4):502-6.
- **Nelson K and Emery E (1993):** Birth asphyxia and the neonatal brain: What do we know and when do we know it? In: *Clinics in Perinatology*. Shankaran S. (ed), E. B. Saunders, Philadelphia, 20(2): 327.
- **Nelson KB and Leviton A (1991):** How much of neonatal encephalopathy is due to birth asphyxia? *Am. J.Dis. Child.*, 145, 1325–31.

Neonate.; 89(3):194–6.
- **Nioka S, Izzetoglu M and Michael J etal (2005):** Fetal transabdominal pulse oximeter studies using a hypoxic sheep model *J Mater Fet Neonat Med*; 17(6): 393-99.

- **Parer JT and King T (2000):** Fetal heart rate monitoring: Is it salvageable? *Am J Obstet Gynecol*: 182:982-87.
- **Peebles DM (1993):** Near infrared spectroscopy and intrapartum fetal surveillance. Abstract page 5, 2nd World Congress of Perinatology Medicine.
- **Perlman J, Tack E, and Martin R (1989):** Acute systemic organ injury in term infants after asphyxia. *Am J. Dis Child*;143:617.
- **Perlman JM (1997):** Intrapartum hypoxic ischemic cerebral injury and subsequent cerebral palsy. *Pediatrics*;99: 851.
- **Perlman JM (2006):** Summary proceedings from the neurology group on hypoxic-ischemic encephalopathy. *Pediatrics*; 117(3):28-33.
- **Perlman JM (2006):** Summary proceedings from the neurology group on hypoxic ischemic encephalopathy. *Pediatrics*;117:S28-33.
- **Pignotti MS, Indolfi G and Ciuti R et al (2005):** Perinatal asphyxia and inadvertent neonatal intoxication from local anaesthetics given to the mother during labour. *BMJ*:330:34-35.
- **Plebani M. and Zaniomotto M (1999):** Cardiac markers, present and Feature. *Int. Clin. Lab*, 20: 56-63.
- **Raju TN, Rosenkrantz T and Windle ML et al (2006):** Hypoxic Ischemic Encephalopathy. *e-medicine, pediatrics, Neonatology*. Available at <http://e.medicine.com/ped/topic 149.htm>.

- **Renato D, Procinoy S, and Olvereiva L (1988):** Neonatal cortical adrenal function in birth asphyxia. *Acta Pediatr Scand*; 77: 671.
- **Rennie J (1998):** Neonatal neurology. In: Forfar and Arneil's textbook of pediatrics (5th edition), by A. G. M. Compbell and Neil. McIntoch (eds.), Churchill Living Store; pp249:52.131.
- **Rivkin M, Anderson M and Kaye E (1992):** Neonatal idiopathic cerebral venous thrombosis. *Ann Neurology*; 32-51.
- **Robertson NRC (1993):** Resuscitation of the newborn. In: Manual of neonatal Intensive Care. Arnold E (ed), 3rd edition. Little Brown Co, London; 7:49-67.
- **Rohl L, Ostergaard L and Simonsen Z et al (2001):** Viability thresholds of ischemic penumbra of hyperacute stroke defined by perfusion-weighted MRI and apparent diffusion coefficient. *Stroke*; 32:1140.
- **Rortocil A, Miklica J and Kucera M (2000):** Continuous monitoring of fetal oxygen saturation (FSpO₂) using intrapartum fetal pulse oximetry (IFPO) in the diagnosis of acute fetal hypoxia. *Cesk Gynekol*; 65: 224-30.
- **Sarnat HB and Sarnat MS (1976):** Neonatal encephalopathy following fetal distress. A clinical and electroencephalographic study *Arch Neurol*; 33(10): 696-705.
- **Sasidharan P (1992):** Breathing pattern abnormalities in full term asphyxiated newborn infants. *Arch. Dis. Child.*; 67: 440-442.

- **Saugstad OD, Ramji S and Vento M (2005):** Resuscitation of depressed newborn infants with ambient air or pure oxygen: A meta-analysis. *Biol Neonate*; 87(1):27-34.
- **Scher M (2001):** Perinatal asphyxia: Timing and mechanisms of injury in neonatal encephalopathy. *Cure Neurol Neurosci Rep*; 1:175-84.
- **Scher MS (2003):** Uncoupling of EEG-clinical neonatal seizures after antiepileptic drug use. *Pediatr Neurol* 2003; 28:277.
- **Seibert JJ, McCowan TC. Chaddock WM (1989):** Duplex pulsed Doppler US versus intracranial pressure in the neonate: Clinical and experimental studies. *Radiology* 171:155-159, 1989.
- **Shah P, Riphagen, S and Beyene J (2004):** Multiorgan dysfunction in infants with post-asphyxial hypoxic-ischemic encephalopathy. *Arch. Dis. Child Fet. Neon. Ed*; 89(2):152-155.
- **Shalak LF, Laptook AR, Velophi SC, and Perlman M (2003):** a-EEG coupled with an early neurologic examination enhances prediction of term infants at risk for persistent encephalopathy.
- **Shankaran S (2002):** The postnatal management of the asphyxiated term infant. *Clin perinatol*; 29:675-92.
- **Siegel MJ, Shackelford GD, Perlman JM, (1984):** Hypoxic-ischemic encephalopathy in term infants: Diagnosis and prognosis evaluated by ultrasound. *Radiology* 152:395-399.

- **Snyder EY and Cloherty JP (1998):** Perinatal asphyxia. In: Manual of neonatal Care. Cloherty J. P. (eds); 4th edition, Little Brown and Lippincott-Raven Co., Boston, pp: 515.
- **Stills Jack (2004):** Perinatal asphyxia. In: Management, Procedures, On call problems, Disease & Drugs in Neonatology. Gomella TL., Cunningham MD., Eyal FG and Zenk KE (eds); 5th edition, The McGraw Hill, USA, 73: 512-523.
- **Stoll BJ and Kliegman RM (2007):** Hypoxia-Ischemia **In:** Kliegman RM, Behrman RE. Jenson HB and Stanton BF (eds). Nelson Textbook of Pediatrics. 18th ed. Philadelphia. WB Saunders, P718.
- **Stoll JB and Kliegman MR (2004):** Hypoxia ischemia. In: Nelson textbook of pediatrics. Behrman R., Robert M., Kleigman RM and Jenson HB (eds); 17th edition, W.B. Saunders Philadelphia 88: 561-568.
- **Strachan B, Sahota D and Wijngaarden WJ et al (2000):** The fetal electro-cardiogram: relationship with acidemia at delivery. Am J Obstet Gynecol-, 182: 603-6.
- **Thompson DG (1994):**Consequences of perinatal asphyxia. AACN Clinical Issues; 5(3): 242-245.
- **Thoresen M, Penrice J, Lorek A, Cady EB, Wylezinska M, Kirkbride V, Cooper CE, Brown GC, Edwards AD, Wyatt JS, Reynolds EOR(1995):** Mild hypothermia after severe transient hypoxia-ischemia ameliorates delayed cerebral energy failure in newborn piglets. Pediatr Res;37:667-670.

- **Toet MC, Lemmers PM, van-Schelwn LJ et al (2006):** **Cerebral** oxygenation and electrical activity after birth asphyxia: Their relation to outcome. *Pediatrics*; 117 (2): 333-39.
- **Toet MC, Mrij VW, de Vries LS, Uiterwaal CS, and Van Huffelen kC (2002):** Comparison between simultaneously recorded a EEG in neonates. *Pediatr*; 109: 772-79.
- **Toet MC, (1999):** Amplitude integrated EEG at 3 and 6 hours after birth in full term neonates with hypoxic ischemic encephalopathy. *Arch Dis Child* ; 81:F19.
- **Trevisanuto D , Pitton M, Doglioni N, Altinier S, Zaninotto M, Plebani M, Zanardo V (2004):** Cord blood cardiac troponin T and troponin I levels in multiple-gestation neonates. *Biol Neonate* 2004;85:269-272.
- **Trevisanuto D, Picco G, Golin R, , Doglioni N, Altinier S, Zaninotto M, Zanardo V. (2006)** Cardiac troponin I in asphyxiated neonates. *Biology of the Neonate.*; 89(3):190–3
- **Turker G, Babaoglu K, Go" kalp AS, Sarper N, Zengin E, Arisoy AE(2004):** Cord blood cardiac Troponin I as an early predictor of short term outcome in perinatal hypoxia. *Biol Neonate*;86:131–137.
- **Verdu PA, Falero MP and Arroyos A et al (2001):** Blood neuronal specific enolase in newborns with perinatal asphyxia. *Rev. Neurok*; 32 (8) 714-17.

- **Vintzileos A M (2000):** Antenatal assessment for the detection of fetal asphyxia: An evidence- based approach using indication-specific testing. *Annals N Y Acad Science*; 900:137-50.
- **Volpe JJ (2001):** Hypoxic ischemic encephalopathy (Clinical aspects). In: *Neurology of the Newborn*. Volpe J (ed), 3rd edition, W.B.Saunders Co, Philadelphia, pp:250.
- **Wahlin A, Herbst A and Lindoff C et al (2001):** Brain-specific NSE and S-100 proteins in umbilical blood after normal delivery. *Clinic Chim Acta*: 304:57-63.
- **Watanabe K & Takeuchi T (1980):** Behavioral state cycles, background EEGs and prognosis of newborns with perinatal hypoxia. *Electroencephalogr Clin Neurophysiol* ; 49:618.
- **Wavenberg JL, Valente F, and Bormans J (1994):** Arterial postnatal base deficit evaluation of birth asphyxia. Abstract 296, 14th European Congress.
- **Wijaberger LD, Nikkels PG VanDongen AJ et al (2002):** Expression in the placenta of neuronal markers for perinatal brain damage. *Pediatric Res*; 51:492-96.
- **Wiswell TE (2000):** Delivery room management of the apparently vigorous meconium-stained neonate: Results of the multicenter, international collaborative trial. *Pediatrics*: 105:1-7.
- **Wyckoff MH, Perlman J and Laptook A (2005):** Use of volume expansion during delivery room resuscitation in near-term

and term infants. Pediatrics-, 115(4): 950-55.

- **Zimmerman KL (1994):** Uric acid correlation with Apgar Score Abstract 53, 14th European Congress.