

REFERENCES

Adams, J.H.; Corsellis, A.N. and Duchen, L.W. (1984):

Greenfield's neuropathology. 4th edition. New York. wiley,
pp. 385-449.

Adam, R. and Greenberg, J.O. (1978):

The mega cisterna magna. J. Neurosurg, 48 : 190 - 192.

Altman, N.R.; Naidich, T.P. and Braffman, B.H. (1992):

Posterior fossa malformations. A.J.N.R., 13 : 691 - 724.

Anderson, R.E. and Osborn, A.G. (1977).

Efficacy of simple sedation for pediatric computed tomography.
Radiology, 124 : 739 - 742.

Archer, C.R.; Darwish, H. and Smith, K. (1978):

Enlarged cisternae magna and posterior fossa cysts simulating
Dandy - Walker syndrome. Radiology, 127 : 681-686.

Babcock, D.S. (1981):

Cranial ultrasonography of infants .

The williams & wilkins Co., Baltimore.

Babcock, D.S. and Hand, B.K. (1981):

The accuracy of high Resolution, Real - time ultrasonography of the
head in infancy. Radiology 139 : 665 - 676.

Bachman, D.S.; Hodges, F.J.; Freeman, J.M. (1977):

Computerized axial tomography in neurologic disorders of children.
Pediatrics, 59 : 352 - 363.

Barkovich, A.J.; Kjos, B.O.; Norman, D. et al., (1989):

Revised classification of posterior fossa cysts and cystlike malformations based on the results of multiplanar M.R. imaging.
A.J.N.R., 10 : 977 - 988.

Banna, M. (1976):

Arachnoid cysts on computed tomography.
A.J.R., 127 : 979 - 982.

Bejar, R. and Coen, R. (1985):

Normal cranial ultrasonography in neonates. In Brain insults in infants and children edited by James, H., Anas, O., Perkin, R. Grune & Stratton Co., Orlando, Fla.

Benjamin, C.P.; Lee, J.B.; Maichael, D.F. et al. (1984):

Posterior fossa lesions : Magnetic Resonance imaging. Radiology, 153 : 137 - 143.

Bindal, A.K.; Storrs, B.B. and McLone, D.G. (1990):

Occipital meningoceles in patients with the Dandy - Walker syndrome. J. Neurosurgery, 28 : 844 - 847.

Bydder, G.M.; Steiner, R.E.; Thomas, D.J. et al. (1983):

Nuclear magnetic resonance imaging of the posterior fossa : 50 cases. Clin. Radiology, 34 : 173 - 188.

Byrd, S.E. and Naidich, T.P. (1988):

Common congenital brain anomalies. Radiologic clinic of north Am., 26 : 755 - 772.

Chiou, Y.M. and Tsai, C.H. (1992):

Sonographic features of Dandy - Walker variant with occipital cephalocele. J.C.U., 20 : 139 - 141.

Comstock, C.H. and Boal, D.B. (1985):

Enlarged fetal cisterna magna : Appearance and significance. J. Obstetrics & Gynecology, 66 : 25 - 28.

Cremin, B.N.; Chilton, S.J. and Peacock, W.J. (1980):

Anatomical landmarks in anterior fontanelle ultrasonography. B.J.R., 56 : 517 - 526.

Daniels, D.L. (1985):

Normal cerebral anatomy. In cranial computed tomography. Edited by Williams, A.L. and Houghton, V.M. 4th edition. The C.V. Mosby company St. Louis - Toronto - Princeton.

Davis, R.L. and Robertson, D.M. (1985):

Text book of neuropathology. 5th edition pp : 208 - 242.
Williams & Wilkins. Baltimore.

Diebler, C. and Dulac, O. (1983):

Cephaloceles : Clinical and neuroradiological appearance.
Neuroradiology, 25 : 199 - 216.

Estroff, J.A.; Scott, M.R. and Benacerraf, B.R. (1992):

Dandy - Walker variant : prenatal sonographic features and clinical outcome. Radiology, 185 : 755 - 758.

Faerber, E.N. (1986):

Cranial computed tomography in infants and children. 4th edition, Philadelphia, Lippincott.

Filly, R.A. (1988):

Ultrasound evaluation of the fetal neural axis. In ultrasonography in obstetrics and Gynecology. edited by Callen, P.W. 2nd edition. pp. 83 - 135. W.B. Saunders. company. Philadelphia - London - Toronto - Montreal - Sydney - Tokyo.

Fitz, C.R. (1982):

Midline anomalies of the brain and spine. Radiologic clinics of North Am. vol. 20 : 95 - 104.

Friede, R.L. (1975):

Developmental Neuropathology. First edition, Springer, Vienna.

Goodwin, L. and Quisling, R.G. (1982):

The neonatal cisterna magna : ultrasonic evaluation. Radiology, 149 : 691 - 697.

Grant, E.; Schellinger, D. and Richardson, J. (1983):

Ultrasonography of the posterior fossa, J.U. Med. 2 : 73 - 93.

Grant, E.G.; Tessler, F. and Perrella, R. (1988):

Infant cranial sonography. Radiologic clinics of North America. vol. 26 : 1089 - 1110.

Harwood - Nash, D.C. and Fitz, C.R. (1976):

Neuroradiology in infants and children. 4th edition. C.V. Mosby company. st. Louis.

Harwood - Nash, D.C. and Breckbill, D. (1976):

Computed tomography in children : a new diagnostic technique. J. pediatrics, vol.

Haughton, V.M. (1985):

Scanning techniques. In cranial computed tomography. edited by williams, A.L. and Haughton, V.M. 4th edition. The C.V. Mosby company. St. Louis - Toronto - Perinceton. pp. 1 -11.

Hidalgo, H.; Bowie, J. and Rosenberg, E.R. et al. (1982):

In utero sonographic Diagnosis of fetal cerebral Anomalies. A.J.R. vol. 139 : 143 - 148.

Hirschi, J.F.; Pierre - Kahn, A. and Renier, D. et al. (1984):

The Dandy - Walker malformation: A review of 40 cases. J. Neurosurgery 61 : 515 - 522.

Hutchinson, J.W.; Stoving, J. and Turner, P.T. (1979):

Occipital encephalocele with holoprosencephaly and aqueduct stenosis. Surg. Neurol, 12 : 331 - 335.

Kendal, B.E. and Kingsley, D. (1978):

The value of computerized axial tomography (CAT) in cranio - cerebral malformations. B.J.R. 51 : 171 - 190.

Kendall, B. and Stevens, J. (1993):

Intracranial lesions (1): Text book of Radiology and Medical imaging
vol (2) edated by Sutton, D. fifth edition. Churchill livingstone.
Edinburgh - London - Madrid - Melbourne - New York and Tokyo.

Kirkinene, P.; Jouppila, P. and Valkeakali, T. et al. (1982):

Ultrasonic evaluation of the Dandy - Walker syndrome. J. obstet.
Gynecol, 59 : 185 - 192.

Latorre, E.; Fortuna, A. and Occhipinti, E. (1973):

Angiographic differentiation between Dandy - Walker cyst and
arachnoid cyst of the posterior fossa in new born infants and children.
J. Neurosury, 38 : 298 - 305 .

Lehman, R.M. (1981):

Dandy - Walker syndrome in consecutive siblings : Familial hindbrain
malformation. J. Neurosurgery, 8 : 717 - 719.

Lemire, R.J.; Loeser, J.D. and Leech, R.W. et al. (1975):

Normal and Abnormal Development of the human nervous system.
Harper and Row, Hagerstown.

Leo, J.S.; pinto, R.S. and Hulvat, G.F. et al. (1979):

Computed tomography of arachnoid cysts. Radiology, 130 : 675 - 680.

Ludwin, S.K. and Malamud, N. (1977):

Pathology of congenital anomalies of the brain. In Radiology of the
skull and Brain. edited by Newton, T.H. and Potts, D.G., C.V. Mosby
Co. St. Louis. vol. 3, pp. 2979 - 3015.

Maggio, M. (1991):

Neonatal intracranial problems. in clinical sonography edited by sanders, R.C. 2nd edition. Little, Brown and company, Boston - Toronto - London pp. 421 - 446.

Maklad, N.F. (1986):

Ultrasound in perinatology. Clinics in Diagnostic ultrasound vol. 19 : 129 - 156. Churchill livingstone. New York - Edinburgh - London - Melbourne.

Marin - Padilla, M. (1978):

Clinical and experimental rachischisis. In Handbook of clinical neurology. edited by virken, P.J. and Bruyn, G.W. Elsevier / North - holland, Amsterdam. pp. 159 - 191.

Masdeu, J.C.; Dobben, G.D. and Kia, B.A. (1983):

Dandy - Walker syndrome studied by computed tomography and pneumoencephalography. Radiology, 147 : 109 - 114.

Masters, C.L. (1978):

Pathogenesis of the Arnold - Chiari malformation and aqueduct stenosis. J. Neuropath. Exp. Neurol, 37 : 56 - 74.

McLeary, R.D.; Kuhns, L.R. and Barr, M.Jr. (1984):

Ultrasonography of the fetal cerebellum. Radiology. 151 : 439 - 442.

Moseley, I.M. (1988):

Magnetic resonance imaging in diseases of nervous system. First edition. Blackwell, Oxford, pp 78 - 118.

Naidich, T.P. (1981):

Cranial C.T. signs of the Chiari II malformation. *J. Neuroradiology*,
8 : 207 - 227.

Naidich, T.P. and Fulling, K.H. (1983):

Congenital malformations of the brain. *Neuroradiology*, 25 : 277 - 304.

Naidich, T.P.; Mclone, D.C. and Fulling, K.H. (1983):

The chiari II malformation : Part IV. The hindbrain deformity. *J. Neuro
radiology* 25 : 179 - 197.

Naidich, T.P.; Pudlowski, R. M. and Naidich, J.B. et al. (1980):

Computed tomographic signs of the chiari II malformation. Part I :
Skull and dural partitions. *Radiology*, 134 : 65 - 71.

Naidich, T.P.; Pudlowski, R.M. and Naidich, J.B. et al. (1980):

Computed tomographic signs of the chiari II malformation. Part II :
Midbrain and cerebellum. *Radiology*, 134 : 391 - 398.

Naidich, T.P.; Pudlowski, R.M. and Naidich, J.B. et al. (1980):

Computed tomographic signs of the chiari II malformation. Part III.
Ventricles and cisterns. *Radiology*, 134 : 657 - 663.

Naidich, T.P.; Radowski, M.A. and Berstein, R.A. et al. (1988):

Congenital malformation of the posterior fossa. In : Taveras, J. M.
and Ferrucci, J.T. editors *Radiology* vol 3 philadelphia Lippincott.
pp. 1 - 17.

Naidich, T.P.; Yousefzadeh, D.K. and Gusnard, D.A. (1986):

Sonography of the normal neonatal head.

J. Neuroradiology, 28 : 408 - 427.

Naidich, T.P. and Zimmerman, R.A. (1987):

Common congenital malformations of the brain. In : Magnetic Resonance imaging of the central nervous system. edited by Brandt-Zawedzhi, M. and Norman, D. New York : Raven Press. pp. 131 - 151.

Newman, G.C.; Buschi, A.I. and Kelly, T.E. et al. (1982):

Dandy - Walker syndrome diagnosed in utero by ultrasonography. Neurology (NY) 32 : 180 - 184.

Newton, T.H. and Potts, D.G. (1981):

Radiology of the skull and Brain : Technical Aspects of computed Tomography. The C.V. Mosby Co. St. Louis, pp. 3956 - 3992.

Nishimaki, S.; Yoda, H. and Seki, K. et al. (1990):

A case of Dandy - Walker malformation associated with occipital meningocele. Pediatr - radiology 20 : 608 - 609.

Pigadas, A.; Thompson, J.R. and Grube, G.L. (1981):

Normal infant brain anatomy : Correlated real - time sonograms and brain specimens. A.J.R. 137 : 815 - 820.

Poe, L.B.; Coleman, L.L. and Mahmud, F. (1989):

Congenital central nervous system anomalies. Radiographics vol. 9 (5): 801 - 826.

Raybaud, C.P. (1982):

Cyst malformations of the posterior fossa abnormalities associated with the development of the roof of the fourth ventricle and adjacent meningeal structures. *J. Neuroradiology*, 9 : 103 - 133.

Ruge, J.R.; Masciopinto, J. and Storrs, B.B. et al. (1992):

Anatomical progression of the Chiari II malformation. *Childs - Nerve - syst.* 8 : 86 - 91.

Russ, P.D.; Pretorius, D.H. and Johnson, M.J. (1989):

Dandy - Walker syndrome : a review of fifteen cases evaluated by pre natal sonography. *Am. J. obstet. Gynecol*, 161 : 401 - 406.

Sabbagha, R.E.; Tamura, R.K. and Compo, S.D. (1980):

Fetal cranial and cranio-cervical mass : ultrasound characteristics and differential diagnosis. *Am. J. Obstet. Gynecol*, 138 : 511 - 517.

Sauerbrei, E.E. and Cooperberg, P.L. (1981):

Neonatal Brain : Sonography of congenital abnormalities. *A.J.R.*, 136 : 1167 - 1170.

Scotti, G.; Musgrave, M.A. and Fitz, C.R. et al. (1980):

The isolated fourth ventricle in children : C.T. and clinical review in 16 cases. *A.J.N.R.*, 1 : 419 - 424.

Slovis, T.L. and Kuhns, L.R. (1981):

Real - time sonography of the brain through the anterior fontanelle. *A.J.R.*, 136 : 277 - 285.

Strother, C.M. and Harwood-Nash, D.C. (1978):

Congenital malformations. In : Radiology of the skull and brain. edited by Newton, T.H. and Potts, D.G. C.V. Mosby company, St. Louis, pp. 3712 - 3748.

Tal, Y.; Freigang, B. and Dunne, H. et al. (1980):

Dandy - Walker syndrome. Analysis of 21 caess. Devel. Med. Child. Neurol, 22 : 189 - 198.

Taverase, J.M. and Wood, E.H. (1976):

Diagnostic neuroradiology. 2nd edition. Williams & Wilkins Baltimore.

Taylor, G.A. and Sanders, R.C. (1983):

Dandy - Walker : recognition by sonography. A.J.N.R. 4 : 1203 - 1209.

Thompson, J.R.; Schneider, S. and Ashwal, S. et al. (1982):

The choice of sedation for computed tomography in children : A prospective Evaluation. Radiology, 143 : 475 - 479.

Weisberg, L.A.; Nice, C. and Katz, M. (1978):

Cerebral computed tomography. Second edition. Saunders, Philadelphia.

Williams, A.L. (1985):

Congenital anomalies. In cranial computed tomography. Edited by Williams, A.L. and Haughton, V.M. 4th edition. The C.V. Mosby company. St. Louis. Toronto - Princeton, pp 316 - 349.

Wolfson, B.J.; Faerber, E.N. and Truex, R.C. (1987):

"The keyhole" sign of herniation of a trapped fourth ventricle and other posterior fossa cyst. *A.J.N.R.* 8 : 473 - 477.

Wolpert, S.M.; Anderson, M. and Scott, R.M. et al. (1987):

Chiari II malformation : M.R. imaging evaluation. *A.J.N.R.* 8:783 - 790.

Wolpert, S.M. (1974):

Vascular studies of congenital anomalies. in *Radiology of the skull and brain* edited by Newton, T.H. and Potts, D.G. vol. 2 C.V. Mosby Co. St. Louis.

Venes, J.L.; Black, K.L. and Latack, J.T. (1986):

Pre operative evaluation and surgical management of the Arnold - Chiari II malformation. *J. Neurosurg.* 64 : 363 - 370.

Vaquero, J.; Carillo, R. and Cabezudo, J. et al. (1981):

Arachnoid cyst of the posterior fossa. *Surg. Neurol.* 16 : 117 - 121.

Yousefzadeh, D.K. and Naidich, T.P. (1985):

U.S. anatomy of the posterior fossa in children : Correlation with brain sections. *Radiology*, 156 : 353 - 359.

Zimmerman, R.A. and Bilanink, L.T. (1988):

Pediatric central nervous system. In *Magnetic resonance imaging*. edited by Stark, D.D. and Bradley, W.G., C.V. Mosby. Co. St. Louis. pp. 132 - 152.

Zimmerman, R.D.; Breckbill, D. and Dennis, M.W. et al. (1979):
Cranial C.T. Findings in patients with meningocele. A.J.R.,
132 : 623 - 629.