

REFERENCES

- Adamis, M.K., Goldszer R.C., pulde, M.F., Sax, E.J and Edelman, R.R (1995) : Renal vasculature in potential renal transplant donors : Comparison of MR imaging and digital subtraction angiography. Radiology, 197 : 467-472.
- Amplatz, K. (1962): Percutaneous arterial catheterization and its applications. Am. J. roentgenol. Radium ther . Nucl. med., 87: 265.
- Anderson, C.M., and Lee, R.E., (1993) : Time-of-flight techniques : Pulse sequences and clinical protocols. MRI clinics of North America, vol (1) No (2), 217-227.
- Aoki, T., Houkin, K., Kamiyama, H., Iwasaki, Y., Abe, M., Kawaguchi, S., and Kashiwaba, T., (1993): Analysis of cerebral blood flow dynamics and evaluation of extracranial-intracranial arterial bypass function using MR angiography with presaturation pulse sequences. No. shinkei-geka, sep, vol(21), No(9), 793-798.
- Baert, A.L., Gryspeerdt, S. and van Hoe, L. (1996): Spiral C.T in non invasive evaluation of hypertensive renal vascular disease international Uro. radiology ed. by Willi, U.V., Kenny, Ph. J. and Thomsen H.S, P. 60-80, FADL publisher.

Barry J.N. (1996):- Donor Nephrectomy. In Marshall F.F., Kavoussi L. R., McAninch. J. W., and peters.G A., (eds): Operative urology 235-247. W.B. Saunders company, Philadelphia, Pennsylvania.

Biojsen, E. (1959): Angiographic studies of the anatomy of single and multiple renal arteries. Acta Radiology, 183: 1.

Borrello, J.A., Hylton, N.M., DULN (1992): Basics of MR angiography in Gutierrez F,R., J.J., Mirowitz, S.A. (eds) cardiovascular. Magnetic resonance imaging st. Louis, Mosby year book.

Borrello, J.A. (1997): renal MR Angiography . MRI Clinics of North America. vol(5), No(1) p. 83-93.

Bowen, B.C., quencer, R.M., Margosian, P., and pattany, P.M., (1994): MR angiography of occlusion disease of the arteries in the head and neck: current concepts. AJR, 162: 9-18.

Bradely, W.G., and Waluch, V. (1985): Blood flow: Magnetic resonance imaging. radiolgy, 154: 443-450.

Braun, B., Weilemann, L.S., and Weigand, W. (1981): Ultrasound demonstration of renal vein thrombosis. Rad., 138: 157.

- Buonocore, M.H. (1994): Algorithms for improving calculated streamlines in 3-D phase contrast angiography. *Mag. Reson. Med.* 31:22 -30.
- Caro, C.G., pedley, J.G., Schroter R.C.,Seed, W.A. (1978): The mechanisms of criculation. OX ford; Oxoford university Press.
- Cecka J.M., Terasaki P.I. (1995):- The UNOS scientific renal registry. in:- Clinical transplants (1995). Cecka J.M, terasaki P.I (eds) los Angles: UCLA tissue typing laboratory . P.P 1-18.
- Change, J.M., Friese, K.,and Caputo, G.R.,(1991): MR measurement of blood flow in the true and false channel in chronic aortic dissection. *J. comput. assist. tomogr.* 15: 418-423.
- Chien, D., Goldmann, A., and Edelman, R.R. (1992): High speed black blood imaging of vessel stenosis in the presence of pulsatile flow. *j. Mag. reson. Imaging* 2: 437-441.
- Cho. W.H., jacobs, S.C. and Lawson. R.K (1987): Digital subtraction angiography for donor evaluation. *urol.*, 30: 289-291.
- Cigarroa, J.E., Isselbacher, E.M., Desanctis, R.W., and Eagle K.A., (1992): Diagnostic imaging in the evaluation of suspected aortic dissection: Old standards and new directions. *N. Engl. J. Med*, 328: 35-43.

Cochlin, D.L., Dubbins P.A., Goldberg, B.B. and Alexander, A.A. (1994): The kidney in urogenital ultrasound ed. by Cochlin, D.L, Dubbins, P.A., Goldberg, B.B. and Alexander, A.A ch.2: 4-113 by Martin Danitz.

Cochran, S.T., Krasny, R.M., and Danovitch, G.M., (1997): Helical C.T angiography for examination of living renal donors. AJR 168: 1569-1573.

Cohen, M.S., (1993): Echo planar Magnetic resonance angiography. MRI Clinics of North America vol(1), No(2), December P. 359-365.

Copper, T.G.,and Gift, D.A. (1992): prospects for the future: Magnetic Resonance fast scanning and flow quantitation seminars in ultrasound, C.T and MRI, vol(13), No(4), Pp.303-310.

Crummy, A.B.; Stieghorst, M.F.; Turski, P.A., strother, ch.M., Lieberman, R.P., Sackett, F.J., Turnipsed, W.D., and Detmer, E.D. (1982): D.S.A.: Current status and use of intra-arterial injection. Rad., 145: 303-307.

Dachman, A.H., Newmark, G.M., Mitchell, M.T., and woodle, E.S (1998) : Helical CT examination of potential kidney donors. AJR 171 : 193-200.

Dalla-palma, L., Bazzocchi, M., Cressa, C., and Tommasini, G (1990):-Radiological anatomy of the kidney. Brit .J of Radiology 63,680-690.

Davis, C.P., Ladd, M.E., Romanowski, B.J., Wildermuth, S., Knoploch, J.F., and Debatin, J.F. (1996) : Human aorta : preliminary results with virtual endoscopy based on three dimensional MR imaging data sets. Radiology 199 : 37-40.

De-cobelli, F., Vanzulli, A., Sironi, S., Mellone, R., Angelli, E., Venturini, M., Garancini, M.P. and Quartagno, R., (1997): renal artery stenosis: evaluation with Breath, hold, three dimesional Dynamic, Gadolinum-enhanced versus three-dimensional phase contrast MR angiography radiology. 205: 689-695.

De-haan, M.W., Kouwenhoven, M.,and Thelissen, R.P.,(1996): Renovascular disease in patients with hypertension: detection with systolic and diastolic gating in three-dimensional-phase contrast MR angiography. Radiology, 198: 449-456.

Debatin, J.F, Ting, R.H., Wegmuller, H., Sommer, F.G., Fredrickson, J.O., Brosnan, T.J., Bowman, B.S., Mayers, B.D., Herfkens, R.J., and Pelc, N.J.(1994); renal artery blood flow quantification with phase contrast MR imaging with and without breath holding. radiology, 190: 371-378.

Drayer, B.P., (1992): Vascular localization of brain infarction. Neuroimaging Clinics of North America, vol (2). No(4), P.801-817.

Duda, S.H., Schick, F., Teufl, F., Muller, M., Erley, C., Schneider, W, Miller, S., and Claussen, C.D.(1997): Phase contrast MR angiography for detection of arteriosclerotic renal artery stenosis. Acta Radiologica 38: 287-291.

Dumoulin, C.L., Cline. H.E., Souza, S.P, Wagle, W.,and Walker, M.F.(1989): Three dimensional time of flight Magnetic Resonance angiography using spin saturation. mag. reson. Med., 11: 35-46.

Dumoulin, C.L., (1992): Phase contrast magnetic Resonance angiography. Neuroimaging Clinics of North American, vol(2), No(4), P. 657-675.

Earls, J.P., Rofsky, N.M., Decorato, D.R., Krinsky, G.A., and Weinreb, J.C. (1996): Breath hold single dose Gadolinum enhanced three-dimensional MR aortography usefulness of a timing examination and MR power injector. Radiology 201: 705-710.

Ecklund, D., Hartnell, G.G., Hughes, L.A., and Strokes, K.P., (1994): MR angiography as the sole method in evaluating abdominal aortic aneurysms. Correlation with conventional techniques and surgery. Radiology, 192: 345.

- Edelman, R.R., Wentz, K.U., Mattle, H.P., O'Reilly, G.V., Candia, G., Liu, C., and Zahao, B., (1989): Intracranial arterio-venous malformations: Evaluation with selective MR angiography and venography. *Radiology*, 173: 831-837.
- Edelman, R.R., Mattle, H.P., and Wallner, B., (1990): Extra cranial Carotid arteries: evaluation with "Black blood" MR angiography *Radiology* 177: 45-50.
- Edelman, R.R., Mattle, H.P., and O'Reilly, G.V., (1990) : Magnetic resonance imaging of flow dynamics in the circle of Willis. *Stroke* 21 : 56-65.
- Edelman, R.R., Chien, D., and Kim, D., (1991): Fast selective black blood MR imaging. *Radiology* 181: 655-660.
- Edelman, R.R., Sungkee, S.A., and Chein, O., (1992): Improved time of flight MRA of the brain with magnetization transfer contrast. *radiology*, 25: 372-392.
- Edelman, R.R. (1993): Magnetic Resonance angiography. An overview. *invest Radiol.* 28(supply 4): 543-546.
- Edelman, R.R. (1993): MRA angiography; present and future. *American journal of radiology* 161:1-11.

Edelman, R.R., Sieuert, B., and Adamis, M., (1994): Signal targeting with alternating radiofrequency (STAR) sequences: Applications to MR angiography, Mag. reson. Med 31: 233-238.

Edelman, R.R., Adamis, M.K., Müller, M., and Hochman, M., (1997): MR angiography of the body in : Edelman, R.R., Hesselink, J.R., and Zlatkin, M.B., (eds): Clinical Magnetic Resonance imaging 2nd edition, vol(2), W.B., Saunders company, Philadelphia, P. 1305-1329.

Eggers, P.W. (1988):- Effect of transplantation on the medical end stage renal disease program. New Engl. J. Med., 318: 223.

Ehman, R.L., and Felmlee, J.P., (1990) : Flow artifact reduction in MRI : a review of the roles of Gradient moment nulling and spatial presaturation. Magn. Reson. Med. 14 : 293-307.

Ekelund, L., Sjoqvist, L., Thuomas, K.A., and Asberg, B., (1996): MR angiography of abdominal and peripheral arteries; Techniques and Clinical applications. Acta Radiologica 37: 3-13.

El-Azab, M., Mohsen, T., El-Diasty, T.A. and Shokeir, A.A. (1996): Doppler ultrasonography in evaluation of potential live kidney donors: Aprospective study J. Urol., Vol. 156: 818-880.

Foly, W.D. (1989): Dynamic hepatic C.T Rad. 110: 617-622.

Foo, T.F., Manojkumar, S., prince, M.R., and Chenevert, T.L., (1996): Smart prep: an automated method for detecting the bolus arrival time and initiating data acquisition in fast 3-D gadolinum enhanced MR angiography (abst). in : proceedings of the society of Magnetic Resonance. Berkely C.A.: Society of Magnetic Resonance 1996: 453.

Furya, Y., Isoda, H., Hasegawa, S., Takahash., Kaneko, M., and Uemura, K., (1992) : Magnetic Resonance angiography of extracranial carotid and vertebral arteries, including their origins : Comparison with digital subtraction angiography. Neuroradiology 35 : 42-45.

Goldberg, M.A., Yucel, E.K.,and Saini, S.,(1993) : MRangiography of the portal and hepatic venous systems : preliminary experience with echo planar imaging. AJR 160 : 35-40.

Graves, F.T. (1954): The anatomy of intrarenal arteries and its application to segmental resection of the kidney. Brit. J. Surg., 42: 32.

Graves, M.J.(1997): Magnetic Resonance angiography. the British. Journal of Radiol. 70: 6-28.

- Grist, T.M., Sproat, I.A., Kennel, T.W., Korosec, F.R., and Swan, J.S., (1996) : MR angiography of the renal arteries during breath-hold using gadolinium enhanced 3D TOF with K. space Zero filling and a contrast timing scan (abst) in proceedings of the international society of Magnetic Resonance in Medicine 1 : 163.
- Gullberg, G.T. wehrli. F.W., Shimaka, A., and Simons, M.A., (1987): MR vascular imaging with a fast gradient refocusing pulse sequence and reformed images from tranaxial sections Radiology 165: 241-246.
- Haacke, E.M., Li.D., Mugler, J.P (1994): Three dimensional Time of flight MR angiography using selective inversion Recovery RAGE with fat saturation and E.C.G triggering: Application to renal arteries. Mag. Reson. Med. 31: 414-422.
- Hany, T.F., Debatin, J.F., Leung; D.A. and Pfammatter, T., (1997): Evaluation of the aorto-iliac and renal arteries: Comparison of breath-hold, contrast-enhanced, three-dimensional MR angiography with conventional catheter Angiography. Radiology. 204: 357-362.
- Hany, T.F., Schmidt, M., Davis, C.P., Göhde, S.C and Debatin, J.F., (1998): Diagnostic impact of four postprocessing techniques in evaluating contrast-enhanced three-dimensional MR angiography. AJR, 170: 907-912.

- Hessel, S.J.(1983): Complications of angiography and other catheter procedures in vascular and interventional radiology ed. by Abrams, H.L>, 3rd edition., volume 11: 1041-1051, little Brown and company Boston.
- Holland, G.A., Dougherty, I.,and Carpenter, J.P., (1996): Breath hold ultrafast three dimensional gadolinum enhanced MR angiography of the aorta, renal and other visceral abdominal arteries AJR 166: 971-981.
- Kaatee, R., Frederik, J.A., Beek, A., De-lange, E.E., Leeuwen M.S., Smith, F.M., and Mali, P.T., (1997): Renal artery stenosis: detection and Quantification with spiral C.T angiography versus optimized Digital subtraction angiography. radiology, 205: 121-127.
- Kabalin, J.N. (1997) : Surgical anatomy of the retroperitoneum, kidney and ureters. In Walsh, P., C., Retik, A.B., and vaughan, E.D.(eds): Campbell's urology, 7th edition p.P49-88. W.B. Saunders company, Philadelphia, Pennsylvania.
- Kadota, T., Kuriyama, K., Inoue, E., Fuiita, M., Nakagawa, H., and Kuroda, C., (1993): MR angiography of Meningioma. Magnetic Resonance Imaging, vol(11), No(4), P. 473-483.

Kandarpa, K., Sandor, T., Tieman, J., Hooshmand, R., Chopra, P.S., and Chakrabati, J., (1993): Rapid three-dimensional surface reconstruction of magnetic resonance images of large arteries and veins. A preliminary evaluation of clinical utility. Cardiovasc. Intervent. Radiol. 16: 25.

Kaufman S.L., Change, R., Radir., S, Mitchell, S.E. and Whit, R.I (1984): Intra arterial digital subtraction angiography in diagnostic arteriography. Rad., 151: 323-327.

Kaufmann, J.A., Geller S.C, and Petersen, M.J., (1994) : MRImaging (including MRA) of abdominal aortic aneurysm : Comparison with conventional angiography. AJR 163 : 203-210.

Keller, P.J., (1992): Time of flight magnetic resonance angiography. Neuro-imaging Clinics of North America, vol(2) No(4), P. 639-656.

Khauli, R.B. (1994):- Defining the role of renal angiography in the diagnosis of renal artery disease. American journal of kidney diseases, 24, 679-684.

Kim, D., Porter, D.H. and crivello, M.S.(1991): Renal artery imaging : a prospective comparison of intra arterial digital subtraction angiography. Angiology, 42: 345-357.

- Kopka, L., Vosshenrich, R., Muller, D., Fischer, U., Rodenwaldt, J. and Grabbe, E., (1997): result of a contrast enhanced three dimensional MR angiography in a single breath hold after optimization of the contrast material bolus. *fortscher. Rontgen.* 166: 15-20.
- Kopka, L., Vosshenrich, R., Rodenwaldt, J., and Grabbe, E., (1998): Differences in injection rates on contrast enhanced breath hold three dimensional MR angiography. *AJR.* 170: 345-348.
- Kouwenhoven, M., Bakker, C.J., Hartkamp, M.J., and Mali, W.P., (1994): Current MR angiographic imaging techniques, *Asurvey vascular diagnostics*, Springer-verlag, heidelberg.
- Laub, G., and Kaiser, W, (1988): MR angiography with gradient motion refocusing . *J. comput. Assist Tomogr.* 12: 377-382.
- Leung, D.A., McKinnon, G.C, Davis, C.P., plammeter, T., and Krestin, G, P. (1996): Breath hold contrast enhanced three dimensional MR angiography. *Radiology* 201: 569-571.
- Leung, D.A., and Debatin, J.F. (1998) : Contrast-enhanced 3D MRA: Impact on thoracic and abdominal vascular imaging. In *Advances in MRI 5* : 28-43. Kluwer Academic publishers. Printed in Great Britain.

Levy, C., Laissy, J.P., Raveau, V., Amarenco, P., Servois, V., Bousser, M.G., and Tubiana, J.M., (1994) : Carotid and vertebral artery dissections : Three-dimensional Time-of Flight MRangiography and MR imaging versus conventional angiography. Radiology, 190 : 97-103.

Lipchik, E.O. and Rogoff, S.M. (1983): Abdominal aortography: Translumbar, femoral and axillary artery catheterization techniques in vascular and interventional radiology ed. by Abrams, H.L., 3rd edition, volume 11: 1029-1039, Little, Brown and Company Boston.

Lowell, J.A., Taylor R.J. (1994): The evaluation of the living renal donor, surgical techniques and results. Semin Urol. 27: 102-107.

Maki, J.H, Prince, M.R., Londy, F.J., and Chenevert, T.L., (1996): The effects of time varying intravascular signal intensity and K-space acquisition order on three-dimensional MR angiography image quality. JMRI 6: 642-651.

Martin , R.P. (1994): Gadolinum enhanced MR aortography. radiology 191: 155.

McDonald, D.A. (1974): Blood flow in arteries. London Edward Arnold.

McDonnel, C.H., Li-Kcp, and Whitney, W.s., (1994): Chronic mesentric ischemia: Evaluation with phase-contrast cine MR imaging. radiology 190: 175-179.

Mcelroy. J, Novick. A.C., streem S.B., Steinmuller. D, Risius B, Buszta. C and Stein Hilber D. (1986): Aprospectve Analysis of the Accuracy and cost effectiveness of digital subtraction Angiography for living-related renal donor evaluation., transplantation 42: 23-27.

Meany J.F., Prince, M.R., Nostrant, T.T and Standey, J.C. (1997) : Gadolinum-enhanced MRA of the visceral arteries in patients with suspected chronic mesentric Ischemia. J. Magn. Reson. Imaging 7 : 171-176.

Meyer, C., Nishimura, D., and Macouski, A., (1992) : Fast spiral Coronary artery imaging. Magn Reson. Med. 28 : 202-213.

Meyers, S.P., Talagala, S.L., Totterman, S., Azodo, M.V., Kwok, E., Shapiro, L., Shapiro, R., Pabico, R.C and Applegate, G.R (1995) : Evaluation of the renal arteries in kidney donors : Value of three-dimensional phase - Contrast MR angiography with maximum-intensity projection or surface rendering. AJR 164 : 117-121.

Milnor, W.R. (1989): Haemodynamics Baltimore: Wiliams & Wilkins. P. 35.

Mitchell, D.G., Stolpen, A.H., Siegelman, E.S., Bolinger, I., and Outwater, E.K., (1996): Fatty tissue on opposed phased MR images paradoxical suppression of signal intensity by paramagnetic contrast agents. radiology 198: 351-357.

Murakami, T., Kashiwagi, T., Nakamura, H., et al., (1993): Display of MRangiograms: Maximum intensity projection versus three-dimensional rendering. Eur. J. Radiol. 17: 95-100.

Netter, F.H. (1980): Kidney, ureters and urinary bladder in the CIBA collection of medical illustrations, ed. by Hapter, R.K. and Yonkman, F.F., 6:, 14-17, published by CIBA.

Nishimura, D.G., (1993): Magnetic Resonance of the body. MRI Clinics of North America, vol(1), No(2), December P. 203-215.

Owen, R.S., Carpenter, J.P., and Baum, R.A., (1992): Magnetic Resonance imaging of angiographically occult run off vessels in peripheral arterial occlusive disease. N. Eng. J. Med, 326: 1577-1581.

parker, D.L., and Blatter, D.D., (1992): Multiple thin slab magnetic Resonance angiography. Neuroimaging clinics of North America, vol(2), No(4) P. 677-692.

Petty. W, Spigos, D.S, Abjo R, Mozes. M, Tan, W. S, Wilbur, A.
(1986) :- Arterial digital angiography in the evaluation of
potential renal donors. (1986) invest. Radiol . 21: 122-
124.

Platt, J.F., Ellis, J.H., Korobkin, M., and reige, K. (1997): Helical C.T
angiography evaluation of potential kidney donors:
finidings in 154 subjects. AJR 169: 1325-1330.

Pozniak, M.A.; Madison, W.I.; Tambeaux, R.11; Gist, M.J., Uehling,
D.T., Moon, T. and Salison, D.J. (1996): C.T
angiography of renal transplant donors (abst).
proceedings of 82 ND scientific assembly and annual
meeting, RSNA, Chicago.

Prince, M.R., Narasimhan, D.L., Stanley, J.C.,and Chenevert, T.L.,
(1995) : Breath hold Gadolinum MRangiography of the
abdominal aorta and its major branches. Radiology,
197 : 785-792.

Prince, M.R., Narasimhan, D.L.,and Jacoby, W.T.,(1996) : Three-
dimensional gadolinum enhanced MRA of the thoracic
aorta AJR 166 : 1387-1397.

Prince, M.R., Schoenberg, S.O., Ward, J.S.,and Londy, F.J., (1997) :
Haemodynamically significant Atherosclerotic Renal
artery stenosis : MR Angiographic features. Radiology,
205 : 128-136.

Prince, M.R., Chenevert, T.L., Thomas, K.F., Londy, F.J., Waard, J.S., and Maki, J.H., (1997): Contrast enhanced Abdominal MR aortography: optimization of imaging delay time by automating the detection of contrast material arrival in the aorta. Radiology, 203: 109-114.

Quinn, S.F., DemLow, T.A., and Hallin, R.W., (1993): Femoral MR angiography versus conventional angiography: Preliminary results. radiology 189: 181-184.

Riehle, R.A., Steckler, R., Naslund, B, E., Riggio, R., Cheigh, J. and suben Bord, W, (1990): Selection criteria for evaluation of the living related renal donors. Urol., 144: 845-848.

Rieumont, M.J, Kaufman, J.A., Geller, S.C., Yucel, E.K., Cambria, R.P., Fang, L.S., Bazari, H., and waltman, A.C. (1997) : Evaluation of Renal artery stenosis with Dynamic Gadolinum-Enhanced MRAngiography. AJR 169 : 39-44.

Rittgers, S.E. Norris, C.S., and Barnes, R.W.(1985): Detection of renal artery stenosis: expermintal and clinical analysis of velocity wave forms. Ultrasound Med. Biol., 11: 523.

Robe, F.E., Smith, E.J.; Yune, H.Y.; E.C., Leapman, S. and Filo, R.S (1983): Limitations of D.S.A. in evaluation of potential renal donors A.J.R., 141: 91.

Robertson R., Murphy, A. and Dubbins, P.A. (1988): renal artery stenosis: The use of duplex ultrasonography as a screening technique B.JR., 61: 196.

Rosenfield, A.T., and Smith, R.C. (1996): Non contrast Helical (spiral) C.T in the evaluation of flank pain: in urology of emergencies international urology 96 ed by Willi, U.V., Kenney, Ph. J. and Thomson, H.S, P. 37-41 FADL publishers.

Rubin, D.G., Dake, M.D.,and Napel, S., (1994) : Spiral CT of Renal artery stenosis : Comparison of three dimensional rendering techniques Radiology 190 : 181-189.

Rubin, D.G. and silverman, S.G. (1995): helical C.T of the retroperitoneum in Helical computed tomography ed. by silverman, P.M., Vol. 33, Number5, P. 913-930, by Saunders, W.B. Company.

Rubin, D.G., Alfery, E.J., Dake, M.D., Semba Ch. P., Sommer, F.G., Kuo, P.C., Dafoe, D.C., Waskerwitz, A. and Jeffrey, R.B. (1995): assesment of living renal donors with spiral C.T. Radiology, 195: 457-462.

Ruggierri, P.M, laub, G.A, Masaryk, T.J, and Medic, M.T: (1989): Intracranial Circulation: pulse -sequence considerations in three dimensional MR angiography. Radiology 171: 785-791.

Ruggieri, P.M., Masary K, T.J., Ross, J.S., and Modic, M.T., (1992): Intracranial Magnetic Resonance angiography. Cardiovascular and interventional Radiology, 15: 71-81.

Runge, V.M., Kirsch, J.E., Lee, C., Garnean, R.A., and Brack, M.A., (1992) : MRAngiography of cerebral disease. MRI Decisions, september/October, P : 2-18.

Sackell, J.f. and mann, F.A. (1982): Contrast media for digital subtraction arteriography of cerebral arteries in : digital subtraction arteriography. ed. by mistretta, C.A., Crummy, A.B., Strother, C.M. and sackell, J.F., 23: 27 Chicago-London: year book medical Publishers.

Saloner, D. (1995): An introduction to MR angiography. radiographics ; 15: 453-465.

Saloner, D. (1997): MRA principles and display in Higgins, C.B., Hricak, H. and Helms, C.A (eds) : Magnetic Resonance imaging of the body 1345-1368. Lippin cott. ravaen publishers, Philadelphia. New York.

Schild, H.H. and Eckmann, A. (1992) : C.T anatomy in computed Tomography in Urology ed. by Schild, H.H., Schweden, F.T. and Lang E.K. Vol. I., chapter 3; 15-18, Thieme Medical publishers, Inc. New York.

Schwerk, W.B., Schwerk, W.N., and Rodeck, G. (1985): Venous renal tuomer extension: a prospective ultrasound evaluation Rad., 156: 491.

Scott, S., (1995): basic concepts in magnetic Resonance angiography. radiologic Clinics of North America. vol(33), No(1) P. 91-112.

Sherwood, T., Rautu, M., and Chisholm, G.D. (1978): Renal angiography problems in live kidney donors. BJR, 51: 99-105.

Shokeir, A.A., Gad, H.M., Shaaban, A.A. El-Kenawy, M.R., El-Sherif. A, Shamaa, M.A., Bakr, M.A. and Ghoneim, M.A. (1993): Differential kidney scans in pre-operative evaluation of kidney donors. transplantation proceedings, 25: No.3 (June): 2327-2329.

Shokeir, A.A.; El-Diasty, T.A.; Shaaban, A.A.; El-Kenaway, M.; Eraky, I. and Ghoneim, M.A.(1993) :evaluation of kidney donors using D.S.A . transplantation proceedings, 25: No.3 (June); 2273-2275.

Sibert, J.E., Pernico, J.R., and Potchen, E.J., (1992): Physical principles and application of Magnetic resonance angiography: Seminar in ultrasond, C.T and MRI, vol(13), No(4) P. 227-245.

- Simonetti, O.P., Wielopolski., P., and Durek, J.L., (1994) : Experimental evaluation of flow effects in echo planar imaging in : Proceedings of the society of Magnetic Resonance, San Francisco, August 6-12, Vol 1. Berkeley, C.A., : Society of Magnetic Resonance P. 460.
- Snidow, J.J., Johnson, M.S, and harris, V.T., (1996): Three dimensional gadolinum enhanced MR angiography for aorto-iliac inflow assesment plus renal artery screening in a single breath hold. Radiology 198: 725-732.
- Spiral, D.B., Salvalierra, O., Palubinskas, A.J., Amend, W.J.C, Vincenti, F.G. and Feduska, N.J.(1979) : Results and significance of angiography in potential kidney donors Rad, 133: 45-47.
- Steiner, P., McKinnon G.C.,and Romanowski, B., (1997) : Contrast enhanced, ultrafast 3D pulmonary MRA in a single breath hold : Initial assesment of imaging performance. J. Magn. Reson. Imaging 7: 177-182.
- Strouse, P.J., Prince, M.R., and Chenevert, T.I., (1996): Effect of the rate of gadopentate dimeglumine adminstration on abdominal vascular and soft tissue MR imaging enhancement patterns. Radiology 201: 809-816.
- Suss-man, S.K., Weinerth, J.L; Braan, S.D., Cohan, R.H., Newman, G.E., perlmutt , L.M. and Dunnick (1987): I.V. DSA in the evaluation of potential renal donors J. Urol., 138: 28-32.

Suto, Y., Ohuchi, Y., and Kimura, T., (1994): Three dimensional black blood MR angiography of the liver during breath hold. Acomparison with two-dimensional time of flight MR angiography. Acta Radiol. 35: 131-134.

Svetky, L.P., Dunnick, N.R., Colfman, T.M., Himmelstien, S.I. et al., (1988): Comparison of intravenous digital subtraction angiography and conventional arteriography in defining renal anatomy. Transplantation, 45: 65-68.

Tempany, C.M., Morton, R.A., and Marshall, F.F., (1992): MR imaging of the renal veins: Assesment of non-neoplastic venous thrombosis. J. Comput. Assist. Tomogr. 16: 929-934.

Tindill, B. (1985): Arteriography and angio-cardiography in : Aguide to radiological procedures ed. by chapman, S. and Nakiely,R. 2nd edition 123-173.

Turski, P. and Korosec, F. (1992): Technical features and emerging clinical applications of phase contrast magnetic resonance angiography. Neuroimaging Clinics of north America, vol(2), No(4) P.785-800.

Vogl, T.J., Bergman, C., Villringer, A., Einhaupl, K., Lissner, J., and Felix, R., (1994): dural sinus thrombosis, value of venous MR venography for diagnosis and follow up. AJT, 162: 1191-1198.

Walker, T.G., Geller, C.S., Delmonico, F.L., Waltman, A. C. and Athanasoulis C.A. (1988): Donor renal angiography: its influence and the decision to use the right or left kidney A.j.R., 151: 1149-1151.

Waltzer, W.C., Engen, D.E., Stanson, A.W., Sterio IF, S., et al., (1985): Use of Radiological abnormal kidneys in living related donors transplantation. Nephron 39: 302.

Wasser, M.N., Westenberg, J., Victor, P.M., Baalen, J.V., Bockel, J.H., and pattynama, P.M., (1997): Haemodynamic significance of renal artery stenosis: Digital subtraction angiography versus systolically gated three dimensional phase contrast MR angiography. radiology 202: 333-338.

Wehrli, F. W., (1990): Time of flight effects in MR imaging of flow. mag. reson. med. 14: 187-193.

Wentz, K.U., Rother, J., Schwartz, A., Mattle, H.P., Suchalla, R., and edelman, R.R., (1994): Intracranial vertebro-basilar system: MR angiography . Radiology, 190: 105-110.

Wielopolski, P.A., and Edelman, R.R., (1994) : Ultrafast, high resolution 3D STAR MRangiography using interleaved segmented echo planar readouts in : Proceedings of society of Magnetic Resonance. San Francisco. August 6-12, volume 2. Berkely, C.A., Society of Magnetic resonance P. 948.

Wielopolski, P.A., Adamis, M., and Prasad, P. (1995): Breath hold 3-D STAR MR Angiography of the renal arteries using segmented echo. planar imaging. *Mag. reson. Med.* 33: 432-438.

Yanaka, K., Shirai, s., Shibata, Y., Kamezaki, T., Matsumura, A., and Nose, T., (1993): Gadolinum-enhanced Magnetic Resonance Angiography in the planning of supratentorial glioma surgery. *Neuro. Med. Chir-Tokyo*, Jul, vol(33), No(7), 439-443.

Yano, T., Kodama, T, Suzuki, Y., and Watanabe, K., (1997): Gadolinum enhanced 3-D Time-of-flight MR angiography. *Acta. Radiologica* 38: 47-54.

Yucel, E.K., Kaufman, J.A., Prince, M., et al., (1993): Time of flight renal MR angiography; utility in patients with renal insufficiency. *Magn. Reson. Imaging*, 11: 925-930.

Yucel, E.K., (1997): MRA of the aorta, Visceral and peripheral arteries in: Higgins, C.B., Hricak, H., and Helm, C.A. (eds): *Magnetic Resonance Imaging of the body*. 3rd edition, Loppincott-Raven press. New York, P. 1383-1399.

Zimmerman, R.A., Bogdan, A.R., and Gusnars, D.A., (1992): Paediatric Magnetic Resonance angiography. *Cardiovascular and interventional Radiology*, 15: 60-64.