

REFERENCES

- 1- Cow IJ. Development and applications of in vivo clinical magnetic resonance spectroscopy. Prog Biophys Mol Biol. 1996;65 (1-2): 45-81.
- 2- Rand SD, Prost R, Hanthton V et al. Accuracy of single voxel proton MR spectroscopy in distinguishing Neoplastic from non-neoplastic lesions AJNR 1994;18: 1695-74.
- 3- Forham E M et al. the utility of MRI in suspected MS. Neurology. Report of the therapeutics & technology Assessment subcommittee of the American Academy of Neurology. Sept 9-2003;61 (5).
- 4- Kent. Braun JA, Miller RG, Weiner MW. Magnetic resonance spectroscopy studies of human muscle. Radiol Clin North Am 1993;32 (2): 313-35.
- 5- Nelson SJ, Vignerson DB. Magnetic resonance spectroscopy. In MRI of Musculoskeletal system. 1st ed. Philadelphia, London, Toronto, Montreal, Sydney, Tokyo: WB Saunders Company, 1994; 69-79.
- 6- Fawcett DW. Muscular tissue, in Fawcett DW: A text Book of Histology, Champan& Hall 1994,260-308.

- 7- Cormack DH, Muscle tissue. In: Cormack DH. Hams histology. J.B. Lippincott company. London, Mexico City, New York, St. Louis, Sao Paulo, Sydney 1987;15:388-420.
- 8- Patten RM, Ovalle WE. Muscle spindle ultrastructure revealed by conventional and high resolution scanning microscopy. Anat. Rec 1991; 230:183-9.
- 9- Page S. Structure of the sarcoplasmic reticulum in vertebrate muscle. Br. Med. Bull. 1968;24:170-5.
- 10- Ogata T. Morphological and cytochemical features of the fiber types in vertebrate skeletal muscle. CRC Crit. Rev. Anat. Cell Biol. 1988;1:299-303.
- 11- MacLennan DH. Wong PTS. Isolation of a calcium sequestering protein from the sarcoplasmic reticulum. Proc. Nat. Acad. Sci USA 1971;68:1231-6.
- 12- Somlyo AV. Bridging structures spanning the junction gap at the triad of skeletal muscle. J cell Biol. 1979;80:743-7.
- 13- Burke RE. Motor units: anatomy, physiology and functional organization. In Handbook of physiology, The nervous system II 1981; 345-52. American physiology society, Bethesda, MD.
- 14- Eisenberg BR, Kuda AM: Discrimination between fiber populations in mammalian skeletal muscle by using ultrastructural parameters. J Ultrastruct Res 1976;54:76-88.

- 15- Durham RM. Muscles and movements in: Durham RM. Human Physiology. Functions of the Human Body. 1st ed. Wm. C. Brown Publishers 1989; 11:352-86.
- 16- Ganong WF. Excitable tissue: Muscle in: Ganong WF. Review of Medical Physiology. 7th ed. Appleton & Lange. Norwalk, Connecticut 1995;3:56-67.
- 17- Murphy RA. Contraction of muscle cells in: Berne RM, Levy MN. Physiology. 1st ed. C.V. Mosby Company St. Louis, Toronto 1983;22:387.
- 18- Guyton AC. Contraction of skeletal muscle in: Guyton AC: Text Book of Medical Physiology. 6th ed. W.B. Saunders company. Philadelphia, London, Toronto, 1981;11:122-36.
- 19- Kienlin M. Von (1999): Basic magnetic resonance spectroscopy and localization methods. 16th annual scientific meeting of Europ.
- 20- Rudin M, Sauter A. Measurement of reaction rates in vivo using Magnetization transfer techniques. In: Diehl P, Fluck E, Cunther H, Kosfeld R, Seelig J, editors. NMR basic principles and progress, Vol. 27. Berlin: springer; 1992 P. 257-93.
- 21- Danielsen E.R. and Ross B. (1999): Magnetic resonance spectroscopy diagnosis of neurological diseases. New York: marcel Dekker, -5-22.

- 22- Bloch F, Hansen WW, Packard M. The nuclear induction experiment. *Phys Rev* 1946;70:474-485.
- 23- Purcell EM, Torrey HC, Pound RV. Resonance absorption by nuclear magnetic moments in a solid. *Phys Rev* 1946;69:37-38.
- 24- Proctor WG, Yu FC. The dependence of nuclear magnetic resonance frequency upon chemical compound. *Phys Rev* 1950; 70:717-9.
- 25- Bottomely, P.A., 1987. Spatial localization in NMR spectroscopy in vivo. *Ann. NY Acad. Sci.* 508, 333-348.
- 26- Brown, T.R., 1992, Practical applications of chemical shift imaging. *NMR Biomed.* 5(5), 238-243.
- 27- Hornak J.P. (2002): The basics of MRI, XII, Chap 4,5 & 12.
- 28- Cousins J.P. (1995): Clinical MR spectroscopy: Fundamentals, current applications, and future potential. *Am J Roentgenol*; 164: 1337-1347.
- 29- Castillo M., Kwok L., Scalfiff J. and Mukherji S. (1996): Clinical applications of proton MR spectroscopy. *Am J Neuroradiol*, Jan; 17:1-15.
- 30- Gonen O., Gruber S., Li B.S.Y., et al (2001): Multivoxel 3D proton spectroscopy at 1.5 versus 3.0 T: Signal-to-noise ratio and resolution comparison. *Am J Neuroradiol*, Oct; 22(9): 1727-31.

- 31- Gruetter R., Weisdorf S.A., Rajanayagan v., et al. (1998): Resolution improvements in invivo ^1H -NMR spectra with increased magnetic field strength. *J Magn Reson.* 135:260-264.
- 32- Bachelard, H.S., Cox, D.W., Morris, P.G., 1987. Nuclear magnetic resonance as a tool to study brain metabolism. *Gerontology* 33 (3-4), 235-246.
- 33- Matthaei, D., Frahm, J., Haase, A., Schuster, R., Bomsdorf, H., 1985. Chemical-shift-selective magnetic-resonance imaging of avascular necrosis of the femoral head. *Lancet* 1 (8425), 370-371.
- 34- Frahm, H., Haase, A., Hanicke, W., Matthaei, D., Bomsdorf, H., Helzel, T., 1985. Chemical shift selective MR imaging using a whole-body magnet. *Radiology* 156 (2), 441-444.
- 35- Frahm, J., Bruhn, H., Gyngell, M.L., Merboldt, K.D., Hanicke, W., Sauter, R., 1989. Localized high-resolution proton NMR spectroscopy using stimulated echoes: initial applications to human brain in vivo. *Magn. Reson. Med.* 9 (1), 79-93.
- 36- Aue WP, Mueller S, Cross TA. Volume selective excitation. A novel approach to topical NMR. *J Magn Res* 1984;56:350-4.
- 37- Haase A. Localization of unaffected spins in NMR imaging and spectroscopy (LOCUS) Spectroscopy. *Magn Res Med* 1986;3:963-9.

- 38- Bottomley PA, Foster TB, Darrow RB. Depth-resolved surface-coil spectroscopy (DRESS) for in vivo H-1, P-31 and C-13 NMR. *J Magn Res* 1984;59:338-42.
- 39- Ordridge RJ, Connelly A, Lohman JAB. Image-selected in vivo spectroscopy (ISIS). A new technique for spatially selective NMR spectroscopy. *J Magn Res* 1986;66:283-94.
- 40- Riddle WR, Gibbs SJ, Willcott MR. Dissecting and implementing STEAM spectroscopy. *Magn Reson Med* 1993;29:378-380.
- 41- Webb PG, Sailasuta N, Kohler SJ, Raidy T, Moats RA, Hurd RE. Auto-mated single-voxel proton MRS: technical development and multisite verification. *Magn Reson Med* 1994;31:365-373.
- 42- Ordridge RJ, Connelly A, Lohman JAB. Image-selected in vivo spectroscopy (ISIS); a new technique for spatially selective NMR spectroscopy. *J Magn Reson* 1986;66:283-294.
- 43- Garwood M. (1997): Principles of spectroscopic localization. Fifth scientific meeting of International Society for Magnetic Resonance in Medicine; Apr 12-18, (Vancouver BC, Canada): ISMRM Educational Course Syllabus, 250-259.
- 44- Robert Lenkinski (1997). Personal communication.
- 45- Lim KO, Pauly J, Webb P, Hurd R, Macovski A. Short TE phosphorous spectroscopy using a spin-echo pulse. *Mag Reson Med* 1994;32:98-103.

- 46- Narayana PA, Jackson EF, Buttlar IJ. ^1H -MRS of muscle physiology and pathophysiology. In: Fleckenstein JL, Crues III JV, Reimer CD, eds. Muscle imaging in health and disease. 1st ed. Springer-Verlag, New York, Berlin· Heidelberg 1996; 1: 133-48.
- 47- Martin PA, Gibson H, Edwards RHT. MRS of muscle. In: Young, IR, Charles, HC, editors. MR spectroscopy clinical applications and techniques. London: Martin Dunitz Ltd.; 1996. p. 55-73.
- 48- Yoshida T. The rate of phosphocreatine hydrolysis and resynthesis in exercising muscle in humans using ^{31}P -MRS. J Physiol Anthropol Appl Hum Sci 2002;21:247-55.
- 49- Boska M. (1991) Estimating the ATP cost of force production in the human gastrocnemius/soleus muscle group using ^{31}P MRS and ^1H MRI. NMR Biomed., 4: 173-181.
- 50- Kemp G.J., C.H. Thompson. P.R.J. Barnes and G.K. Radda (1994a) Comparisons of ATP turnover in human muscle during ischaemic and aerobic exercise using ^{31}P magnetic resonance spectroscopy. Magn. Reson. Med., 31: 248-258.
- 51- Houtman CJ, Heerschap A, Zwarts MJ, Stegeman DF. An additional phase in PCr use during sustained isometric exercise at 30% MVC in the tibialis anterior muscle. NMR Biomed 2002; 15:270-7.

- 52- Street D, Bangsbo J, Juel C. Interstitial PH in human skeletal muscle during and after dynamic graded exercise. *J Physiol* 2001;537:993-8.
- 53- Clark MG, Rattigan S, Clerk LH, et al. Nutritive and non-nutritive blood flow: rest and exercise. *Acta Physiol Scand* 2000;168:519-30.
- 54- Russ DW, Vandenborne K, Walter GA, Elliott M, Binder-Macleod SA. Effects of muscle activation on fatigue and metabolism in human skeletal muscle. *J Appl Physiol* 2002;92:1978-86.
- 55- Lenkinski RE, Schnall MD. Magnetic resonance spectroscopy. In: Kricun ME. *Imaging of bone tumours*. 1st ed. W.B. Saunders Company, Philadelphia, London, Toronto, Montreal, Sydney, Tokyo 1993; 9:447-73.
- 56- Cohen JS. Phospholipid and energy metabolism of cancer cells monitored by ³¹P magnetic resonance spectroscopy: Possible clinical significance. *Mayo Clin Proc* 1988;63: 1199-207.
- 57- Clarke GD, Bonte FJ, Fleckenstein JL. Ionizing radiation imaging techniques in the evaluation of muscle disorders. In: Fleckenstein JL, Crues III JV, Reimers CD, eds. *New York, Berlin, Heidelberg* 1996;1:3-12.

- 58- Bertocci LA, Haller RG, Lewis SF. Abnormal high energy phosphate metabolism in human muscle phosphofructokinase deficiency. *J Appl Physiol* 1991;70:1201-7.
- 59- Blei ML, Conley KE, Kushmerick MJ, Separate measures of ATP utilization and recovery in human skeletal muscle. *J Physiol (Lond)* 1993;456:203-22.
- 60- Kemp GJ, Radda GK. Quantitative interpretation of bioenergetic data from ^{31}P and ^1H magnetic resonance spectroscopic studies of skeletal muscle: an analytical review. *Mag Res Quart* 1994;10:43-63.
- 61- Taylor DJ, Bore PJ, Styles P, Gadian DG, Radda GK. Bioenergetics of intact human muscle. A ^{31}P nuclear magnetic resonance study. *Mol Biol Med* 1983;1:77-94.
- 62- Iotti S., Lodi R., Frassinetti C., Zaniol P., Barbiroli B., In vivo assessment of mitochondrial functionality in human gastrocnemius muscle by ^{31}P MRS. *NMR Biomed.* 6 (1993) 248-253.
- 63- Hutson S.M., Williams G.D., Berkich D.A., LaNoue K.F., Briggs R.W., A ^{31}P NMR study of mitochondrial inorganic phosphate visibility: effects of Ca^{2+} , Mn^{2+} , and the pH gradient, *Biochemistry* 31 (1992) 1322-1330.
- 64- Wohlrab H., Flowers N., pH gradient-dependent phosphate transport catalyzed by the purified mitochondrial phosphate transport protein, *J. Biol. Chem.* 257 (1982) 28-31.

- 65- Barbiroli B., Iotti S., Iodi R., In vivo assessment of human skeletal muscle mitochondria respiration in health and disease, *Mol. Cell Biochem.* 174 (1997) 11-15.
- 66- Fabrizi, G.M., Lodi, R., D'Ettorre, M., Malandrini, A., Cavallaro, T., Rimoldi, M., Zaniol, P., Barbiroli, B., Guazzi, G., 1996. Autosomal dominant limb girdle myopathy with ragged-red fibers and cardiomyopathy. A pedigree study by in vivo ^{31}P -MR spectroscopy indicating a multisystem mitochondrial defect. *J. Neurol. Sci.* 137 (1), 20-27.
- 67- Argov, Z., De Stefano, N., Arnold, D.L., 1996. ADP recovery after a brief ischemic exercise in normal and diseased human muscle-a ^{31}P MRS study. *NMR Biomed.* 9(4), 165-172.
- 68- Simon, D.K., Johns, D.R., 1999. Mitochondrial disorders: clinical and genetic features. *Annu. Rev. Med.* 50, 111-127.
- 69- Zeviani, M., Tiranti, V., Piantadosi, C., 1998. Mitochondrial disorders. *Medicine (Baltimore)* 77 (1), 59-72.
- 70- Adams GR, Foley JM, Meyer RA. Muscle buffer capacity estimated from pH changes during rest-to-work transitions. *J Appl Physiol* 1990;69:968-72.
- 71- McCully K, Vandenborne K, De Meirleir K. Muscle metabolism in track athletes using ^{31}P NMR spectroscopy. *Can J Physiol Pharmacol* 1992;70:1353-9.

References

- 72- Barbiroli B, Funicello R, Iotti S. ^{31}P NMR spectroscopy of skeletal muscle in Becker dystrophy and DMD/BMD carriers: altered rate of phosphate transport. *J Neurol Sci* 1992;109:188-95.
- 73- Agrov Z, Bank WJ, Maris J, Chance B. Muscle energy metabolism in McArdle's syndrome by in vivo phosphorous magnetic resonance spectroscopy. *Neurology* 1987;37: 1720-4.
- 74- Agrov Z, Bank WJ, Boden B, Young R, Chance B. ^{31}P magnetic resonance spectroscopy of partially blocked muscle glycolysis. *Arch Neurol* 1987;44:614-7.
- 75- Chance B, Younkin DP, Kelley R. Magnetic resonance spectroscopy of normal and diseased muscles. *Am J Med Genet* 1986;25:659-79.
- 76- Radda GK, Bore PJ, Gadian DG. ^{31}P NMR examination of two patients with NADH-CoQ reductase deficiency. *Nature* 1982, 295:608-9.
- 77- Arnold DL, Taylor DJ, Radda GK. Investigation of human mitochondrial myopathies by phosphorous magnetic resonance spectroscopy. *Ann Neurol* 1985; 18:189-96.
- 78- Matthews PM, Allaire C, Shoubridge EA, Karpati G, Carpenter S, Arnold DL. In vivo muscle magnetic resonance spectroscopy in the clinical investigation of mitochondrial disease. *Neurology* 1991;41:114-20.

- 79- Agrov Z, Bank WJ, Maris J. Bioenergetic heterogeneity of human mitochondrial myopathies: phosphorus magnetic resonance spectroscopy study. *Neurology* 1987; 37:257-62.
- 80- Kuhl CK, Layer G, Traber F, Zierz S, Block W, Reiser M. Mitochondrial encephalomyopathy: correlation of P-31 exercise MR spectroscopy with clinical findings. *Radiology* 1994;192:223-30.
- 81- Lewis SF, Haller RG, Cook JD, Nunnally RL. Muscle fatigue in McArdle's disease studied by ^{31}P -NMR: effect of glucose infusion. *J Appl Physiol* 1985;59:1991-4.
- 82- Chance B, Eleff S, Bank W, Leigh JS, Warnell R. ^{31}P NMR studies of control of mitochondrial function in phosphofructokinase-deficient human skeletal muscle. *Proc Natl Acad Sci USA* 1982;79:7714-8.
- 83- Chinnery PF, Turnbull DM. Epidemiology and treatment of mitochondrial disorders. *Am J Med Genet* 2001;106:94-101.
- 84- Barbiroli, B., Frassinetti, C., Martinelli, P., Iotti, S., Lodi, R., Cortelli, P., Montagna, P., 1997. Coenzyme Q10 improves mitochondrial respiration in patients with mitochondrial cytopathies. An in vivo study on brain and skeletal muscle by phosphorous magnetic resonance spectroscopy. *Cell Mol. Biol. (Noisy-le-grand)* 43 (5), 741-749.

- 85- Barbiroli, B., Iotti, S., Lodi, R., 1999b. Improved brain and muscle mitochondrial respiration with CoQ. An in vivo study by ³¹P-MR spectroscopy in patients with mitochondrial cytopathies. *Biofactors* 9 (2-4), 253-260.
- 86- Arahata K, Ishiura S, Ishiguro T, Suhara Y, Eguchi C, Ishihara T, Nonaka I, Ogawa E, Sugita H. Immuno staining of skeletal and cardiac muscle surface membrane with antibody against Duchenne muscular dystrophy at the muscle cell membrane. *Nature* 1988;333:861-3.
- 87- Hoffman EP, Brown RL Jr, Kunkel LM. Dystrophin the protein product of Ducheene muscular dystrophy locus. *Cell* 1987;51:919-28.
- 88- Ervasti JM, Campbell KP. Membrane organization of the dystrophin-glycoprotein complex. *Cell* 1991;66: 1121-31.
- 89- Ervasti JM, Kahl SD, Campbell KP. Purification of dystrophin from skeletal muscle. *J Biol Chem* 1991;266:9161-5.
- 90- Matsumura K, Campbell KP. Dystrophin-glycoprotein complex: its role in the molecular pathogenesis of muscular dystrophies. *Muscle and Nerve* 1994;17:2-15.
- 91- Fong P, Turner PR, Denetclaw WF, Steinhardt RA. Increased activity of calcium channels in myotubes of Duchenne human and mdx mouse origin. *Science* 1990;250:673-6.

- 92- Franco A, Lansman JB. Calcium entry through stretch inactivated ion - channels in mdx myotubes. *Nature* 1990;344:670-3.
- 93- Willig, T.N.; Carlier, L.; Legrand, M.; Riviere, H.; Navarro, J. Nutritional assessment in Duchenne muscular dystroph. *Dev. Med. Child. Neurol.* 35:1074-1082; 1993.
- 94- Argov Z, Bank WJ. Phosphorus magnetic resonance spectroscopy (^{31}P MRS) in neuromuscular disorders. *Ann Neurol* 1991;30:90-7.
- 95- Argov Z, Lofberg M, Arnold DL. Insights into muscle diseases gained by phosphorus magnetic resonance spectroscopy. *Muscle and Nerve* 2000;23:1316-34.
- 96- Kemp GJ, Taylor DJ, Dunn JF, Frostick SP, Radda GK. Cellular energetics of dystrophic muscle. *J Neurol Sci* 1998;116:201-6.
- 97- Lodi R, Kemp GJ, Muntoni F, Thompson CH, Rae C, Taylor J, Styles P, Taylor DJ. Reduced cytosolic acidification during exercise suggests defective glycolytic activity in skeletal muscle of patients with Becker muscular dystrophy. An in-vivo ^{31}P magnetic resonance spectroscopy study. *Brain* 1999;122:121-30.
- 98- Barnes PR, Kemp GJ, Taylor DJ, Radda GK. Skeletal muscle metabolism in myotonic dystrophy. A ^{31}P magnetic resonance spectroscopy study. *Brain* 1997; 120:1699-711.

- 99- Taylor DJ, Kemp GJ, Woods CG, Edwards JH, Radda GK. Skeletal muscle bioenergetics in myotonic dystrophy. *J Neurol Sci* 1993; 116: 193-200.
- 100- Younkin DP, Berman P, Sladky J, Chee e, Bank W, Chance B. ³¹P NMR studies in Duchenne muscular dystrophy: Age-related metabolic changes. *Neurology* 1987;37: 165-9.
- 101- Bchark L, Den Hollander JA, Stromski ME, Ogino T, Shulman RG, Perrott OAC, Prichard JW. High resolution ¹H nuclear magnetic resonance study of cerebral hypoxia in-vivo. *Proc Natl Acad Sci USA* 1983;80.
- 102- Mole PA, Chung Y, Tran TK, Sailasuta N, Hurd R, Jue T. Myoglobin desaturation with exercise intensity in human gastrocnemius muscle. *Am J Physiol* 1999; 277(pt 2):R173-R180.
- 103- Wang Z, Noyszewski EA, Leigh JS. In vivo MRS measurement of deoxymyoglobin in human forearms. *Magn Reson Med* 1990; 14:562-567.
- 104- Brillault-Salvat C, Giacomini E, Jouvensal L, Wary C, Bloch G, Carrier PG. Simultaneous determination of muscle perfusion and oxygenation by interleaved NMR plethysmography and deoxymyoglobin spectroscopy. *NMR Biomed* 1997;10:315-323.

- 105-Kreis R, Bruegger K, Skjelsvik C, et al. Quantitative ^1H magnetic resonance spectroscopy of myoglobin de-and re-oxygenation in skeletal muscle: reproducibility and effects of location and disease. *Magn Reson Med* 2001;46:240-248.
- 106- Argov Z, Lofberg M, Arnold DL. Insights into muscle diseases gained by phosphorus magnetic resonance. *Muscle Nerve* 2000; 23:1316-1334.
- 107-Holmes GP, Kaplan JE, Gantz NM. Chronic fatigue syndrome: a working case definition. *Ann Intern Med* 1988;108:387-9.
- 108-Stokes MJ, Cooper RG, Edwards RHT. Normal muscle strength and fatigability in patients with effort syndromes. *Br Med J* 1988;297:1014-7.
- 109-Arnold DL, Radda GK, Bore PJ, Styles P. Excessive intracellular acidosis of skeletal muscle on exercise in a patient with a post-viral exhaustion/fatigue syndrome. *Lancet* 1984;2:1367-9.
- 110-Kent-Braun JA, Sharma KR, Weiner MW, Massie B, Miller RG. Central basis of muscle fatigue in chronic fatigue syndrome. *Neurology* 1993;43: 125-31.
- 111-Lloyd AR, Gandevia SC, Hales JP. Muscle performance, voluntary activation, twitch properties and perceived effort in normal subjects and patients with the chronic fatigue syndrome. *Brain* 1991; 114:85-98.

- 112-Jamal GA, Hansen S, Apartopoulos F, Peden A. The “Gulf War syndrome”. Is there is evidence of dysfunction in the nervous system?. *J Neurol Neurosurg Psychiatry* 1996;60:449-51.
- 113-Hubert M, Lison D. Study of muscular effects of short term pyridostigmine treatment in resting and exercising rats. *Human Exp Toxicol* 1995;14:49-54.
- 114-Yeckel CW, Weiss R, Dziura J, et al. Validation of insulin sensitivity indices from oral glucose tolerance test parameters in obese children and adolescents. *J Clin Endocrinol Metab* 2004; 89:1096-1101.
- 115-Decombaz J, Schmitt B, Ith M, et al. Post-exercise fat intake repletes intramyocellular lipids but no faster in trained than in sedentary subjects. *Am J Physiol Regul Integr Comp Physiol* 2001;281:R760-R769.
- 116-Howald H, Boesch C, Kreis R, et al. Content of intramyocellular lipids derived by electron microscopy, biochemical assays, and (1)H-MR spectroscopy. *J Appl Physiol* 2002; 92:2264-2272.
- 117-Boesch C, Slotboom J, Hoppeler H, Kreis R. In vivo determination of intra-myocellular lipids in human muscle by means of localized ¹H-MR-spectroscopy. *Magn Reson Med* 1997; 37:484-493.

- 118-Szczepaniak LS, Babcock EE, Schick F, et al. Measurement of intracellular triglyceride, stores by H spectroscopy: validation in vivo .Am J Physiol1999; 276(5 pt 1):E977-E989.
- 119-Schick F, Eismann B, Jung WI, Bongers H, Bunse M, Lutz O. Comparison of localized proton NMR signals of skeletal muscle and fat tissue in vivo: two lipid compartments in muscle tissue. Magn Reson Med 1993;29:158-167.
- 120-Sinha R, Dufour S, Petersen KF, et al. Assessment of skeletal muscle triglyceride content by (1) H nuclear magnetic resonance spectroscopy in lean and obese adolescents: relationships to insulin sensitivity, total body fat, and central adiposity. Diabetes 2002; 51:1022-1027.
- 121-Krssak M, Falk Petersen K, Dresner A, et al. Intramyocellular lipid concentrations are correlated with insulin sensitivity in humans: a ¹H NMR spectroscopy study. Diabetologia 1999; 42:113-116.
- 122-Jacob S, Machann J, Rett K, et al. Association of increased intramyocellular lipid content with insulin resistance in lean nondiabetic offspring of type 2 diabetic subjects. Diabetes 1999; 48:1113-1119.
- 123-Levin K, Daa Schroeder H, Alford FP, Beck Nielsen H. Morphometric documentation of abnormal intramyocellular fat storage, and reduced glycogen in obese patients with type II diabetes. Diabetologia 2001;44:824-833.

- 124-Rico-Sanz J, Hajnal JV, Thomas EL, Mierisova S, Ala-Korpela M, Bell JD. Intracellular and extracellular skeletal muscle triglyceride metabolism during alternating intensity exercise in humans. *J Physiol* 1998; 510(pt 2):615-622.
- 125-Schrauwen-Hinderling VB, van Loon LJ, Koopman R, Nicolay K, Saris WH, Kooi ME. Intramyocellular lipid content is increased after exercise in nonexercising human skeletal muscle. *J Appl Physiol* 2003;95:2328-2332.
- 126-Luzi L, Perseghin G, Tambussi G, et al. Intramyocellular lipid accumulation and reduced whole body lipid oxidation in HIV lipodystrophy. *Am J Physiol Endocrinol Metab* 2003;284:E274-E280.
- 127-K. Brechtel, A. Niess, 1. Machann, S. Jacob, H.-U. Haring, H. H. Dickhuth, C. D. Claussen, and F. Schick, Direct assessment of intramyocellular lipids (IMCL) during exercise in well trained male runners. A ¹H-MRS study, *Horm. Metab. Res.* 33,63-66 (2001).
- 128-M. Krssak, K. F. Petersen, R. Bergeron, T. Price, D. Laurent, D. L. Rothman, M. Roden, and G. I. Shulman, Intramuscular glycogen and intramyocellular lipid utilization during prolonged exercise and recovery in man: A ¹²C and ¹H nuclear magnetic resonance spectroscopy study, *J. Clin. Endocrinol. Metab.* 85(2),748-754 (2000).

- 129-Ng TC, Evanochko WT, Hiramoto RN, et al. ^{31}P NMR spectroscopy of in vivo tumors. *J Magn Reson* 1982; 49:271-286.
- 130-Sijens, PE, Bovee WMMJ, Seijkens D. Los G, Rutgers DH. In vivo ^{31}P -nuclear magnetic resonance study of the response of a murine mammary tumor to different doses of γ -radiation. *Cancer Res* 1986;46:1427-1432.
- 131-Semmler W. Gademann G. van Kaick G, Zabel H-J, Lorenz WJ. In vivo ^{31}P -spectroscopy of human tumors and their response to therapy using a 1.5 Tesla whole body scanner (abstr.). In: Book of abstracts: Society of Magnetic Resonance in Medicine. Vol. 1. Berkeley. Calif.: Society of Magnetic Resonance in Medicine. 1986;39.
- 132-Ross B, Marshall V, Smith M, Bartlett S, Freeman D. Monitoring response to chemotherapy of intact human tumors by ^{31}P nuclear magnetic resonance. *Lancet* 1984;1:641-646.
- 133-Singer S. New diagnostic modalities in soft tissue sarcoma. *Semin Surg Oncol* 1999; 17:11-22.
- 134-Mukherji SK, Schiro S, Castillo M, et al. Proton MR spectroscopy of squamous cell carcinoma of the upper aerodigestive tract: in vitro characteristics. *AJNR Am J Neuroradiol* 1996; 17:1485-1490.

- 135-Mackinnon WB, Barry PA, Malycha PL, et al. Fine-needle biopsy specimens of benign breast lesions distinguished from invasive cancer ex vivo with proton MR spectroscopy. *Radiology* 1997; 204:661-666.
- 136-Millis K, Weybright P, Campbell N, et al. Classification of human liposarcoma and lipoma using ex vivo proton NMR spectroscopy. *Magn Reson Med* 1999; 41:257-267.
- 137-Oya N, Aoki J, Shinozaki T, et al. Preliminary study of proton magnetic resonance spectroscopy in bone and soft tissue tumors: an unassigned signal at 2.0-2.1 ppm may be a possible indicator of malignant neuroectodermal tumor. *Radiat Med* 2000; 18:193-198.
- 138-Negendank W. Studies of human tumors by MRS: a review. *NMR Biomed* 1992;5:303-324.
- 139-Maheshwari SR, Mukherji SK, Neelon B, et al. The choline/creatine ratio in five benign neoplasms: comparison with squamous cell carcinoma by use of in vitro MR spectroscopy. *AJNR Am J Neuroradiol* 2000; 21:1930-1935.
- 140-Sharma U, Atri S, Sharma MC, Sarkar C, Jagannathan NR. Skeletal muscle metabolism in Duchenne muscular dystrophy (DMD): an in-vitro proton NMR spectroscopy study. *Magn Reson Imaging* 2003;21:145-153.