

References

- Abou Steit M.H., El Shiaty A.F., El Desouki M.M and Said D.G., (2002):** “use of corneal topography in postoperative assessment of LASIK for myopia and myopic astigmatism” Bull. Ophthalmol. Soc. Egypt; 95(1):65-72.
- Agarwal A. (2000):** Excimer laser: fundamental concepts” In “Refractive surgery” Agarwal S., Agarwal A., Pallikaris I.G., Neuhaun T.H., Knorz MC, and Agarwal A., Ch., 12, page. 124-128, Jaypee Bros. Medical publishers. Eds.
- Aramberri J., (2003):** “Intraocular lens power calculation after corneal refractive surgery: Double-K method” J. Cataract Refract. Surge.; 29:2063-2068.
- Binkhorst R.D (1975):** “The optical design of IOL implant” In “Ophthalmic surgery” Vol.6 ch.3, page. 17-31.
- Bogan J., Waring G., Ibrahim O., Drews C., and Crutis L., (1990):** “Classification of normal corneal topography based on computer assisted videokeratography” Arch. Ophthalmol.; 108:945-949.
- Borasio E., Stevens J., and Smith G.T., (2006):** “Estimation of true corneal power after keratorefractive surgery in eyes requiring cataract surgery: BESSt formula” J. Cataract Refract. Surge.; 32:2004-2014.
- Bores L.D (2001):** “Topography the fine art of corneal surface measurements” in “Refractive eye surgery” Bores L.D, Smith R. and Barraquer J. 2nd edition, ch.6 page. 126-140, Blackwell Scientific Publications, London, Edinburgh, Paris.

Byrne S.F., and Green R.L., (2002): “Ultrasound of the eye and orbit”
Byrne S.F., and Green R.L., 2nd edition Ch.1, P.1-8,
Mosby, Inc. USA.

**Cairns G.C, McGhee C.N., Collins M.J., Owens H., and Gamble G.D.,
(2002):** “Accuracy of orbscan II slit-scanning elevation
topography” J. Cataract Refract. Surge.; 28:2181-2187.

Chalita M.R, Chavala S., Meng XU., and Krueger R.R., (2004): “Wave
front analysis in post-LASIK eyes and its correlation with
visual symptoms, refraction and topography”
Ophthalmology; 111:447-453.

Chan C.C, Hodge C., and Lawless M., (2006): “Calculation of
intraocular lens power after corneal refractive surgery”
Clin. Experiment Ophthalmol.; 34(7):640-644.

Cho HJ., Koo OS., (2005): “comparison of refraction –derived
keratometric value and orbscan corneal power after laser in
situ keratomileusis” J. Korean Ophthalmol. Soc.;
46(4):623-628.

Cho Hj., Oh CH., and Koo OS., (2005): “Intraocular lens power
calculation in the absence of pre-PRK data in moderate
myopic eyes” J. Korean Ophthalmol. Soc.; 46(4):643-648.

Coleman D.J, Lizzi F.L., and Jack R.L., (1977): “Ultrasonography of
the eye and orbit” Lea and Febiger, Philadelphia, P.7-11.

Colendrander M.C (1973): “Calculation of the power of an iris clip lens
for distant vision” Br. J. Ophthalmol. 57:735-740.

Corbett M.C., Rosen E.S., and O’Brart D.P. (1999): “Normal
topography” In “Corneal topography Principles and

Applications” Corbett M.C., ch.5, Page. 63-79, the BMJ Books, London.

Dingeldein S.A and Klyce S.D (1989): “The topography of normal corneas” Arch. Ophthalmology, 107:512-518.

Dowidar M., (2003): “Refractive and corneal topographic stabilization after discontinuation of contact lenses” Bull. Ophthalmol., Soc. Egypt; 96(1):81-87.

Eghbali F. Yeungh K.K and Maloney R.K (1995): “Togographic determination of corneal asphericity and its lack of effect on the outcome of radial keratotomy” Am. J. ophthalmol., 119:275-280.

Feiz V., Mannis M.J., Garcia F.F., Kandavel G., Darlington J.K., Kim E., Caspar J., Wang J.L., and Wang W., (2001): “Intraocular lens power calculation after laser in situ keratomileusis for myopia and hyperopia: a standardized approach” Cornea; 20:792-797.

Fishman G.R., Pans M.E., Seedor J.A., Liebmann J.M., and Ritch R., (2005): “Assessment of central corneal thickness using OCT” J. Cataract Refract. Surg.; 31:707-711.

Fredrick S., Wilson G., (1983): “The relation between eyelid tension, corneal toricity and age” Invest. Ophthalmol. Vis. Sci.; 24:1367-1373.

Gobbi P.G., Carones F., and Brancuto R. (1998): “Keratometric index, videokeratography and refractive surgery” J. Cataract Refract. Surg.; 24:202-211.

Haigis W. (2003): “Corneal power after refractive surgery for myopia: contact lens method” J. Cataract Refract. Surg., 29:1397-1411.

Haigis W., Lege B., Miller N, and Schneider B., (2000): “Comparison of immersion ultrasound biometry and partial coherence interferometry for intraocular lens calculation according to Haigis” Graefes Arch. Clin. Exp. Ophthalmol.; 238:765-773.

Hamed A.M, Wang L., Misra M., and Koch D.D (2002): “A comparative analysis of five methods of determining corneal refractive power in eyes that have undergone myopic laser in situ keratomileusis” Ophthalmology; 109:651-658.

Hamilton D.R., and Hardten D.R., (2003): “Cataract surgery in patients with prior refractive surgery” Curr. Opin. Ophthalmol.; 14:44-53.

Helena M.C., Meisler D., Wilson SE., (1997): “Epithelial growth within the lamellar interface after laser in situ keratomileusis (LASIK)”Cornea;3:300-305.

Hersh P.S., Scher K.S., and Irani R., (1998): “Corneal topography of photorefractive keratectomy versus laser in situ keratomileusis; Summit PRK-LASIK study group” Ophthalmology; 105:612-619.

Hill W.E., (2003): “IOL master” Techniques in Ophthalmology; 1:62-67.

Hill W.E., (2005): “Understanding the Haigis formula”
<http://www.doctor-hill.com.Haigisformula.htm>.

- Hoffer K.J., (2000):** “Intraocular lens power calculation” In “Phacoemulsification, laser cataract surgery and foldable IOL” Agrawal S., Agrawal A., Sachde M.S., Mehta K.R., Fine I.H., and Agrawal A., 2nd edition, Ch.3, P.17-25. JAYPEE Brothers Medical Publishers New Delhi.
- Hoffer K.J., (2002):** “Calculating intraocular lens power after refractive corneal surgery” Arch. Ophthalmol.; 120:500-501.
- Holladay J.T, Praeger T.C, Chandler T.Y, Masgrove K.H, Lewis J.W, and Ruiz R.S. (1988):** “A three part system for refining intraocular lens power calculation” J. Cataract Refract. Surg. 14:17-24.
- Holladay J.T., (1989):** “Consultations in refractive surgery” Refract. Corneal Surg.; 5-203.
- Holladay J.T., (1993):** “Refractive power calculation for intraocular lenses in the phakic eye” Am. J. Ophthalmol.; 116:63-66.
- Holladay J.T., (1997):** “Corneal topography using the Holladay Diagnostic Summary” J. Cataract Refract. Surg.; 23:209-221.
- Holladay J.T., (1997):** “Standardizing constants for ultrasonic biometry, keratometry and intraocular lens power calculation” J. Cataract Refract. Surg.; 23:1356-1370.
- Hugger P., Kohmen T., LaRosa F.A., (2000):** “Comparison of changes in manifest refraction and corneal power after photorefractive keratectomy: Am. J. Ophthalmol.; 129:68-75.

- Jarade E., and Tabbara F., (2001):** “Keratometry reading following laser in situ keratomileusis” *Invest. Ophthalmol. Vis. Scien.*; 42:604.
- Jarade E.F., and Tabbara K.F., (2002):** “Intraocular lens calculation after corneal refractive surgery” *Middle East J. of Ophthalmology*; 10(2):106-111.
- Jonathan D.C., Stulting R.D., Thompson K.P. and Waring G.O. (2003):** “Laser in situ keratomileusis, surgical technique” In “Ophthalmic Surgery: Principles and Practice” George L. Spaeth, 3rd edition, ch.19, P.177-184, SAUNDERS, USA.
- Jonsson M., and Behndig A., (2005):** “Pachymetric evaluation prior to laser in situ keratomileusis” *J. Cataract Refract. Surg.*; 31:701-706.
- Joslin C.E., Koster J., and TU., E.Y., (2005):** “Contact lens overrefraction variability in corneal power estimation after refractive surgery” *J. Cataract Refract. Surge.*; 31:2287-2292.
- Kalski RS., Danyoux JP., Franenkel GE., Lawless M.A., and Rogers C., (1997):** “Intraocular lens power calculation for cataract surgery after photorefractive keratectomy for high myopia” *J. Refr. Surg.*; 13(4):362-366.
- Kawana K., Tokunaga T., Miyato K., Okomoto F., Kiuchi T., and Oshiks T., (2004):** “Comparison of corneal thickness measurements using Orbscan 11, non-contact specular microscopy, and ultersonic pachymetry in eyes after laser in situ keratomileusis” *Br. J. Ophthalmol.*; 88:466-468.

- Kim J.H., Lee D.H., and Joo C.K., (2002):** “Measuring corneal power for intraocular lens power calculation after refractive surgery” J. Cataract Refract. Surg.; 28:1932-1938.
- Koch D.D., and Wang L., (2003):** “Calculating intraocular lens power in eyes that have had refractive surgery: J. Cataract refract. Surge.; 29:2039-2042.
- Krone S.S., Moreno G.L., Balbi O.V.B., Arce C.G., Schor P., and Campos M., (2004):** “A direct method to measure the power of the central cornea after myopic laser in situ keratomileusis” Arch. Ophthalmol.; 122:159-166.
- Ku HC., Cho SH., Kim HJ., Kim JH., and Joo CK., (2005):** “Relationship between clinical history method and Orbscan 11 for measuring corneal power after LASIK” J. Korean Ophthalmol. Soc.; 46(4):581-588.
- Langenbucher A., Torres F., Behrens A., Suarez E., Haigis W., and Seitz B., (2004):** “Consideration of the posterior corneal curvature for assessment of corneal power after myopic LASIK” Acta Ophthalmologica Scand.; 82:264-269.
- Liesegang T.J., Skuta G.L., and Cantor L.B (2004):** “Refractive surgery” In “Basic and clinical science course in American academy of ophthalmology” Sec 14, ch.11, p:199-209.
- Liu Z., Huang A.J., and Pflugfelder S.C., (1999):** “Evaluation of corneal thickness and topography in normal eyes using the Orbscan corneal topography system” Br. J. Ophthalmol.; 83:774-778.

- Mackool R.J., and Mackool R., (2006):** “Intraocular lens power calculation after laser in situ keratomileusis: Aphakic refraction technique” J. Cataract Refract. Surg.; 32(3):371-372.
- Maeda N., Klyce SD., Smolek MK., and McDonald MB., (1997):** “Disparity between keratometry – style readings and corneal power within the pupil after refractive surgery for myopia” Cornea; 16:517-524.
- Mandell R.B (1994):** “Corneal power correction factor for photorefractive keratectomy” J. Refract. Corneal Surg.; 10:125-128.
- Mccarey B.E. (1997):** “Corneal optics, nutrition and alloplastic material dynamics” In “Principles and Practice of refractive surgery” Elander R., Rich F.L., and Robin B.J., ch.3 page. 27-37, W.B. SAUNDERS, Philadelphia.
- McGhee CNJ, and Bryce IG., (1996):** “Natural history of central topographic islands following excimer laser potorefractive keratectomy” J. Cataract Referact. Surg.; 22:1151-1158.
- Merlin, U., (1996):** "Corneal topography & astigmatism" in "Corneal Topography: the clinical Atlas" Buratto Lucio (ed.), SLACK Incorporated, New Jersey, Ch.4, P. 137-213.
- Mohrman R., (1985):** “The keratometer” In “Clinical Ophthalmology,” Vol. (1), Ch.60, P.1-12., Done, T.D., Harper and Row Publishers, Philadelphia, London.
- Mundt G.H., and Hughes W.F., (1956):** “Ultrasonics in ocular diagnosis” Am. J. Ophthalmol.; 41:488-498.

- Munnerlyn CR., Kooms S.J., and Marshall J., (1988):** “Photorefractive keratectomy: A technique for laser refractive surgery” J. Cataract Refract. Surg.; 14(1):46-52.
- Odenthal M.T., Eggink C.A., Melles G., Pameyer J.H., Geerards A.J.M., and Beekhuis H., (2002):** “Clinical and theoretical results of intraocular lens power calculation for cataract surgery after photorefractive keratectomy for myopia” Arch. Ophthalmol.; 120:431-438.
- Olsen T. (1987):** “Theoretical approach to intraocular lens calculation using Gaussian optics” J. Cataract Refract. Surg. 13:141-145.
- Olsen T. (1992):** “Sources of error in intraocular lens power calculation” J. Cataract Refract. Surg.; 18:125-129.
- Olsen T., (1986):** “On the calculation of power from curvature of the cornea” British J. Ophthalmol.; 70:152-154.
- Olsen T., Olsen H., Theim K., and Corydon L., (1992):** “Prediction of pseudophakic anterior chamber depth with the newer IOL calculation formulas” J. Cataract refract. Surg.; 18:280-285.
- Packer M., Brown L.K., Hoffman R.S., and Fine I.H., (2004):** “Intraocular lens power calculation after incisional and thermal keratorefractive surgery” J. Cataract Refract. Surg.; 30:1430-1434.
- Pallikaris I., (2006):** "Epi-LASIK Vs LASEK for treatment of myopia". Highlights of ophthalmology; 34(1):2-3.
- Patel S., and Alio JL., (2002):** “Factors in a hyperopic shift after LASIK (Letter)” Ophthalmology; 109:407-408.

- Pedrotti L.S and Pedrotti F.L (1998):*** “Optics and vision” ch.10, page 194-223, Prentice HALL, Inc.
- Preussner P.R., Wahl J., and Weitzel D., (2005):*** “Topography – based intraocular lens power selection” J. Cataract Refract. Surg.; 31:525-533.
- Probst L.E, Machat J.J, (1998):*** “Mathematics of laser in situ keratomileusis for high myopia” J. Cataract Refract. Surg.; 24(2):190-195.
- Rabinowitz Y., Yang H., Brickman Y., Akkina J., Riley C., Rotter J., and Elashoff J., (1996):*** “Videokeratography database of normal human corneas” Br. J. Ophthalmol.; 80:610-616.
- Randleman J.B, Loupe D.N, Song C.D, Waring G.O. and Stulting R.D. (2002):*** "Intraocular lens power calculations after laser insitu keratomileusis" Cornea; 21(8):751-752.
- Retzlaff J., and Prust J., (1995):*** “Intraocular lens implant power calculation, selection and ocular biometry” In “Cataract surgery: Technique, complications and management” Steinert R.F, Fine H., Gimbel H.V., Lindstrom R.L., Neubann T.F., and Osher R.H., Ch.3, P.22-33. W.B. Saunders Company.
- Retzlaff J.A., Sander D.R, and Kraff M.C.(1990):*** “Development of SRK/T intraocular lens implant power calculation formula” J. Cataract Refract. Surg. 16:333-340.
- Rosa N., Capasso L., Lanza M., Iaccarino G., and Romano A., (2005):*** “Reliability of a new correcting factor in calculating intraocular lens power after refractive corneal surgery” J. Cataract Refract. Surg.; 31:1020-1024.

- Rowsey J., Reynolds A., and Brown S., (1981):** “Corneal topography” Arch., Ophthalmol.; 99:1093-1100.
- Sarayba MA., Ignacio TS., Binder PS., Tran DB., (2007):** "Comparative study of stromal bed quality by using mechanical, IntraLase femtosecond laser 15- and 30-kHz microkeratomes" Cornea;4:446-451.
- Schultze R., (1998):** “Accuracy of corneal elevation with four corneal topography Systems” J. Refract. Surg.; 14(2):100-104.
- Seiler T., Koufala K., and Richter G., (1998):** “Iatraogenic keratectasia after laser in situ keratomileusis” J. Refract Surg.; 14:312-317.
- Seitz B., and Langenbucher A., (2000):** “Intraocular lens power calculation in eyes after corneal refractive surgery” J. Refract. Surg.; 16:349-361.
- Seitz B., Langenbucher A., and Haigis W., (2002):** “Pitfalls of IOL power prediction after photorefractive keratectomy for high myopia” Klin Monatsbl Augenheilkd; 219(12):840-850.
- Seitz B., Langenbucher A., Nguyen N., and Kus M.M., (1999):** “Underestimation of intraocular lens power for cataract surgery after myopic photorefractive keratectomy” Ophthalmology; 106:693-702.
- Seitz B., Torres F., Langenbucher A., Behrens A., and Suarez E., (2001):** “Posterior corneal curvature changes after myopic laser in situ keratomileusis” Ophthalmology; 108:666-673.

- Shammas H.J., (1984):** “Basic physics, technology and instrument of ultrasound and axial length measurement” In “Atlas of Ophthalmic Ultrasonography and Biometry; the C.V. Mosby Co., St., Louis Toronto.; P.2-10.
- Shammas H.J., (2001):** “Intraocular lens power calculations” In “Intraocular lenses in cataract and refractive surgery” Dimitri T. Azar, Ch.4, P.51-64, W.B. Saunders Company, Philadelphia.
- Shawky M.M., Abd El-Azeem A.A., and El-Brombly T.H., (2002):** “Intraocular lens power calculation after myopic laser in situ keratomileusis” Bull. Ophthalmol. Soc. Egypt.; 95(1):73-79.
- Shiotani Y., Maeda N., Inoue T., Watanabe H., Inoue Y., Shimomura Y., and Tano Y., (2000):** “Comparison of topographic indices that correlate with visual acuity in videokeratography” Ophthalmology; 107:559-564.
- Skridge T.B., Amos J.F., and Bartlett J.D., (1991):** “Keratometry” In “Clinical procedures in optometry” Ch.15, P.135-153. J.B Lippincott Company, Philadelphia, New York, London, Hagerstown.
- Smolek M., Klyce S., and Hovis J., (2002):** “The universal standard scale proposed improvements to the American National Standards Institute (ANSI) scale for corneal topography” Ophthalmology;109:361-369.
- Speicher L., (2001):** “Intraocular lens calculation status after corneal refractive surgery” Curr. Opin. Ophthalmol.; 12:17-29.

- Srivannaboon S., Reinstein D.Z., Sutton H.F., and Holland S.P., (1999):** “Accuracy of Orbscan total optical power maps in detecting refractive changes after myopic laser in situ keratomileusis” J. Cataract Refract. Surg.; 25:1596-1599.
- Topuz H., Ozdemir M., Cinal A., and Gumusalan Y., (2004):** “Age related differences in normal corneal topography” Ophthalmic surgery Lasers and Imaging; 35(4):298-303.
- Trokel S.L, Srinivasan R., and Braven B. (1983):** “Excimer laser surgery of the cornea” Am. J. Ophthalmol.; 96:710-715.
- Ucakhan O.O., Ozkan M., and Kanpolat A., (2006):** “Corneal thickness measurements in normal and keratoconic eyes: Pentacam comprehensive eye scanner versus noncontact specular microscopy and ultrasound pachymetry” J.Cataract Refract. Surg.;32:970-977.
- Wallace R.B. III (2005):** “Refractive cataract surgery” <http://www.opthalmichyperguides.com/tutorials/cataract.refractivecatarat/tutorial.asp>
- Walter K.A., Gagnon M.R., Hoopes PC., and Dickinson PJ., (2006):** “Accurate intraocular lens power calculation after myopic laser in situ keratomileusis, by passing corneal power J. Cataract Ref. Surg. 32(3):425-429.
- Wang L., Booth M.A., Koch D.D., (2004):** “Comparison of intraocular lens power calculation methods in eyes that have undergone LASIK” Ophthalmology; 111:1825-131.
- Waring G.O., III (1992):** “Refractive keratotomy for myopia and astigmatism” P.77-89., Mosby yearbook 1st ed., Louis Baltimore, London.

References

- Wilson S., Klyce S., and Hussein Z., (1993):*** “Standardized colour coded maps for corneal topography” *Ophthalmology*; 100:1723-1727.
- Young J., Siegel I (1995):*** “Three-dimensional digital subtraction modeling of corneal topography” *J. Refract. Surg.*; 11:188-193.