

References

- **Abraham T, Laskowski C, Zhan W, et al. (2003):** Myocardial contractility by strain echocardiography: comparison with physiological measurements in an in vitro model. *Am J Physiol Heart Circ Physiol*; 285:H2599–H2604.
- **Abdelmoniem A, Awad M, Neama A, et al. (1995):** Predictors of exercise tolerance after acute uncomplicated myocardial infarction, role of left ventricular diastolic function. Thesis.
- **Adams KF Jr, Lindenfeld J, Arnold JM, et al. (2006):** Executive summary: HFSA 2006 comprehensive heart failure practice guideline. *J Card Fail*; 12:10–38.
- **Adams KF, Lindenfeld J, Arnold JMO, et al. (2006):** HFSA 2006 comprehensive heart failure practice guideline. *J Cardiac Fail*; 12:e1–e122.
- **Akera T, Baskin SI, Tobin T, et al. (1973):** Temporal relationship between the inotropic effect and the in vitro binding to, and dissociation from, (Na + + K +)-activated ATPase. *Naunyn Schmiedebergs Arch Pharmacol*; 277:151-62.
- **Alam M and Hoglund C. (1992):** Assessment by echocardiogram of left ventricular diastolic function in healthy subjects using the atrioventricular plane displacement. *Am J Cardiol*; 69:565-8.
- **Alam M, Wardell J, Andersson E, et al. (2000):** Effects of first myocardial infarction on left ventricular systolic and diastolic function with the use of mitral annular velocity determined by pulsed wave Doppler tissue imaging. *J Am Soc Echocardiogr*; 13:343–52.
- **Alam M, Wardell J, Andersson E, et al. (2003):** Assessment of left ventricular function using mitral annular velocities in patients with congestive heart failure with or without the presence of significant mitral regurgitation. *J Am Soc Echocardiogr*; 16:240-245.
- **American College of Sports Medicine. (2000):** Guidelines for Exercise Testing and Prescription, 6th edn. Pennsylvania: Williams and Wilkins.

-
- **American Heart Association. (2003):** Heart disease and stroke statistics: Dallas, TX: American Heart Association.
 - **American Heart Association. (2005):** Heart disease and stroke statistics: 2005 update. Dallas, TX: American Heart Association. In: Zipes DP, Libby P, Bonow RO, Braunwald E, ed. *Braunwald's Heart Disease*, 7th ed.. Philadelphia: Elsevier; 539-568.
 - **Anand IS, Liu D, Chugh SS, et al. (1997):** Isolated myocyte contractile function is normal in post-infarct remodeled rat heart with systolic dysfunction. *Circulation*; 96:3974-3984.
 - **Appleton C, Hatle L, and Popp R. (1988):** Relation of transmitral flow velocity pattern to left ventricular diastolic function: new insights from a combined hemodynamic and Doppler echocardiographic study. *J Am Coll Cardiol*; 12:426-440.
 - **Appleton CP. (1993):** Doppler assessment of left ventricular diastolic function: The refinements continue (Editorial). *J Am Coll Cardiol*; 21(7):1697-700.
 - **Appleton C, Hatle L, Nishimura RA, et al. (1997):** The non invasive assessment of left ventricular diastolic function with two dimensional echocardiography. *J Am Soc Echocardiogr*; 10:246-270.
 - **Awan NA, Amsterdam EA, Hermanovich J, et al. (1982):** Long-term hemodynamic and clinical efficacy of captopril therapy in ambulatory management of chronic congestive heart failure. *Am Heart J*; 103:474-479. [PMID: 6801943]
 - **Bach D S, Armstrong W F, Donovan C L, et al. (1996):** Quantitive Doppler tissue imaging for assessment of regional myocardial velocity during transient ischemia and reperfusion. *Am Heart J*; 132:721-5.
 - **Bader H, Garrigue S, Lafitte S, et al. (2004):** Intra-left ventricular electromechanical asynchrony. A new independent predictor of severe cardiac events in heart failure patients. *J Am Coll Cardiol*; 43:248-56.
 - **Bakris GL, Fonseca V, Katholi RE, et al. (2004):** Metabolic effects of carvedilol vs metoprolol in patients with type 2 diabetes mellitus and
-

- hypertension: a randomized controlled trial. *JAMA*;292:2227–2236. [PMID: 15536109]
- **Bardy GH, Lee KL, Mark DB, et al. (2005):** Amiodarone or an implantable cardioverter-defibrillator for congestive heart failure. *N Engl J Med*; 352:225.
 - **Bax JJ, Abraham T, Barold SS, et al. (2005):** Cardiac resynchronisation therapy: Part 1—issues before device implantation. *J Am Coll Cardiol*; 46:2153–67.
 - **Bednarz J, Vignon P, Mor-Avi V, et al. (1998):** Color kinesis: Principles of operation and technical guidelines. *Echocardiography*; 15: 21–34.
 - **Belohlavek M, Bartleson VB, and Zobitz ME. (2001):** Real-time strain rate imaging: validation of peak compression and expansion rates by a tissue mimicking phantom. *Echocardiography*; 18:565–571.
 - **Bittner V, Weiner DH, Yusuf S, et al. (1993):** Prediction of mortality and morbidity with a 6-minute walk test in patients with left ventricular dysfunction: SOLVD Investigators. *JAMA*; 270:1702–1707.
 - **Bleeker GB, Yu CM, Nihoyannopoulos P, et al. (2007):** Optimal use of echocardiography in cardiac resynchronisation therapy. *Heart*; 93:1339–50.
 - **Bolling SF, Pagani FD, Deeb GM, et al. (1998):** Intermediate-term outcome of mitral reconstruction in cardiomyopathy. *J Thorac Cardiovasc Surg*; 115: 381–386.
 - **Bonow RO. (1991):** Radionuclide angiographic evaluation of left ventricular diastolic function. *Circulation*; 84: 1208-15.
 - **Bouchard A, Blumlein S, Schiller NB, et al. (1987):** Measurement of LV stroke volume using continuous wave Doppler echocardiography of the ascending aorta and M-mode echocardiography of the aortic valve. *J Am Coll Cardiol*; 1:75–83.
 - **Bountiukos M, Schinkel A, Bax J, et al. (2004):** Pulsed-wave tissue Doppler quantification of systolic and diastolic function of viable and nonviable myocardium in patients with ischemic cardiomyopathy. *Am Heart J*; 148:1079–84.

- **Braunwald, Ott M, and John D. (2008):** Clinical assessment of heart failure. Braunwald's Heart diseases, 8th ed; 561-580.
- **Bruch C, Stypmann J, Gradaus R, et al. (2004):** Usefulness of tissue Doppler imaging for estimation of filling pressures in patients with primary or secondary pure mitral regurgitation. *Am J Cardiol*; 93:324–8.
- **Burgess MI, Jenkins C, Chan J, et al. (2007):** Measurement of left ventricular dyssynchrony in patients with ischaemic cardiomyopathy: a comparison of real-time threedimensional and tissue Doppler echocardiography. *Heart*; 93:1191–6.
- **Burnett Jr JC, Costello-Boerrigter L, and Boerrigter G. (2003):** Alterations in the kidney in heart failure: The cardiorenal axis in the regulation of sodium homeostasis. In: Mann DL, ed. *Heart Failure: A Companion to Braunwald's Heart Disease*, Philadelphia: Saunders; 279-289.
- **Butler J, Forman DE, Abraham WT, et al. (2004):** Relationship between heart failure treatment and development of worsening renal function among hospitalized patients. *Am Heart J*; 147:331-338.
- **Calvo F, Teijeira S, Fernandez JM, et al. (2000):** Evaluation of heart involvement in gammasarcoglycanopathy (LGMD2C). A study of ten patients. *Neuromuscul Disord*; 10:560–565.
- **Cardim N, Perrot A, Ferreira T, et al. (2002):** Usefulness of Doppler myocardial imaging for identification of mutation carriers of familial hypertrophic cardiomyopathy. *Am J Cardiol*; 90:128–32.
- **Carr KW, Engler RL, Forsythe JR, et al. (1979):** Measurement of left ventricular ejection fraction by mechanical cross-sectional echocardiography. *Circulation*; 59:1196–1206.
- **Ceia F, Fonseca C, Mota T, et al. (2002):** Prevalence of chronic heart failure in Southwestern Europe: The EPICA study. *Eur J Heart Fail*; 4:531.
- **Chati Z, Zannad F, Jeandel C, et al. (1996):** Physical deconditioning may be a mechanism for the skeletal muscle energy phosphate metabolism abnormalities in chronic heart failure. *Am Heart J*; 131:560-566.

-
- **Chen Y., Sheirid M.V., and Duryes E.M. (1985):** Value of two-dimensional echocardiography in evaluating coronary artery disease. A randomized blinded analysis. *J Am Coll Cardiol*; 5(4) : 911-7.
 - **Choong CY, Herrmann HC, Weymann AE, et al. (1987):** Preload dependence of Doppler-derived indexes of left ventricular diastolic function in humans. *JAMA*; 10:800–8.
 - **Choong CY. (1994):** Left ventricle: diastolic function- its principles and evaluation. In: principles and practice of echocardiography. Weyman, AE (ed). Second edition, Lea and Febiger, Philadelphia, Bultimore, Hong Kong, London. Anonymous. Assessment of diastolic function. In: Oh JK, Seward JB, Tajik AJ, eds. The echo manual. Philadelphia: Lipincott Williams & Wilkins, 1999:45–57.
 - **Clark AL, Sparrow JL, and Coats AJ. (1995):** Muscle fatigue and dyspnoea in chronic heart failure: two sides of the same coin? *Eur Heart J*;16:49-52.
 - **Cleland JG, Pennell DJ, Ray SG, et al. (2003):** Myocardial viability as a determinant of the ejection fraction response to carvedilol in patients with heart failure (CHRISTMAS trial): randomised controlled trial. *Lancet*; 362: 14–21.
 - **Cohn JN, Ferrari R, Sharpe N, et al. (2000):** Cardiac remodeling: concepts and clinical implications: a consensus paper from an international forum on cardiac remodeling. *J Am Coll Cardiol*; 35:569-582.
 - **Cohn JN, Johnson G, Ziesche S, et al. (1991):** A comparison of enalapril with hydralazine-isosorbide dinitrate in the treatment of chronic congestive heart failure. *N Engl J Med*; 325: 303–310.
 - **Cohn J, Johnson GR, Shabetai R et al. (1993):** Ejection fraction, peak oxygen consumption, cardiothoracic ratio, ventricular arrhythmias, and plasma norepinephrine as determinants of prognosis in heart failure. *Circulation*; 87: V15–16.
 - **Colucci WS, Packer M, Bristow MR, et al, for the US Carvedilol Heart Failure Study Group. (1996):** Carvedilol inhibits clinical progression in patients with mild symptoms of heart failure. *Circulation*; 94: 2800–2806.
-

- **Costill DL. (1972):** Physiology of marathon running. *JAMA*; 221:1024–1029.
- **Cowie MR, Jourdain P, Maisel A, et al. (2003):** Clinical applications of B-type natriuretic peptide (BNP) testing. *Eur Heart J*;24:1710–1718. [PMID: 14522565]
- **Cowie MR, Mosterd A, Wood DA, et al. (1997):** The epidemiology of heart failure. *Eur Heart J*; 18:208-225.
- **Cowie MR, Struthers AD, Wood DA, et al. (1997):** Value of natriuretic peptides in assessment of patients with possible new heart failure in primary care. *Lancet*; 350:1349–1353. [PMID: 9365448]
- **Curtis JP, Sokol SI, Wang Y, et al. (2003):** The association of left ventricular ejection fraction, mortality, and cause of death in stable outpatients with heart failure. *J Am Coll Cardiol*; 42:736–742.
- **D’Hooge J, Bijmens B, and Jamal F. (2000):** High frame rate, myocardial integrated back scatter. Does this change our understanding of this acoustic parameter? *Eur J Echocardiography*, 1:32-41.
- **D’Hooge J, Heimdal A, Jamal F, et al. (2000):** Regional strain and strain rate measurements by cardiac ultrasound: principles, implementation and limitations. *Eur J Echocardiogr*; 1:154 –170.
- **Davis JA, Vodak P, Wilmore JH, et al. (1976):** Anaerobic threshold and maximal aerobic power for three modes of exercise. *J Appl Physiol*; 41:544 - 550.
- **De Backer J, Matthys D, Gillebert TC, et al. (2005):** The use of TDI for the assessment of changes in myocardial structure and function in inherited cardiomyopathies. *Eur J Echocardiogr*; 6:243–50.
- **De Sutter J, De Backer J, Van de Veire N, et al. (2005):** Effects of age, gender and left ventricular mass on septal mitral annulus velocity (Ea) and the ratio of transmitral early peak velocity to Ea (E/Ea). *Am J Cardiol*; 95:1020–3.

-
- **Dell'Italia L, Sabri A. (2004):** Activation of the renin-angiotensin system in hypertrophy and heart failure. In: Mann DL, ed. *Heart Failure: Companion to Braunwald's Heart Disease*, Philadelphia: Saunders: 129-143.
 - **Derumeaux G, Michel O, Loufoua J, et al. (2000):** Assessment of non uniformity of transmural myocardial velocities by color-coded tissue Doppler imaging. Characterization of normal, ischemic and stunned myocardium. *Circulation* ;101:1390-1395.
 - **Derumeaux G, Ovisze M, Loufova J, et al. (1998):** Doppler tissue imaging quantitative regional wall motion during myocardial ischemia and reperfusion. *Circulation*; 97:1970-1997.
 - **Devereux RB, Roman MJ, Liu JE, et al. (2003):** An appraisal of echocardiography as an epidemiological tool. The Strong Heart Study. *Ann Epidemiol*; 13:238–244. [PMID: 12684189]
 - **Donal E, Raynier P, Coisne D, et al. (2005):** Tissue Doppler echocardiographic quantification. Comparison to coronary angiography results in acute coronary syndrome patients. *Cardiovascular Ultrasound*; 3:10.
 - **Duzenli MA, Ozdemir K, Aygul N, et al. (2008):** The role of tissue Doppler echocardiography in the evaluation of functional capacity in patients of heart failure. *Turk Kardiyol Dern Ars*; 36(3):143-9.
 - **Edmonds PM, Rogers A, Addington-Hall JM, et al. (2005):** Patient descriptions of breathlessness in heart failure. *Int J Cardiol*; 98:61-66.
 - **Edvardsen T, Aakhus S, Enderson E, et al. (2000):** Effect of first myocardial infarction on left ventricular systolic and diastolic function with the use of mitral annular velocity determined by pulsed wave Doppler tissue imaging. *J Am Soc Echocardiogr*; 13:343-52.
 - **Edvardsen T, Gerber BL, Garot J, et al. (2002):** Lima JA, Smiseth OA. Quantitative assessment of intrinsic regional myocardial deformation by Doppler strain rate echocardiography in humans: validation against three-dimensional tagged magnetic resonance imaging. *Circulation*; 106:50 –56.
-

-
- **Elliott P. (2000):** Cardiomyopathy: diagnosis and management of DCM. *Heart*. 84(1):106-112.
 - **Erbel R, Wallbridge DR, and Zammarano J. (1999):** Tissue Doppler echocardiography. *Heart*, 76:193-196.
 - **Eriksson H. (1995):** Heart failure: a growing public health problem. *J Intern Med*; 237:135-141.
 - **Fang ZY, Leano R, and Marwick TH. (2004):** Relationship between longitudinal and radial contractility in subclinical diabetic heart disease. *Clin Sci (Lond)*; 106:53– 60.
 - **Farrias CA, Radriguez L, Garcia MJ, et al. (1999):** Assessment of diastolic function by tissue Doppler echocardiography: comparison with standard transmitral and pulmonary venous flow. *J Am Soc Echocardiogr*;12:609-617.
 - **Feigenbaum H. (1994):** Hemodynamic information derived from echocardiography. Echocardiographic evaluation of cardiac chambers. In echocardiography, fifth edition. Lea and Febiger . Philadelphia, Bultimore, Hong Kong; p 134 and 204.
 - **Feigenbaum, Harvey, Armstrong, et al. (2005):** Evaluation of Systolic and Diastolic Function of the Left Ventricle. *Feigenbaum's Echocardiography*, 6th Edition 138-180.
 - **Flather MD, Yusuf S, Kober L, et al. (2000):** Long-term ACE-inhibitor therapy in patients with heart failure or left-ventricular dysfunction: A systematic overview of data from individual patients. ACE-Inhibitor Myocardial Infarction Collaborative Group. *Lancet*; 355:1575.
 - **Fleg JL, Lakatta EG. (1988):** Role of muscle loss in the age-associated reduction in $\dot{V}O_2$ max. *J Appl Physiol*; 65:1147–1151.
 - **Fletcher GF, Balady G, Froelicher VF et al. (1995):** Exercise standards: a statement for healthcare professionals from the American Heart Association Writing Group. Special Report. *Circulation*; 91: 580–615.
-

-
- **Floras JS. (2003):** Sympathetic activation in human heart failure: Diverse mechanisms, therapeutic opportunities. *Acta Physiol Scand*; 177:391.
 - **Foster C, Pollock ML, Rod JL, et al. (1983):** Evaluation of functional capacity during exercise radionuclide angiography. *Cardiology*; 70:85–93.
 - **Fox KF, Cowie MR, Wood DA, et al. (2001):** Coronary artery disease as the cause of incident heart failure in the population. *Eur Heart J*; 22:228–236. [PMID: 11161934]
 - **Fox KF, Cowie MR, Wood DA, et al. (2001):** Coronary artery disease as the cause of incident heart failure in the population. *Eur Heart J*; 22:228–236. [PMID: 11161934]
 - **Francesco F, Paolo T, Guiseppina M, et al. (1998):** New modalities of regional and global left ventricular function analysis: state of Art. *Am J Cardiol*; 81:49-57.
 - **Francis GS, and Chu C. (1995):** Post-infarction myocardial remodeling: why does it happen? *Eur Heart J*; 16:31-36.
 - **Freidman MJ, Roeske WR, Sahn DJ, et al. (1982):** Accuracy of M-mode echocardiographic measurements of the left ventricle. *Am J Cardiol*; 49:416-423.
 - **Fujii J, Yazaki Y, Sauada H, et al. (1985):** Non invasive assessment of left and right ventricular filling in myocardial infarction with a two-dimensional Doppler echocardiographic method. *J Am Coll Cardiol*; 5(5):1155-60.
 - **Fukuta H, Sane DC, Brucks S, et al. (2005):** Statin therapy may be associated with lower mortality in patients with diastolic heart failure: A preliminary report. *Circulation*; 112:357.
 - **Gaasch WH, Zile MR. (2004):** Left ventricular diastolic dysfunction and diastolic heart failure. *Annu Rev Med*; 55:373-94.
 - **Garcia M J, Rodriguez L, Ares M, et al. (1996):** Myocardial wall velocity assessment by pulsed wave Doppler tissue imaging: characteristic findings in normal subjects. *Am Heart J*; 132:648-656.
-

-
- **Garcia M, and Thomas J. (1999):** Tissue Doppler to assess diastolic left ventricular function, *Echocardiography*; 165:501-508.
 - **Garcia MJ, Ares MA, Asher C, et al. (1997):** An index of early left ventricular filling that combined with pulsed Doppler peak E velocity may estimate capillary wedge pressure. *J Am Coll Cardiol* 29:448–454
 - **Garcia MJ, Rodriguez L, Ares M, et al. (1996):** Differentiation of constrictive pericarditis from restrictive cardiomyopathy: assessment of left ventricular diastolic velocities in longitudinal axis by Doppler tissue imaging. *J Am Coll Cardiol*; 27:108–14.
 - **Garcia-Fernandez M A, Zamorano J, and Azevedo J. (1998):** Normal pattern with Doppler tissue imaging. In Garcia-Fernandez M A, Zamorano J, and Azevedo J (eds): *Doppler tissue imaging echocardiography*; Mc Grow-Hill Madrid. pp 23-44.
 - **Garot J, Dermeaux GA, Monin GL, et al. (1999):** Quantitative systolic and diastolic transmyocardial velocity gradients assessed by M-mode color Doppler tissue imaging as reliable indicators of regional left ventricular function after acute myocardial infarction. *Eur Heart J*; 20:593-603.
 - **Gattis WA, O'Connor CM, Gallup DS, et al. (2004):** Predischage initiation of carvedilol in patients hospitalized for decompensated heart failure: results of the Initiation Management Predischage: Process for Assessment of Carvedilol Therapy in Heart Failure (IMPACT-HF) trial. *J Am Coll Cardiol*; 43: 1534–1541.
 - **Gheorghiade M, Zannad F, Sopko G, et al. (2005):** Acute heart failure syndromes: current state and framework for future research. *Circulation*; 112:3958-3968.
 - **Giannuzzi P, Temporelli PL, Bosimini E, et al. (1996):** Independent and incremental prognostic value of Doppler-derived mitral deceleration time of early filling in both symptomatic and asymptomatic patients with left ventricular dysfunction. *JAMA*; 28:383–90.
 - **Gibson DG, and Francis DP. (2003):** Clinical assessment of left ventricular diastolic function. *Heart*; 89:231-238.
-

-
- **Givertz M, Colucci WS, and Braunwald E. (2005):** *Clinical aspects of heart failure: High-output failure, pulmonary edema.*
 - **Goldberger Ary L. (2006):** Clinical electrocardiography: a simplified approach. 7th edition.
 - **Gorascan J, Gulati VK, Mandarino WA, et al. (1996):** Color-coded measures of myocardial velocity throughout the cardiac cycle by tissue Doppler imaging to quantify left ventricular function. *Am Heart J*; 131:1203-1213.
 - **Gorscan J III, Kanzaki H, Bazaz R, et al. (2004):** Usefulness of echocardiographic tissue synchronization imaging to predict acute response to cardiac resynchronization therapy. *Am J Cardiol*; 93:1178–81
 - **Gorscan J, Gulati V, Manriano W, et al. (1999):** Color-coded measures of myocardial velocity throughout the cardiac cycle by Doppler tissue imaging to quantify regional left ventricular function. *Am Heart J*; 131:1203-13.
 - **Granger CB, McMurray JJ, Yusuf S, et al. (2003):** Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function intolerant to angiotensin-converting-enzyme inhibitors: the CHARM-Alternative trial. *Lancet*; 362: 772–776.
 - **Grinstead WC, Francis MJ, Marks GF, et al. (1994):** Discontinuation of chronic diuretic therapy in stable congestive heart failure secondary to coronary artery disease or to idiopathic dilated cardiomyopathy. *Am J Cardiol*; 73:881-886.
 - **Gulati V, Katz W, Follansbee W, et al. (1996):** Mitral annular descent velocity by Tissue Doppler echocardiography as an index of global left ventricular function. *Am J Cardiol*; 77:979-84.
 - **Ha JW, Lee HC, Kang ES, et al. (2007):** Abnormal left ventricular longitudinal functional reserve in patients with diabetes mellitus: implication for detecting subclinical myocardial dysfunction using exercise tissue Doppler echocardiography. *Heart*; 93:1571–6.
 - **Ha J-W, Ommen SR, Tajik AJ, et al. (2004):** Differentiation of constrictive pericarditis from restrictive cardiomyopathy using mitral
-

annular velocities by tissue Doppler echocardiography. *Am J Cardiol*; 94:316–9.

- **Harlan WR, Oberman A, Grimm R, et al. (1977):** Chronic congestive heart failure in coronary artery disease. Clinical criteria. *Ann Intern Med*; 86:133.
- **Hasegawa H, Little WC, and Ohno M. (2003):** Diastolic mitral annular velocity during the development of heart failure. *J Am Coll Cardiol* 41:1590–1597
- **Haulica I, Petrescu G, Slatineanu SM, et al. (2004):** New bioactive angiotensins formation pathways and functional involvements. *Rom J Intern Med*; 42:27.
- **Henein MY, Gibson DG. (1999):** Long axis function in disease. *Heart*; 81: 229 –31.
- **Hillege H, Girbes AR, de Kann PJ, et al. (2000):** Renal function, neurohormonal activation, and survival in patients with chronic heart failure. *Circulation*; 102:205.
- **Hiller HS, Moller JE, Pellikka PA, et al. (2004):** Noninvasive estimation of left ventricular filling pressure by E/E9 is a powerful predictor of survival after acute myocardial infarction. *J Am Coll Cardiol*; 43:360–7.
- **Hjalmarson A, Goldstein S, Fagerberg B, et al, for the MERIT-HF Study Group. (2000):** Effects of controlled-release metoprolol on total mortality, hospitalizations, and well-being in patients with heart failure: the Metoprolol CR/XL Randomized Intervention Trial in congestive heart failure (MERIT-HF). *JAMA*; 283: 1295–1302.
- **Ho C, and Solomon S.D. (2006):** A Clinician’s guide to tissue Doppler imaging. *Circulation*; 113:396-398.
- **Ho KKL, Pinsky JL, Kannel WB, et al. (1993):** The epidemiology of heart failure: the Framingham study. *J Am Coll Cardiol*; 22(suppl A):6A-13A.
- **Hope MD, de la Pena E, Yang PC, et al. (2003):** A visual approach for the accurate determination of echocardiographic left ventricular ejection fraction by medical students. *J Am Soc Echocardiogr*; 16:824–831.

-
- **Hoskins PR, and Mc Dicken WN. (1997):** Color ultrasound imaging of blood flow and tissue motion. *British J Radiol.* 70:878-890.
 - **Huang CH, Tsai MS, Hsieh CC, et al. (2006):** Diagnostic accuracy of tissue Doppler echocardiography for patients with acute heart failure. *Heart;* 92:1790–4.
 - **Hundley W, Kizilbash A, Afridi I, et al. (1998):** Administration of an intravenous perfluorocarbon contrast agent improves echocardiographic determination of left ventricular volumes and ejection fraction: Comparison with cine MRI. *J Am Coll Cardiol;* 32:1426–1432.
 - **Hunt SA, Abraham WT, Chin MH, et al. (2005):** ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult: summary article. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to revise the 2001 Guidelines for the Evaluation and Management of Heart Failure). *J Am Coll Cardiol* ; 46:1116–1143.
 - **Hunt SA, Abraham WT, Chin MH, et al. (2005):** ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure): developed in collaboration with the American College of Chest Physicians and the International Society for Heart and Lung Transplantation: endorsed by the Heart Rhythm Society. *Circulation;* 112(12):e154–235.
 - **Hunt SA, Baker DW, Chin MH, et al. (2001):** Guidelines for the evaluation and management of chronic heart failure in the adult: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to revise the 1995 Guidelines for the Evaluation and Management of Heart Failure). *J Am Coll Cardiol;* 38:2101-13.
 - **Hunt SA; American College of Cardiology; American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure).**
-

- (2005): ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*; 46:e1-82.
- **Isaaz K, Thompson A, Thevenot G, et al. (1989):** Doppler echocardiographic measurement of low velocity motion of left ventricular posterior wall. *Am J Cardiol*; 64:66-75.
 - **Isaaz K. (2002):** Tissue Doppler imaging for the assessment of systolic and diastolic functions. *Current opinion in cardiology*; 17(5) j:431-42.
 - **Israili ZH, Hall WD. (1992):** Cough and angioneurotic edema associated with angiotensin-converting enzyme inhibitor therapy: a review of the literature and pathophysiology. *Ann Intern Med*; 117:234-42.
 - **J Am Coll Cardiol (2007);** 49:1903–14.
 - **J De Sutter, N R Van de Veire, J J Bax, et al. (2008):** Technological advances in Tissue Doppler Imaging echocardiography. *Heart*; 94:1065-1074.
 - **Javaheri S. (1999):** A mechanism of central sleep apnea in patients with heart failure. *N Engl J Med*; 341:949-954.
 - **Jerome L Fleg, and Ileana L Piña. (2000):** Assessment of Functional Capacity in Clinical and Research Applications: *Circulation*; 102:1591.
 - **John Gorscan, Warren J Manning, and Susan B Yeon. (2008):** Tissue Doppler echocardiography. *Up to date*.
 - **Johnson W, Omland T, Hall C, et al. (2002):** Neurohormonal activation rapidly decreases after intravenous therapy with diuretics and vasodilators for class IV heart failure. *J Am Coll Cardiol*; 39:1623-1629.
 - **Jones CJ, Raposo L, and Gibson DG. (1990):** Functional importance of the long axis dynamics of the human left ventricle. *Br Heart J*; 63:215–220.
 - **Joshi N. (1999):** The third heart sound. *South Med J*; 92:756.

-
- **Juurlink DN, Mamdani M, Kopp A, et al. (2003):** Drug-drug interactions among elderly patients hospitalized for drug toxicity. *JAMA*; 289: 1652–1658.
 - **Kannel WB, Belanger AJ. (1991):** Epidemiology of heart failure. *Am Heart J*; 121:951–957. [PMID: 2000773]
 - **Kannel WB, Ho K, Thom T. (1994):** Changing epidemiological features of cardiac failure. *Br Heart J*; 72:S3–S9.
 - **Kasper DL, Braunwald E, Fauci AS, et al (2007):** Harrison's Principles of Internal Medicine. 17th ed. New York, McGraw-Hill, Anonymous. Assessment of diastolic function. In: Oh JK, Seward JB, Tajik AJ, eds. The echo manual. Philadelphia: Lipincott Williams & Wilkins, 1999:45–57.
 - **Katz WE, Gulati VK, Mahler CM, et al. (1997):** Quantitative evaluation of the segmental left ventricular response to dobutamine stress by tissue Doppler echocardiography. *Am J Cardiol*; 79:1036–42.
 - **Kitzman DW, Little WC, Brubaker PH, et al. (2002):** Pathophysiological characterization of isolated diastolic heart failure in comparison to systolic heart failure. *JAMA*; 288:2144.
 - **Kornbluth M, Liang DH, Brown P, et al. (2000):** Contrast echocardiography is superior to tissue harmonics for assessment of left ventricular function in mechanically ventilated patients. *Am Heart J*; 140:291–296.
 - **Kostis J.B, Mavrogeorgis E, Slater A, et al. (1972):** Use of a range-gated, pulsed ultrasonic Doppler technique for continuous measurement of velocity of the posterior wall. *Chest*; 62; 597-604.
 - **Lang RM, Bierig M, Devereux RB, et al. (2005):** Recommendations for chamber quantification: A report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *J Am Soc Echocardiogr*; 18:1440–1463.
-

-
- **Lee KI, Pryor DB, Pieper KS, et al. (1990):** Prognostic value of radionuclide angiography in medically treated patients with coronary artery disease. A comparison with clinical and catheterization variables. *Circulation*; 82:1706.
 - **Levy D, Kenchaiah, S, Larson MG, et al. (2002):** Long-term trends in the incidence of and survival with heart failure. *N Engl J Med* ;347:1397-1402.
 - **Lipkin DP, Canepa-Anson R, Stephens MR, et al. (1986):** Factors determining symptoms in heart failure: comparison of fast and slow exercise tests. *Br Heart J*; 55:439-445.
 - **Lisauskas J, Singh J, and Courtois M. (2001):** The relation of the peak Doppler E-wave to peak mitral annulus velocity ratio to diastolic function. *Ultrasound Med Biol* 27:499–507
 - **Little WC. (2001):** Assessment of normal and abnormal cardiac function. In Braunwald E, Zipes, Libby: Heart diseases 6th edition.
 - **Lloyd-Jones DM, Larson MG, Leip EP, et al. (2002):** Lifetime risk for developing congestive heart failure: the Framingham Heart Study. *Circulation*; 106:3068-3072.
 - **Lundback S. (1986):** Cardiac pumping and function of the ventricular septum. *Acta Physiol Scand Suppl*; 550:1–101.
 - **Malm S, Frigstad S, Sagberg E, et al. (2004):** Accurate and reproducible measurement of left ventricular volume and ejection fraction by contrast echocardiography: A comparison with MRI. *J Am Coll Cardiol*; 44: 1030–1035.
 - **Mandinov L, Ebreli FR, Seiler C, et al. (2000):** Diastolic heart failure. *Cardiovasc Res*; 45:813–25.
 - **Mann DL, Bristow MR. (2005):** Mechanisms and models in heart failure: The biomechanical model and beyond. *Circulation*; 111:2837.
 - **Marwick TH. (2006):** Measurement of strain and strain rate by echocardiography. Ready for prime time? *J Am Coll Cardiol*; 47:1313–27.
 - **McDicken WN, Sutherland GR, Moran CM, et al. (1992):** Color Doppler velocity imaging of myocardium. *Ultrasound Med Biol.*; 18:651-654.
-

-
- **McCullough PA, Nowak RM, McCord J, et al. (2002):** B-type natriuretic peptide and clinical judgment in emergency diagnosis of heart failure: analysis from Breathing Not Properly (BNP) Multinational Study. *Circulation*; 106:416–422. [PMID: 12135939]
 - **Mclean A, Needham P, Stewart D, et al. (1996):** Estimation of cardiac output in critically ill subject by noninvasive echocardiographic techniques. (Abstract). *Anaesth. Intensive care*, (in press).
 - **McMurray JJ, Ostergren J, Swedberg K, et al. (2003):** Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function taking angiotensin-converting-enzyme inhibitors: the CHARM-Added trial. *Lancet*; 362: 767–771.
 - **Mehmet Akif, Kurtulus Ozdemir, Nazif Aygul, et al. (2008):** Relationship between Systolic Myocardial Velocity Obtained by Tissue Doppler Imaging and Left Ventricular Ejection Fraction: Systolic Myocardial Velocity Predicts the Degree of Left Ventricular Dysfunction in Heart Failure. *ECHOCARDIOGRAPHY: A Jnl. of CV Ultrasound & Allied Tech*; 856-863.
 - **Mehra M, and Greenberg B. (2004):** Cardiac resynchronisation therapy: Caveat medicus! *J Am Coll Cardiol*; 43:1145–8.
 - **Meluzin J, Spirnarova L, Bakala J, et al.** Pulsed Doppler tissue imaging of the velocity of tricuspid annular motion: a new, rapid, and non-invasive method of evaluating right ventricular systolic function. *Eur Heart J*; 22:340–8.
 - **Mendez GF, and Cowie MR. (2001):** The epidemiological features of heart failure in developing countries: a review of the literature. *Int J Cardiol.*; 80:213–219. [PMID: 11578717]
 - **Meyers J, Salleh A, Buchanan N, et al. (1992):** Ventilatory mechanisms of exercise intolerance in chronic heart failure. *Am Heart J*; 124:7-10.
 - **Miller LW. (1998):** Listing criteria for cardiac transplantation: results of an American Society of Transplant Physicians-National Institutes of Health conference. *Transplantation*; 66: 947–951.
-

-
- **Miyamura M, Honda Y. (1972):** Oxygen intake and cardiac output during peak treadmill and bicycle exercise. *J Appl Physiol*; 32:185–188.
 - **Miyatake K, Yamagashi M, Tanaka N, et al. (2000):** Analysis of ventricular wall motion using color-coded tissue Doppler imaging system. *Eur Heart J*; 21:1337-57.
 - **Miyatake K, Yamagishi M, Tanaka N et al (1995):** New method of evaluating left ventricular wall motion by color coded tissue Doppler imaging in vitro and in vivo studies. *J Am Coll Cardiol*. 25:717-724.
 - **Mock MB, Ringqvist I, Fisher LD, et al. (1982):** Survival of medically treated patients in the Coronary Artery Surgery Study (CASS) registry. *Circulation*; 66:562–568.
 - **Mokdad A, Herdula M, and Deitz W. (1999):** The spread of obesity in the united states. *JAMA*; 282:1519-1522.
 - **Mottram PM, and Marwick TH. (2005):** Assessment of diastolic function: What the general cardiologist needs to know. *Heart*; 91(5):681-695.
 - **Mulvagh SL, DeMaria AN, Feinstein SB, et al. (2000):** Contrast echocardiography: Current and future applications. *J Am Soc Echocardiogr*; 13:331–342.
 - **Myers J, Buchanan N, Walsh D, et al. (1991):** Comparison of the ramp versus standard exercise protocols. *J Am Coll Cardiol*; 17:1334–1342.
 - **Nagueh SF, Middleton K, Kopelen H, et al. (1997):** Doppler tissue imaging: a noninvasive technique for evaluation of left ventricular relaxation and estimation of filling pressure. *J Am Coll Cardiol* 30:1527–1533
 - **Nagueh SF, Middleton KJ, Kopelen HA, et al. (1997):** Doppler tissue imaging: a non-invasive technique for evaluation of left ventricular relaxation and estimation of filling pressures. *J Am Coll Cardiol*; 30:1527–33.
 - **Nagueh SF, Mitaki I, Kopelen HA, et al. (1998):** Doppler estimation of left ventricular filling pressure in sinus tachycardia. A new application of tissue Doppler imaging. *Circulation*; 98:1644–50.
-

-
- **Nagueh SF, Sun HB, Kopelen HA, et al. (2001):** Haemodynamic determinants of the mitral annulus diastolic velocities by tissue Doppler. *JAMA*; 37:278–85.
 - **Nakayama K, Miyatake MD, Uematsu M, et al. (1998):** Application of tissue Doppler technique in evaluating early ventricular contraction associated with accessory A-V pathways in Wolff- Parkinson White syndrome. *Am-Heart-J*; 35(1):99-106.
 - **Neil Smart, Brian Haluska, Philip M, et al. (2005):** Determinants of functional capacity in patients with chronic heart failure: Role of filling pressure and systolic and diastolic function. *AHJ*; 152-158.
 - **Neskovic AN, and Otasevic P. (2005):** Stress echocardiography in idiopathic dilated cardiomyopathy: instructions for use. *Cardiovasc Ultrasound*; 3:3.
 - **Nishimura RA, Housmans PR, Hatle, et al. (1989):** Assessment of diastolic function of the heart. Background and current applications of Doppler echocardiography part II: Clinical studies. *Mayo Clin. Proc*; 64:181-204.
 - **Nishimura RA, and Tajik AJ. (1994):** Quantitative hemodynamics by Doppler echocardiography: A noninvasive alternative to cardiac catheterization. *Prog Cardiovasc Dis*; 4:309–342.
 - **Nishimura R, and Tajik A.** Evaluation of diastolic filling of left ventricle in health and disease: Doppler echocardiography is the clinician's Rosett stone. *J Am Coll Cardiol* 1997; 30:8 –18.
 - **Nixon JV, Murray RG, Leonard PD, et al. (1982):** Effect of large variations in preload on left ventricular performance characteristics in normal subjects. *Circulation*; 4:698–703.
 - **Nonogi H, Hess OM, Ritter M, et al. (1988):** Diastolic properties of the normal left ventricle during supine exercise. *Br Heart J* ; 60:30.
 - **Oh JK, Appleton CP, Hatle LK, et al. (1997):** The noninvasive assessment of left ventricular diastolic function with two-dimensional and Doppler echocardiography. *J Am Echocardiogr* 10:246–270
-

-
- **Oh JK, Hatle L, Tajik AJ, et al. (2006):** Diastolic heart failure can be diagnosed by comprehensive two-dimensional and Doppler echocardiography. *J Am Coll Cardiol*; 47:500–506.
 - **Ohno M, Cheng CP, Little WC, et al. (1994):** Mechanism of altered patterns of left ventricular filling during the development of congestive heart failure. *Circulation*; 89:2241.
 - **Ohte N, Narita H, and Kimura O. (1999):** Evaluation of cardiac function using Tissue Doppler Imaging, *Medical review*; 30-35.
 - **Oki T, Tabata T, Yamada H, et al. (1997):** Clinical application of pulsed Doppler tissue imaging for assessing abnormal left ventricular relaxation. *Am J Cardiol*; 79:921-8.
 - **Okura H, Takada Y, Kubo T, et al. (2006):** Tissue Doppler-derived index of left ventricular filling pressure, E/Ea, predicts survival of patients with non-valvular atrial fibrillation. *Heart*; 92:1248–52.
 - **Ommen SR, Nishimura RA, Appleton CP, et al. (2000):** Clinical utility of Doppler echocardiography and tissue Doppler imaging in the estimation of left ventricular filling pressures: a comparative simultaneous Doppler-catheterization study. *Circulation* 102:1788–1794
 - **Ommen SR, Nishimura RA, Appleton CP, et al. (2000):** Clinical utility of Doppler echocardiography and tissue Doppler imaging in the estimation of left ventricular filling pressures: a comparative simultaneous Doppler-catheterization study. *Circulation*; 102:1788–94.
 - **Opasich C, Pinna GD, Mazza A et al. (1998):** Reproducibility of the six-minute walking test in chronic congestive heart failure patients: practical implications. *Am J Cardiol*; 81: 1487–500.
 - **Packer M, Coats AJ, Fowler MB, et al. (2001):** Effect of carvedilol on survival in severe chronic heart failure. *N Engl J Med*; 344: 1651–1658.
 - **Packer M, Lee WH, and Kessler PD. (1986):** Preservation of glomerular filtration rate in human heart failure by activation of the renin-angiotensin system. *Circulation*; 74:766-74.
-

-
- **Packer M, Lee WH, and Kessler PD. (1987):** Identification of hyponatremia as a risk factor for the development of functional renal insufficiency during converting enzyme inhibition in severe chronic heart failure. *J Am Coll Cardiol*;10:837-44.
 - **Packer M, Lee WH, Medina N, et al. (1987):** Functional renal insufficiency during long-term therapy with captopril and enalapril in severe chronic heart failure. *Ann Intern Med*; 106:346-54.
 - **Packer M, Lee WH, Medina N, et al. (1987):** Influence of diabetes mellitus on changes in left ventricular performance and renal function produced by converting enzyme inhibition in patients with severe chronic heart failure. *Am J Med*; 82:1119-26.
 - **Packer M, Medina N, and Yushak M. (1984):** Relation between serum sodium concentration and the hemodynamic and clinical responses to converting enzyme inhibition with captopril in severe heart failure. *J Am Coll Cardiol*; 3:1035-43.
 - **Pai RG, Bodenheimer MM, Pai SM, et al. (1991):** Usefulness of systolic excursion of the mitral annulus as an index of left ventricular systolic function. *Am J Cardiol*; 67:222– 4.
 - **Pai RG, and Buech GC. (1996):** New Doppler measures of left ventricular function. *Clin. Cardiol*; 19:277:88.
 - **Page E, Cohen-Solal A, Jondeau G et al. (1994):** Comparison of treadmill and bicycle exercise in patients with chronic heart failure. *Chest*; 106: 1002– 6.
 - **Park TH, Nagueh SF, Khourey DS, et al. (2006):** Impact of myocardial structure and function postinfarction on diastolic strain measurements: implications for assessment of myocardial viability. *Am J Physiol Heart Circ Physiol*; 290:H724–31.
 - **Paulus WJ, Tschoepke C, Sanderson JE, et al. (2007):** How to diagnose heart failure: a consensus statement on the diagnosis of heart failure with normal left ventricular ejection fraction by the Heart Failure and Echocardiography Associations of the European Society of Cardiology. *Eur Heart J*; 28:2539–50.
-

- **Penicka M, Bartunek J, De Bruyne B, et al. (2004):** Improvement of left ventricular function after cardiac resynchronization therapy is predicted by tissue Doppler imaging echocardiography. *Circulation*; 109:978–83.
- **Pfeffer MA, and Braunwald E. (1990):** Ventricular remodeling after myocardial infarction. *Circulation*; 81:1161-1172.
- **Piña IL, Karalis DG. (1990):** Comparison of four exercise protocols using anaerobic threshold measurement of functional capacity in congestive heart failure. *Am J Cardiol*; 65:1269–1271.
- **Pitt B, Poole-Wilson PA, Segal R, et al. (2000):** Effect of losartan compared with captopril on mortality in patients with symptomatic heart failure: randomised trial: the Losartan Heart Failure Survival Study ELITE II. *Lancet*; 355: 1582–1587.
- **Pitt B, Williams G, Remme W, et al. The EPHESUS trial. (2001):** eplerenone in patients with heart failure due to systolic dysfunction complicating acute myocardial infarction: Eplerenone Post-AMI Heart Failure Efficacy and Survival Study. *Cardiovasc Drugs Ther*; 15: 79–87.
- **Pitt B, Zannad F, Remme WJ, et al for the Randomized Aldactone Evaluation Study Investigators. (1999):** The effect of spironolactone on morbidity and mortality in patients with severe heart failure. *N Engl J Med*; 341: 709–717.
- **Pitt B, Remme W, and Zannad F. (2003):** Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med*; 348:1309.
- **Pitt B, Zannad F, Remme WJ, et al. (1999):** The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized Aldactone Evaluation Study Investigators. *N Engl J Med*; 341:709.
- **Quinones MA, Gaasch WH, Alexander JK, et al. (1974):** Echocardiographic assessment of left ventricular function: With special reference to normalized velocities. *Circulation*; 50(1):42-51.

-
- **R C Jones, G S Francis and M S Lauer. (2004):** Predictors of mortality in patients with heart failure and preserved systolic function in digitalis investigation group trial, *J.Am.Coll.Cardiol.*44, P 1025-1029.
 - **Rassi A Jr, Rassi A, and Little WC. (2000):** Chagas' heart disease. *Clin Cardiol.* ;23:883–889. [PMID: 11129673]
 - **Redfield MM, Jacobsen SJ, Borlaug BA, et al. (2005):** Age- and gender-related ventricular-vascular stiffening: A community-based study. *Circulation*; 112:2254.
 - **Redfield MM, Jacobsen SJ, Burnett Jr JC, et al. (2003):** Burden of systolic and diastolic ventricular dysfunction in the community: appreciating the scope of the heart failure epidemic. *JAMA*; 289:194.
 - **Remme W J and Swedberg K. (2001):** Task force for the diagnosis and treatment of chronic heart failure, European Society of cardiology. Guidelines for the diagnosis and treatment of chronic heart failure. *Eur. Heart J.*, 22:1527-1560.
 - **Riegger GAJ. (1991):** Effects of quinapril on exercise tolerance in patients with mild to moderate heart failure. *Eur Heart J* 1991; 12: 705–11.
 - **Rihal CS, Nishimura RA, Hatle LK, et al. (1994):** Systolic and diastolic dysfunction in patients clinical diagnosis of dilated cardiomyopathy; relation to symptoms and prognosis. *Circulation* 90:2772–2779
 - **Rihal CS, Nishimura RA, Hatle LK. et al. (1994):** Systolic and diastolic dysfunction in patients with clinical diagnosis of dilated cardiomyopathy. Relation to symptoms and prognosis. *Circulation* 90:2772–9.
 - **Ross J Jr. (1976):** Afterload mismatch and preload reserve: A conceptual framework for the analysis of ventricular function. *Prog Cardiovasc Dis*; 18:255–264.
 - **Ross J Jr, Braunwald E, and Sonnenblick EH. (1979):** Mechanisms of Contraction of the Normal and Failing Heart. Boston, Little Brown.
 - **Rockville, Rodeheffer RJ, Lerman A, et al (1994):** Increased plasma concentrations of endothelin in congestive heart failure in humans. *Mayo Clin Proc*;67:719-724.
-

-
- **Rowell LB. (1988):** Muscle blood flow in humans: how high can it go? *Med Sci Sports Exerc*; 20:S97–S103.
 - **Sait Terzi, Nurten Sayar , Tuba Bilsel, et al. (2007):** Tissue Doppler imaging adds incremental value in predicting exercise capacity in patients with congestive heart failure. *Heart Vessels*; 22:237–244
 - **Sanderson JE, Wang M, and Yu CM. (2004):** Tissue Doppler imaging for predicting outcome in patients with cardiovascular disease. *Curr Opin Cardiol*; 19:458–63.
 - **Sanderson JE. (2007):** Heart failure with a normal ejection fraction. *Heart*; 93:155– 8.
 - **Scherrer-Crosbie M, Ullrich R, Bloch KD, et al. (2001):** Endothelial nitric oxide synthase limits left ventricular remodeling after myocardial infarction in mice. *Circulation*; 104:1286.
 - **Schiller N.B., Shan P.M., Crawford M., et al. (1989):** Recommendations for quantitations of the left ventricle by two-dimensional echocardiography. *J Am Soc Echocardiogr*; 2:358-367.
 - **Schrier RW, and Abraham WT. (1999):** Hormones and hemodynamics in heart failure. *N Engl J Med*; 341:577.
 - **Sengupta P, Mohan J, and Pandean N. (2002):** Tissue Doppler echocardiography; principles and applications. *Indian Heart J*; 54:368-378.
 - **Shalaby S, Reda A, Khalil T, et al. (1998):** Age related changes in myocardial contraction and relaxation velocities. Doppler tissue imaging study. *Menoufya Med J*; Jan (10)1:257-73.
 - **Sharir T, Germano G, Kavanagh PB, et al. (1999):** Incremental prognostic value of post-stress left ventricular ejection fraction and volume by gated myocardial perfusion single photon emission computed tomography. *Circulation*; 100:1035–1042.
 - **Silverberg DS, Wexler D, Sheps D, et al. (2001):** The effect of correction of mild anemia in severe, resistant congestive heart failure using subcutaneous erythropoietin and intravenous iron: A randomized controlled study. *J Am Coll Cardiol*; 37:1775.
-

-
- **Simpson J A. (1997):** Echocardiographic assessment of long axis functions: a simple solution to a complex problem? *Heart*; 78:211-212.
 - **Smith GL, Masoudi FA, Vaccarino V, et al. (2003):** Outcomes in heart failure patients with preserved ejection fraction: Mortality, readmission, and functional decline. *J Am Coll Cardiol*; 41:1510.
 - **Sohn DW, Chai IH, Lee DJ, et al. (1997):** Assessment of mitral annulus velocity by Doppler tissue imaging in the evaluation of left ventricular diastolic function. *J Am Coll Cardiol*; 30:474–80.
 - **Solal AC, and Gourgon R. (1991):** Assessment of exercise tolerance in chronic congestive heart failure. *Am J Cardiol*; 67:36C-40C.
 - **Solin P, Bergin P, Richardson M, et al. (1999):** Influence of pulmonary capillary wedge pressure on central apnea in heart failure. *Circulation*; 99: 157-1579.
 - **Spirito BB, Brecker SD, Ziao HB, et al. (1992):** Left ventricular filling characteristic in pulmonary hypertension: A new mode of ventricular interaction. *Br Heart J*; 68:16.
 - **Spain MC, Smith MD, Grayburn PA, et al. (1989):** Qualitative assessment of mitral regurgitation by Doppler color flow imaging: angiographic and haemodynamic correlations. *J Am coll Cardiol*; 13:585-90.
 - **Steele B. (1996):** Timed walking tests of exercise capacity in chronic cardiopulmonary illness. *J Cardiopulm Rehabil*; 16:25–33.
subcutaneous erythropoietin and intravenous iron: A randomized controlled study. *J Am Coll Cardiol*; 37:1775.
 - **Stevenson LW, Couper G, Natterson B et al. (1995):** Target heart failure populations for newer therapies. *Circulation*; 92 (Suppl II): II-174–II-181.
 - **Stevenson LW, Massie BM, and Francis G. (1998):** Optimizing therapy for complex or refractory heart failure: a management algorithm. *Am Heart J* ; 135: S293–S309.
 - **Sutherland G.R, Bijnen B, and Mc Dicken W.N. (1999):** Tissue Doppler echocardiography. Historical prospective and technological consideration. *Echocardiography*, 16:445-453.
-

-
- **Sutherland G.R, Stewart M.J. Groundstroem K.W, et al. (1994):** Color Doppler myocardial imaging: A new technique for the assessment of myocardial imaging. *J-Am-Soc-Echocardiogr*; 7:441-458.
 - **Sutherland GR, Bijmens B, and McDicken WN. (1999):** Tissue Doppler echocardiography. Historical perspective and technological considerations. *Echocardiography*; 16:445–453.
 - **Sutherland GR, Di SG, Claus P, et al. (2004):** Strain and strain rate imaging: a new clinical approach to quantifying regional myocardial function. *J Am Soc Echocardiogr*; 17:788–802.
 - **Sutherland GR, Hatle L. (2004):** Normal data. In: Sutherland GR, Hatle L, Rademakers FE, et al, eds. *Myocardial imaging*. Leuven: Leuven University Press, 53–108.
 - **Swedberg K. (1994):** Exercise testing in heart failure. *Drugs* ; 47 (Suppl 4): 14–24.
 - **Swedberg K, Cleland J, Dargie H, et al. (2005):** Guidelines for the diagnosis and treatment of chronic heart failure: executive summary (update 2005): the Task Force for the Diagnosis and Treatment of Chronic Heart Failure of the European Society of Cardiology. *Eur Heart J*; 26:1115–1140. [PMID: 15901669]
 - **Tang WHW, Girod JP, Lee MJ, et al. (2003):** Plasma $\hat{\text{I}}^2$ -type natriuretic peptide levels in ambulatory patients with established chronic symptomatic systolic heart failure. *Circulation*; 108:2964-2966.
 - **Task Force for the Diagnosis and Treatment of Chronic Heart Failure of the European Society of Cardiology. (2005):** Guidelines for the diagnosis and treatment of chronic heart failure: Executive summary (update 2005). *Eur Heart J*; 26:1115.
 - **Taylor AL, Ziesche S, Yancy C, et al. (2004):** Combination of isosorbide dinitrate and hydralazine in blacks with heart failure. *N Engl J Med*; 351:2049–2057. [PMID: 15533851]
-

-
- **Teichholz LE, Kreulen T, Herman MV, et al. (1976):** Problems in echocardiographic volume determinations: Echocardiographic-angiographic correlations in the presence of absence of asynergy. *Am J Cardiol*; 37: 7–11.
 - **Tretjak M, and Kozely M. (2004):** Assessment of left ventricular function using tissue Doppler imaging. *Zdrav Vestn*; 73:663-6.
 - **The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). (2002):** Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium channel blocker vs diuretic. *JAMA*; 288:2981-2997.
 - **The Cardiac Insufficiency Bisoprolol Study II (CIBIS-II). (1999):** a randomised trial. *Lancet*; 353: 9–13.
 - **The Digitalis Investigation Group. (1997):** The effect of digoxin on mortality and morbidity in patients with heart failure. *N Engl J Med*; 336:525–533.
 - **The SOLVD Investigators. (1991):** Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. *N Engl J Med*; 325: 293–302.
 - **Urheim S, Edvardsen T, Torp H, et al. (2000) :** OA. Myocardial strain by Doppler echocardiography: validation of a new method to quantify regional myocardial function. *Circulation*; 102:1158–1164.
 - **Van de Veire N, Bleeker GB, De Sutter J, et al. (2007):** Tissue synchronization imaging accurately measures left ventricular dyssynchrony and predicts response to cardiac resynchronization therapy. *Heart* ; 93:1034–9.
 - **Van de Veire N, De Backer J, Ascoop A-K, et al. (2006):** Echocardiographically estimated left ventricular end-diastolic and right ventricular systolic pressure in normotensive healthy individuals. *Int J Cardiovasc Imaging*; 22: 633–41.
 - **Van de Veire N, Yu C-M, Ajmone-Marsan N, et al. (2008):** Tri-plane tissue Doppler imaging: a novel 3-dimensional imaging modality that
-

- predicts reverse left ventricular remodeling after cardiac resynchronisation therapy. *Heart*; 94:e9.
- **Vinereanu D, Lim PO, Frenneaux MP, et al (2005):** Reduced myocardial velocities of left ventricular longaxis contraction identify both systolic and diastolic heart failure-a comparison with brain natriuretic peptide. *Eur J Heart Fail*;7:512–519.
 - **Volpi A. (1993):** Determinants of 6 months mortality in survivors of myocardial infarction after thrombolysis results of Gruppo Italiano Per Lo studio della nellinfarto miocardia (GISSI-2) database. *Circulation*; 88:419 – 429.
 - **Von Bibra H, Tchnitz A, Klein A, et al. (2000):** Regional diastolic function by pulsed Doppler myocardial mapping for the detection of left ventricular ischemia during pharmacologic stress testing. *J Am Coll Cardiol*; 36:444–52.
 - **Waggoner A, and Michelle Dierig S. (2001):** Tissue Doppler imaging: A useful echocardiographic method for cardiac sonographer to assess systolic and diastolic ventricular function. *J Am Soc Echocardiogr*; 14:1143-1152.
 - **Wang M, Yip GM, Wang AY, et al. (2003):** Peak early diastolic mitral annulus velocity by tissue Doppler imaging adds independent and incremental prognostic value. *JAMA*; 41:820–26.
 - **Wasserman K, Hansen JE, Sue DY, et al. (1994):** Principles of exercise testing and interpretation, 2nd edn. Philadelphia: Lea & Febiger.
 - **Weber KT, Janicki JS. (1985):** Cardiopulmonary exercise testing for evaluation of chronic heart failure. *Am J Cardiol*; 55:22A–31A.
 - **Weber KT, Kinasewitz GT, Janicki JS, et al. (1982):** Oxygen utilization and ventilation during exercise in patients with chronic cardiac failure. *Circulation*; 65:1213-23.
 - **Weisman IM, Zeballos RJ. (1994):** An integrated approach to the interpretation of cardiopulmonary exercise testing. *Clinics in Chest Medicine*; 15: 423–45.

-
- **Weyman AE. (1994):** Appendix A: Normal cross-sectional echocardiographic measurements. In. principles and practice of echocardiography. Philadelphia. Lea& Febiger.
 - **White HD, Norris RM, Brown MA, et al. (1987):** Left ventricular end-systolic volume as the major determinant of survival after recovery from myocardial infarction. *Circulation*; 76:44-51.
 - **Wilson JR, Hanamanthu S, Chomsky DB, et al. (1999):** Relationship between exertional symptoms and functional capacity in patients with heart failure. *J Am Coll Cardiol*; 33:1943-7.
 - **Wilson JR, Martin JL, Ferraro N, et al. (1983):** Effect of hydralazine on perfusion and metabolism in the leg during upright bicycle exercise in patients with heart failure. *Circulation*; 68:425-432.
 - **Witte KK, Nikitin NP, De Silva R, et al. (2004):** Exercise capacity and cardiac function assessed by tissue Doppler imaging in chronic heart failure. *Heart* 90:1144–1150
 - **Wong M, Staszewsky L, Latini R, et al. (2002):** Valsartan benefits left ventricular structure and function in heart failure: Val-HeFT echocardiographic study. *J Am Coll Cardiol*; 40: 970–975.
 - **Xie GY, Berk MR, Smith MD, et al. (1994).** Prognostic value of Doppler transmitral flow patterns in patients with congestive heart failure. *JAMA*; 274:132–9.
 - **Yamamoto T, Oki T, Yamada H, et al. (2003):** Prognostic value of the atrial systolic mitral annular motion velocity in patients with left ventricular systolic dysfunction. *J Am Soc Echocardiogr*; 16:333–9.
 - **Yaron Shapira, a Tuvia Bengal, a Mali Mansur, et al. (2006):** Tissue Doppler Imaging in Patients with Advanced Heart Failure: Relation to Functional Class and Prognosis. *The Journal of Heart and Lung Transplantation*; 214-218.
-

- **Yoshida T, Mori M, Nimura Y, et al. (1961):** Analysis of heart motion with ultrasonic Doppler method and its clinical application. *Am heart J*; 61:61-75.
- **Yu CM, Chau E, Sanderson JE, et al. (2002):** Tissue Doppler echocardiographic evidence of reverse remodeling and improved synchronicity by simultaneously delaying regional contraction after biventricular pacing therapy in heart failure. *Circulation*; 105:438–45.
- **Yvorchuk KJ, Davies RA, and Chan KL. (1994):** Measurement of left ventricular ejection fraction by acoustic quantification and comparison with radionuclide angiography. *Am J Cardiol*; 74:1052–1056.
- **Zamorano J, Rodriguez L, Ares M, et al. (1997):** Assessment of cardiac physiology by tissue Doppler echocardiography. *Eur Heart J*; 18:330-9.
- **Zaphiriou A, Robb S, Murray-Thomas T, et al. (2005):** The diagnostic accuracy of plasma BNP and NT-proBNP in patients referred from primary care with suspected heart failure: results of the UK natriuretic peptide study. *Eur J Heart Fail*; 7:537–41. [PMID: 15921792]

This document was created with Win2PDF available at <http://www.win2pdf.com>.
The unregistered version of Win2PDF is for evaluation or non-commercial use only.
This page will not be added after purchasing Win2PDF.