

REFERENCES

- Abd Ellah, R.G. (1994): Heat, salt and water budget of Lake Nasser water. M.Sc. Assiut Univ., 110 pp.
- Abd Ellah, R.G. (2004): Temperature distributions in Lake Nasser water, Egypt. Asw. Sc. Tec. Bull., 22 : 57- 66.
- Abdallah, M.S. (1972): Taxonomical studies in the flora of Egypt. III. Comparative studies in the natural flora of Egyptian Nubia, Egypt. Egypt. J. Bot., 15: 265-281.
- Abdel-Monem, AM.A. (1995): Spatial distribution of phytoplankton and primary productivity in Lake Nasser, Ph.D Thesis, University Collage for Girls, Ain Sham Univ., 161pp.
- Abdel-Rahman, A. and Goma, R.H. (1992): Monthly variation of air and water temperature, pH, transparency and water color in 1984. Work. Rep. Fish. Mang.. Cent., High Dam Lake Develop. Auth., Aswan, 1: 191-212.
- Abdel-Satar, A.M. (1994): Studies on Some environmental factors affecting the distribution of some chemical elements and productivity of River Nile at El-Kanater El-Khyria”, M.Sc. Thesis, Faculty of Science, Cairo Univ., Egypt.
- Abdel-Satar, A. M. (1998): Distribution of some elements in River Nile environment at Great Cairo region. Ph.D. Thesis, Fac. of Sci. Cairo Univ., Egypt.
- Abouel Kheir, W.; Khalil, M.M.; Heikal, M.T. and Yousry, K.M. (2003): Assessment of Physico-chemical characteristics and phytoplankton growth in Lake Nasser. J. Environ. Sci., 7 (1): 59-84.
- Ali, M.M. (1987): Studies on the shoreline vegetation of Aswan High Dam Lake (Lake Nasser) and impact of the Lake on the desert. M.Sc. Thesis, Assiut Univ., 253pp.
- Ali, M.M. (1992): Ecological studies on freshwater macrophyte in regulated water bodies in Egypt and U.K. Ph.D. Thesis, Faculty of Sci., Aswan, Assiut Univ.
- Ali, M.M. (1998): Aquatic macrophytes in Egyptian Nubia Pre-and Post-formation of Lake Nasser in relation to environmental factors. ICLARM Symposium on Ecological Basis and Management Policy for sustainable Fish Production in Lake Nasser, High Dam Lake Development Authority, June 20-23, Aswan, Egypt.

- Ali, M. A.; Mageed, A. A. and Heikal, M. (2007): Importance of Aquatic macrophyte for invertebrate diversity in large subtropical reservoir. *Limnolgia* 37: 155- 169.
- Al-Zahaby, A.S.; Mageed, A.A.; Barbary, M.S.; Tawfik, T.A and Bedair, S.M. (2004): Feeding habits of fish from the River Nile in Shoubra El-Khima Region, Egypt. Second Regional Conference on Arab Water 2004 Action Plans for Integrated Development. 1-14.
- American Public Health Association (APHA) (1985): Standard methods of the examination of water and waste water. 15th edition, AWWA, WPCF, 1015Pp.
- American Public Health Association (APHA) (1992): Standard methods of the examination of water and waste water. 17th ed., AWWA, WPCF, 1015Pp.
- Arawomo, G.A.O. and Fawole, O.O. (1997): The food and feeding habits of *Sarotherodon galilaeus* (Artedi) in Opa reservoir of Obafemi Awolowo University, Ile-Ife, Nigeria. *Biosci. Res. Commun.*, 9: 15-20.
- Arora, J. and Mehra, N.K. (2003): Species diversity of planktonic and Epiphytic Rotifers in the Backwaters of the Delhi Segment of the Yamuna River, with remarks on new Records from India. *Zoological Studies*, 42 (2):239-247.
- Aruga, Y. and Monsi, M. (1963): Chlorophyll amount as an indicator of matter productivity in biocommunities. *Plant Cell Physiol.*, 4: 29-39.
- Azim, E. (1982): Biological studies of some Cyprinids from Lake Nasser. Ph.D. Thesis, Faculty of Science, Alexandria University, 172pp.
- Azim, E. (1991): Food and feeding habits of *Oreochromis niloticus* from different localities in Lake Nasser in spring and late summer. *Asw. Sc. Tech. Bull.*, 12:115-142.
- Belal, A.E. (1992) (ed.): Final report on Wadi Allaqi Eco-development project: Sustainable development: An experiment in Wadi Allaqi, Aswan High Dam Lake region, Egypt project number: FP/9101-88-91, 78pp.
- Bishai, H.M.; Abdel-Malek, S.A. and Khalil, M.T. (2000): Lake Nasser, Egyptian Environmental affairs Agency (EEAA), Publication of National Biodiversity Unit, 577pp.
- Boulos, L. (1986): Flora of the Nile region in Egyptian Nubia. *Feddes Repertorium*, 38 (3): 183-215.

- Bowen, S.H. (1982): Feeding digestion and growth-qualitative consideration. In. R. S. Pullin & R.H. Lowe-McConnel (Eds), The Biology and culture of Tilapias. ICLARM. Metro. Manila, Philippines, 141-156.
- Broberg, O. and Persson, G. (1988): Particulate and dissolved phosphorus forms in freshwater: Composition and analysis. *Hydrobiologia*, 170: 61 – 90.
- Campbell J.M.; Clark, J.J. and Kosinski, R. (1982): A technique for examining microspatial distribution of Cladocera associated with shallow water macrophytes. *Hydrobiologia* 97: 225-232.
- Cheruvelil K.S.; Soranno P.A. and Serbin R.D. (2002): Macroinvertebrate associated with submerged macrophyte: sample size and power to detect effects. *Hydrobiologia*, 441: 133-139.
- Clark, B. G. and Micheal, A. U. (1972): Waste water engineering in the water Resources and Environmental Engineering, McGraw-Hill New York.
- Clay, D. (1979): Population biology. Growth and feeding of African catfish (*Clarias gariepinus*) with special reference to juveniles and their importance in fish culture. *Arch. Hydrobiologia*, 87 (4) : 453-482.
- Delince, G. (1992): The ecology of the fish pond ecosystem with special reference to Africa. Text book, Kluwer Academic Publishers 230pp.
- Difonzo, C.D. and Campbell, J.M. (1988): Spatial partitioning of microhabitats in littoral cladocern communities. *Journal of Freshwater Ecology*, 4, 303-313.
- Downing J.A. and Cyr, H. (1985): Quantitative estimation of epiphytic invertebrate populations. *Canadian journal of Fisheries and Aquatic Science*, 42, 1570-1579.
- Drenner, R.W.; Vinyard, G.L; Gophen M. and McComass, S.R. (1982): Feeding Behavior of the cichlid *Sarotherodon galilous*: selective predation on Lake Kinneret zooplankton. *Hydrobiologia*, 87: 17-20.
- Edmondson. W.T. (1966): *Freshwater Biology*. 2nd ed. John Wiley and Sons. Inc. New York & London, 12248 pp.
- Edmondson. W.T. and Lill, A.H. (1987): *Conochillus* in Lake Washington. *Hydrobiologia*, 147: 157-162.
- El Shabrawy, G.M., & Dumont, H.J.: 2003, Spatial and seasonal variation of the zooplankton in the coastal zone and main khors of Lake Nasser (Egypt). *Hydrobiologia*, 491, 119–132.
- El Shahawy, M.: 1975, Studies on the physical properties of Lake Nasser. M.Sc. Thesis, Cairo Univ., Cairo. 128 pp.

- El-Enany, H.R.A. (2004): Ecological and biological studies on Lake El-Manzalah with special reference to their water quality and sediment productivity. M.Sc. Thesis Faculty of Science, Al-Azhar University. 386pp.
- Elewa, A.A. (1976): Chemical analysis for Lake Nasser, M.Sc. Thesis, Al-Azhar Univ., Cairo 220pp.
- Elewa, A.A. (1980): Studies on the distribution of some elements in water and sediments in the Aswan High Dam Reservoir, Ph.D. Thesis, Al-Azhar Univ., Cairo, 274pp.
- Elewa, A.A. (1985): Preliminary analysis of khor water in Lake Nasser, Bull. Inst. Oceanogr. & Fish., ARE, 2: 43-52.
- Elewa, A.A. (1987): Physical and chemical studies on some khors of Lake Nasser, Egypt J. Geol., 31(1-2): 253-265.
- Elewa, A.A.; Sayyah, S.M.; Latif, A.F.A. and Touffek, M.F. (1988): Nutrient status in lake Nasser and River Nile at Aswan, Egypt. Bull. Inst. Oceanogr. & Fish., ARE, 14(3): 177-188.
- El-Hadidi, M.N. (1976): The riverian flora in Nubia. In: the Nile, Biology of an Ancient River. Rzoska, J. (edit.) pp. 87-91. Dr. W. Junk B.V. Publishers, The Hague, Netherlands.
- El-Hawet, A.; Adam, E.; Sangq, Y. and El-Far, A. (2008): Assessment of Lake Nasser Fisheries. Egyptian Journal of Aquatic Research. 43(2): 285-298.
- El-Ngar, H.A.M.S (2008): Distribution and abundance of planktonic rotifers of aquatic habitats of the Nile Delta, Egypt. M.Sc. Thesis Faculty of Science, Al-Azhar University. 216pp.
- El-Otify A.M. (1985): Studies on phytoplankton of Aswan High Dam Lake M. Sc. Thesis Assiut Univeristy. Egypt.
- Entz, B.G.A. (1972): Comparison of the physical and chemical environments of Lake Volta and Lake Nasser. In: Proceeding of Freshwaters. UNESCO symposium in productivity problems of freshwaters. Kazimierz: Dolny: 883-891.
- Entz, B.G.A. (1974): Liminology, In: "Report on survey to Lake Nasser and Lake Nubia (1972-1973)", Entz, B.G.A & Latif, A.F.A (eds.), pp 4-63, Lake Nasser Development Center, Working paper No. 6.
- Entz, B.G.A. (1976): Lake Nasser and Lake Nubia. In:" The Nile Biology of an Ancient River", Rzoska, J. (ed.) pp 271-298, Dr W. Junk B.V., Publishers. The Hauge, Netherlands.

- Entz, B.G.A. (1980): Sedimentation process in the Reservoir Lake Nasser and Lake Nubia during 1965-1974 and future aspects. In: Water supply and management, Vol. 4, Biswas, A.K. (ed.) pp. 53-66. Pergamon press Oxford.
- Fead, E.M. (1980): Primary production of Lake Nasser. Thesis to School of public Health, Michigan Univ.
- Fernando, C.H., (1994): Zooplankton, fish and fisheries in fresh water. *Hydrobiologia*, 272: 105-123.
- Fernando, C.H., J.C. Paggi and R. Rajapaksa (1987): *Daphnia* in tropical low lands. *Mem. 1st. Ital. Hyd.*, 45:107-141.
- Fishar, M.R.A. (1995): Studies on bottom fauna in Lake Nasser, Egypt. Ph.D. Thesis, Suez Canal Univ., 267pp.
- Gaber, M.A. (1981): Report on zooplankton investigations in Lake Nasser (during August and September). Report, Aswan Regional Planning. Water Resources Department. 60 pp.
- Gaber, M.A. (1982): Report on phytoplankton investigations in Lake Nasser during October 1979. Report, Aswan Regional Planning. Water Resources Department. 56pp.
- Ghallab, M.H.M. (2000): Some physical and chemical changes on River Nile downstream of Delta Barrage at El-Rahawy drain. M. Sc. Thesis Fac. of Sci. Ain Shams Univ. Cairo, Egypt.
- Gharib, S.M. (1991): Study of biological productivity of Nozha Hydrodrom As a model of artificially fertilized fish farm. Ph.D Thesis, Fac. Sci., Mansoura Uni., 453pp.
- Gindy, A.R. (1991): Geochemical, mineralogical and sedimentological studies on some khor sediments of Lake Nasser. M.Sc. Thesis, Fac, Sci., Assiut Univ. 183pp.
- Goher, M.A. (1998): Factors affecting the precipitation and dissolution of some chemical elements in River Nile at Damietta Branch. M. Sc. Thesis, Fac. of Sci. Menofiya Univ., Egypt.
- Goldman, C. R. and Heron, A.J. (1983): *Limnology*. McGraw-Hill, New York, 464pp.
- Goma, R.H. and Abdel-Rahman, A. (1996): Preliminary report on long-term dissolved oxygen variation at El-Ramla 1985-1989. Work. Rep. Fish. Mang. Cent., Nasser Lake Overlap. Auth., Aswan 5: 53-60.

- Green, J. (1965): Zooplankton of Lake Mutanda, Bunyoyi and Mulehe, Proc. Zool. Soc. London, 144 (3): 383-404.
- Green, J. (2003): Association of planktonic and periphytic rotifers in a tropical swamp, the Okavango Delta, Southern Africa. *Hydrobiologia*, 490: 197-209.
- Habib, O.A. (1984): Distribution of chlorophyll *a* along the main channel of the High Dam Lake as an Index of productivity of phytoplankton. Work. Rep. Fish. Mang. Cent., High Dam Lake Development Auth., Aswan, 8pp.
- Habib, O.A. (1992): Distribution of chlorophyll *a* along the main channel of the High Dam Lake as an Index of primary productivity of phytoplankton. Work. Rep. Fish. Mang. Cent., High Dam Lake Development Auth., Aswan, 1: 127-134.
- Habib, O.A. (1995): Seasonal variations of the standing crop of phytoplankton in the northern part of the High Dam Lake, Egypt. Work. Rep. Fish. Mang. Cent., High Dam Lake Development Auth., Aswan, 4: 95-105.
- Habib, O.A. (1997): Limnological studies on Nasser Lake, Egypt, with special reference to phytoplankton and zooplankton. Ph.D. Thesis, Tokyo University of Fisheries, Japan.
- Habib, O.A. and Aruga, Y. (1988): Changes of the distribution of phytoplankton chlorophyll *a* in the main channel of the High Dam Lake, Egypt, J. Tokyo Univ. Fish., 75(2): 343-352.
- Habib, O.A.; Ioriya, T. and Aruga, Y. (1987): The distribution of chlorophyll *a* as an index of primary productivity of phytoplankton in Khor El-Ramla of the high Dam Lake, Egypt. J. Tokyo Univ. Fish., 76 (2): 145-157.
- Habib, O.A.; Mohamed, I.O.; Mohamed, M.S. and Aruga, Y. (1996): Seasonal changes of the Secchi disc depth and suspended solids at six stations along the main channel of the high Dam Lake Egypt, Work. Rep. Fish. Mang. Cent., Nasser Lake Development, 5: 169-184.
- Hammer, U.T. (1964): The succession of bloom species of blue-green algae and some casual factors. *Verh. Int. Ver. Limnol.*, 15: 829-836.
- Haney, J.F. and Buchanan, C. (1987): Distribution and Biogeography of *Daphnia* in the Arctic. *Men. 1st. Ital. Hydrobiol.* 45: 77-107.
- Hann, B.J. (1995): Invertebrates associations with submerged Aquatic plants in a Prairie wetland. USF (Delta Marsh). Annual report, Vol. 30.

- Hart, R.C. (1990): Zooplankton distribution in relation to turbidity and related environmental gradients in a large subtropical reservoir: Patterns and implications. *Freshwat. Biol.*, 24:241-263.
- Havens, K.E. (1991): Summer zooplankton dynamics in the limnetic and littoral zones of a humic acid lake. *Hydrobiologia*, 215:21-29.
- Heikal, M. T. and Abdel Bary, M.R. (1999): Stratification variability and trophic state along Lake Nasser. In Lake 99, Eighth International Conference on the Conservation and Management of Lakes, Copenhagen.
- Hureau, J.C. (1970): Biologie compare de quelques poisons antarctiques. (nototheniidae). *Bull. Inst. Oceanogr. Monac.*, 68: 1-250.
- Iskaros I.A. (1993): Biological studies on bottom fauna of Lake Nasser and adjacent water. M.Sc. Thesis. Faculty of Science. Alex. Univ. 184pp.
- Iskaros, I.A.; Bishai, R. M. and Mokhtar, F. M. (2008a): Seasonal variations of zooplankton and physico-chemical factors in Lake Nasser, Egypt. *Egypt J. Aquat. Res.* 34 (2): 232-259.
- Iskaros, I.A.; Bishai, R. M. and Mokhtar, F. M. (2008b): Comparative study of Zooplankton in Aswan Reservoir, and the River Nile at Aswan, Egypt. *Egypt. J. of Aquat. Res.* Vol. 34, No. (2) 260-284.
- Jeje, C.Y. (1988): A revision of the Nigerian species of the genera *Mesocyclops* Sars, 1914 and *Thermocyclops* Kiefer, 1927 (Copepoda: Cyclopoida). *Hydrobiologia*, 164: 171-184.
- Khalifa U.S.A.; Agaypi, M.Z. and Adam, H.A. (2000): Population Dynamics of *Oreochromis niloticus* L and *Sarotherodon galilaeus* Art. 87-90. in J. F: Craig (ed.) Sustainable fish production in Lake Nasser: ecological basis and management policy. ICLARM Conf. Proc. 61, 184 pp.
- Khalil, M.T. (1990): The physical and chemical environment of Lake Manzalah, Egypt. *Hydrobiologia*, 196: 193-199.
- Khallaf, E.A. and Alne-na-ei, A.A. (1987): Feeding ecology of *Oreochromis niloticus* (Linnaeus) and *Tilapia zilli* (Gervias) in a Nile Canal. *Hydrobiologia*, 146: 57-62.
- Krzyzanek, E. (1986): Zoobenthos of the small eheolimnic Wilsa-Czarre Dam Reservoir (south Poland) in the period 1975-1984. *Acta Hydrobiol.*, 28, 414-427.

- Latif, A. F. A. (1974): Fisheries of Lake Nasser and Lake Nubia. In report on Trip to Lake Nasser and Lake Nubia, by Entz, B. and Latif, A.F.A. Aswan regional planning. Lake Nasser development center. A.R.E., 64-137.
- Latif, A.F.A. (1978): Effect of impoundment on Nile biota in Lake Nasser. In: Proceeding international Symposium. Environmental effects of hydraulic engineering works, pp. 435-445, Tennessee Valley Authority, Knoxville, Tennessee, USA.
- Latif, A. F. A. (1981): Management of fish resources of Aswan High Dam Reservoir: A Joint study between the University of Michigan and Egyptian Academy of Scientific Research and Technology. In: Proceeding of the second Conference on Water Resources Management in Egypt. L-Assiouti, I. & Marks, D. (edits) pp. 516-642.
- Latif, A.F.A (1984): Lake Nasser. The New man-made lake in Egypt (with reference to Lake Nubia). In: "Lakes and Reservoirs", Taub, F.B. (ed.), pp 385-410, Elsevier Science Publishers, Amsterdam, Netherlands.
- Latif, A.F.A. & Co-Worker. (1982): water quality studies on the River Nile and Lake Nasser-Lake studies- Report on surveys to Lake Nasser and River Nile project, Acad. Sci. Res. Technol., Cairo.
- Latif, A.F.A. & Elewa, A.A. (1977): Physico-chemical characteristics of Lake Nasser, with special reference to Lake Nubia: In. Report on surveys to Lake Nasser and River Nile project, Latif, A.F.A. (ed.) pp. 19-91, Acad. Sci. Res. Technol., Cairo, 105pp.
- Latif, A.F.A. & Elewa, A.A. (1980): Lake studies, Physico-chemical characteristics of Lake Nasser and Lake Nubia, In. Report on surveys to Lake Nasser and River Nile project, Latif, A.F.A. (ed.) pp. 19-91, Acad. Sci. Res. Technol., Cairo, 105pp.
- Latif, A.F.A; El-Etreby, S.G. and Agaiby, I (1979): Bottom fauna of High Dam Reservoir. Report, River and Lake Nasser project. Academy of Scientific Research and Technology, Cairo, Egypt.
- Lauridsen, T.L .and Buenk I. (1996): Dial changes in the horizontal distribution of zooplankton in the littoral zone of two shallow eutrophic lakes. Arch Hydrobiologia, 137: 161-176.
- Lepš, J. & Šmilauer, P. (1999): Multivariate Analysis of Ecological Data. Faculty of Biological Sciences, University of South Bohemia, České Budejovice, 110pp.

- Mageed, A.A. (1992): Ecological studies on zooplankton from Khor El-Ramla (Lake Nasser), M.Sc. Thesis, Fac. Sci., Al-Azhar Univ., Cairo, 219.
- Mageed, A.A. (1995): Studies on zooplankton from Lake Nasser (Egypt), Ph.D. Thesis, Fac. Sci., Al-Azhar Univ., Cairo, 243.
- Mageed, A.A. (2004): Species composition, abundance and biomass of zooplankton with their importance to fishes in Wadi El Raiyan Lakes, Western Desert, Egypt. J. Egypt. Acad. Environ. Develop., (D-Environmental studies) 5 (2): 43-68.
- Mageed, A.A. and Heikal, M.T. (2005): Water quality and zooplankton dynamic in Lake Nasser, Egypt. Egypt J. Aquat. Biol. & Fish., Vol. 9, No. 3: 1-24.
- Mageed, A.A.; Abdel-Rahman, S.H.; Heikal, M.T.; Salem T.A and Salama E. (2008): Response of plankton prevalence to water quality variables in Lake Nasser during turnover period. Egyptian Journal of aquatic Research. 34 (2): 217-231.
- Malik, C.P. and Serivastava, A.K. (1982): Plant Physiology, Kalyant Publishers, New Delhi, 110002.
- Mohamed, I.O. (1992): Distribution of chlorophyll a concentration along the main channel of the High Dam Lake from January to December 1986. Work. Rep. Fish. Mang. Cent., High Dam Lake Develop. Auth., Aswan, 1: 213-236.
- Mohamed, I.O. (1993): Seasonal variations of chlorophyll a in the main channel of the High Dam Lake in 1986-1987. Work. Rep. Fish. Mang. Cent., High Dam Lake Develop. Auth., Aswan, 2: 133-154.
- Mohamed, I.O. (1995): Chlorophyll a concentration at six stations in the main channel of the High Dam Lake in 1990. Work. Rep. Fish. Mang. Cent., High Dam Lake Develop. Auth., Aswan, 4: 309-328.
- Mohamed, I.O. (1996): Vertical distribution of chlorophyll a concentration, water temperature and transparency in main channel of Lake Nasser. Work. Rep. Fish. Mang. Cent., High Dam Lake Develop. Auth., Aswan, 5: 83-99.
- Mohamed, I.O. and Ioriya (1998): *Microcystis aeruginosa* water blooms in Lake Nasser. ICLARM symposium on the Ecological Basis and Management Policy for Sustainable Fish Production in Lake Nasser. High Dam Lake Develop. Auth. June 20-23, 1998, Aswan, Egypt.

- Mokhtar, F.M. (2003): Ecological studies on the distribution of zooplankton in Lake Nasser and Adjacent waters. Ph.D. Thesis, Fac. Sci., Mansoura Univ., Egypt 361pp.
- Nessim, R.B. (1972): Liminological study of Lake Nasser. M.Sc. Thesis. Alex Univ., Alexandria, 295pp.
- Nielsen, D. L.; Hillman, T.J.; Smith, F.J. and Shile, R. J. (2000): The influence of a planktivors fish on zooplankton assemblages in experimental billabongs. *Hydrobiologia*, 434, 1-9.
- Ogutu-Ohwaya, R. (1984): The effect of predation by the Nile Perch. *Lates niloticus* (Linne) introduced onto Lake Kariba in relation to the fisheries of Lake Kyoga and Lake Victoria FHO. Report, 335: 18-39.
- Oronsaye, C.G. and Nakpodia, F.A. (2005): A comparative study of the food and feeding habits of *Chrysichthys nigrodigitatus* and *Brycinus nurse* in a tropical river. *Pak. J. Sci. Ind. Res.*, 48: 118-121.
- Oso, J.A.; Ayodele, I.A. and Fagbuaro, O. (2006): Food and Feeding habits of *Oreochromis niloticus* (L.) in a tropical Reservoir. *World Journal of Zoology*. 1 (2): 118-121.
- Paasche, E. (1980): Silicon. In I. Morris (ed.). *The physiological ecology of phytoplankton*. Blackwell Scientific Publication, Oxford. *Studies in Ecology* volume 7: 259 – 284.
- Pennak, R.W. (1978): *Freshwater invertebrates of the United States*. 2nd ed. John Wiley & Sonns. New York, 803 pp.
- Phillippart, J.Cl and Ruwet, J.Cl. (1982): Ecology and distribution of tilapias. In. R. S.V. Pullin & R.H. Lowe-McConnel (Eds), *The Biology and culture of Tilapias*. ICLARM. Metro. Manila, Philippines, 141-156.
- Pontin, R. (1978): A key TO British freshwater planktonic Rotifer, *Freshwater Biol. Assoc. Sci. Publ.*, 38. Britain 178pp.
- Rey, J. and L. Saint-Jean (1968): Les cladoceres (Crustacea, Branchiopodes) Du Tchao. *Cah. O.R.S.T.O.M., Ser. Hyd.*, II (3-4): 79 118.
- Ruttner- Kolisko, .A. (1974): Planktonic rotifers. *Biology and Taxonomy. Binnengewasser. Suppl.*, 26:146pp.
- Rzoska, J. (1976): Zooplankton of the Nile System. In J. Rzoska (Ed.), *The Nile, biology of an ancient river* (pp. 333–343). The Hague: W. Junk.

- Sabae, Z.S. and Abdel Rahman S.H. (2008): Bacteria as indicator of sewage pollution and as trophic levels in some selected khors of Lake Nasser, Egypt. *Egypt. J. of Aquat. Res.*, 34 (2): 206-216.
- Sader, J.P., (1991): Inorganic chemistry and Drug Design. In *Adv. Inorganic Chem.*, Ed. A.G. Sykes. Academic press, 36: 1- 44.
- Sakuma, M; Hanazato, T. and Nakazato, R. (2002): Abundance of Chydoridae associated with plant surface, water column and bottom sediments in the macrophyte zone of a lake. *Verh. Internat. Verein. Limnol.* 28.
- Saleh, A.M. (1976): A study on the phenomenon of taste and odor in Lake Nasser. Report. Academy of Science Research and Technology. Cairo. Egypt. 67pp.
- Samaan, A.A. (1971): Report on trip to Lake Nasser to investigate its primary productivity during March 1971, Rep LUNDC., Aswan, 11pp.
- Samaan, A.A, and Gaber, M.A. (1976): Report on plankton investigation in Lake Nasser during March and August, 1976. In: AFA Latif (ED.). Reports on surveys to Lake Nasser & River Nile Project. Acad. Sci. Res. & Tech., Cairo, Egypt. 13pp.
- Schmid-Araya, J.M. and Zuniga, L.R. (1992): zooplankton community structure in two Chilean reservoirs. *Arch. Hydr.*, 123(3): 309-335.
- Shehata, K.K; Shehata, S.M.A.; Hussien, M.M. and Abd El-Mageed, A. (1996a): Taxonomical and ecological studies on some Zooplankton species; Ciliata, Rhizopoda, Turbellaria and Crustaceae of high Dam Lake. *Egypt. J. Aquat. Biol. & Fish.*, 2 (1): 37-63.
- Shehata, S.M.A.; Shehata, K.K; Hussien, M.M. and Abd El-Mageed, A. (1996b): Ecological aspects of high Dam Lake with special reference to the seasonal distribution of the fresh water copepods. *Al-Azhar Bull. Sci.*, 7 (1): 731-761.
- Shehata, S.M.A.; Shehata, K.K; Hussien, M.M. and Abd El-Mageed, A. (1998a): Taxonomy, population structure, and species diversity of Rotifera in the high Dam Lake. *Egypt. J. Biol. And Fish.*, 2(1):1-36.
- Shehata, S.M.A.; Shehata, K.K; Hussien, M.M. and Abd El-Mageed, A. :(1998b): Taxonomical and ecological studies on some zooplankton species: Ciliata, Rhizopoda, Turbellaria and Crustacea of the High Dam Lake. *Egypt. J. Biol. And Fish.*, 2(1): 37-63.

- Shoreit, A.A.M.; Abd-Alla, M.H. and Shabel, M.S.A. (1992): Acetylene reduction by Rhodospirillaceae from the Aswan High Dam Lake, World J. Microbiology. & Biotechnology, 8: 151-154.
- Siliem, T.A.A. (1984): Chemical studies on pollution in the Damietta Nile branch between Faraskour Dam and Ras El Bar Outlet Ph.D. Thesis submitted to Fac. Sci. Alex. Univ., 253pp.
- Springuel, I.; Ali, M.M. and Murphy, K.J. (1990): Aquatic macrophyte growth in relation to water level regime in the River Nile and its impoundments in Upper Egypt. Proceedings EWRS 8th Symposium on Aquatic weeds: 199-201.
- Taha, O.E. and Mageed, A.A. (2002): Spatial distribution and relationship between phytoplankton and zooplankton in Lake Nasser (Egypt) after the flood season. Egypt J. Aquat. Biol. & Fish., 6 (4): 265-281.
- Talling, J.F., & Razoska, J.: (1967): The development of plankton in relation to hydrological regime in the Blue Nile. J. Ecol., 55: 637-662.
- Talling, J.F., & Talling, T.B.: 1965. The chemical composition of African lake waters. Internationale Revue der Gesamten Hydrobiologie, 50(3), 421–463.
- Train, R.E. (1979): Quality criteria for water. Castle House Publication LTD., London, 256pp.
- Vijverberg J. and Boersma M. (1997): Long-term Dynamics of small-bodied and large bodied Cladocerans during the eutrophication of a shallow reservoir, with special attention for *Chydorus sphaericus*. Hydrobiologia, 360 :233-242.
- Windell, J. T. and Bowen, S.H., (1978): Methods for study of fish diets based on analysis of stomach contents. In Methods for the Assessment of Fish Production in Fresh Waters. 3rd edn. (T. Bagnel. Ed.) pp. 219-226. Oxford: Blackwell Scientific Publications.
- Zafar, A. R. (1964): On the ecology of the algae in certain fresh ponds of Hyderabad, India. Hydrobiologia., 23: 179 – 195.
- Zaghloul, F.A. (1985): Seasonal variations of plankton in Lake Nasser, Ph.D. thesis, Fac. Sci., Suez Canal Univ., 364pp.
- Zahran, M.A. and Smith, A.J. (2003): Plant life in the River Nile in Egypt. Mars Publishing House. 518pp.
- Zrum, L. and Hann J.B. (1997): Planktonic microinvertebrate community structure in a prairie wetland in response to additional inorganic nutrient and organ phosphorus insecticide. UFS (Delta March) Annual Report, 32: 56-67.
- Zyadah, M.A. (1995): Environmental impact assessment of pollution in Lake Manzalah and its effect on fish. Ph.D. Thesis, Fac. Sci., El-Mansoura Univ.127p.