

REFERENCES

- Abdulrahman, A., K. and Nasimuzzaman, M. (2001):**Antimicrobial resistance patterns of Gram –negative bacteria isolated from urine culutures in almanac general hospital . Annals of Saudia Med. ; 21: 51-52.
- Adedeji , B. A. M. ,and Abdulkadir, O. A.(2009):** Etiology and antimicrobial resistance pattern of bacterial agents of urinary tract infections in students of tertiary institutions in Yola Metropolis . Department of Microbiology, Federal University of Technology, P.M.B. 2076. Advances in Biological Research 3 (3-4): 67-70.
- Ahmad ,S. and Ahmad ,F.(1995):**Urinary tract infection at a specialist hospital in Saudia Arabia. Bangladesh Med. Res. Counc .Bull , 21:95-98.
- Ahmad, I. and Aqil , F. (2007):** In vitro efficacy of bioactive extracts of 15 medicinal plants against ESbL-producing multidrug-resistant enteric bacteria. Microbiol. Res., 162: 264-275.
- Ahmed, F.M. (2004):** Antimicrobial studies of some essential oils on some bacteria; Master Degree; Botany Department; Faculty of Science; Zagazig University; Egypt.
- Aibinu ,I.; Adenipekun ,T.; Adelowotan ,T.; Ogunsanya ,T. and Odugbemi, T.(2007):** Evaluation of the antimicrobial properties of different parts of *Citrus aurantifolia* (Lime fruit) as used locally. Afr. J. Trad. CAM. 4:185-190.
- Alanis, A.D.; Calzada, F. ;Cerrantes, J .A. ;Torres, J. and Ceballos, G.M.(2005):** Antibacterial properties of some plants used in Mexican traditional medicine for the treatment of gastrointestinal disorders. J Ethnopharmacology , 22;100(1-2):153-157.

- Al-Haddad, A.M (2005):**Urinary tract infection among pregnant women in Al-Mukalla district, Yemen Eastern Mediterranean Health Journal;11 (3) : 505-510.
- Allan ,R. (1984):**Current concept in the management of urinary tract infection in adults . Medical Clinic of north America; 68(2):59-76.
- Allan ,R. (2003):**The etiology of urinary tract infection. Tradition and Emerging pathogens. Dis Mon ,49:71-82.
- Amit ,R. and Shailendra, S. (2006):**Ethnomedicinal approach in biological and chemical investigation of phytochemicals as antimicrobials. Available at <http://www.phamainfo.net>.
- Andrade, S.S; Sader, H.S.; Jones, R.N. and Pereira, A.S. (2006):** Increased resistance to first-line agents among bacterial pathogens isolated from urinary tract infections in Latin America. Mem. Inst. Oswaldo. Cruz, Rio de Janeiro, Vol. 101 (7):741-748.
- Ayad(2009):** Evaluation of antibacterial efficiency of some plant extracts to control urinary tract system pathogens. Master Degree, Benha University, Faculty of Science, Botany Department.
- Ayana, R.; Remya ,R. and Deepthi, S. R.(2008):** Antimicrobial activity studies on *Cissus quadrangularis* . Indian Journal of Botanical Research, 4 (2): 201-204.
- Ayoola, G. A. ; Lawore, F. M. ; Adelowotan , T. , Aibinu , I. E. ; Adenipekun , E. ; Coker H. A. B.; and Odugbemi , T. O.(2008):** Chemical analysis and antimicrobial activity of the essential oil of *Syzigium aromaticum* (clove).African Journal of Microbiology Research. Vol.(2): 162-166 .
- Bariska, J.; Valko,K.; Takacs-Novak, K. and Kalasz, H. (1999):**Planar chromatography, J. Planar Chromatogr. 12:46-50.

- Baron, E.J. and Finegold ,S.M.(1990):**General microbiology; Baily and Scotts Diagnostic Microbiology. 8th. Ed. Mosby Co. St. Louis. Balitimore.
- Barry, L.B. and Thrnsberry, Y.(1987):**Inhibition of growth and aflatoxin production by cinnamon and clove oils, cinnamic aldhyle and eugenol. Food sci. 42:1107-1109.
- Barrett ,S.P; Savage ,M.A; Rebec ,M.P ;Guyot ,A ;Andrews ,N and Shrimpton ,S.B(1999):**Antibiotic sensitivity of bacteria associated with community –acquired urinary tract infection in Britain Journal of Antimicrobial Chemotherapy ;44: 359-365.
- Bauer, A. W.; Kirby, W. M.; Sherris, .J. C. and Truck, M. (1986):** Antibiotic susceptibility testing by the standard single disk method. J. Clane. Pathol., 45: 493-496.
- Becerril R; Gómez-Lus, R.; Goñi, P.; López, P. and Nerín, C. (2007):** Combination of analytical and microbiological techniques to study the antimicrobial activity of a new active food packaging containing cinnamon or oregano against *E. coli* and *Staph. aureus*. Anal Bioanal Chem. Jul;388(5-6):1003-1011.
- Bengtsson ,C.; Bengtsson ,V. and Lincoin ,K. (1980):**Bacteriuria in a population sample of women :Prevalence ,characteristics ,results of treatment ,and prognosis , Acta Med SCi.,208:417-423.
- Bethesda, M.D (1990):**The National Kidney and urologic Diseases. National institutes of Health, 90:583 .
- Bethesda, M.D. (2005):**National kidney and urologic Diseases Information Clearing house .NIH Publication , No. 08–3925.
- Betoni, J.E.C.; Mantovani, R.P.P.; Barbosa, L.N.; Di Stasi, L.C. and Fernandes Junior, A. (2006):** Synergism between plant extract and antimicrobial drugs used on *Staphylococcus aureus* diseases. Mem. Inst. Oswaldo Cruz., 101: 387-390.

- Betty ,A.F. ; Daniel , F.S. and Alice ,S.W.(1998):**Diagnostic microbiology. Tenth Edition .Moshy publishing.
- Beyer ,I. ; Mergan ,A ;Benoit ,F. ;Theunissen ,C. and Peper sack ,T. (2001):**Management of urinary tract infections in the elderly .Z. Gerontol Geriatr ,34:153-157.
- Bigelow, J. (1918):** American medicinat botany". Vol 1, 2 and 3 (1948) screening methods for detemining antibiotic activity of higher plants . J. Bact. 55: 235- 240.
- Boon, E.M.; Downs, A. and A. and Marcey, D. (2007):**Catalase:H₂O₂ oxidoreductase. Catalase structural tutorial text. Retrived on 2007-02-11.
- Boyd , L.B. ; Maynard M.J.; Morgan-Linnell , S.K., Horton , L.B.; Sucgang , R. ; Hamill ,R.J.; Jimenez, J.R.; Versalovic ,J.; Steffen D., and Zechiedrich , L. (2009) :**Relationships among ciprofloxacin, gatifloxacin, levofloxacin, and norfloxacin MICs for fluoroquinolone-resistant *Escherichia coli* clinical isolates . Antimicrob Agents Chemother. Jan;53(1):229-234.
- Boyko ,E. J. ;Fihn, S.D and Scholes, D. (2002):** Diabetes and the risk of acute urinary tract infection among postmenopausal women. Diabetes care ,25:1778-1783.
- Brandberg ,A. ;Herner ,B. ; Swartling ,P.G. and Vahlne ,G.(1970):**Urine vagsinfektion hosden vuxna befolkningen i en svensk stand .La Kartidningen ,67:2655-2662.
- Brandberg ,A ;Herner ,B; Kaijser ,B.; Kjerru , IFL ;Lindblom ,G.B and Swarting P.G(1978):**Skall asymtomatisk bakteriuri behandlas? Lakartidningen ,75:2115-2118.
- Brocklehurst ,J.C ;Dillane ,J.B;Griffithl and Fry ,J (1968):**The prevalence and symptomatology of urinary infection in an agent population .Gerontol clin (Basel),10:242-253.

- Cai, L. and Wu, C.D. (1996):** Compounds from *Syzygium aromaticum* possessing growth inhibitory activity against oral pathogens. J. Nat. Prod., 59(10): 987-990.
- Campo ,J. ; Amiot, M.J. and Nguyen-The, C. (2000):**Antimicrobial effect of rosemary extracts. J. Food Prot.;63(10):1359-1368.
- Carlini, E. A.; Contor, J. P. and Silva filho (1986):** Pharmacology of lemon grass (*Cymbopogon citrates*). J. of Ethnopharmacology 17: 37- 64.
- Carlson, H. J.; Bissell, H. D. and Muller, M. G. (1948):** Antimalarial and antibacterial substances separated form higher plants. Bact., 52: 155-168.
- Chaieb, K.; Hajlaoui, H.; Zmantar, T. and Nakbi, K.A.B.(2007a):**The chemical composition and biological activity of essential oil, *Eugenia cryophyllata* (*Syzygium aromaticum* L. Myrtaceae): A short review. *Phytother Res.*, 21(6): 501-506.
- Chaieb, K.; Zmantar T.; Ksouri, R. and Hajlaoui (2007b):**Antioxidant properties of essential oil of *Eugenia caryophyllata* and its antifungal activity against a large number of clinical *Candida species*. *Mycosis*, 50(5): 403-406.
- Cheesbrough ,M.(1993):**Collection, transport and examination of specimens. In: Cheesbrough ,M.(ed).Medical laboratory Manual for Wort- Heinemann Ltd., Oxford.
- Cheesbrough ,M.(2000):**Biochemical test to identify bacteria .In Cheesbrough ,M.(ed). Laboratory practice in Tropical Countries , Vol. 2:63-70.Cambridge University press ,U.k.

- Clem ,M. ; Allan , D.; and Iain, H. (2006) :** A study into the antimicrobial effects of cloves (*Syzygium aromaticum*) and cinnamon (*Cinnamomum zeylanicum*) using disc-diffusion assay. Journal: Nutrition & Food Science ;36 (4): 225 – 230
- Collee ,J.G. ; and Marr ,W.(1989):** Culture containers and culture media .In practical medical microbiology 13th ed. Vol. (2).Collee ,J.G ; Duguid, J.P.; Fraser ,A.G. and Marmion ,B.P; (eds); Churchill. Livingstone .New York .PP.100-120.
- D'Amato, R.F.; Isenberg, H.D. and Mckinley, G.A. (1985) :** Collaborative evaluation of the UniScept qualitative antimicrobial susceptibility test. J Clin Microbiol 21: 293-297.
- Dalia, A. (2009):** Isolation,characterization and control of some clinical *Candida albicans*. Ph.D. Degree, Zagazig University. Faculty of Science. Botany Department.
- Dan, B. ; Steven ,C.; Erich, S. and Andrew ,G. (2004):**Chinese herbal medicine: Materia Medica, Third Edition.
- David ,S.H. ;Joseph ,A.S. ;Francisco, T.; Richard, S. ;John ,D.H. and Pamela ,L.D. (2008):**Urinary tract infections ,Female. e medicine from web MD. Comprehensive Study Guide. 6th ed. New York: 606-612.
- Drago, L.; Nicola , I.; Mattina ,R. and Vecch, E.D.(2010) :** In vitro selection of resistance in *Escherichia coli* and *Klebsiella spp.* at in vivo fluoroquinolone concentrations. B.M.C. Microbiology, 1471-2180.
- Dontas ,A.S. ;Kasviki-charvati ,P. ;Papa nayiotou ,P.C. and Marketos ,S.G. (1981):-**Bacteriuria and survival in old age. N Engl J Med,304:939-943.

- Ehrlich, G. ;Hu, F. ;Shen, K. ;Stoodley, P. and Post, J. (2005):** Bacterial plurality as general mechanism driving persistence in chronic infections .Clin . Orthop. Related RES:20-4 PMID 16056021.
- Eiros, B. J.M.; Ochoa, S. C.; and Grupo I. (2007):** Etiological profile of urinary tract infections and antimicrobial susceptibility of urinary pathogens An Pediatr (Barc).;67(5):461-468. PMID: 17991366.
- El-Mishad, J.(1991):**Microorganisms isolated from urinary tract infections and their beta lactamase production and evalution of clinical efficacy of sulperazone .Hinyokika Kivo 34 :8:1503-1514.
- Emeka, I. and Elizabeth ,E.(2009):** Justification for the use of *Ocimum gratissimum* L in herbal medicineand its interaction with disc antibiotics. B.M.C. Complementary and Alternative Medicine, 28, 9:37. Available on line at <http://www.biomedcentral.com /1472-6882/9/37>.
- Evans, D.A. ;Kass, E.H ;Hennekens, C.H. and *etal* (1982):-**Bacteriuria and subsequent mortality in women. Lancet,1:156-158.
- Faro, S. and Fenner ,D.E. (1998):**Urinary tract infections.Clin obstet Gynecol. 41:744-754.
- Farrell, D. and Qlobinson, M. (1972):**Nutrient and blood agar.J. Appl.Bact.,25:625-630.
- Fihn ,S. D. ;Boyko ,E.J. and Normad ,E.H. (1996):**Association between use of spermicide –coated condms and *Escherichia coli* urinary tract infections in young women . AM J Epidemiol,144:512-520.

- Fihn ,S.D. ;Boyko ,E.J. and Chen ,C. L. (1998):**Use of permicide – coated condoms and other risk factors for urinary tract infection caused by *Staphylococcus saprophyticus*. Arch Intern Med ,158:281-287.
- Fihn ,S.D. ;latham ,R.H. ;Roberts ,P. ;Running ,K. and Stamm ,W.E. (1985):**Association between diaphragm use and urinary tract infection ,JA MA,254:240-245.
- Foxman, B.(1990):**Recurring urinary tract infection :incidence and risk factors .AMJ public Health ,80:331-333.
- Foxman ,B. ;Barlow ,R. ; Arcy ,H. ;Gillespie ,B. and Sobel ,J.D. (2000):**Urinary tract infection :self reperted incidence and associated costs. Ann Epidermoil ,10:509-515.
- Friedman , M.; Henika, P.R. and Mandrell, R.E. (2002)** Bactericidal activities of plant essential oils and some of their isolated constituents against *Campylobacter jejuni*, *Escherichia coli*, *Listeria monocytogenes*, and *Salmonella enterica*. Journal of Food Protection 65, 1545-1560.
- Gary, K. (2007):**Gram stain and capsule stain. Biol. 230 Microbiology Laboratory Manual.
- Gebre- Selassie, S. (1998):**Asymptomatic bacteriuria in pregnancy: epidiomology, clinical and microbiological approach. Ethiop.Med.J.,36:185-192.
- Geo, F. ;Janets ;Butel ;Stephan ,A. and Morse (2004):**Medical Microbiology .Twenty –third edition. Published by the Mc Graw-Hill companies.
- George ,A. and Max, D. (1988):**Biology ,fifth edition ,813-818.Macmillan publishing company .New York.

- Gerand ,J. ;Berdell ,R.; Funke and Christinel ,L. (2007):**Microbiology. ninth edition .Published by person Benjumin Cummings.
- Ghaleb, A. ; Abu-Shanab, B. A. and Adwan, K. M.(2008):-** *In vitro* activity of certain drugs in Combination with plant extracts against *Staphylococcus aureus* infections Pak.J.Med Sci ,24(4)541-4.
- Ghosh ,R.; Nadiminty, N.; Fitzpatrick, J.E. ;Alworth, W.L. ;Slaga, T. J. and Kumar, A.P. (2005):** Eugenol causes melanoma growth suppression through inhibition of E2F1 transcriptional activityJ.Biol. Chem., 280, 5812.
- Goetz, A. and Yu ,V.L.(1997):**Intensive care unit :the hottest zone .Curr Opin Infect Dis ,10:319-323.
- Goldstein ,F.W. (2000):**The Multicentre study Group.Antibiotic susceptibility of bacterial strains isolated from patients with community-acquired urinary tract infections in France .Eur J. Microbiol. Infect Dis. ,19:112-117.
- Gottlieb ,P.L.(1995):**Comparison of enxacin versus trimethoprim – sulfamethoxazole in the treatment of patients with complicated urinary tract infection Clin .ther ,17:493-502.
- Guidoni , E.B.; Berezin, E.N.; Nigro, S. and Santiago, N.A. (2008):** Antibiotic resistance patterns of pediatric community-acquired urinary infections .J.Braz. J. Infect Dis.;12(4):321-323. PMID: 19030734.
- Gunnar, K. (2000):**The Eco .SENS P roject :apropective ,multinational, multicenter epidermiological survey of the prevalence and antimicrobial susceptibility of urinary tract pathogens –interim report .Journal of antimicrobial chemotherapy 46,suppl.s1 ,15-22.

- Gunnar ,R.; Brandberg, C. and Anders, S.(1986):**Bacteriuria and mortality in an elderly population. The New England and Journal of Medicine ; 314 (18): 1152-1156.
- Gupta, K. ; Stamm ,W.E. and Facp, M.D. (2005):** Urinary tract infections ACP medicine. Available at:<http://www.medscape.com/viewarticle/505095>
- Gupta, K. ;Scholes ,D. and Stamm ,W.E. (1999):**Increasing prevalence of antimicrobial resistance among uropathogens causing acute uncomplicated cystitis in women .J. am. Med. Assoc ,281:736-738.
- Gupta, K ; Hooton ,T.M. and Stamm ,W.E (2001):**-Increasing resistance to fluoroquinolones in *Escherichia coli* from urinary tract infections in the Netherlands .J. Antimicrob. Chemother,46:223-228.
- Hammer, K. A.; Carson C. F.and Riley T. V. (1999):** Antimicrobial activity of essential oils and other plant extracts Journal of Applied Microbiology ;86 (6) :985–990 .
- Henrique, D.M. ; José, G.M.; Edeltrudes, O. ; Vivyanne S. and José, P. (2008).:** Enhancement of the Antibiotic Activity against a Multiresistant *Escherichia coli* by *Mentha arvensis* L. and Chlorpromazine. *Chemotherapy*;54:328-330.
- Hilf ,M.; Yu, V.L. and Sharp ,J. (1989):**Antibiotic therapy for *Pseudomonas aeruginosa* bacteremia :outcome correlation a prospective study of 200 patients. Am .J. Med. ,87:540-546.
- Holt, J.G. ; Krieg, N.R. ; Sneath ,P .H. ; Staley ,J. T. ; Williams ,T. and Hensyl, W.R.(1994):**"Bergy's manual of determination bacteriological" :9th ed Williams and Wilkins, Baltimore, Meryland.

- Hooton ,T. M. ;Latham, R. H. ;Wong, E. S.; Roberts, P.L. and Stamm, W.E.(1989):**Ofloxacin versus trimethoprim-sulfamethoxazole for treatment of acute cystitis . Antimicrobial-Agents chemother,33:1308-1312.
- Hooton ,T. M ;Johnson,C. and Winter, C. (1991):**Single –dose and three –day regimens of loxacin versus trimethoprim – sulfamethoxazole for acute cystitis in women .Antimicrob. Agents chemther. ,35:1479-1483. (<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?>
- Hooton, T.M.; Roberts, P.L. and Stamm, W.E. (1994):** Effects of recent sexual activity and use of a diaphragm on the vaginal microflora. Clin Infect. Dis. 19:274-278.
- Hooton ,T. M.; Scholes ,D. and Hughes ,J .P. (1996):**A prospective study of risk factors for symptomatic urinary tract infection in young women . Engl. J. Med. ,335:468-474.
- Huang, S., G. Contronic and W. Khan(1999):** Resistance of *shigella* to ampicillin and other antibiotics . Clinical and epidemiological application. JAMA., 221: 45-47.
- Husnu P. S. ; Aydemlr ,M. I. ; Feriha C. ; Alper, T. ,and Sercan U.(2008):** Susceptibility of Extended-Spectrum Beta-Lactamase-Producing *Escherichia coli* Urine Isolates to Fosfomycin, Ciprofloxacin, Amikacin and Trimethoprim- Sulfamethoxazole. Turk J Med Sci; 38 (2): 175-180.
- Ikram, M. and Inamul- Hag. (1980):** Screening of medicinal plants for antimicrobial activity. Part 1 Fitoterapia, 51 (6): 231- 235.

- Iravani ,A. ;Klimberg ,L.; Briefer ,C. ;Munera ,C. ;Kowalsky ,S.F. and Echols ,R.M. (1999):**A trial comparing low –dose ,short – course ciprofloxacin and standard 7 day therapy with co-trimoxazole or nitrofurantoin in treatment of uncomplicated urinary tract infection .J. Antimicrob chemother ,43 (suppl A):s 67-s75.
- Iroha, I.R. ; Adikwu , M.U. ; Esimone , C.O. ;Aibinu ,I. and Amadi ,E.S. (2009) :**Extended spectrum Beta-Lactamase (ESBL) in *E.coli* isolated from a tertiary hospital in Enugu state ,Nigeria .Pak J Med Sci ; Vol. 25 No. 2: 279-282.
- Isogai, E.; Isogai, H.; Hirose, K.; Hayashi, S., and Oguma, K.(2001):**In vivo synergy between green tea extract and levofloxacin against enterohemorrhagic *Escherichia coli* O157 infection._Curr Microbiol.;42(4):248-51PMID: 11178724.
- Jackie, R.(2005):**Oxidase test, Richland College 12800 Abrams RD Dallas, T X, 75243US 972.238.6100.
- James, K. (2008):**Nitrofurantoin should become the first- line therapy of choice. B .J.U. Int.;102:1634-1637.
- Jepson ,R.; Mihaljevic ,L. and Craig ,J.(2000):**Cranberries for preventing urinary tract infections .Cochrane Database Syst. Rev. :CD001321.PMID 14973968 (<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?>
- John, P. (2008):** Studies on Antimicrobial Efficiency of *Citrullus colocynthis* (L.) Schrad: A Medicinal Plant. Ethnobotanical Leaflets 12: 944-947.
- Johnson, J.R.; Kuskowski ,M.A.; Menard, M.; Gajewski, A.; Xercavins, M.; Garau ,J.(2006):-** Similarity between human and chicken *Escherichia coli* isolates in relation to ciprofloxacin resistance status. J Infect Dis. 1;194(1):8-10.

- Johnson ,J.R and Stamm ,W.E (1989):**Urinary tract infections in women :Diagnosis and treatment .Ann inter Med ,111:906-917.
- Judith ,A.O. ;Steven ,P. and Elius, A.(2002):**Selecting drug regimens for urinary tract infections :Current recomendations .Infect Med 19(1):14-22.
- Justice, S. ,Hunstad, D., Seed, P. and Hultgren, S. (2006):** Filamentation by *Escherichia coli* subverts innate defenses during urinary tract infection Proc Natl Acad Sci .USA 103(52):19884-9.
- Kader ,A.A.; Kumar ,A. and Dass ,S.M.(2004):**Antimicrobial Resistance Patterns of Gram –negative Bacteria Isolated from Urine Culutures at a General Hospital .Saudi J Kidney Dis. ;15:135-9 .
- Kahlmeter ,G.(2003):**An international survey of the antimicrobial susceptibility of pathogens from uncomplicated urinary tract infections :the Eco .SENS project .Journal of antimicrobial chemotherapy ,51:69-76.
- Kang, D. H. and Fung, Y.C. (1999) :**Effect of diacetyl on controlling *Escherichia coli* 0157: H7 and *Salmonella Typhimurium* in the presence of starter culture in a laboratory medium and during meat fermentation. Journal of Food Protection 62, 975-979.
- Kass, E.H. (1957):-**Bacteriuria and the diagnosis of infections of urinary tract .Arch Intern Med ;100:709-714.
- Kaur, G. J. and Daljit ,S. A.(2010):** Bioactive potential of *Anethum graveolens*, *Foeniculum vulgare* and *Trachyspermum ammi* belonging to the family Umbelliferae - Current status .J. of Medicinal Plants ; 4(2): 087-094.

- Kim, J.M.; Marshall, M.R. and Wei, C.I. (1995)** :Antimicrobial activity of some essential oils components against five food borne pathogens. *Journal of Agriculture and Food Chemistry* 43, 2839-2845.
- Kim, M.E. and Cho, YH. (2008)**:Prevalence of antimicrobial resistance among uropathogens causing acute uncomplicated cystitis in female outpatients in South Korea: a multicentre study in 2006. *Int.J. Antimicrob. Agents*; 31 (1): 515-518.
- Kim, C. S.; Kim, M. E.; Yong-Hyun, C.; Rae, C. and Gilho, L.(2010)**:Virulence Characteristics and Phylogenetic Background of Ciprofloxacin Resistant *Escherichia coli* in the Urine Samples from Korean Women with Acute Uncomplicated Cystitis. *J. Korean Med. Sci.*; 25: 602-607.
- Koch, F. E.; Zentralbl, B. P. and Abt (1942)**:Mannitol salt agar. 1 Orig. 149.
- Komaroff A.L.(1986)**: Urinalsis and urine culuture in women with dysuria .*Ann Intern Med* ;104:212-218.
- Koneman ,E.W ;Allen ,S.D; Dowell, V.R. and Cho, Y.(1983)**: Colour atlas and text book of diagnostic microbiology.2nd ed.J.P. Lippincott Company; N.Y.;London. 157-175.
- Krieg and Holt (1994)**:Bergy's manual of systematic bacteriology, Vol. 1. Williams& Wilkins, Baltimore, Md.
- Kumar, P.; Bhatt R.P. ; Singh, L. ;Shailja; Chandra, H. and Prasad, R.(2010)**: Identification of phytochemical content andantibacterial activity of *Juniperus communis* leaves. *International Journal of Biotechnology and Biochemistry*. Volume 6 Number 1 pp. 87–91.

- Kunin ,C .M. (1978):**Sexual intercourse and urinary tract infections .N Engl Med ,298:336-337.
- Kunin ,C .M. ,White L.V., Hua T.H. (1993):**A reassessment of the importance of low –count in young women with acute urinary symptoms .Ann Intern. Med. ;119:454-460.
- Kunin ,C.M.(1994):**Urinary tract infections in females .Clin .Infect .Dis ;18 (1):1–12.
- Kunin ,C.M.(2001):**Nosocomial urinary tract infections and the indwelling catheter: What is new and what is true? Chest; 120:10-12.
- Laemmli, V. K. (1973):** Cleavage of structural proteins during the assembly of the head of bacteriophage T₄. Nature (London), 227 : 680-685.
- Latiful, B. ; Hoque, M.M.; Inatsu,Y. And Kawamoto, S.(2007):**Antibacterial activity of cloves and cinnamon extracts against food borne pathogens and spoilage microorganisms. National food research Institute , Japan. Journal of Food Science and Technology. 72:9-21.
- Lau, W.S.(1999):**Infrared characterization for microelectronics. World Scientific.
- Lau, S.M.; Peng, M.Y. and Chang, F.Y. (2004):** Among pathogens of both bacteremic and non- bacteremic community –acquired urinary tract infection.

- Leonid ,S. and Vladimir ,V. (2006):**Antimicrobial susceptibility of pathogens isolated from adult patients with uncomplicated community –acquired urinary tract infections in the Russian Federation: two multicentre studies ,UTIAP -1and UTIAP-2 International Journal of Antimicrobial Agents ,Volume 28,Supplement 1.pages 4-9.
- Lerner, G.R. (1994):**Urinary tract infections in children *Pediatr. Ann.*, 23;463-473.
- Liza D. ;Scott H. ;Francisco T. and Michael D. (2003):**Urinary tract infection .e Medicine world Medical library from Web MD.
- Lowry ,S.J.(1975):**Diarrheal disease :Current concepts and future challenges .Enteroadherent *E.coli* :A heterogenous group of *E.coli* implication as diarrheal pathogens .*Trans .R. Soc. Trop. Med .Hyg.* 87:549-553.
- Lucas, E. H. and Lewis, R. W. (1944):** Antibacterial substances in organs of higher plants. *Science* 100: 597- 599.
- Lul, R. ;Gjyle M. ;Lule B. ;Lumturije B. ;Shyhreta O. and Linda P. (2004):**Etiology and susceptibility of urinary tract isolates in Kosova .*International Journal of Antimicrobial Agents* ,23(1):2-5.
- MacDonald, R. E. and Bishop, C. J. (1952):** A further survey of plant for antibacterial substances. *Can- Journal of Botany* 31: 123- 131.
- Mackey, J.P. and Sandys, J.H. (1966):**A trial of cystine- lactose electrolyte deficient medium for urinary bacteriology.*Br.Med.J.*;1:1173.
- Maruzzella, j. C. and Henry, P.A. (1958):** The in vitro antibacterial activity of essential oils and oil combinations. *J. Amer. Pharm. Assoc.*; 47: 294-296.
- Maruzzella, J. C. and Sicurella, N. A. (1960):** Antibacteial activity of essential oil vapors. *J. Amer. Pharm. Assoc.*; 49:692-694.

- Mazzulli ,T.(2001):**Susceptibility of community Gram-negative urinary tract isolates to mecillinam and other oral agents. *Can J Infect Dis*;12(5):289-292.
- Meyhoff ,H; Nordling ,J; Gammelgaard ,P and Vejlsgaard ,R (1981):**Dose antibacterial ointment applied to urethral meatus in women prevent recurrent cystitis ? .*Scand J Urol Nephrol* 15(2) :81-83. PMID 7036332.
- Miraqliotta G; Di pierro, M.N. ; Miraqliotta, L. ;Mosca, A.(2008):-**Antimicrobial resistance among uropathogens responsible for community acquired urinary tract infections in an Italian community.
- Monica, C. (1985):**Medical laboratory manual for tropical countries. ELBS. Ed. English Languag Book Society; tropical health. Technology. Buttrworth. Vol. II. Enteric Gram- negative rods.Chap.42:250-255.
- Munoz ,P. ;Lian C.A. and Rodriguez C. M. (1997):**Group B *Streptococcus* bacteremia in non pregnant adults .*Arch Intern Med*,157:213-216.
- National Institutes of Health (2005):**National Kidney and Urologic Diseases information clearing house (NKUDIC).Accessed on line at <http://Kidney.niddk.nih.gov/>.
- NCCLS(National Committee For Clinical Laboratory Standard) (2001):**Approved guideline, 2nd ed. Routine urinalysis and collection, transport and preservation of urine specimens. Document GP16-A2, Nov.2001.
- Nelson C. A.; Onyeagba, R. A.(2007):** Antimicrobial Properties Of Extracts Of *Allium cepa* (Onions) And *Zingiber officinale* (Ginger) On *Escherichia coli*, *Salmonella typhi* And *Bacillus subtilis* . The Internet Journal of Tropical Medicine. ; 3 (2). ISSN: 1540-2681

- Niamsa ,N. and Sitiwet ,C.(2009):** Antimicrobial Activity of *Curcuma longa* Aqueous Extract. Journal of Pharmacology and Toxicology. Volume (4) Issue(4) Pp.173-177.
- Nicolle ,L.E ;Harding ,G.K.M. ;Preiksaitis ,J. and Ronald ,A.R. (1982):**The association of urinary tract infection with sexual intercourse . J infect Dis,146:574-583.
- Nicolle ,L.E. ; Hoban, S.A. and Harding ,G.K.(1983):**Characterization of coagulase negative Staphylococci from urinary isolates. J. Clin. Microbiol. , Vol. 17 (2)P p. 267-271.
- Nicolle ,L.E(2001):**A practical guide to antimicrobial management of complicated urinary tract Infection. Drug Aging ,18(4)243-254.
- Nicolle, L. E.(2003):** Urinary Tract Infection . traditional pharmacologic therapies. Dis. Mon,49:111-128.
- Nunezsanchez ,J.C ;Catala, B.M.T. ;Navarro, O. I. and Balaquer M.J.V.(1999):**Tratamiento delas infecciones urinarias en pacientes con factors deriesgo .Med .Clin ,24:558-559.
- O'' Donnell ,J.A. and Gelone ,S.P. (2000):**The fluoroquinolones .Infect Dis. clin. North Am ,14:489-513.
- Odugbemi, TO. (2006) :**Outlines and pictures of medicinal plants from Nigeria , University of Lagos Press, Lagos, Nigeria. pp. 91.
- Okada ,K. ;Usui ,Y. and Abe ,T. (1994):**Statistics studies on bacteria isolated from urinary tract infections .Hinyokika Kiyo ,40:175-185.
- Orret ,F.A. and Shurland ,S.M.(1998):**The changing patients of antimicrobial susceptibility of urinary pathogens in Trinidad .Singapore Med .J ,39:256-259.
- Osman S.; Alper K.; Musa O. and Sami O.(2001):** Effects of Turkish spice extracts at various concentrations on the growth of *Escherichia coli* O157:H7 Food Microbiology Vol 19, Issue 5, Pages 473-480.

- Oyedemi , S. O.; Okoh ,A. I.; Mabinya ,L. V.; Pirochenva, G. and Afolayan, A. J. (2009):-** The proposed mechanism of bactericidal action of eugenol, α -terpineol and α -terpinene against *Listeria monocytogenes*, *Streptococcus pyogenes*, *Proteus vulgaris* and *Escherichia coli*. African Journal of Biotechnology Vol. 8 (7), pp. 1280–1286.
- Palac, D.M. (1986):**Urinary tract infections in women : a physician 's perspective .lab .Med. , pg 17:25.
- Park ,N.; Juven, B.J.; Shaaya, E.; Menasherov,M.; Nitzan, R.; Weiss, H and Ravid, V.(1984):** Inhibitory effect of oregano and thyme essential oils on moulds and food borne bacteria .Letters in Applied Microbiology; 11(1):33-37
- Park, J.; Min, K. and Kang, D.(2007):** The efficacy and safety of a once-daily extended- release ciprofloxacin tablet for the empirical treatment of symptomatic uncomplicated cystitis in Korean women. Korean J. Urol.; 48: 35-39.
- Patricia, S.(2006):** Mannitol salt agar .Department of cell biology and molecular genetics University of Maryland College Park, MD 20742 USA.
- Patton ,J .P. ;Nash ,D .B. and Abrutyn ,E. (1991):**Urinary tract infection :economic considerations .Med clin North Am ,75:495-513.
- Perfetto ,E.M. and Gondek ,K. (2002):***Escherichia coli* resistance in uncomplicated urinary tract infection: A model for determining when to change first –line empirical antibiotic choice .Manag Care Interface ,15:35-42.
- Platt ,R.; Polk ,B.F.; Murdock ,B. and Rosner ,B. (1982):**Mortality associated with nosocomial urinary tract infection .N Engl Med,307:637-642.

- Powers ,J.S.; Billings ,F.T. ;Behrend ,T.D. and Burger ,M.C. (1988):**Antecedent factors in urinary tract infections among nursing home patients .South Med J ,81(6), 734-735.
- Prescott ,M.L.; Harley, P.J. and Klein, A.D. (2008):** Microbiology. (7th Edition).Wm. C. Brown Publishers. U.S.A. pp 557-561.
- Priscila, I. U.; Mariama T. N.; Luiz C. D.; Luciano B. and Ary Fernandes J.(2007):** Antibacterial activity of medicinal plant extracts. Braz. J. Microbiol. ; 38 (4) : 717-719.
- Rebecca, B. (2005):**Blood agar plates and hemolysis: Streptococcus and other catalase negative Gram-positive cocci University of Utah, Department of Pathology, Salt Lake City, UT84132.USA.
- Robert, C.R.; Valkenburg, H.A.; Haverkon,M.J. and Gosling ,W.H. (1991):**Aprospective study of urinary tract infection in Dutch general practice. Lancet 2:674-677.
- Roller, S. and Seedhar, P. (2002):** Carvacrol and cinnamic acid inhibit microbial growth in fresh-cut melon and kiwifruit at 4° and 8°C. Letters in Applied Microbiology 35, 390-394.
- Ronald ,A. and Ludwig ,E. (2001):**Urinary tract infections in adults with diabetes .International Journal of Antimicrobial Agents ,17:287-292.
- Ronald ,A. (2002):**The etiology of urinary tract infection :traditional and emerging pathogens .AM J Med ;113:suppl (1A):145-195.
- Ronald, A.R. and Pattulo, M.S. (1991):**The natural history of urinary infection in adults. Med. Clin. Am., 75:299-312.
- Rosato A.; Vitali C.; De Laurentis, N.; Armenise, D., and Antonietta ,M.(2007):** Antibacterial effect of some essential oils administered alone or in combination with norfloxacin. Phytomedicine, 2007 Nov;14(11):727-732.

- Rosa, D. ;Jose, G. and Gonzalo, P. (2001):**Antibiotic susceptibility of bacterial strains isolated from patients with community –acquired urinary tract infections .International Journal of Antimicrobial Agents ,18(3):211-215.
- Rosemary, X.; Sorna, L. V. and Sivagama, S. M. (2008):**Antimicrobial activity of Selective native medicines. Indian Journal of Botanical Research, 4 (2): 213-222.
- Rula ,M. D. and Talal, A. (2010):** Effect of ethnomedicinal plants used in folklore medicine in Jordan as antibiotic resistant inhibitors on *Escherichia coli*. BMC Complementary and Alternative Medicine. 1472-6882.
- Saeed, S. and Tariq, P. (2007):** Antimicrobial activities of *Embllica officinalis* and *Coriandrum sativum* against Gram-positive bacteria and *Candida albicans*. Pak. J. Bot., 39(3): 913-917.
- Saeed, S.; Naim, A. and Tariq P. (2007):**A study on prevalence of multi-drug-resistant Gram negative bacteria. Int. J. Biol. Biotech., 4(1): 71-74.
- Saeed, S. and Tariq ,P. (2008):**In vitro antibacterial activity of clove against gram negative bacteria. Pak. J. Bot., 40(5): 2157-2160.
- Sahika, E. A. and Mehmet, K. (1986):**Sensitivity of some common food-poisoning bacteria essences against five important food borne pathogens. Letters in Applied Microbiology. 26(2): 118-122.
- Samy, R.P. (2005).** Antimicrobial activity of some medicinal plants from India. Fitoterapia., 76: 697-699.
- Savas, L.; Guvel, S.; Onlen, Y.;Savas, N. and Duran, N.(2006):** Nosocomial urinary tract infections : Micro-organisms, antibiotic sensitivities and risk factors. West Indian Med J., 55(3):188-193.

- Saxena ,D.;Amin ,M. and Khanna ,S. (1996):**Modulation of protein profiles in *Rhizobium* *sp.* Under salt stress.Can.J.Microbiol.Vol.42(6): 617–620
- Schappert ,S.M. (1997):**Ambulatory care visits to physician offices ,hospital outpatients departments ,and emergency departements :united states .vital and health statistics ,series 13-no 143.
- Scholes ,D.; Hooton ,T.M.; Roberts ,P.L. ;Stapleton ,A.E.; Gupta ,K. and Stamm ,W.E. (2000):**Risk factors for recurrent urinary tract infection in young women .J. Infect Dis ,182:1177-1182.
- Shaikh N.; Morone, N.E.; Bost, J.E. and Farrell, M.H.(2008):-**Prevalence of urinary tract infection in childhood: a meta-analysis. *Pediatr. Infect. Dis. J.*;27(4):302-308.
- Shirazi, M.H.; Ranjbar R.; Eshraghi S.; Amin G.; Seyed, N. M. and Bazzaz, N. (2008):** inhibitory effects of sage extract on the growth of enteric bacteria. *Pakistan J. Sci.* 11(3):487-489.
- Shymaa, A. (2004):**Identification and susceptibility testing of non-lactose fermenting Gram-negative bacilli in urinary tract infection. Master Degree, Zagazig University, Faculty of Medicine, Microbiology and Immunology Department.
- Sibanda ,T. and Okoh ,A. (2007):**The challenges of overcoming antibiotic resistance:Plant extracts as potential sources of antimicrobial and resistance modifying agents. *African J. of biotechnology* 6(25):2886-2896.
- Silverstein, R.M.; Bassler, G.C. and Morrill, T.C.(1991):**Spectrometric identification of organic compounds, 5th Ed., Wiley.
- Sinha, A. K.; Mehra, M. S.; Pathec, R. C. and Sinha, J. K. (1977):** Antibactrerial activity of 3 volatile oils from some indigenous plants. *Indian J. Exp. Boil.* 15: 339-340.

- Smith ,P.W. ;Seip ,C.W. ;Schaefer ,S.C. and Bell –Dixon ,C. (2000) :** Microbiologic survey of long –term care facilities .Am J Infect Control,28:8-13.
- Sobel ,J.D. ;Kaye ,D. ;Mandelle ,G.L. ; Bennett ,J.E. and Dolin ,R. (1995):**Urinary tract infections .In :Principles and practice of infectious diseases ,Fourth edition New York ,USA :Churchill livingstone ,662-690.
- Sobel ,J.D. and Kay ,D. (1984):**Host factors in the pathogenesis of urinary tract infections .AM J Med ,76:122-130.
- Sonnenwirth ,A .C. (1980):**Gradwohol's Clinical Laboratory Methods and Diagnosis Vol .2 ;8th ed. Sonnenwirth ;AC and Jarett L. (eds).The CV Mosby Co .London ;Tornto.
- Sourander ,L.B. (1966):**Urinary tract infection in the aged –an epidermiological study .Ann Med Intern fenn , 45:7-55.
- Sourander ,L.B. and Kaslanen ,A.(1972):-** A 5-year follow- up of bacteriuria in the aged. Cerontol Clin. ,14:274-281.
- Soussy C.J.; Lascols C. and Dib-Smahi, C.(2006):**Comparative in vitro activity of levofloxacin against *Escherichia coli* isolated from acute pyelonephritis in france in 2005.16th European Congress of Clinical Microbiolo. And Infectious Diseases Nice, France, April 1-4 .
- Stamm ,W.E and Hooton ,T.M. (1989):**Urinary tract infections :from pathogenesis to treatment .J. Infect .Dis. ,159:400-406.
- Stamm ,W.E. and Hooton ,T.M. (1993):** Mangement of urinary tract infections in adults .N. Engl. J. Med. ,329:1328-1334.
- Stamm, W. E. and Norrby, S. R. (2001):** Urinary tract infections: disease panorama and challenges. J. Infect. Dis. 183(1):51-54.

- Stapleton ,A (1999)** :Prevention of recurrent urinary tract infections in women .Lancet ,353:7-8.
- Stephan ,D. F. (2003)**:Acute uncomplicated urinary tract infection in women .N Engl. J. Med. ,349:259-266.No(3).
- Steven ,G. ; Carol, C. ; Karen, F. ;Van, H. ; Kathy, M. and Lauren ,Z. (1999)**:Urinary tract infection .University of Michigan health system.
- Strom, B.L; Collins ,M. ;West ,S.L. ;Kreisberg ,J. and Wellers (1987)**: Sexual activity ,contraceptive use ,and other risk factors for symptomatic and a symptomatic bacteriuria :a case –control study .Ann Inter Med, 107 :816-823.
- Sule, I.O. and Agbabiaka T.O. (2008)**: Antibacterial Effect of some Plant Extracts on Selected Enterobacteriaceae. Ethnobotanical Leaflets 12: 1035-1042.
- Susan ,A .;Mehnert –Kay,M.D. (2005)**:Diagnosis and management of uncomplicated urinary tract infections .American family physician ;72(3):451-456.
- Suslow, T.V.; Schroth, M.N. and Isaka, M. (1982)**:Application of a rapid method for Gram differentiation of plant pathogenic and saprophytic bacteria without staining. Phytopathology 72:917-918.
- Talan, M.D.(2010)**: National Ciprofloxacin Shortage: What to Do? E.M.N.(Emergency Medicine News) . Volume 32 - Issue 4 - p 10.
- Thomson, K.S.; Esanders ,W. and Sanders ,C.C (1994)**: A resistance patterns among UTI pathogens .J,Antimicrob Chemother ,33:9-15.
- Todar ,k.(2007)**:-Pathogenic *E.coli* .online text book of bacteriology. University of Wisconsin –Madison department of bacteriology. Retrived on 2007-11-30.

- Victor, N. ; Ma. Luisa, V. ; Gabriela, R. and Xavier L.(1996):** Antimicrobial evaluation of some plants used in Mexican traditional medicine for the treatment of infectious diseases. J. of Ethnopharmacology. Volume53, Issue 3, September 1996, Pages 143-147
- Vosti ,K .L. (2002):**Infections of urinary tract in women :a prospective longitudinal study of 235 women observed for 1-19 years .Medicine (Baltimore),81 :369-387.
- Vromen, M. ;Vanderven ,A.J. ;Knolls ,A. and Stobberingh ,E.E. (1999):**Antimicrobial resistance patterns in urinary isolates from nursing home resident .J Antimicrob chemother ,44:113-116.
- Vuddhakul, V.; Bhooponga, P.; Hayeebilana, F. and Subhadhirasakulb, S. (2007).** Inhibitory activity of Thai condiments on pandemic strain of *Vibrio parahaemolyticus*. Food Microbiol., 24: 413-418.
- Walsh, T.J. (1993):**Randomized double- blinded trial of rifampin with either novobiocin or trimethoprim- sulfamethoxazole against methicillin- resistant *Staphylococcus aureus* colonization: Prevention of antimicrobial resistance and effect of host factors on outcome. Antimicrobial agents and chemotherapy, Vol.37 No 6 p. 1334-1342.
- Walsh, S.E.; Maillard, J.Y.; Russell, A.D., Catrenich, C.E., Charbonneau, D.L. and Bartolo, R.G. (2003):** Activity and mechanisms of action of selected biocidal agents on Gram-positive and -negative bacteria. Journal of Applied Microbiology 94, 240-247.

- Warren ,J.W. (1996):**Clinical presentation and epidemiology of urinary tract infections. P.3-27. In H.L.T. Mobley and J.W. Warren (ed.),Urinary Tract infections. Molecular pathogenesis and clinical management. American Society of microbiology, Washington, D.C.
- Warren ,J.W. (2001):**Guidlines for antimicrobial treatment of uncomplicated acute bacterial cystitis and acute pyelonephritis in women .Infectious Diseases Society of America (IDSA).Clin Infect Dis ,33:296-299.
- Watt, K.; Christofi, N. and Young, R.(2007):**The detection of antibacterial actions of whole herb tinctures using luminescent *Escherichia coli*. Phytother Res. ;21(12):1193-9. PMID: 17661335.
- Well, B. and Stood, R.J. (1989):**The genus *shigella*. In Bergy's manual of systematic bacteriology. Vol. I. 8th ed. Krieg, N.R. Holt, J.G. and others (e d s)Wilkins . Hong Kong., pp: 423-427.
- Wikipedia(2007):**Urinary tract infection. http://en.wikipedia.org/wiki/urinary_tract_infection.
- Will L.; Dlawer ,A. and Tim Boswell(2005):** Medical Microbiology section F –Clinical manifestations of infection .F10,Urinary tract infection .P:313-318.Published by Taylor and Francis group.
- Winstanely ,T.G.; Limb ,D.T. ;Eggington ,R. and Hancock ,F. (1997):** A ten year survey of the antimicrobial susceptibility of urinary tract isolates in the UK :Microbe Base project .J.Antimicrob .chemother ,40:591-594.
- Wong ,E.S.(1983):**Guideline to prevention of catheter –associated urinary tract infections AM.J Infect control;11:28-36.

- Yamamoto, S.; Akiyama, K.; Yoshimoto, T. and Kanokogi, M. (2009):** Clinical efficacy of oral administration of 200 mg gatifloxacin once daily for 3 days for the treatment of patients with uncomplicated cystitis. *J. Infect. Chemother.*;15(2):104-107.
- Yang, Y.C.; Lee, S.H.; Lee, W.J.; Choi, D.H. and Ahn, Y.J. (2003):-** Ovicidal and adulticidal effects of *Eugenia cryophyllata* bud and leaf oil compounds on *Pediculus capitis*. *J. Agric. Food Chem.*,51(17): 4884-4888.
- Zahar, J.R.; Lortholary, O.; Martin, C.; Potel, G. and Plesiat, P. (2009):** Addressing the challenge of extended-spectrum beta-lactamases *Curr Opin Investig Drugs.*;10(2):172-80. PMID: 19197.
- Zaidan, M.R.; Noor, R. A.; Badrul, A.R.; Adlin, A.; Norazah, A. and Zakiah, I.(2005):** In vitro screening of five local medicinal plants for antibacterial activity using disc diffusion method. *Trop Biomed* ;22(2):165-70 PMID: 16883283.
- Zhanel, G.G.; Hisanaga T.L.; Lain, N.M.; Decorby, M.R. and Nichol K.A.(2006):**Antibiotic resistance in *Escherichia coli* outpatient urinary isolates: final results from the North American Urinary Tract Infection Collaborative Alliance (NAUTICA) . *J. Antimicrob. Chemother.* 27(6):468-475 . PMID: 16713191.