

References

- Abdel-Aziz, T.A.M.; Fouad, H.H.; Rashed, L.A.; Salama, H.A.S.; Abdel-Wahhab M.A. and Ahmed, T. (2005): Interferon- α gene therapy prevents aflatoxin and carbon tetrachloride promoted hepatic carcinogenesis in rats. *Int. J. Mol. Med.*, 15: 21–26.
- Abdel-Wahhab, M.A.; Nada, S.A.; Farag, I.M.; Abbas, N.F. and Amra, H.A. (1998): Potential protective effect of HSCAS and bentonite against dietary aflatoxicosis in rat: with special reference to chromosomal aberrations. *Nat. Toxins*, 6: 211–218.
- Abdel-Wahhab, M. A.; Nada, S. A. and Amra, H. A. (1999): Effect of alumino-silicates and bentonite on aflatoxin-induced developmental toxicity in the rats. *J Appl. Toxicol.*, (19: 199-204.
- Abdel-Wahhab, M.A.; Nada, S.A. and Khalil, F.A. (2002): Physiological and toxicological responses in rats fed aflatoxin-contaminated diet with or without sorbent materials. *Anim. Feed Sci. Technol.*, 10740: 1–11.
- Abdel-Wahhab, M. A. and Aly, S. E. (2003): Antioxidants and radical scavenging properties of vegetable extracts in rats fed aflatoxin-contaminated diet. *J Agric. Food Chem.*, 51(8):2409-14.
- Abdel-Wahhab, M. A. and Aly, S. E. (2005): Antioxidant property of *Nagilia Sativa* (Black cumin) and *Syzygium Aromaticum* (Clove) in rats during aflatoxicosis. *J. Appl. Toxicol.*, 25: 218-223.
- Abdel-Wahhab, M. A.; Abdel-Galil, M. M. and Hamed, N. F. (2005): The protective effects of hydrated sodium calcium alumino-silicate and the egyptian montmorillonite against aflatoxicosis in rats. *SC. J. AZ. Med. Fac. (Girls)*, 26(3): 243-258.
- Abdel-Wahhab, M.A.; Ahmed, H.H. and Hagazi, M.M. (2006): Prevention of aflatoxin B₁-initiated hepatotoxicity in rat by marine algae extracts. *J. Appl. Toxicol.*, 26: 229-238.

- Abdel-Wahhab, M.A.; Abdel-Galil, M.M.; Hassan, A.M.; Hassan, N.H.; Nada, S.A.; Saeed, A. and El-Sayed, M.M. (2007): *Zizyphus spina-christi* extract protects against aflatoxin B₁-initiated hepatic carcinogenicity. Afr. J. Trad. CAM., 4 (3): 248-256.
- Adams, R.P. (1995): Identification of essential oil components by gas chromatography mass/ spectroscopy. Allured publishing corporation, USA.
- Adams, R. Z. (2002): Indoor environment connections featured public library closes down for mold investigation
www.ieconnections.com/archive/jan_01/jan-01.htm
- Adamson, R. H.; Correa, P. and Dalgard, D.W. (1973): Brief communication: occurrence of primary liver carcinoma in rhesus monkey fed aflatoxin B.J Nath. Cancer Inst., 50: 549-553.
- Afzal, M.; Al-Hadidi, D.; Menon, M.; Pesek, J. and Dhimi, M.S. (2001): Ginger: an ethnomedical, chemical and pharmacological review. Drug Metab. Drug Interact., 18: 159–190.
- Agarawal, M.; Walia, S.; Dhingra, S. and Khambay, B.P. (2001): Insect growth inhibition, antifeedant and antifungal activity of compounds isolated/derived from *Zingiber officinale* Roscoe (ginger) rhizomes. Pest Manage. Sci., 57: 289–300.
- Ahmed, R.S.; Seth, V. and Banerjee, B.D. (2000): Influence of dietary ginger (*Zingiber officinales* Rosc) on antioxidant defense system in rat: comparison with ascorbic acid. Indian J Exp Biol., 38(6):604–6.
- Alam, M.M.; Javed, K. and Jafri, M.A. (2005): Effect of Rheum emodi (*Revand Hindi*) on renal functions in rats. J. Ethnopharmacol., 96: 121–125.
- Angus, J.F.; Gardner, P.A.; Kirkegaard, J.A. and Desmarchelier, J.M. (1994): Isothiocyanates released from brassica roots Inhibit growth of the take-All Fungus. Biofumig. Plant Soil, 162: 107-112.

- Ansar, S.; Iqbal, M. and Athar, M. (1999): Nordihydroguaiaretic acid is a potent inhibitor of ferric-nitritotriacetate-mediated hepatic and renal toxicity, and renal tumour promotion, in mice. *Carcinogenesis*, 20: 599–606.
- A.O.A.C. (2000): Official Methods of Analysis of AOAC International 17th ed., Nature Toxins. AOAC International, Arlington, Virginia, USA, chapter 49.
- Assmann, G.; Jabs, H.U.; Nolte, W. and Schriewer, H. (1984): LDL-cholesterol determination in blood serum following precipitation of LDL with polyvinyl sulfate. *Clinica. Chimica. Acta.*, 140:77-83.
- Atef A. H.; Tawakkol W.M. and Abdel Azeem E. (2010): The hepatoprotective effect of dimethyl 4,4- dimethoxy 5,6,5,6- dimethylene dioxy-biphenyl- dicarboxylate (D.D.B.) on aflatoxin B₁ induced liver injury. *Life Science J.*, 7 (3): 148.
- Athar, M. and Iqbal, M. (1998): Ferric nitritotriacetate promotes N-diethylnitrosamine-induced renal tumorigenesis in the rat: implications for the involvement of oxidative stress. *Carcinogenesis*, 19: 1133–1139.
- Bahk J. and Elmer H. M. (1983): Aflatoxin production is inhibited by selected herbal drugs. *Mycopatholo.*, (83) , 129-134.
- Barillari, J.; Canistro, D.; Paolini, M.; Ferroni, F.; Pedulli, G.F.; Iori, R. and Valgimigli, L. (2005): Direct antioxidant activity of purified glucoerucin, the dietary secondary metabolite contained in rocket (*Eruca sativa Mill.*) seeds and sprouts. *J. Agric. Food Chem.*, 53: 2475–2482.
- Bartles, H.; Bohmer, M. and Heirli, C. (1972): Serum creatinine determination without protein precipitation. *Clin. Chem. Acta.* , 37: 193-197.
- Bartley, J. and Jacobs, A. (2000): Effects of drying on flavour compounds in Australian-grown ginger (*Zingiber officinale*). *Journal of the Science of Food and Agriculture*, 80, 209–215.

- Barton, C. C.; Ganey, P. E. and Roth, R. A. (2000): Lipopolysaccharide augments aflatoxin B₁-induced liver injury through neutrophil-dependent and -independent mechanisms. *Toxicol. Sci.*, 58: 208–215.
- Bennett, J. W. and Klich, M. (2003): Mycotoxins. *Clin. Microbiol. Rev.*, 16: 497 – 516.
- Bergmeyer, H.U. (1978): Principles of enzymatic analysis. Verlag Chemie.
- Betina, V. (1989): Biological effects of mycotoxins. In Betina, V. (Ed.). *Mycotoxins chemical, biological and environmental aspects*. Amsterdam 42-58.
- Bhatnagar, D.; Yu, J. and Ehrlich, K.C. (2002): Toxins of filamentous fungi. In: *Fungal Allergy and Pathogenicity*. Breitenbach M, Cramer R, Lehrer S (Eds.) *Chem. Immunol.*, Vol 81, Basel, Karger, 167-206.
- Bhatnagar, D. and Mc-Cormick, S. P. (1988): The inhibitory effect of neem (*Azadirachta indica*) leaf extracts on aflatoxin synthesis in *Aspergillus parasiticus*. *J. Am. Oil Chem. Soci.*, 65: 1166–1168.
- Blois, M.S. (1958): Antioxidant determinations by the use of a stable free radical. *Nature*, 181: 1199–1200.
- Boer J. G.; Quiney B.; Walter P. B.; Thomas C.; Hodgson K.; Murch S. J. and Saxena P. K. (2005): Protection against aflatoxin-B₁-induced liver mutagenesis by *Scutellaria baicalensis*. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 578: 15-22.
- Buescher, R.; and Yang, L. (2000): Natural Food Colorants. Turmeric. In: Lauro, G.J.; Francis, F.J. (Eds.), *Science and Technology*. Marcel Dekker, NY, 205–226.
- Bull B.S.; D’Onofrio G. and Fujimoto K. (2001): Recommendation for reference method for the packed cell volume (ICSH Standard Lab Hematol., 7:148-170.
- Cai, Y.Z.; Sun, M. and Corke, H. (2003): Antioxidant activity of betalains from plants of the amaranthaceae. *J. of Agri. and Food Chem.*, 51 (8): 2288-2294.
- Cai, Y.; Luo, Q.; Sun, M. and Corke, H. (2004): Antioxidant activity and phenolic

- compounds of 112 traditional Chinese medicinal plants associated with anticancer, Life Sciences., 74: 2157–2184.
- Calabrese, V.; Butterfield, D.A.; and Stella, A. M. (2003): Nutritional antioxidants and the heme oxygenase pathway of stress tolerance: novel targets for neuroprotection in Alzheimer's disease. Ital J Biochem., 52(4):177-81.
- Carolina, A.; Manzan, C.M.; Toniolo, F.S.; Bredow, E. and Povh, N.P. (2003): Extraction of essential oil and pigments from *Curcuma longa* [L.] by steam distillation and extraction with volatile solvents. J. Agric. Food Chem., 51: 6802–6807.
- Casado, J. M.; Theumer, M.; Masih, D. T.; Chulze, S. and Rubenstein, H. R. (2001): Experimental subchronic mycotoxicoses in mice: individual and combined effects of dietary exposure to fumonisins and aflatoxin B₁. Food and Chem. Toxicol., 39: 579-586.
- CAST; (1989): Mycotoxins: Economic and Health Risks. Council for Agricultural Science and Technology, Task Force Report No. 116, Ames, IA.
- CAST; (2003): Mycotoxins: Risks in Plant, Animal and Human Systems. Council for Agricultural Science and Technology Task Force Report No. 139, Ames, IA.
- Chang, P.K.; Cary, J.W.; Yu, J.; Bhatnagar, D. and Cleveland, T.E. (1995): *Aspergillus parasiticus* polyketide synthetase gene, pksA, a homolog of *Aspergillus nidulans*, required for aflatoxin B₁. Mol. Gen. Genet., 248:270–277.
- Chen, C.H.; Kuo, M.; Wu, C.H. and Ho, C.H. (1986): Pungent compounds of ginger (*Zingiber officinale* (L) Rosc) extracted by liquid carbon dioxide. J. of Agri. and Food Chem., 34: 477–480.
- Citil, M.; Gunes, V.; Atakisi, O.; Ozcan, A.; Tuzcu, M. and Dogan, A. (2005): Protective effect of Lcarnitine against oxidative damage caused by experimental chronic aflatoxicosis in quail *Coturnix coturnix*). Acta Vet. Hung., 53(3):319-324.

- Clark, J. D.; Hatch, R. C.; Miller, D. M. and Jain; A. V. (1984): Caprine aflatoxicosis: Experimental disease and clinical pathologic changes. J. Dairy Sci., 88:3063–3069.
- Conner, D. E. (1993): Naturally occurring compounds, In Antimicrobial in Foods, pp.441- 468. Edited by Davidson, P. M. and Brannen, A. L. Marcel Dekker Inc., New York.
- Coulombe, R.A. (1991): Aflatoxin. In: Sharma, R.P. & Salunkhe, D.K. (Eds.), Mycotoxins and Phytoalexins. London: CRC Press, 103-144.
- Creppy; (2002): Update of survey, regulation and toxic effect of mycotoxins in Europe, Toxicol., Review 127, 19–28.
- Cullen, J.M. and Newberne, P.M. (1994): Acute hepatotoxicity of aflatoxins. In: The toxicology of aflatoxins. Eaton D.L, and Groopman J.D (Eds.). New York, Academic Press, 3-26.
- Das, K.C. and Das, C.K. (2002): Curcumin (diferuloylmethane), a singlet oxygen (1O_2) quencher. Biochem. Biophys. Res. Commun., 295: 62–66.
- Depkevicius, A.; Van Beek, T.A.; Lelyveld, G.P.; Van Veldhuizen, A.; de Groot, A.; Linssen, J.P.H. and Venskutonis, R. (2002): Isolation and structural elucidation of radical scavengers from *Thymus vulgaris* leaves. J. Nat. Prod., 65: 892-896.
- Drabkin, D.L. and Austin, J.H. (1935): Spectrophotometric studies. II. Preparations from washed blood cells; nitric oxide hemoglobin and sulfhemoglobin. J. Biol. Chem., 112, 51.
- Duke, J. (2004): USDA, ARS, National Genetic Resources Program. Phytochemical and Ethnobotanical Databases.[Online Database] National Germplasm Resources Laboratory, Beltsville, Maryland,
- Dutton, M.F. and Anderson, M.S. (1980): Inhibition of aflatoxin biosynthesis by organophosphorus compounds. J. Food Prot., (43): 381-384.

- Dwivedi, C.; John, L.M.; Schmidt, D.S. and Engineer, F.N. (1998): Effects of oil- soluble organosulfur compounds from garlic on doxorubicin-induced lipid peroxidation. *Anti Cancer Drugs*, 9 (3): 291-294.
- Eaton, L. and Gallagher, E.P. (1994): Mechanism of aflatoxin Carcinogenesis. *Annu. Rev. Pharmacol. Toxicol.*, 34:135-72.
- Eaton, D.L. and Groopman, J.D. (1994): The toxicology of aflatoxins-human health. Veterinary and agricultural significance. New York, Academic Press.
- Eaton, D. L.: Ramsdell, H. S. and Neal, G. E. (1994): Biotransformation of Aflatoxins. In: *The Toxicology of Aflatoxins: Human Health, Veterinary and Agricultural Significance* (Eaton D. L., Groopman J., eds), 4571. New York: Academic Press.
- Edrington, T.S.; Harvey, R.B. and Rottinghaus. G.E. (1997): Environmental health influence of α super activated charcoal of the toxic effects of aflatoxins or T₂ toxins in broiler chickens. *Poultry Sci.*, 76: 1205 -1211.
- Edrington, T.S.; Kamps-Holtzapple, C.A.; Harvey, R.B.; Kubena, L.F.; Elissalde, M.H. and Rottinghaus, G.E. (1995): Acute hepatic and renal toxicity in lambs dosed with fumonisin containing culture materials. *J. Anim. Sci.*, 73: 508–515.
- (EHC): Environmental Health Criteria, (1979): Environmental Health Criteria 11. Mycotoxins. Geneva.
- El-Agamy, D. S. (2010): Comparative effects of curcumin and resveratrol on aflatoxin B₁ induced liver injury in rats. *rch Toxicol.*, 84(5):389-396.
- El-Ghorab, A.H.; El-Massry, K.F. and Shaaban, H.A. (2007): Effect of drying on the chemical composition of the Egyptian *Cupressus macrocarpa* (Hartw.ex. Gordon) essential oils and their biological characteristics. *J. Essential Oil Bearing Plants.*, 10: 399 – 411.
- Eraslan, G.; Karaoz, E.; Bilgili, A.; Akdogan, M.; Oncu, M. and Essiz, D. (2003): The effects of aflatoxin on kidney function in broiler chicks. *Turk J Vet Anim Sci* 27: 741-749.

- Fejes, S.; Kery, A.; Blazovics, A.; Lugasi, A.; Lemberkovics, E. and Petri, G. (1998): Investigation of the in vitro antioxidant effect of *Petroselinum crispum* (Mill.) Nym. ex A. W. Hill. Acta Pharmac. Hung., 68(3): 150–156.
- Ficker, C.E.; Arnason, J.T.; Vindas, P.S.; Alvarez, L.P.; Akpagana, K.G.M.; DeSouza, C. and Smith, M.L. (2003): Inhibition of human pathogenic fungi by ethnobotanically selected plant extracts. Mycoses, 46: 29-37.
- Finley, P.R. (1978): Clin. Chem., 24: 931-9347.
- Flegg, H.M. (1973): An investigation of determination of serum cholesterol by an enzymatic method. Ann. Clin. Biochem., 10: 79-84.
- Food and Agricultural Organization (FAO), (1997): Worldwide regulations for mycotoxins 1995. FAO Food and Nutrition Paper 64, Rome.
- Fossati P. and Prencipe L. (1980): Clin. Chem., 26: 227.
- Fossati, P. and Prencipe, L. (1982): Serum triglyceride determined colorimetrically with an enzyme that produces hydrogen peroxide. Clin. Chemi., 28: 2077-2088.
- Galvano, F.; Piva, A.; Ritieni, T. and Galvano, G. (2001): Dietary strategies to counteract the effects of mycotoxins: A review .J. Food Prot., 64 (1): 120-131.
- Gamliel, A. and Stapleton, J.J. (1993): Characterization of antifungal volatile compounds evolved from solarized soil amended with cabbage residues. Phytopathology., 83: 899-905.
- Gary, G. A. and Elmer, H. M. (1976): Inhibition of growth and aflatoxin production of *Aspergillus parasiticus* by citrus oils Zeitsch. für Leben. Smitt. Forsch. A, 160, (4): 353-358.
- Gazzani, G. (1994): Anti and prooxidant activity of some dietary vegetables. Rivi. Sci. Aliment., 23(3): 413–420.
- Giri, U.; Iqbal, M. and Athar, M. (1999): Potassium bromate (KBrO₃) induces renal proliferative response and damage by elaborating oxidative stress. Cancer Lett.,

135: 181–188.

- Gonzalez, E.; Felicio, J.D. and Pinto, M.M. (2001): Biflavonoids inhibit the production of aflatoxin by *Aspergillus flavus*. Braz. J. Med. Biol. Res., (34) : 1453-1456.
- Groopman J.D. and Sabbioni G. (1991): Detection of aflatoxin and its metabolites in human biological fluids. Bray, G.A. and Ryan, DH (Eds.) Pennington Center Nutrition Series, Vol. 1. Mycotoxins, cancer and health. Baton Rouge, Louisiana State University Press, 18-31.
- Guenther, E. (1961): The essential oils. D. Van Nostrand Company, New York, London, Vol I,II.
- Guerre, P.; Larrieu, G.; Burgat, V. and Galtier, P. (1999): Cytochrome P450 decreases are correlated to increased microsomal oxidative damage in rabbit liver and primary cultures of rabbit hepatocytes exposed to AFB1. Toxicol Lett., 104(1-2): 117-125.
- Halliwell, B. and Gutteridge, J. M. C. (1998): Free Radicals in Biology and Medicine; Oxford University Press: Oxford.
- Halliwell, B.; Zhao, K. and Whiteman, M. (1999): Nitric oxide and peroxynitrite. The ugly, the uglier and the not so good: a personal view of recent controversies. Free Radic. Res., 31: 651–669.
- Harrigan, W.F. and Nargaret, E.M. (1966): Grain storage studies. The fungus flora of stored wheat. Cereal Chem., 29: 450-462.
- Harvey, R. B.; Huff, W. E.; Kubena, L. F.; Corner, D. E. and Phillips, T. D. (1988): Progression of aflatoxicosis in growing barrows. Am. J. Vet. Res., 49:482.
- Harvey, R.B.; Phillips, T.D.; Ellis, J.A.; Kubena, L.F.; Huff, W.E. and Peterson, H.D. (1991): Effect of aflatoxin M₁ residues in milk by addition of hydrated sodium calcium aluminosilicate to aflatoxin contaminated diets of dairy cows. Am. J.Vet. Res., 52: 1556-1559.
- Helferich, W. G.; Garrett, W. N.; Hsieh, D.P.H. and Baldwin, R. L. (1986): Feed lot performance and tissue residues of cattle consuming diets containing aflatoxins. J.

- him . Sci., 62:691.
- Hempel, J.; Pforte, H.; Raab, B.; Engst, W.; Bohm, H. and Jacobasch, G. (1999): Flavonols and flavones of parsley cell suspension culture change the antioxidative capacity of plasma in rats. *Nahrung*., 43: 201-204.
- Herrmann, K. (1994): Antioxidativ wirksame Pflanzenphenole sowie Carotinoide als wichtige Inhaltsstoffe von Gewürzen. *Gordian*, 94: 113–117.
- Hong, C.H.; Hur, S.K.; Oh, O.; Kim, S.S.; Nam, K.A.; and Lee, S.K. (2002): Evaluation of natural products on inhibition of inducible cyclooxygenase (COX-2) and nitric oxide synthase (iNOS) in cultured mouse macrophage cells. *J. Ethnopharm.*, 83: 153–159.
- Huang, S. W. and Frankel, E. N. (1997): Antioxidant activity of tea catechins in different lipid systems. *J. Agric. Food Chem.*., 5: 3033–3038.
- IARC; (1993): Some naturally occurring substances: Food items and constituents. IARC monographs on Evaluation of carcinogenic risk to humans No. 56.
- IFCC: (the International Federation of Clinical Chemistry) (1977): Section no. 3: Enzymes, III. Revised IFCC Method for aspartate aminotransferase *J. Clin. Chem. Clin. Biochem.*, 15: 719-720.
- Ippoushi, K.; Azuma, K.; Ito, H.; Horie, H. and Higashio, H. (2003): [6]-Gingerol inhibits nitric oxide synthesis in activated J774.1 mouse macrophages and prevents peroxynitrite-induced oxidation and nitration reactions. *Life Sci.*, 73:3427–37.
- Jain, S.K. (1994): Ethnobotany and research in medicinal plants in India. *Ethnobot. Search, New Drugs*., 185: 153 - 168.
- Jeyakumar, S.M.; Nalini, N. and Menon, V.P. (1999): Antioxidant activity of ginger (*Zingiber officinale* Rosc) in rats fed a high fat diet. *Med. Sci. Res.*, 27: 341-344.
- Jian, Y.T.; Mai, G.F.; Wang, J.D.; Zhang, Y.L.; Luo, R.C. and Fang, YX. (2005): Preventive and therapeutic effects of NF-kB inhibitor curcumin in rats colitis induced by trinitrobenzene sulfonic acid. *World J. of Gastro.*, 11:1747–52.

- Jodynis-Liebert, J.; Matlawska, I.; Bylka, W. and Murias, M. (2006); Protective effect of *Aquilegia vulgaris* L on aflatoxin B₁ -induced hepatic damage in rats , Environ. Toxicol. Pharmacol., 22(1): 58-63.
- Joon-Kyoung, L.; Eun, H. C.; Kwang-Geun, L. and Hyang, S. C. (2005): Alleviation of aflatoxin B₁-induced oxidative stress in HepG2 cells by volatile extract from *Allii Fistulosi* Bulbus. Life Sciences 77, (23): 2896-2910
- Juglal, S.; Govinden, R. and Odhav, B. (2002): Spice oils for the control of co-occurring mycotoxin producing fungi. Food Prot., 65(4):683-687.
- Jung, H.A.; Park, J.C.; Chung, H.Y.; Kim, J. and Choi, J.S. (1999): Antioxidant flavonoids and chlorogenic acid from the leaves of *Eriobotrya japonica*. Arch. Pharm. Res., 22(2): 213-218.
- Justesen, U. and Knuthsen, P. (2001): Composition of flavonoids in fresh herbs and calculation of flavonoid intake by use of herbs in traditional Danish dishes. Food Chem., 73: 245–250.
- Kalembe. D. and Kunicka, A. (2003): Antibacterial and antifungal properties of essential oils. Current Medicin. Chem., 10: 813–829.
- Kaneko, J.J. (1989): Clinical chemistry of domestic animals , 4th edition, Academic press, San Diego, CA.
- Kaplan, M.M. (1987): Laboratory Tests. In: Diseases of the Liver. Eds. Schiff, L. & Schiff, E.R., J.B. Lippincott, Philadelphia, USA. p. 219.
- Kapoor, LD. :(1990) CRC Handbook of Ayurvedic Medicinal Plants. CRC Press, Boca Raton, 149-150.
- Kiran, M.M.; Demet, O.; Ortatatli, M., and Oguz, H. (1998): The preventive effect of polyvinyl–polypyrrolidone on aflatoxicosis in broilers. Avian Pathology, 27:250–255.
- Klein, B.; Read, P.A. and Babson, L.A. (1960): Alkaline phosphatase activity measurement. Clin. Chem., 6: 269-275.

- Klein, P. J.; Van Vleet T. R.; Hall J. O. and Coulumbe R. A. (2003): Effects of dietary butylated hydroxytoluene on aflatoxin B₁-relevant metabolic enzymes in turkeys. *Food Chem. Toxicol.* 41: 671 – 678.
- Klich, M.A.; Mullaney, E.J.; Daly, C.B. and Cary, J.W. (2000): Molecular and physiological aspects of aflatoxin and sterigmatocystin biosynthesis by *Aspergillus tamarii* and *A. ochraceoroseus*. *Appl. Microbiol. Biotechnol.*, 53: 605-609.
- Kolb JP. (2000); Mechanisms involved in the pro- and antiapoptotic role of NO in human leukemia. *Leukemia*, 14:1685–1694.
- Krishnamachari, K. A. V. R.; Bhat, R. V.; Nagarajan, V. and Tilnak, T. M. G. (1975): Hepatitis due to aflatoxicosis. An outbreak in Western India. *Lancet* 1, 1061–1063.
- Kuhn, D. M. and Ghannoum, M. A. (2003): Indoor mold, toxigenic fungi, and *Stachybotrys chartarum*: Infectious disease perspective. *Clinical Microbiol. Reviews*, 144: 150-151.
- Kusak, V. (1977): International programme on chemical safety. Institute of Experimental Medicine, Czechoslovak Academy of Sciences, Prague, Czechoslovakia (Vice-Chairman).
- Larsson, P. and Tjalve, H. (1995): Extrahepatic bioactivation of aflatoxin B₁ in fetal, infant and adult rats. *Chem.-Biol. Interact.*, 94: 1–19.
- Lee S.H.; Chang K.S.; Su M.S.; Huang Y.S., and Jang H.D. (2007): Effects of some Chinese medicinal plant extracts on five different fungi. *J. Food Control* doi:10.1016/j.foodcont.2006.12.005.
- Lindemann, M. D.; Blodgett, D. J.; Kornegay, E. T. and Schurig, G. G. (1993): Potential ameliorators of aflatoxicosis in weanling/ growing swine. *J. Anim. Sci.*; 71:171.
- Liu, J.R.; Chen, M.J. and Lin, C.W. (2005): Antimutagenic and antioxidant properties of milk–kefir and soymilk–kefir. *J. Agric. Food Chem.*, 53: 2467-2474.
- Llewellyn, G. C.; Burkett, M. L. and Eadic, T. (1981): Potential mold growth, aflatoxin

- production, and antimycotic activity of selected natural spices and herbs. J. Assoc. Off. Anal. Chem., 64: 995-960.
- Luchese R.H.; and Harrigan W.F. (1993): Biosynthesis of aflatoxin - the role of nutritional factors. J. Appl. Bacteriol., 74: 5-14.
- Mabrouk, S. S. and El-Shayeb, N. M. A. (1980): Inhibition of aflatoxin formation by some spices. Z. Lebensm. Unters. Forsch., 171: 344-347.
- Mari, M.; Iori, R.; Leoni, O. and Marchi, A. (1993): In vitro activity of glucosinolate-derived isothiocyanates against post harvest fruit pathogens. Ann. Appl. Biol., 123: 155–164.
- Martínez-Tomé, M., García-Carmona, F., and Antonia, M. M. (2001): Comparison of the antioxidant and pro-oxidant activities of broccoli amino acids with those of common food additives. J. of the Sci. of Food and Agri., 81: 1019–1026.
- Matasyoh, J.C.; Maiyo, Z.C.; Ngure, R.M. and Chepkorir, R. (2009): Chemical composition and antimicrobial activity of the essential oil of *Coriandrum sativum*. J. Food Chem., 113:526-529.
- Matthaus B. (2002): Antioxidant activity of extracts obtained from residues of different oilseeds. J. Agric. Food Chem., 50: 3444-3452.
- Meki A. R.; Sary, Kh. A. and El-Gibaly, I. (2001): Aflatoxin B₁ induces apoptosis in rat liver protective effect of melatonin. Neuroendocrinology, 22:417–426.
- Meki A. R.; Esmail E. D.; Hussein A. A. and Hassanein. H. M. (2004): Caspase -3 and heat shock protein–70 in rat liver treated with aflatoxin B₁: effect of melatonin. Toxicon.m., 43: 93-100.
- Menut, C.; Lamaty, G.; Bessiire, J. M. and Koudou, J. (1994): Aromatic plants of tropical central Africa.. Rhizomes volatile components of two zingiberales from the Central African Republic. J. Essent. Oil Res., 6: 161-164.
- Microbial Diagnostics, Inc. (2002): Germology.
<http://www.germology.com/mycotoxins.htm>.

- Miliauskasa, G.: Venskutonis, P. R. and van Beek, T. A. (2002): Screening of radical scavenging activity of some medicinal and aromatic plant extracts *Nahrung/ food* (46) 5: 317-320.
- Miliauskas, G.; Venskutonis, P. R. and Beek, T. A. (2004): Screening of radical scavenging activity of some medical and aromatic plant extracts. *Food Chem.*, 85: 231–237.
- Miller, D.M. and Willson, D.M. (1994): Veterinary diseases related to aflatoxins. In *The Toxicology of Aflatoxins*; Eaton, L. D., Groopman, J. D., Eds.; Academic Press: San Diego, CA., 347-364.
- Montgomery, H.A.C.; and Dymock, J.F. (1961): The determination of nitrite in water. *Analyst.*, 86: 414–416.
- Moreno, O. J. and Kang, M. S. (1999): Aflatoxins in maize: The problem and genetic solutions. *Plant Breeding*, 118: 1-16.
- Moss, M. O. (1998): Recent studies of mycotoxins. *J. of Appl. Microbiology Symposium Supplement*. 84: 62S-76S.
- Moure, A.; Cruz, J. M.; Franco, D.; Domínguez, J. M.; Sineiro, J. and Dominguez, H. (2001): Natural antioxidants from residual sources. *Food Chem.*, 72: 145–171.
- Muro-Cacho, C. A.; Stedeford, T.; Banasik, M.; Suchecki, T. T.; and Persad. A. S. (2004): Mycotoxins: mechanisms of toxicity and methods of detection for identifying exposed individuals. *J. Land Use*, 19: 537-549.
- Nielsen, P.V. and Rios, R. (2000): Inhibition of fungal growth on bread by volatile components from spices and herbs and the possible application in active packaging, with special emphases on mustard essential oil. *Inter. J. Food microbial.*, 60: 219-229.
- Nielsen S.E.; Young J.F.; Daneshvar B.; Lauridsen S.T.; Knuthsen P.; Sandstrom B. and Dragsted Lo. (1999): Effect of parsley (*Petroselinum crispum*) intake on urinary apigenin excretion, blood antioxidant enzymes and biomarkers for oxidative stress in human subjects. *Br J Nutr.*, 81: 447-455.

- Nyandieka, H.S.; Wakhis, J. and Kilonzo, M.M. (1990): Association of reduction of AFB₁-induced liver tumours by antioxidants with increased activity of microsomal enzymes. *Indian J. Med. Res.*, 92: 332-336.
- Ohkawa, H.; Ohishi, N. and Yagi, K. (1979): Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analyti. Biochem.*, 95: 351-358.
- Oke, F.; Aslim, B.; Ozturk, S. and Altundag, S. (2009): Essential oil composition, antimicrobial and antioxidant activities of *Satureja cuneifolia* Ten. *Food Chem.*, 112: 874-879.
- Ozer, J.; Ratner, M.; Shaw, M.; Bailey, W. and Schomaker, S. (2008): The current state of serum biomarkers of hepatotoxicity. *Toxicol.*, 245:194-205.
- Papp, E.; H-Otta, K.; Záray, G. and Mincsovcis, E. (2002): Liquid chromatographic determination of aflatoxins. *Microchem. J.*, 73: 39-46.
- Pedraza-Chaverri, J.; Maldonado, P.D.; Barrera, D.; Ceron, A.; Medina-Campos, O.N. and Hernandez-Pando, R. (2003): Protective effect of diallyl sulfide on oxidative stress and nephrotoxicity induced by gentamicin in rats. *Mol. Cell. Biochem.*, 254: 125–130.
- Percy, C.; Pat, B.; Poronnik, P. and Gobe, G. (2005): Role of oxidative stress in age-associated chronic kidney pathologies. *Adv. Chronic Kidney Dis.* 12: 78–83.
- Peterson, S.W.; Ito, Y.; Horn, B.W. and Goto, T. (2001): *Aspergillus bombycis*, a new aflatoxigenic species and genetic variation in its sibling species, *A. nomius*. *Mycologia.*, 93: 689-703.
- Petropoulos, S. A.; Daferera, D.; Akoumianakis, C. A.; Passam, H. C. and Polissiou, M. G. (2004): The effect of sowing date and growth stage on the essential oil composition of three types of parsley *Petroselinum crispum*. *J. of Sci. Food Agri.*, 84: 1606–1610.
- Phillips, D. J. (1980): Aflatoxins in Almonds, USDA, Science and Education Administration, Agricultural Reviews and Manuals.

- Piper, J.T.; Singhal, S.S.; Salameh, S.M.; Torman, R.T.; Awasthi, Y.C. and Awasthi, S. (1998): Mechanisms of anticarcinogenic properties of curcumin: the effect of curcumin on glutathione linked detoxification enzymes in rat liver. *Int. J. Biochem. Cell Biol.*, 30: 445–456.
- Pitt, J.I. (1996): What are mycotoxins, *Australian Mycotoxin Newsletter*, 7:1.
- Raccach, M. (1984): The antimicrobial activity of phenolic antioxidants in foods: a review. *J. of Food Safety*, 6, 141–170.
- Rahman, A.; Ahmed, S.; Vasenwala, S.M. and Athar, M. (2003): Glyceryl trinitrate, a nitric oxide donor, abrogates ferric nitrilotriacetate-induced oxidative stress and renal damage. *Arch. Biochem. Biophys.*, 418: 71–79.
- Ramsdell, H.S. and Eton, D.L. (1988): Modification of aflatoxin B₁ biotransformation in vitro and DNA binding in vivo by dietary broccoli in rats. *J. Toxicol. Environ. Health.*, 25: 269-284.
- Rasooli I. and Abyaneh M. R. (2004): Inhibitory effects of Thyme oils on growth and aflatoxin production by *Aspergillus parasiticus* *Food Control*, 15: 479-483.
- Recknagel, R. O.; Glende, E. A.; Dolak, J. A. and Waller, R. L. (1989): Mechanisms of carbon tetrachloride toxicity. *Pharmacology & Therapeutics*, 43: 139-154.
- Reddy R.V.; Taylor M.J. and Sharma R.P. (1987): Studies of immune function of CD-1 mice exposed to aflatoxin B₁. *Toxicology*, 43, (2):123 -132.
- Reddy, S. V. and Waliyar, F. (2000): Properties of aflatoxin and it producing fungi. URL: <http://www.aflatoxin.info/aflatoxin.asp>.
- Roebuck, M.; Yi-Liang L.; Adrianne E. R.; Groopman, J. D. and Kensler T. W. (1991): Protection against Aflatoxin Bi-induced Hepatocarcinogenesis in F344 Rats by 5-(2-Pyrazinyl)-4-methyl-1,2-dithiole-3-thione (Oltipraz): Predictive Role for Short-Term Molecular Dosimetry B. D. *Cancer Res.*, 51:5501-5506.

- Saad, M. M. M. and Abdel-Fattah, Sh.M. (2008): A food additive formula to minimize the negative effects due to ingesting aflatoxin(s) contaminated food. J. of Saudi Society for Food and Nutrition, 3 (1): 17-31.
- Sabry B.A. (2004): Biological studies on telomerase and other parameters in bladder cancer tissues. M.Sc. Thesis Faculty of Science Helwan univ. (5):139.
- Salbe, A.D. and Bjeldanes, F. (1989): Effect of diet on route of administration on the DNA binding of aflatoxin B₁ in the rat. Carcinogenesis, 10: 629-634.
- Salh, B.; Assi, K. and Templeman, V. (2003): Curcumin attenuates DNB-induced murine colitis, Ameri. J. of Physiol. Gastro. and Liver Physiol., 285: G235–43.
- Sanchez E.; Heredia N. and Garcia S. (2005): Inhibition of growth and mycotoxin production of *Aspergillus flavus* and *Aspergillus parasiticus* by extracts of *Agave species* Int. J. of Food Microbiol., 98 :271– 279.
- Sarwar, M. A.; Kaur, G.; Jabbar, Z.; Javed, K. and Athar, M. (2007): *Eruca sativa* seeds possess antioxidant activity and exert a protective effect on mercuric chloride induced renal toxicity. Food and Chem. Toxicol., (45): 910–920.
- SAS Institute, Inc. (1982): SAS User's Guide: statistics. SAS Institute, Cary, N.C.
- Selvi A. T.; Joseph G.S. and Jayaprakasha G.K. (2003): Inhibition of growth and aflatoxin production in *Aspergillus flavus* by *Garcinia indica* extract and its antioxidant activity Food Microbiol., 20 455–460.
- Schieffer, G.W. (2002): Pressurized liquid extraction of curcuminoids and curcuminoid degradation products from turmeric (*Curcuma longa*) with subsequent HPLC assays. J. Liq. Chromatogr. Related Technol., 25: 3033–3044.
- Schinella, G.R.; Tournier, H.A.; Prieto, J.M.; Mordujovich de Buschiazso, P.; and Rios, J.L. (2002): Antioxidant activity of anti-inflammatory plant extracts. Life Sci., 70: 1023–1033.
- Sell, S.; Xu, L.U. and Huff, W.E. (1998): Aflatoxin exposure produces serum alpha

- fetoprotein elevation and marked oval cell proliferation in young male pekin ducklings. J.of Pathol., 30: 34-39.
- Selvam, R.; Subramanian, L.; Gayathri, R. and Angayarkanni, N. (1995): The anti-oxidant activity of turmeric (*Curcuma longa*). J. Ethnopharm., 47: 59–67.
- Shahidi, F. (2008): Antioxidant factors in plant foods and selected oilseeds. BioFactors, (13), 1-4: 179–185.
- Shen H. M.; Ong. C. N. and Shi, C. Y. (1995): Involvement of reactive oxygen species in aflatoxin B₁-induced cell injury in cultured rat hepatocytes. Toxicol., 99:115-123.
- Shen, H. M.; Shi, C. Y.; Lee, H. P and Ong, C. N. (1994): Aflatoxin B₁ induced lipid peroxidation in rat liver. Toxicol. Appl. Pharmacol., 127:145-150.
- Shreya, R. V.; Rahul, B. P. and Meena, M P. (2009): Protective effect of *petroselinum crispum* (Mill) Nyman ex. A.W. Hill leaf extract on D-galactose-induced oxidative stress in mouse brain. Ind. J. of exper. biol., 47: 338-342.
- Shyamal, S.; Latha, P G.; Suja, S R.; Shine, V J.; Anuja, G I.; Sini, S.; Pradeep, S.; Shikha, P. and Rajasekharan, S. (2010): Hepatoprotective effect of three herbal extracts on aflatoxin B₁-intoxicated rat liver Singapore Med J., 51(4): 326.
- Sies, H. (1991): Oxidative stress: Introduction. In: Oxidative stress, oxidants and antioxidants. (SIES H., Ed.), pp. 15.22, Academic Press, California.
- Simon, J. E.; Chadwick, A. F. and Craker, L. E. (1984): Herbs: an indexed bibliography, 1971–1980. The scientific literature on selected herbs, and aromatic and medicinal plants of the temperate zone. Hamden, CT: Archon Books 770 .
- Simon, J. E., and Quinn, J. (1988): Characterization of essential oil of parsley. Journal of Agricul. Food Chemi., 36, 467–472.
- Simon, Delsaut P.; Lafontain, M.; Morele, R. and Nicot, T. (1998): Automated column-switching HPLC for the determination of Aflatoxin M₁, J. of Chromatography, B 712: 95–104.

- Singh, U.; Devaraj, S. and Jialal, I. (2005): Vitamin E, oxidative stress and inflammation. *Annu. Rev. Nutr.*, 25: 151-174.
- Singh, G.; Kapoor, I.S.; Singh, P.; Heluani, C.S.; Lampasona, M.P. and Catalan, C.A.N. (2008): Chemistry, antioxidant and antimicrobial investigation on essential oil and oleoresine of *Zingiber officinale*. *Food Chem. Toxicol.*, 46: 3295-3302.
- Singh, G.; Kapoor, I.P.S.; Singh, P.; Carola, S.; de Heluani, M. P.; de Lampasona, and Cesar A.N. (2010): Comparative study of chemical composition and antioxidant activity of fresh and dry rhizomes of turmeric (*Curcuma longa* Linn.). *Food and Chemical Toxicol.*, 48: 1026-1031.
- Smela, M. E.; Currier, S. S.; Bailey, E. A. and Essigmann, J. M. (2001): The chemistry and biology of aflatoxin B₁ from mutational spectrometry to carcinogenesis. *Carcinogenesis*, 22: 535-545.
- Srivastava, K.C. and Mustafa, T. (1989): Ginger (*Zingiber officinale*) and rheumatic disorders. *Med Hypoth.*, 29(1):25–8.
- Steinhart, C.E.; Doyle M.E. and Cochrane, B.A. (1996): Mycotoxins. Steinhart, C.E.; Doyle M.E.; Cochrane, B.A. (Eds), *Food safety*. New York: Marcel Dekker, 376-394.
- Stoilova, I.; Krastanov, A.; Stoyanova, A.; Denev, P. and Gargova, S. (2007): Antioxidant activity of a ginger extract (*Zingiber officinale*), *J. of Food Chemi.*, 102 : 764–770.
- Stubblefield, R.D.; Shotwell, O.L.; Hesseltine, C.W.; Smith, M.L. and Hall, H.H. (1967): Production of Aflatoxin on wheat and oats: Measurement with a recording densitometer. *Appli. Microbiol.*, 15(1): 186-190.
- Surh, Y. (2002): Anti-tumor promoting potential of selected spice ingredients with antioxidative and anti-inflammatory activities: a short review. *Food Chem. Toxicol.*, 40: 1091–1097.
- Suttle, NF. (1986): *The Veterinary Record*, 119: 519-522.

- Sweeney, M. J. and Dobson, A. D. W. (1998): Mycotoxin production by *Aspergillus*, *Fusarium* and *Penicillium* species. *International Journal of Food Microbiology*, 43: 141-158.
- Tabacco, A.; Meiattini, F.; Moda, E. and Tarli, P. (1979): Simplified enzymatic/colorimetric serum urea nitrogen determination. *Clin. Chem.*, 25: 336–337.
- Thapliyal, R.; Deshpande, S.S. and Maru, G.B. (2002): Mechanism(s) of turmeric-mediated protective effects against benzo(a)pyrene derived DNA adducts. *Cancer Lett.*, 175: 79–88.
- Thomson, M.; Al-Qattan, K.K.; Al-Sawan, S.M.; Alnaqeeb, M.A.; Khan, I. and Ali, M. (2002): The use of ginger (*Zingiber officinale* Rosc.) as a potential anti-inflammatory and antithrombotic agent. *Prostagl Leukotr Essen Fatty Acids.*, 67: 475–478.
- Trail, F.; Mahanti, N. and Linz, J. E. (1998): Molecular biology of aflatoxin biosynthesis. *Microbiology*, 141: 755–765.
- Tripathi, P. and Dubey, N.K. (2004): Exploitation of natural products as an alternative strategy to control post-harvest fungal rotting of fruit and vegetables. *Post-harvest Biology and Technology*, 32: 235-245.
- Van Rensburg, S. J.; Cook-Mozaffari, P.; Van Schalkwyk, D. J.; Van der Watt, J. J.; Vincent, T. J. and Purchase, I. F. (1985): Hepatocellular carcinoma and dietary aflatoxin in Mozambique and Transkei. *British J. of Cancer*, 51: 713-720.
- Van Vleet, J.F. and Ferrans, V.J. (1992): Etiological factors and pathologic alterations in selenium – vitamin E deficiency and excess in animals and humans. *Biol. Trace Elm., Res.* 33:1-12.
- Vaziri, N.D. (2004): Roles of oxidative stress and antioxidant therapy in chronic kidney disease and hypertension. *Curr. Opin. Nephrol. Hypertens.*, 13: 93–99.
- Verma, S.P.; Salamone, E. and Goldin, B. (1997): Curcumin and genistein, plant natural

- products, show synergistic inhibitory effects on the growth of human breast cancer MCF-7 cells induced by estrogenic pesticides. *Biochem. Biophys. Res. Commun.*, 233: 692–696.
- Visioli, F.; Grande, S.; Bogani, P. and Galli, C. (2004): The role of antioxidants in the Mediterranean diets: focus on cancer *Eur J Cancer Prev.*, 13(4):337-43.
- Waller, R.A. and Duncan, D. B. (1969): A Bayes rule for the symmetric multiple comparison problems. *J. Am. Stat. Assoc.*, 64: 1484-1503.
- Wang, J. S. and Groopman, J. D. (1999): DNA damage by mycotoxins. *Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis* 424: 167-181.
- Wardini, T.H. and Prakoso, B. (1999): Curcuma. In: *Plant Resources of South-East Asia* de Padua, L.S., Bun-yaphrathatsara, N., Lemmens, R.H.M.J. (Eds.), No. 12 (1). Prosea, Bogor, Indonesia, 210–219.
- Wei, A. and Shibamoto, T. (2007): Antioxidant activity and volatile constituents of various essential oils. *J. Agric. and Food Chem.*, 55: 1737– 1742.
- Whitlow, L.W.; Hagler W.M; and Hopkins. B.A. (1998): Mycotoxin occurrence in farmer submitted samples of North Carolina feedstuffs: 1989-1997. *J. Dairy Sci.* 81 (Abstr.):1189.
- Whitty, J.P. and Bjeldanes, L.F. (1987): The effects of dietary cabbage on xenobiotic metabolizing enzymes and the binding of aflatoxin B₁ to hepatic DNA in rats. *Food Chem. Toxicol.*, 25: 581-587.
- W. H. O. (1979): International program on chemical safety. World Health Organization Geneva, Environmental health criteria mycotoxins. ISBN 92 4 154071 0.
- Wild, D. (1994): The immunoassay handbook, Stockton press, 400-402.
- Williams, G. M.; Tanaka, T. and Maeura, Y. (1986): Dose-related inhibition of aflatoxin B₁, induced hepatocarcinogenesis by the phenolic antioxidants, butylated hydroxyanisole and butylated hydroxytoluene. *Carcinogenesis (Lond.)*, 7: 1043-1050.
- Wong, P. Y. Y. and Kitts, D. D. (2006): Studies on the dual antioxidant and bacterial

- properties of parsley (*Petroselinum crispum*) and cilantro (*Coriandrum sativum*) extracts. Food Chem., 97: 505–515.
- Yang, C.F.; Liu, J.; Shen, H.M. and Ong, C.N. (2000): Protective effect of ebselen on aflatoxin B₁-induced cytotoxicity in primary rat hepatocytes. Pharmacol. Toxicol., 86: 156-161.
- Yaniv, Z.; Schafferman, D. and Amar, Z. (1998): Tradition, uses, and biodiversity of rocket (*Eruca sativa*) in israel. Econ. Bot., 52, 394–400.
- Yiannikouris, A.; Andre, G.; Poughon, L.; Francois, J.; Dussap C.G.; Jeminet, G.; Bertin, G. and Jouany, J.P. (2006): Chemical and conformational study of the interactions involved in mycotoxin complexation with beta-D-glucans. Biomacromolecules, 7(4): 1147-1155.
- Yin, M., and Cheng, W. (1998): Anti-oxidant activity of several Allium members. J. Agric. Food Chem., 46: 4097-4101.
- Young D.S. (1975): Clin. Chem., 21:1D.
- Yu J. (2003) : Genetics and biochemistry of mycotoxin synthesis. In: Arora DK (Ed.). Handbook of fungal biotechnology (2 nd Edition. New York, Marcel Dekker.
- Zeringue, H. J. (2000): Identification and effects of maize silk volatiles on cultures of *Aspergillus flavus*. J. Agri. Food Chem., 48: 921–925.
- Zeringue, H. J. and Bhatnagar, D. (1990): Inhibition of aflatoxin production in *Aspergillus flavus* infected cotton bolls after treatment with neem (*Azadirachta indica*) leaf extracts. J. Am. Oil Chemi. Soci., 67: 215–216.
- Zhang, H.; Chen, F.; Wang, X. and Hui-Yuan, Y. (2006): Evaluation of antioxidant activity of parsley (*Petroselinum crispum*) essential oil and identification of its antioxidant constituents. Food Res. Inter., 39: 833–839.
- Zheng, G. Q.; Kenney, P. M.; Zhang, J. and Lam, L. K. T. (1992): Inhibition of benzo [a] pyrene-induced tumorigenesis by myristicin, a volatile aroma constituent of parsley leaf oil. Carcinogenesis, 13(10): 1921–1923.
- Zheng, W. and Wang, S. Y. (2001): Antioxidant activity and phenolic compounds in selected herbs. J. of Agri. Food Chem., 49(11): 5165–5170.

Zou, Y.; Lu, Y. and Wei, D. (2004): Antioxidant activity of a flavonoid-rich extract of *Hypericum perforatum* L. in vitro. J. Agric. and Food Chem., 52: 5032–5039.