

REFERENCES

Abd Alla, G. F. A. (2002): Toxicological and physiological effects of some muscle relaxant drugs on rats.

Ph.D thesis Fac. of Sci., Cairo Univ., Egypt.

Apple, F. S. (1989): Postmortem tricyclic antidepressant concentration assessing cause of death using parent drug to metabolite ratio.

J. Anal. Taxicol., **13 (4):** 197 – 198.

Backman, J. T.; Karjalainen, M. J.; Neuvonen, M.; Laitila, J. and Neuvonen, P. J. (2006): Refecoxib is apotent inhibitor of cytochrome P450 1A2: studies with tizanidine and caffeine in healthy subjects.

Br. J. Clin. Pharmacol., **26 (3):** 345 – 357 .

Bakalli, R. I.; Pesti, G. M.; Ragland, W. L. and Konjufca, V. (1995): Dietary copper in excess of nutritional requirement reduces plasma and breast muscle cholesterol of chickens.

Poultry Sci., **74 (2):** 360 – 365.

Bathory, G.; Sazko, E.; Magyer, K. and Deutsch, T. (1990):

Properties of the melanin binding of p-bromomethyl amphetamine.

Pol. J. Pharmacol. Pharm., **42**: 19 – 27.

Baumgartner, W. A.; Hill, V. A. and Bland, W. H. (1989):

Hair analysis of drugs of abuse.

J. Forensic Sci., **34**: 1433 – 1453.

Berkman, M. Z.; Zirh, T. A.; Berkman, K. and Pamir, M.N.

(1998): Tizanidine is an effective agent in the Prevention of focal cerebral ischemia in rats: An Experimental study.

Surg. Neurol., **50**: 264 – 271.

Burtis, C. A.; Bowers, L. D.; Chatteraj, S. C. and Ullman,

M. D. (1987):" Chromatography" In "Fundamentals of clinical chemistry" 3rd ed.

(N. W. Tietz, ed.), W. B. Saunders company, Philadelphia, London. P. 105 – 124 .

Cain, B. W.; Sileo, L.; Franson, J. C. and Moore, J. (1983):

Effects of dietary cadmium on mallard duckling.

Environ. Res., **32(2)**: 286 – 297 .

- Cameron, M.; Donati, F. and Varin, F. (1995):** *In vitro* plasma protein binding of neuromuscular blocking agent in different subpopulation of patients.
Anesth. Analg., **18(5)**: 286 – 297.
- Casarett, L. J. and Doull, J. (1996):** "The basic science of poisons" 5th ed.
(Curtis D. Klassen, ed.), McGraw – Hill. New York. **P.** 89-186 .
- Chan, E. M. and Chan, S. C. (1984):** Screening for acidic and neutral drugs by HPLC in postmortem blood.
J. Anal. Toxicol., **8(4)**: 173 – 176 .
- Chou, R.; Peterson, K. and Helfand, M. (2004):** Comparative efficacy and safety of skeletal muscle relaxants for spasticity and musculoskeletal conditions: A systematic review.
J. Pain and Symptom Manage., **28(2)**: 140 – 175.
- Clarke, E. G. C. (1986):** "Isolation and identification of drugs"
London the pharmaceutical press., 2nd ed. **P.** 636 .
- Collins, C.; Muto, J. and Spiehler, V. (1992):** Whole blood deproteinization for drug screening using automatic pipettors.

J. Anal. Toxicol., **16(5)**: 354 – 342 .

Cone, E. J.; Darwin, W. D. and Wang, W. L. (1993): The occurrence of cocaine, heroin and metabolites in hair of drug abusers.

Forensic Sci. Int., **63**: 55 – 68 .

De-Graaf, E. M.; Oosterveld, M.; Tjabbea, T. and Strickes, B. H. Ch. (1996): A case of tizanidine-induced hepatic injury.

J. Hepatol., **25**: 772 – 773 .

Devaitova, N. V. and Buravkova, L. B. (1999): The individual and integral effects of brief simulated and anesthetic pharmacological preparations on human metabolism.

Aviakosm. Ekolog. Med., **33(2)**: 26 – 30 .

Elliott, S. P. and Hale, K. A. (1999): Analysis of etorphine in postmortem samples by HPLC with UV diode-array detection.

Forensic Sci. Int., **12; 101(1)**: 9 – 16 .

Emre, M.; Leslie, G. C.; Muir, C.; Part, N. J.; Pokorny, R. and Roberts, R. C. (1994): Correlations between dose, plasma concentrations, and antispastic action of tizanidine (sirdalud).

J. Neurol. Neurosurg. Psychiatry, **57**: 1355 – 1359 .

Freitag, F. G. (2003): Preventative treatment for migraine and tension-type headaches: do drugs having effects on muscle spasm and tone have a role?
C N S Drugs, **17**: 373 – 381.

Gallichio, J. E. (2004): Pharmacologic management of spasticity following stroke.
Physical Therapy, **84 (10)**: 973 – 981.

Garriott, J. C. (1991): Skeletal muscle as a specimen for alcohol and drugs analysis.
J. Forensic Sci., **36 (1)**: 60 – 69.

Giorgi, M.; Bertini, S.; Soldani, G. and Giusiani, M. (2000):
Comparison of HPLC and GC-MS methods for determination of embutramide (A component of tanax R or T-61 R) in biological specimens.
J. Anal. Toxicol., **24 (5)**: 233 – 326.

Gomal, A. G.; Bardawil, G. M. and David, M. M. (1946):
Determination of serum protein by means of the biuret reaction. J. Bio. Chem., 175 – 177.

- Goodman, H. M. (2003):** "Basic medical endocrinology" 3rd ed.
Elsevier Science. San Diego. California. USA. **P:** 297 – 299.
- Granfors, M. T.; Backman, J. T.; Laitila, J. and Newvonen, P. J. (2003):** Tizanidine is mainly metabolized by cytochrome P450 1A2 in vitro.
Br. J. Clin. Pharmacol., **57 (3):** 349 – 353.
- Harper, H. A.; Rodwell, V. W. and Mayes, P. A. (1977):**
"Review of physiological chemistry" 16th ed.
Loss Ailos, California , **P.** 67.
- Heazlewood, V.; Symoniw, P.; Maruff, P. and Eadie, M. J. (1983):** Tizanidine initial pharmacokinetic studies in patients with spasticity.
Eur. J. Clin. Pharmacol., **25 (1):** 65 – 67.
- Henderson, G. L. (1993):** Mechanisms of drug incorporation into hair.
Forensic Sci. Int., **63:** 19 – 29.
- Henney, H. R.; Fitzpatrick, A.; Stewer, J. and Runyan, J. D. (2008):** Relative bioavailability of tizanidine hydrochloride capsule formation compared with capsule contents administered in applesauce: A single-dose, open-

label, randomized , two-way, crossover in fasted healthy adult sublecls.

Clin. Therape., **30 (12)**: 2263 – 2271.

Hilberg, T.; Rogde, S. and Morland, J. (1999): Postmortem drug redistribution-human case related to results in experimental animals.

J. Forensic Sci., **44 (1)**: 3 – 9.

Holtebeck, B. (2000): Tizanidine an alternative in the pharmacologic treatment of spasticity.

J. Pharm. Soc. Wisc. Mar. / Ap. : 26 – 30.

Honda, M.; Sekiguchi, Y.; Sato, N. and Ono, H. (2002): Involvement of imidazoline receptors in centrally acting muscle-relaxant effects of tizanidine.

Eur. J. Pharmacol., **445**: 187 – 193.

Howanitz, P. J. and Howanitz, J. H. (1984): In Clinical "Diagnosis and Management by laboratory methods" 17th ed.

(J. B. Henry. Ed.). W. B. Saunders, Philadelphia, **P.168**.

Ito, S.; Kudo, K.; Imamura, T.; Jitsufuchi, N. and Kimura,K. (1997): Detection of drugs and poisons in

postmortem tissues determination of paraquat in tissues of rabbits buried under ground.

Nippon Hoigaku Zasshi, **51 (2)**: 83 – 88.

Katzung, B. G. (2001): "Basic and clinical pharmacology" 8th ed.

McGaw-Hill Companiea, Inc. USA. Chapter 11, **P. 161**.

Kaul, N.; Dhaneshwar, S. R.; Agrawal, H.; Kakad, A. and Patil, B. (2005): Application of HPLC and HPTLC for the simultaneous determination of tizanidine and rofecoxib in pharmaceutical dosage form .

J. Pharmac. And Biome. Anal., **37**: 27 – 38 .

Klys, M.; Klementowicz, W. and Trela, F. (1997): Selected medicolegal problems in case of poisoning with substances of abuse.

Prezegl-Lek., **54 (6)**: 404 – 409.

Knight, J. A.; Andreson, S. and Rawle, J. M. (1972): Chemical basis of sulfophospho-vanillin reaction in estimating total lipid. Clin. Chem., **18**: 199 – 204.

Koch, P.; Hirst, D. R. and Von-Watburg, B. R. (1989): Biological fate of sirdalud in animals and man.

Xenobiotica, **19 (11)**: 1255 – 1265.

Koves, E. M. (1995): Use of high performance liquid chromatography-diode array detection in forensic toxicology.

J. Chromatogr. A., **692 (1-2):** 103 – 119.

Koves, E. M. and Wells, J. (1992): Evaluation of a photodiode array / HPLC-base system for the detection and quantitation of basic drugs in postmortem blood.

J. Forensic Sic., **37 (1):** 42 – 60.

Krishna, D. and Klotz, U. (1994): Extrahepatic metabolism of drugs in human.

Clin. Pharmacokinet., **26:** 144 – 160.

Lake, A. E. and Saper, J. R. (2002): Chronic headache: New advances in treatment strategies.

Neurology, **59 (2):** 8 – 13.

Lawson, K. (1998): Tizanidine a therapeutic weapon for spasticity .

Physiotherapy, **84 (9):** 418 – 420.

Logan, B. K. and Stafford, D. T. (1989): Liquid /solid extraction on diatomaceous earth for drug analysis in postmortem blood.

J. Forensic Sci., **34(3):** 553 – 564 .

Lukas, G.; Brindle, S. D. and Greengard, P. (1971): The route of absorption of interperitoneally administration compounds.

J. Pharmacol. Exp. THER., **178**: 562 – 564.

Lukin, C. L.; Hein, H. A.; Swygert, T. H.; Gunning, T. C. 3rd ; Valek, T. R.; Donica, S. K.; Nelson, R. B. 3rd and Ramsay, M. A. (1995): Duration of vacuronium induced neuromuscular block as a predictor of liver allograft dysfunction.

Anesth. Analg., **80 (3)**: 526 – 533.

Mahadik, K. R.; Paradkar, A. R.; Agrawal, H. and Kaul, N. (2003): Stability-indicating HPTLC dertermination of tizanidine hydrochloride in bulk drug formulations.

J. Pharmac. and Biome. Anal., **33**: 545 – 552.

Malanga, G. A.; Gwynn, M. W.; Smith, R. and Miller, D. (2002): Tizanidine is effective in the treatment of myofascial pain syndrome.

Pain Pysician., **5 (4)**: 422 – 432 .

Martindale (1998): "The extrapharmacopia" 31st ed. London the pharmaceutical Press, **P. 1205.**

Mathias, C. J.; Luckitt, J. Desai, P.; Barker, H.; El Masri, W. and Frankel, H. L. (1989): Pharmacodynamics and pharmacokinetics of the oral antispastic agent tizanidine in patients with spinal cord injury.
J. Rehabil. Res. Dev. **26**: 9- 16.

Meyer, U. A. (1996): Overview of enzymes of drug metabolism.
J. Pharmacokinet. Biopharm., **24**: 449 – 459.

Meythaler, J. M.; Guin-Renfroe, S.; Johnson, A. and Brunner, R. M. (2001): Prospective assessment of tizanidine for spasticity due to aquired brain injury.
Arch. Phys. Med. Rehabil., **82**: 1155 – 1163.

Miettinen, T. J.; Kanto, J. H.; Salonen, M. A. and Scheinin, M. (1996): The sedative and sympatholytic effects of oral tizanidine in healthy volunteers.
Anesth. Analg., **82 (4)**: 817 – 820.

Momo, K.; Homma, M.; Abei, M.; Hyodo, I. and Kohda, Y. (2008): Tizanidine-induced hypotension in patients with liver cirrhosis.
Eur. J. Clin. Pharmacol., **64**: 647 – 648.

Moor, K. A.; Cina, S. J.; Jones, R.; Selby, D. M.; Levine, B. and Smith, M. L. (1999): Tissue distribution of tramadol and metabolites in an overdose fatality.

Am. Forensic Med. Pathol., **20 (1):** 98 – 100 .

Moriya, F. and Hashimoto, Y. (1997): Evaluation of triage screening for drug of abuse in postmortem blood and urine samples.

Nippon. Hoigaku Zasshi, **51 (3):** 214 – 219.

Mourier, P. A. (1989): Determination of gallamine and its impurities by reversed phase ion-pair HPLC and comparison with thin layer chromatography.

J. Chromatogr., **13 (462):** 281 – 292.

Mule, S. J. (1971): Routine identification of drug of abuse in human urine.

J. Chromatogr., **55:** 255 – 257 .

Nakahara, Y. U. J.; Kikura, P. U. R. I. and Takahashi, K. (1995): Effect of structural factors on incorporation of drug into hair . The incorporation rates of amphetamine analogs. Pro. of TIAFT Int.

Conference and workshop for hair analysis in Forensic Toxicology., **P. 28 – 49.**

Nakamura, G. R.; Liu, Y. and Noguchi, T. T. (1981): A method for the separation and determination of neutral compounds in postmortem tissues.

J. Anal. Toxicol., **5 (4):** 162 – 164.

Nakamura, T.; Ono, H.; Fukuda, H.; Tanabe, M.; Kozuk, M. and Iwata, N. (1993): Local cerebral glucose utilization in rats with decerbrate rigidity and effects of centrally acting muscle relaxants, diazepam and tizanidine.

Biol. Pharm. Bull. Jan., **16 (1):** 33 – 35.

Nance, P. W.; Bugaresti, J. and Shellenberger, K. (1994): Efficacy and safety of tizanidine in the treatment of spasticity in patients with spinal cord injury.

Neurology, **44 (suppl 9):** 544 – 552.

Neumann, P. (1937): Preparation and antitremor activity.

Neth. Pa. Apple., **7(306):** 228-230.

Nickolls, L. C. (1956): "The scientific investigation of crime"
1st ed.

London, Butterworth, **P.** 382 – 383.

- Ochs, G.; Loew, M.; Tonn, J. and Toyka, K. (1998):**
Distribution, tolerability and tissue compatibility of
intrathecal tizanidine in sheep.
Acta. Anaesthesiol. Scand., **42 (7)**: 786 – 793.
- Ono, H.; Fukushima, C. and Fukuda, H. (1993):** Effects of
centrally acting muscle relaxant tizanidine on spinal
reflexes: Involvement of descending noradrenergic
system.
Jpn. J. Pharmacol., **62**: 357 – 362.
- Pipkin, F. B. (1984):** "Medical statistics made easy". Churchill
Livingston, London, **P.** 1 – 138 .
- Pounder, D. J.; Adams, E.; Fuke, C. and Longford, A. M.**
(1996): Variability of postmortem drug concentrations in
liver and lung.
J. Forensic Sci., **41 (6)**: 927 – 932.
- Puranik, M.; Wadher, S. J.; Dhole, S. and Yeale, P. G.**
(2006): Simultaneous estimation of valdecoxib and
tizanidine hydrochloride in tablets by RP-HPLC .
Ind. J. Pharmace. Sci., **68 (5)**: 670 – 672 .
- Püschel, K.; Thomasch, P. and Arnold, W. (1983):** Opiates
levels in hair.

Forensic Sci. Int., **21**: 181 – 186 .

Qi, M. L.; Wang, P. and Wang, L. (2003): Validated liquid chromatography method for assay of tizanidine in drug substance and formulated products.

Analytica. Chimica. Acta., **478**: 171 – 177.

Reitman, S. and Frankel, S. (1957): A colorimetric method for the determination of serum glutamic oxalo acetic and glutamic pyruvic transaminases.

Am. J. Clin. Patho., **28**: 56 – 60.

Satia, M. C.; Damani, R.; Shukla, M. L.; Gandhi, T. P. and Goyal, R. K. (1995): Clonidine still a better option for hypertension associated with diabetes mellitus.

Indi. J. Pharmacol., **27**: 45 – 48.

Sato, O. (1999): "Pain drugs hand book" 2nd ed. State of the art technology, Inc.

Information system for health care. **P.** 5 – 6.

Schlesinger, H. L. (1985): Topics in the chemistry of cocaine.

ODCCP. Bulletin on narcotics. Issue, **1**: 63 – 70 .

Shah, J.; Wesnes, K. A.; Kovelesky, R. A. and Henney, H. R.

(2006): Effects of food on the single-dose

pharmacokinetics /pharmacodynamics of tizanidine capsules and tablets in healthy volunteers.

Clin. Therape., **28 (9)**: 1308 – 1317.

Shellenberger, M. K.; Groves, L.; Shah, J. and Novack, G.

D. (1999): A controlled pharmacokinetic evaluation of tizanidine and baclofen at steady state.

Am. Soci. Pharmacol. Experi. Therape., **27 (2)**: 201-204.

Singh, R. M.; Singh, A.; Qardri, A. and Fahmy, N. (2002):

Acute liver failure in a patient prescribed tizanidine (zanaflex) who successfully recovered with high dose steroids.

Am. J. Gastroent., **97 (9)**: 149.

Sivasubramanian, L. and Devarajan. (2009):

Spectrophotometric and HPLC methods for simultaneous estimation of tizanidine and valdecoxib from tablets.

Int. J. Chem. Tech. Res., **1 (1)**: 96 – 102.

Smith, C.; Brnbaum, G. and Carter, J. L. (1994): Tizanidine

treatment of spasticity caused by multiple sclerosis, results of a double blind placebo controlled trial.

Neurology, **44 (suppl 9)**: 534-543.

Spratt, E. (1977): Interfering compound in urine screening for drugs.

Excerpta Medical, Drug Dependence, **5**: 2121.

Steinert, L. and Coffman, N. (1992): "Techniques of drug analysis . In clinical laboratory medicine".

(Edited by Tilton, R. C.; Balows, A.; Hohnadel, D. C. and Reiss, R. F.), Mosby Year Book, Inc., **P.** 346.

Stein, E. A. (1986): "In textbook of clinical chemistry".

(Tietz, N.W. and Saunders, W. B. ed.), Philadelphia. **P.** 879 – 886, 1818 – 1829.

Sykes, A. R.; Coop, R. L. and Robinson, M. C. (1980):

Chronic subclinical ovine fascioliasis: Plasma glutamate dehydrogenase gamma glutamyl transaminase activities and their significance as diagnostic aids.

Res. Vet. Sci., **28**: 71 – 75 .

Takenaka, M.; Iida, H.; Kasamatsu, M.; Katoh, H.; Tashiro, T. and Dohi, S. (1996): Tizanidine for pre-anesthetic medication .

Masui., **45 (8)**: 970 – 975 .

- Tilton, R. C.; Balows, A.; Hohnadel, D.C. and Reiss, R.F. (1992):** "Laboratory medicine" Mosby-year Book, Inc., London, Philadelphia. Chapter 24 .
Techniques of drug analysis. **P.** 346 – 362 .
- Toosy, A. and Thompson, A. (2000):** Multiple sclerosis: The disease and its treatment.
The Pharmaceutical J., **264:** 694 – 700.
- Traqui, A.; Kintz, P. and Mangin, P. (1995):** Hair analysis a worthless tool for therapeutic compliance monitoring.
Forensic Sci. Int., **70:** 183 – 189 .
- Trevelyan, W. E. and Harrison, J. S. (1952):** Fractionation and micro-determination of cell carbohydrates.
Biochem. J., **50:** 298 – 303.
- Tse, F. L.; Jaffe, J. M. and Bhuta, S. (1987):** Pharmacokinetics of orally administrated tizanidine in healthy volunteers.
Fundam. Clin. Pharmacol., **1 (6):** 479 - 488.
- Uematoso, T.; Sato, R.; Fujimori, O. and Nakashima, M. (1990):** Human scalp hair as evidence of individual dosage history of haloperidol: A possible linkage of haloperidol excretion into hair with hair pigment.

Arch. Derm. Res., **P.** 120 -125.

Uematsu, M.; Nakano, M.; Akiyama, H. and Najkashima, M. (1992): Possible effect of pigment on the pharmacokinetics of ofloxacin and its excretion in hair.
J. Pharm. Sci., **81**: 45-48.

Valsala, P. and Kurup, P. A. (1987): Investigations on the mechanism of hypercholesterolemia observed in copper deficiency in rats.
J. Biosciences (Bangalore), **12**: 137-142.

Vearanjaneyulu, A.; Sridhar, N. and Sharma, A. (1995): Alpha-adreno-receptor mediated hyperglycemic effect of clonidine and 1st 606 in wister rats.
Indian J. of pharmacol., **27**: 45-48.

Wagstaff, A. J. and Bryson, H. M. (1997): Tizanidine: A review of its pharmacology, clinical efficacy and tolerability in the management of spasticity associated with cerebral and spinal disorders.
Drugs, **53**: 435-452.

Wahlefeld, A. W. (1974): "In methods of enzymatic analysis"
Bermeyer, H. U. ed.
Academic Press, New York, **5**: 1831-1835.



Wallace, J. D. (1994): Summary of combined clinical analysis of controlled clinical trials with tizanidine.
Neurology, **44 (9):** 553 – 559.

Whitter, P. D. and Cary, P. L. (1986): Postmortem liver specimen by toxi-lab.
J. Anal. Toxicol., **10 (2):** 68-71.

Wilson, J. F.; Green, S.; Dunstan, D.G.; Wicks, J. F. C.; Brewer, C. and Perfect, N. (1995): Relationship between methadone dose, hair methadone concentration and hair color in man.
Ther. Drug Monit., **17:** 419 - 421.



