

REFERENCES

- Agarwal A, Gautam S, Gupta D.** Evaluation of a single preoperative dose of pregabalin for attenuation of postoperative pain after laparoscopic cholecystectomy. *Br J Anaesth* **2008**; 101:700–704.
- Ahmadi S, Lippross S, Neuhuber WL.** PGE(2) selectively blocks inhibitory glycinergic neurotransmission onto rat superficial dorsal horn neurons. *Nature Neuroscience* **2002**; 5: 34–40.
- Andrieu G, Roth B, Ousmane L.** The efficacy of intrathecal morphine with or without clonidine for postoperative analgesia after radical prostatectomy. *Anesth Analg* **2009**; 108:1954–1957.
- Apkarian AV, Bushnell MC, Treede RD.** Human brain mechanisms of pain perception and regulation in health and disease. *Eur J Pain* **2005**; 9(4):463–84.
- Ashburn MA, Caplan RA, Carr DB.** Practice guidelines for acute pain management in the perioperative setting. An updated report by the American Society of Anesthesiologists task force on acute pain management. *Anesthesiology* **2004**; 100:1573–1581.
- Aubrun F, Gaillat C, Rosenthal D.** Effect of a low-dose ketamine regimen on pain, mood, cognitive function and memory after major gynaecological surgery: a randomized, double-blind, placebo-controlled trial. *Eur J Anaesthesiol* **2008**; 25:97–105.
- Aveline C, Gautier JF, Vautier P.** Postoperative analgesia and early rehabilitation after total knee replacement: a comparison of continuous low-dose intravenous ketamine versus nefopam. *Eur J Pain* **2009**; 13:613–619.
- Axelsson K, Nordenson U, Lidegran G, Gupta A.** Patient-controlled regional analgesia (PCRA) with ropivacaine after arthroscopic subacromial decompression. *Acta Anaesthesiol Scand* **2003**; 47:993-1000.
- Backhaus J, Junghanns K, Broocks A, Riemann D, Hohagen F.** Test-retest reliability and validity of the Pittsburgh sleep quality index in primary insomnia. *J Psychosom Res.* **2002**; 53:737–740.
- Barash PG, Cullen BF and Stoelting RK:** Clinical anesthesia, 4th edition, Philadelphia, Pa: Lippincott Williams & Wilkins; 929-967, **2001**.
- Barbosa MA.** Pain assessment in postoperative cardiac surgery. *Acta Paul Enferm* **2010**; 23(2):224-9.

References

- Barry BW.** Novel mechanisms and devices to enable successful transdermal drug delivery. *Eur J Pharm Sci* **2001**; 14: 101–14.
- Basse L, Hjort Jakobsen D, Billesbolle P.** Clinical pathway to accelerate recovery after colonic resection. *Annals of Surgery* **2000**; 232: 51-57.
- Beattie WS, Badner NH, Choi P.** Epidural analgesia reduces postoperative myocardial infarction: a meta-analysis. *Anesthesia and Analgesia* **2001**; 93: 853-858.
- Beattie WS, Badner NH, Choi P.** Epidural analgesia reduces postoperative myocardial infarction: a meta-analysis. *Anesth Analg* **2001**; 93: 853–858.
- Beecher HK.** The Measurement of Subjective Responses. Oxford: Oxford University Press; **1959**.
- Bell RF, Dahl JB, Moore RA.** Perioperative ketamine for acute postoperative pain. *Cochrane Database of Systematic Reviews* **2006**; (1): CD004603.
- Beloeil H, Ababneh Z, Chung R.** Effects of Bupivacaine and Tetrodotoxin on Carrageenan-induced hind paw inflammation in rats: hyperalgesia, edema, and systemic cytokines. *Anesthesiology* **2006**; 105: 128-138.
- Bian F, Li Z, Offord J.** Calcium channel alpha2-delta type 1 subunit is the major binding protein for pregabalin in neocortex, hippocampus, amygdala, and spinal cord: an ex vivo autoradiographic study in alpha2-delta type 1 genetically modified mice. *Brain Res* **2006**; 1075:68–80.
- Bjune K, Stubhaug A, Dodgson MS, Breivik H.** Additive analgesic effect of codeine and paracetamol can be detected in strong, but not moderate, pain after Caesarean section. Baseline pain-intensity is a determinant of assay-sensitivity in a postoperative analgesic trial. *Acta Anaesthesiol Scand* **1998**; 40: 399–407
- Bombardier C.** Outcome assessments in the evaluation of treatment of spinal disorders. *Spine*. **2000**; 25:3100–3103.
- Bonnet F.** Postoperative pain management and outcome after surgery. *Best Pract Res Clin Anaesthesiol* **2007**; 21 (1): 99- 107
- Bowdle T.** Nocturnal arterial oxygen desaturation and episodic airway obstruction after ambulatory surgery. *Anesth Analg* **2004**; 99:70-6.
- Bowler I, Djaiani G, Abel R.** A combination of intrathecal morphine and remifentanyl anesthesia for fast track cardiac anesthesia and surgery. *J Cardiothorac Vasc Anesth* **2002**; 16: 709–714.

References

- Breivik EK, Barkvoll P, Skovlund E.** Combining ciclofenac with acetaminophen or acetaminophen–codeine after oral surgery: a randomized, double-blind single-dose study. *Clin Pharmacol Ther* **1999**; 66: 624–35
- Breivik EK, Bjornsson GA, Skovlund E.** A comparison of pain rating scales by sampling from clinical trial data. *Clin J Pain* **2000**; 16: 22–8
- Breivik H, Borchgrevink PC, Allen SM, Rosseland LA.** Assessment of pain. *Br J Anaesth* **2008**; 101: 17–24
- Breivik H, Collett B, Ventafridda V, Cohen R, Gallacher D.** Survey of chronic pain in Europe: prevalence, impact on daily life, and treatment. *Eur J Pain* **2006**; 10: 287–333
- Breivik H, Stubhaug A.** Management of acute postoperative pain: still a long way to go! *Pain*. **2008**;137(2):233-4.
- Brennan TJ.** Frontiers in translational research: the etiology of incisional and postoperative pain. *Anesthesiology* **2002**; 97: 535–537.
- Buvanendran A, Kroin JS, Berger RA.** Upregulation of prostaglandins E2 and interleukin in the central nervous system and peripheral tissue during and after surgery in humans. *Anesthesiology* **2006**; 104:403–410.
- Buvanendran A, Kroin JS, Tuman KJ.** Effects of perioperative administration of selective cyclooxygenase 2 inhibitor of pain management and recovery of function after knee replacement. *JAMA* **2003**; 290:2411–2418.
- Buvanendran A, Kroin JS.** Multimodal analgesia for controlling acute postoperative pain. *Curr Opin Anaesthesiol* **2009**, 22:588–593.
- Byers MR, Bonica JJ.** Peripheral pain mechanisms and nociceptor plasticity. In: Loeser JD, editor. *Management of pain*. New York: Lippincott Williams & Wilkins; **2001**. p. 26–72.
- Callesen T, Kehlet H.** Postherniorrhaphy pain. *Anesthesiology* **1997**; 87: 1219-1230.
- Capdevila X, Barthelet Y, Biboulet P.** Effects of perioperative analgesic technique on the surgical outcome and duration of rehabilitation after major knee surgery. *Anesthesiology* **1999**; 91: 8-15.
- Capdevila X, Dadure C, Bringuier S, Bernard N, Biboulet P, Gaertner E, Macaire P.** Effect of patient controlled perineural analgesia on rehabilitation and pain after ambulatory orthopedic surgery: A multicenter randomized trial. *Anesthesiology* **2006**; 105:566-573.

- Carli F, Mayo N, Kluben K.** Epidural analgesia enhances functional exercise capacity and health-related quality of life after colonic surgery: results of a randomized trial. *Anesthesiology* **2002**; 97: 540-549.
- Carney J, McDonnell JG, Ochana A.** The transversus abdominis plane block provides effective postoperative analgesia in patients undergoing total abdominal hysterectomy. *Anesth Analg* **2008**; 107:2056– 2060.
- Carr DB, Goudas LC, Albin R, Hamilton D, Rogers MC, Firestone L, Lavin PT, Mermelstein F.** Safety and efficacy of intranasal ketamine for the treatment of breakthrough pain in patients with chronic pain: A randomized, double-blind, placebo controlled, crossover study. *Pain* **2004**; 108:17-27.
- Cashman JN, Dolin SJ.** Respiratory and haemodynamic effects of acute postoperative pain management: Evidence from published data. *Br J Anaesth* **2004**; 93:212-223.
- Chen PP, Cheam EW, Ma M, Lam KK, Ngan Kee WD, Gin T.** Patient-controlled pethidine after major upper abdominal surgery: Comparison of the epidural and intravenous routes. *Anaesthesia* **2001**; 56: 1106-1112.
- Cherny NI.** Opioid analgesics: Comparative features and prescribing guidelines. *Drugs* **1996**; 51:713-737.
- Coghill RC, Talbot JD, Evans AC.** Distributed processing of pain and vibration by the human brain. *J Neurosci* **1994**;14(7):4095–108.
- Coull JA, Boudreau D, Bachand K.** Trans-synaptic shift in anion gradient in spinal lamina I neurons as a mechanism of neuropathic pain. *Nature* **2003**; 424(6951):938–42.
- Cowan P, Kelley N.** Quality of Life Scale: a Measure of Function for People with Pain. Rocklin, CA: American Chronic Pain Association; **2003**.
- Curdy C, Kalia YN, Guy RH.** Non-invasive assessment of the effects of iontophoresis on human skin in-vivo. *J Pharm Pharmacol* **2001**; 53: 769–77
- Dal D, Kanbak M, Caglar M, Aypar U.** A background infusion of morphine does not enhance postoperative analgesia after cardiac surgery. *Can J Anaesth* **2003**; 50: 476–479.
- Dale O, Hjortkjaer R, Kharasch ED.** Nasal administration of opioids for pain management in adults. *Acta Anaesthesiol Scand* **2002**; 46:759-770.
- Davis MP, Shaiova LA, Angst MS.** When opioids cause pain. *J Clin Oncol* **2007**; 25(28): 4497–8.

- DeWeese FT, Akbari Z, Carline E.** Pain control after knee arthroplasty: Intraarticular versus epidural anesthesia. *Clin Orthop Relat Res* **2001**; 392:226-231.
- di Benedetto P, Casati A, Bertini L.** Continuous subgluteus sciatic nerve block after orthopedic foot and ankle surgery: Comparison of two infusion techniques. *Reg Anesth Pain Med* **2002**; 27:168-172.
- Dirks J, Moiniche S, Hilsted KL, Dahl JB.** Mechanisms of postoperative pain: clinical indications for a contribution of central neuronal sensitization. *Anesthesiology* **2002**; 97: 1591-1596.
- Dolin SJ, Cashman JN, Bland JM.** Effectiveness of acute postoperative pain management: I. Evidence from published data. *Br J Anaesth* **2002**; 89:409-423.
- Dowling R, Thielmeier K, Ghaly A.** Improved pain control after cardiac surgery: results of a randomized, double-blind, clinical trial. *J Thorac Cardiovasc Surg* **2003**; 126: 1271–1278.
- Dracup K, Bryan-Brown CW.** Pain in the ICU: fact or fiction? *Am J Critical Care* **1995**; 4: 337–339.
- Eide PK.** Wind-up and the NMDA receptor complex from a clinical perspective. *Eur J Pain* **2000**; 4(1):5–15.
- Eledjam JJ, Cuvillon P, Capdevila X, Macaire P, Serri S, Gaertner E, Jochum D.** Postoperative analgesia by femoral nerve block with ropivacaine 0.2% after major knee surgery: Continuous versus patient- controlled techniques. *Reg Anesth Pain Med* **2002**; 27:604-611.
- Engelhardt T, Zaarour C, Naser B.** Intraoperative low-dose ketamine does not prevent a remifentanyl-induced increase in morphine requirement after pediatric scoliosis surgery. *Anesth Analg* **2008**; 107:1170–1175.
- Farmery AD, Wilson-MacDonald J.** The analgesic effect of epidural clonidine after spinal surgery: a randomized placebo-controlled trial. *Anesth Analg* **2009**; 108:631–634.
- Fassoulaki A, Triga A, Melemenis A, Sarantopoulos C.** Multimodal analgesia with gabapentin and local anesthetics prevents acute and chronic pain after breast surgery for cancer. *Anesthesia and Analgesia* **2005**; 101: 1427-1432.
- Fayaz MH, Abel RJ, Pugh SC.** Opioid-sparing effects of diclofenac and paracetamol lead to improved outcomes after cardiac surgery. *J Cardiothorac Vasc Anesth* **2004**; 18: 742–747.

References

- Feldt K:** Pain measurement, Present concerns and future directions. *Pain Med*; 8 (7): 541-543, **2007**.
- Feng Y, Ju H, Yang B, An H.** Effects of a selective cyclooxygenase-2 inhibitor on postoperative inflammatory reaction and pain after total knee replacement. *J Pain* **2008**; 9:45–52.
- Fields HL, Malick A, Burstein R.** Dorsal horn projection targets of ON and OFF cells in the rostral ventromedial medulla. *J Neurophysiol* **1995**; 74(4):1742–59.
- Fisher K, Coderre TJ, Hagen NA.** Targeting the N-methyl-D-aspartate receptor for chronic pain management. Preclinical animal studies, recent clinical experience and future research directions. *Journal of Pain and Symptom Management* **2000**; 20: 358–373.
- Fredman B, Shapiro A, Zohar E, Feldman E, Rawal N, Jedeikin R.** The analgesic efficacy of patient-controlled ropivacaine instillation after Cesarean delivery. *Anesth Analg* **2000**; 91:1436-1440.
- Freye E, Anderson-Hillemacher A, Ritzdorf I, Victor Levy J.** Opioid Rotation from High-Dose Morphine to Transdermal Buprenorphine (Transtec®) in Chronic Pain Patients. *Pain Practice*, **2007**; 7 (2): 123–129.
- Ganapathy S, Amendola A, Lichfield R, Fowler PJ, Ling E.** Elastomeric pumps for ambulatory patient controlled regional analgesia. *Can J Anaesth* **2000**; 47:897-902.
- Ganapathy S, Choi PT, Chisholm KC, Sutton I, Chan VWS.** Inhaled liposome-encapsulated fentanyl (AeroLEF) for post-operative pain after orthopedic surgery. *Anesthesiology* **2006**; 105:A1191.
- Goffaux P, Redmond WJ, Rainville P.** Descending analgesia when the spine echoes what the brain expects. *Pain* **2007**; 130(1–2):137–43.
- Granot M, Granovsky Y, Sprecher E.** Contact heat-evoked temporal summation: tonic versus repetitive-phasic stimulation. *Pain* **2006**; 122(3):295–305.
- Grape S, Tramer MR.** Do we need preemptive analgesia for the treatment of postoperative pain? *Best Practice & Research Clinical Anaesthesiology* **2007**; 21 (1):51–63.
- Grass JA.** Patient-controlled analgesia. *Anesth Analg* **2005**; 101:S44-S61.
- Griffin M.** Con: non-steroidal anti-inflammatory drugs should not be routinely administered for postoperative analgesia after cardiac surgery. *J Cardiothorac Vasc Anesth* **2000**; 14: 735–738.

- Grond S, Radbruch L, Lehmann KA.** Clinical pharmacokinetics of transdermal opioids: focus on transdermal fentanyl. *Clin Pharmacokinet* **2000**; 38: 59–89
- Groudine SB, Fisher HAG, Kaufman RP.** Intravenous lidocaine speeds the return of bowel function, decreases postoperative pain, and shortens hospital stay in patients undergoing radical retropubic prostatectomy. *Anesth Analg* **1998**; 86: 235–239.
- Guilbaud G, Besson J-M.** Physiologie du circuit de la douleur. In: Brasseur L, Chauvin M, Guilbaud G, editors. *Douleurs*. Paris: Maloine; **1997**. p. 7–22.
- Guindon J, Walezak JS and Beaulieu P:** Recent advances in the pharmacological management pain. *Drugs*; 67 (15): 2121-2133, **2007**.
- Guler T, Unlugenc H, Gundogan Z.** A background infusion of morphine enhances patient-controlled analgesia after cardiac surgery. *Can J Anaesth* **2004**; 51: 718–722.
- Gupta R, Raja SN.** Chronic pain: pathophysiology and its therapeutic implications. *Current Review of Pain*. **1996**; 1:1–9.
- Gust R, Pecher S, Gust A.** Effect of patient-controlled analgesia on pulmonary complications after coronary artery bypass grafting. *Crit Care Med* **1999**; 27: 2218–2223.
- Gutstein H, Akil H.** Opioid analgesics. In: Hardman J, Limbird L, editors. *Goodman and Gilman's: The pharmacological basis of therapeutics*. New York: McGraw- Hill; **2001**. p. 569-619.
- Habib AS, White WD, El Gasim MA.** Transdermal nicotine for analgesia after radical retropubic prostatectomy. *Anesth Analg* **2008**; 107:999–1004.
- Hall R, Adderley N, MacLaren C.** Does intrathecal morphine alter the stress response following coronary artery bypass grafting surgery? *Can J Anaesth* **2000**; 47: 463–466.
- Hallett A, O'Higgins F, Francis V, Cook TM.** Patient controlled intranasal diamorphine for postoperative pain: An acceptability study. *Anaesthesia* **2000**; 55: 532-539.
- Hariharan S, Moseley H, Kumar A, Raju S.** The effect of preemptive analgesia in postoperative pain relief: a prospective double-blind randomized study. *Pain Med* **2009**; 10:49–53.
- Haythornthwaite JA, Raja SN, Fisher B.** Pain and quality of life following radical retropubic prostatectomy. *J Urol* **1998**; 160: 1761-1764.

- Hicks CL, von Baeyer CL, Spafford PA.** The faces pain scale-revised: toward a common metric in pediatric pain measurement. *Pain* **2001**; 93: 173–83
- Hoar PF, Hickey RF, Ulliyot DJ.** Systemic hypertension following myocardial revascularization: a method of treatment using epidural anesthesia. *J Thorac Cardiovasc Surg* **1976**; 71: 859–864.
- Horlocker TT, Heit JA, Mulroy MF, Rosenquist RW, Rowlingson J, Tryba M, Yuan CS.** Regional anesthesia in the anticoagulated patient: Defining the risks. *Reg Anesth Pain Med* **2004**; 29:1-12.
- Huang YM, Wang CM, Wang CT.** Perioperative celecoxib administration for pain management after total knee arthroplasty: a randomized, controlled study. *BMC Musculoskelet Disord* **2008**; 9:77.
- Hurley RW, Cohen SP, Williams KA.** The analgesic effects of perioperative gabapentin on postoperative pain: a meta-analysis. *Regional Anesthesia and Pain Medicine* **2006**; 31: 237-247.
- Hynninen MS, Cheng DC, Hossian IJ.** Non-steroidal anti-inflammatory drugs in treatment of postoperative pain after cardiac surgery. *Can J Anaesth* **2000**; 47: 1182–1187.
- Ilfeld BM, Morey TE, Wright TW, Chidgey LK, Enneking FK.** Continuous interscalene brachial plexus block for postoperative pain control at home: A randomized, double-blinded, placebo-controlled study. *Anesth Analg* **2003**; 96:1089-1095.
- Immer FF, Immer-Bansi AS, Trachsel N.** Pain treatment with a COX-2 inhibitor after coronary artery bypass operation: a randomized trial. *Ann Thorac Surg* **2003**; 75: 490–495.
- Insler SR, O'Connor M, Samonte AF, Bazaral MG.** Lidocaine and the inhibition of postoperative pain in coronary artery bypass patients. *J Cardiothorac Vasc Anesth* **1995**; 9: 541–546.
- International Association for the Study of Pain.** Pain terms: a list with definitions and notes on usage. *Pain*. **1979**; 6:249.
- Ji RR, Strichartz G.** Cell signaling and the genesis of neuropathic pain. *Science* **2004**; 252: STKE re 14
- Jokela R, Ahonen J, Tallgren M.** A randomized controlled trial of perioperative administration of pregabalin for pain after laparoscopic hysterectomy. *Pain* **2008**; 134:106–112.

- Jokela R, Ahonen J, Tallgren M.** Premedication with pregabalin 75 or 150mg with ibuprofen to control pain after day-case gynaecological laparoscopic surgery. *Br J Anaesth* **2008**; 100:834–840.
- Kalia YN, Naik A, Garrison J, Guy RH.** Iontophoretic drug delivery. *Adv Drug Deliv Rev* **2004**; 56: 619–58
- Kardash KJ, Sarrazin F, Tessler MJ.** Single-dose dexamethasone reduces dynamic pain after total hip. *Anesth Analg* **2008**;106:1253–57.
- Katz J.** George Washington Crile, anoci-association and pre-emptive analgesia. *Pain* **1993**; 53: 243–245.
- Kearns GL, Heacock J, Daly SJ, Singh H, Qu S.** Percutaneous lidocaine administration via a new iontophoresis system in children: tolerability and absence of systemic bioavailability. *Pediatrics* **2003**; 112: 578–82
- Kehlet H, Jensen TS & Woolf C.** Persistent postsurgical pain: risk factors and prevention. *Lancet* **2006**; 367: 1618-1625.
- Kelly DJ, Ahmad M & Brull SJ.** Preemptive analgesia II: recent advances and current trends. *Canadian Journal of Anaesthesia* **2001**; 48: 1091–1101.
- Kissin I.** Preemptive analgesia: terminology and clinical relevance. *Anesthesia and Analgesia* **1994**; 79: 809–810.
- Kulika A, Ruela M, Bourke ME.** Postoperative naproxen after coronary artery bypass surgery: a double blind randomized controlled trial. *Eur J Cardiothorac Surg* **2004**; 26: 694–700.
- Kuperberg KG, Grubbs L.** Coronary artery bypass patients' perceptions of acute postoperative pain. *Clin Nurse Specialist* **1997**; 11: 116–122
- Lavand'homme PM, Roelants F, Waterloos H.** An evaluation of the postoperative antihyperalgesic and analgesic effects of intrathecal clonidine administered during elective cesarean delivery. *Anesth Analg* **2008**; 107: 948–955.
- Lebovits AH, Zenetos P, O'Neill DK, Cox D, Dubois MY, Jansen LA, Turndorf H.** Satisfaction with epidural and intravenous patient-controlled analgesia. *Pain Med* **2001**; 2:280-286.
- Lehmann LJ, DeSio JM, Radvany T, Bikhazi GB.** Transdermal fentanyl in postoperative pain. *Reg Anesth* **1997**; 22: 24–8
- Lenz FA, Gracely RH, Romanoski AJ.** Stimulation in the human somatosensory thalamus can reproduce both the affective and sensory dimensions of previously experienced pain. *Nat Med* **1995**; 1(9):910–3.

References

- Li J, Simone DA, Larson AA.** Windup leads to characteristics of central sensitization. *Pain* **1999**; 79(1):75–82.
- Lin TF, Yeh YC, Lin FS.** Effect of combining dexmedetomidine and morphine for intravenous patient-controlled analgesia. *Br J Anaesth* **2009**; 102:117–122.
- Liu SS, Allen HW, Olsson GL.** Patient-controlled epidural analgesia with bupivacaine and fentanyl on hospital wards: Prospective experience with 1,030 surgical patients. *Anesthesiology* **1998**; 88:688–695.
- Liu SS, Block BM, Wu CL.** Effects of perioperative central neuraxial analgesia on outcome after coronary artery bypass surgery. A Meta-analysis. *Anesthesiology* **2004**; 101: 153–161.
- Macrae WA.** Chronic pain after surgery. *British Journal of anaesthesia* **2001**; 87(1): 88–98.
- Marchand S, Arsenault P.** Spatial summation for pain perception: interaction of inhibitory and excitatory mechanisms. *Pain* **2002**; 95(3):201–6.
- Marchand S.** The Physiology of Pain Mechanisms: From the Periphery to the Brain. *Rheum Dis Clin N Am* **2008**; 34:285–309
- Marret E, Flahault A, Samama CM.** Effects of postoperative, nonsteroidal, anti-inflammatory drugs on bleeding risk after tonsillectomy: metaanalysis of randomized, controlled trials. *Anesthesiology* **2003**; 98:1497–1502.
- Mathews ET, Abrams LD.** Intrathecal morphine in open heart surgery [Correspondence]. *Lancet* **1980**; 2: 543.
- Mathiesen O, Jacobsen LS, Holm HE.** Pregabalin and dexamethasone for postoperative pain control: a randomized controlled study in hip arthroplasty. *Br J Anaesth* **2008**; 101:535–541.
- McCaffery M, Beebe A.** *Pain: Clinical Manual for Nursing Practice*. Philadelphia, PA: Mosby; **1989**, 7.
- McCarberg B, Stanos S.** Key Patient Assessment Tools and Treatment Strategies for Pain Management. *Pain Practice* **2008**; 8 (6): 423–432
- McCartney CJ, Sinha A, Katz J.** A qualitative systematic review of the role of N-methyl-D-aspartate receptor antagonists in preventive analgesia. *Anesthesia and Analgesia* **2004**; 98: 1385–1400.
- McConnell S, Kolopack P, Davis AM.** The Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC): a review of its utility and measurement properties. *Arthritis Rheum.* **2001**; 45:453–461.

- McDonald SB, Jacobsohn E, Kopacz DJ.** Parasternal block and local anesthetic infiltration with levobupivacaine after cardiac surgery with desflurane: the effect on postoperative pain, pulmonary function, and tracheal extubation times. *Anesth Analg* **2005**; 100: 25–32.
- McQuay HJ.** Pre-emptive analgesia: a systematic review of clinical studies. *Annals of Medicine* **1995**; 27: 249–256.
- Melzack R, Wall PD.** Pain mechanisms: a new theory. *Science* **1965**; 150:971–9.
- Meyer RA, Ringkamp M, Campbell JN.** Peripheral mechanisms of nociception. In: McMahon SB, Koltzenburg M, editors. *Wall and Melzack's textbook of pain*. 5th edition. Philadelphia: Elsevier; **2006**. p. 3–34.
- Millan MJ.** Descending control of pain. *Prog Neurobiol* **2002**; 66(6):355–474.
- Millan MJ.** The induction of pain: an integrative review. *Prog Neurobiol* **1999**; 57(1):1–164.
- Modig J, Borg T, Kalstrom G.** Thromboembolism after total hip replacement: role of epidural and general anesthesia. *Anesthesia and Analgesia* **1983**; 62: 174-180.
- Møiniche S, Kehlet H, Dahl JB.** A qualitative and quantitative systematic review of preemptive analgesia for postoperative pain relief: the role of timing of analgesia. *Anesthesiology* **2002**; 96: 725–741.
- Moodie JE, Brown CR, Bisley EJ.** The safety and analgesic efficacy of intranasal ketorolac in patients with postoperative pain. *Anesth Analg* **2008**; 107:2025–2031.
- Munro AJ, Long GT, Sleight JW.** Nurse-administered subcutaneous morphine is a satisfactory alternative to intravenous patient-controlled analgesia morphine after cardiac surgery. *Anesth Analg* **1998**; 87: 11–15.
- Mystakidou K, Katsouda E, Parpa E, Tsiatas ML, Vlahos L.** Oral transmucosal fentanyl citrate for the treatment of breakthrough pain in cancer patients: An overview of its pharmacological and clinical characteristics. *Am J Hosp Palliat Care* **2005**; 22:228- 232.
- Nielsen KC, Greengrass RA, Steele SM.** Continuous interscalene brachial plexus blockade provides good analgesia at home after major shoulder surgery – Report of four cases. *Can J Anaesth* **2003**; 50:57-61.
- Niemi G, Breivik H.** The minimally effective concentration of adrenaline in a low-concentration thoracic epidural analgesic infusion of bupivacaine, fentanyl and adrenaline after major surgery. A randomized, double blind dose-finding study. *Acta Anaesthesiol Scand* **2003**; 47: 439–50

- Nirschl RP, Rodin DM, Ochiai DH, Maartmann-Moe C.** Iontophoretic administration of dexamethasone sodium phosphate for acute epicondylitis. A randomized, double-blinded, placebo controlled study. *Am J Sports Med* **2003**; 31: 189–95
- Norris EJ, Beattie C, Perler BA.** Double-masked randomized trial comparing alternate combinations of intraoperative anesthesia and postoperative analgesia in abdominal aortic surgery. *Anesthesiology* **2001**; 95: 1051–1053.
- Nussmeier NA, Whelton AA, Brown MT.** Complications of the COX-2 inhibitors parecoxib and valdecoxib after cardiac surgery. *New Eng J Med* **2005**; 352: 1081–1091.
- Ong CKS, Philipp L, Robin AS.** The efficacy of preemptive analgesia for acute postoperative pain management: a meta-analysis. *Anesthesia and Analgesia* **2005**; 100: 757–773.
- Ott E, Nussmeier NA, Duke PC.** Efficacy and safety of the cyclooxygenase 2 inhibitors parecoxib and valdecoxib in patients undergoing coronary artery bypass surgery. *J Thorac Cardiovasc Surg* **2003**; 125: 1481–92.
- Paech MJ, Lim CB, Banks SL, Rucklidge MW, Doherty DA.** A new formulation of nasal fentanyl spray for postoperative analgesia: A pilot study. *Anaesthesia* **2003**; 58:740–744.
- Palecek J, Paleckova V, Willis WD.** Fos expression in spinothalamic and postsynaptic dorsal column neurons following noxious visceral and cutaneous stimuli. *Pain* **2003**; 104(1–2): 249–57.
- Palecek J, Willis WD.** The dorsal column pathway facilitates visceromotor responses to colorectal distention after colon inflammation in rats. *Pain* **2003**; 104(3):501–7.
- Park WY, Thompson JS, Lee K.** Effect of epidural anesthesia and analgesia on perioperative outcome. A randomized, controlled Veteran Affairs Cooperative study. *Annals of Surgery* **2001**; 234: 560–571.
- Parker MJ, Handoll HH, Griffiths R.** Anesthesia for hip fracture surgery in adults. *Cochrane Database of Systematic Reviews* **2001**; (4): CD000521.
- Perttunen K, Tasmuth T, Kalso E.** Chronic pain after thoracic surgery: a follow-up study. *Acta Anaesthesiologica Scandinavica* **1999**; 43: 563–567.
- Pettersson PH, Lindskog EA, Owall A.** Patient-controlled vs. nurse-controlled pain treatment after coronary artery bypass surgery. *Acta Anaesth Scand* **2000**; 44: 43–47.

References

- Peyton PJ, Myles PS, Silbert BS.** Perioperative epidural analgesia and outcome after major abdominal surgery in high-risk patients. *Anesthesia and Analgesia* **2003**; 96: 548-554.
- Pogatzki EM, Vandermeulen EP, Brennan TJ.** Effect of plantar local anesthetic injection on dorsal horn neuron activity and pain behaviors caused by incision. *Pain* **2002**; 97:151-161.
- Pollard CA.** Preliminary validity study of the pain disability index. *Percept Mot Skills*. **1984**; 59:974.
- Price DD.** Psychological and neural mechanisms of the affective dimension of pain. *Science* **2000**; 288(5472):1769-72.
- Priestley MC, Cope L, Halliwell R.** Thoracic epidural anesthesia for cardiac surgery: the effects on tracheal intubation time and length of hospital stay. *Anesth Analg* **2002**; 94: 275-282.
- Rainville P, Carrier B, Hofbauer RK.** Dissociation of pain sensory and affective dimensions using hypnotic modulation. *Pain* **1999**; 82(2):159-71.
- Rainville P, Duncan GH, Price DD.** Pain affect encoded in human anterior cingulate but not somatosensory cortex. *Science* **1997**; 277:968-71.
- Randinitis EJ, Posvar EL, Alvey CW.** Pharmacokinetics of pregabalin in subjects with various degrees of renal function. *J Clin Pharmacol* **2003**; 43:277-283.
- Rapanos T, Murphy P, Szalai JP.** Rectal indomethacin reduces postoperative pain and morphine use after cardiac surgery. *Can J Anaesth* **1999**; 46: 725-730.
- Rawal N, Allvin R, Axelsson K, Hallen J, Ekback G, Ohlsson T, Amilon A.** Patient-controlled regional analgesia (PCRA) at home: Controlled comparison between bupivacaine and ropivacaine brachial plexus analgesia. *Anesthesiology* **2002**; 96:1290-1296.
- Reynolds DV.** Surgery in the rat during electrical analgesia. *Science* **1969**; 164(878):444-5.
- Richman JM, Liu SS, Courpas G, Wong R, Rowlingson AJ, McGready J, Cohen SR, Wu CL.** Does continuous peripheral nerve block provide superior pain control to opioids? A meta-analysis. *Anesth Analg* **2006**; 102:248-257.
- Roberge CW, McEwen M.** The effects of local anesthetics on postoperative pain. *AORN J* **1998**; 68: 1003-1012.

- Rodgers A, Walker N, Schug S.** Reduction of postoperative mortality and morbidity with epidural or spinal anaesthesia: results from overview of randomised trials. *British Medical Journal* **2000**; 321: 1493-1497.
- Roediger L, Larbuisson R, Lamy M.** New approaches and old controversies to postoperative pain control following cardiac surgery. *European Journal of Anaesthesiology* **2006**; 23: 539–550
- Roehrs T, Hyde M, Blaisdell B, Greenwald M, Roth T.** Sleep loss and REM sleep loss are hyperalgesic. *Sleep*. **2006**; 29:145–151.
- Romundstad L, Breivik H, Roald H.** Chronic pain and sensory changes after augmentation mammoplasty: long term effects of preincisional administration of methylprednisolone. *Pain* **2006**; 124: 92–9
- Rosseland LA, Stubhaug A, Sandberg L, Breivik H.** Intraarticular (IA) catheter administration of postoperative analgesics. A new trial design allows evaluation of baseline pain, demonstrates large variation in need of analgesics, and finds no analgesic effect of IA ketamine compared with IA saline. *Pain* **2003**; 104: 25–34
- Rosseland LA, Stubhaug A.** Gender is a confounding factor in pain trials: women report more pain than men after arthroscopic surgery. *Pain* **2004**; 112: 248–53
- Rosseland LA.** No evidence for analgesic effect of intra-articular morphine after knee arthroscopy: a qualitative systematic review. *Reg Anesth Pain Med* **2005**; 30: 83–98
- Rothschild JM, Orav EJ, Burdick E, Thompson S, Hayes J, Bates DW.** A controlled trial of smart infusion pumps to improve medication safety in critically ill patients. *Crit Care Med* **2005**; 33:533-540.
- Royse C, Royse A, Soeding P.** Prospective randomized trial of high thoracic epidural analgesia for coronary artery bypass surgery. *Ann Thorac Surg* **2003**; 75: 93–100.
- Ryu JH, Kang MH, Park KS, Do SH.** Effects of magnesium sulphate on intraoperative anaesthetic requirements and postoperative analgesia in gynaecology patients receiving total intravenous anaesthesia. *Br J Anaesth* **2008**; 100:397–403.
- Saito M, Okutomi T, Kanai Y, Mochizuki J, Tani A, Amano K, Hoka S.** Patient-controlled epidural analgesia during labor using ropivacaine and fentanyl provides better maternal satisfaction with less local anesthetic requirement. *J Anesth* **2005**; 19:208-212.

- Salerno A, Hermann R.** Efficacy and safety of steroid use for postoperative pain relief. Update and review of the medical literature. *J Bone Joint Surg Am* **2006**; 88:1361–1372.
- Samad TA, Moore KA, Sapirstein A.** Interleukin-1beta-mediated induction of Cox-2 in the CNS contributes to inflammatory pain hypersensitivity. *Nature* **2001**; 410: 471-475.
- Scholten W, Nygren-Krug H, Zucker HA.** The World Health Organization paves the way for action to free people from the shackles of pain. *Anesth Analg* **2007**; 105: 1–4
- Scott NB, Turfrey DJ, Ray DAA.** A prospective randomized study of the potential benefits of thoracic epidural anesthesia and analgesia in patients undergoing coronary artery bypass grafting. *Anesth Analg* **2001**; 93: 528–535.
- Senturk M, Ozcan PE, Talu GK.** The effects of three different analgesia techniques on long-term postthoracotomy pain. *Anesthesia and Analgesia* **2002**; 94: 11-15.
- Silvasti M, Pitkanen M.** Patient-controlled epidural analgesia versus continuous epidural analgesia after total knee arthroplasty. *Acta Anaesthesiol Scand* **2001**; 45:471-476.
- Simonnet G, Rivat C.** Opioid-induced hyperalgesia: abnormal or normal pain? *Neuroreport* **2003**; 14(1):1–7.
- Sinatra RS, Torres J, Bustos AM.** Pain management after major orthopaedic surgery: Current strategies and new concepts. *J Am Acad Orthop Surg* **2002**; 10: 117-129
- Singelyn FJ, Seguy S, Gouverneur JM.** Interscalene brachial plexus analgesia after open shoulder surgery: Continuous versus patient-controlled infusion. *Anesth Analg* **1999**; 89:1216-1220.
- Smirnov G, Terava M, Tuomilehto H.** Etoricoxib for pain management during thyroid surgery: a prospective, placebo-controlled study. *Otolaryngol Head Neck Surg* **2008**; 138:92–97.
- Smythe M, Loughlin K, Schad RF.** Patient-controlled analgesia versus intramuscular analgesic therapy. *Am J Hosp Pharm* **1994**; 51:1433- 40.
- Stein C.** The control of pain in peripheral tissue by opioids. *N Engl J Med* **1995**; 332:1685-90.
- Stoelting R.** Pharmacology and physiology in anesthetic practice. 3rd ed. Philadelphia: Lippincott Williams and Wilkins; **1999**. p.78-112.

- Strichartz GR.** Novel ideas of local anaesthetic actions on various ion channels to ameliorate postoperative pain. *Br J Anaesth* **2008**; 101: 45–7
- Stubhaug A, Breivik H, Kreunen M.** Mapping of punctuates hyperalgesia around a surgical incision demonstrates that ketamine is a powerful suppressor of central sensitization to pain following surgery. *Acta Anaesthesiol Scand* **1997**; 41: 1124–32
- Stubhaug A, Breivik H.** Clinical trials—acute and chronic pain. In: Breivik H, Campbell W, Nicholas M, eds. *Clinical Management of Pain—Practice and Procedures*, 2nd Edn. London: Arnold, **2008**; pp 1562- 1591.
- Stubhaug A, Breivik H.** Prevention and treatment of hyperalgesia and persistent pain after surgery. In: Breivik H, Shipley M, eds. *Pain Best Practice and Research Compendium*. London: Elsevier, **2007**; 281–8
- Subramaniam K, Subramaniam B, Steinbrook RA.** Ketamine as adjuvant analgesic to opioids: a quantitative and qualitative systematic review. *Anesth Analg* **2004**; 99:482–495.
- Terman GW, Bonica JJ.** Spinal mechanisms and their modulation. In: Loeser JD, editor. *Management of pain*. 3rd edition. New York: Lippincott Williams & Wilkins; **2001**. p. 73–152.
- Thipphawong JB, Babul N, Otulana BA.** Analgesic efficacy of inhaled morphine in patients after bunionectomy surgery. *Anesthesiology* **2003**; 99:693-700.
- Toussaint S, Maidl J, Striebel HW.** Patient-controlled intranasal analgesia: Effective alternative to intravenous PCA for postoperative pain relief. *Can J Anesth* **2000**; 47:299-302.
- Tsang J, Brush B.** Patient-controlled analgesia in postoperative cardiac surgery. *Anaesth Intensive Care* **1999**; 27: 464–470.
- Tufanogullari B, White PF, Peixoto MP.** Dexmedetomidine infusion during laparoscopic bariatric surgery: the effect on recovery outcome variables. *Anesth Analg* **2008**; 106:1741–1748.
- Venkateswaran R and Prasad KN:** Management of post operative pain. *Indian J Anesth*; 50 (5): 345-354, **2006**.
- Vicente KJ, Hillel G, Cassano A, Orser BA.** Programming errors contribute to death from patient-controlled analgesia: Case report and estimate of probability. *Can J Anaesth* **2003**; 50:328- 332.

- Vierck CJ Jr, Cannon RL, Fry G.** Characteristics of temporal summation of second pain sensations elicited by brief contact of glabrous skin by a preheated thermode. *J Neurophysiol* **1997**; 78(2):992–1002.
- Vintar N, Pozlep G, Rawal N, Godec M, Rakovec S.** Incisional self-administration of bupivacaine or ropivacaine provides effective analgesia after inguinal hernia repair. *Can J Anaesth* **2002**; 49:481-486.
- Vintar N, Rawal N, Veselko M.** Intraarticular patient- controlled regional anesthesia after arthroscopically assisted anterior cruciate ligament reconstruction: Ropivacaine/morphine/ketorolac versus ropivacaine/morphine. *Anesth Analg* **2005**; 101:573- 578.
- Viscusi ER.** Patient-Controlled Drug Delivery for Acute Postoperative Pain Management: A Review of Current and Emerging Technologies. *Reg Anesth Pain Med* **2008**; 33:146-158.
- Walder B, Schafer M, Henzi I, Tramer MR.** Efficacy and safety of patient-controlled opioid analgesia for acute postoperative pain. *Acta Anaesthesiol Scand* **2001**; 45:795-804.
- Walder B, Schafer M, Henzi I, Tramer MR.** Efficacy and safety of patient-controlled opioid analgesia for acute postoperative pain. A quantitative systematic review. *Acta Anaesthesiol Scand* **2001**; 45:795-804.
- Wallace MS, Wallace AM, Lee J, Dobke MK.** Pain after breast surgery: a survey of 282 women. *Pain* **1996**; 66: 195-205.
- Wang JJ, Sung KC, Huang JF, Yeh CH, Fang JY.** Ester prodrugs of morphine improve transdermal drug delivery: a mechanistic study. *J Pharm Pharmacol.* **2007** Jul;59(7):917-25.
- Webster LR.** Fentanyl buccal tablets. *Expert Opin Investig Drugs* **2006**; 15:1469-1473.
- Wei F, Dubner R, Ren K.** Dorsolateral funiculus-lesions unmask inhibitory or disfacilitatory mechanisms which modulate the effects of innocuous mechanical stimulation on spinal Fos expression after inflammation. *Brain Res* **1999**; 820(1–2):112–6.
- Wermeling DP, Miller JL, Rudy AC.** Systematic intranasal drug delivery: Concepts and applications. *Drug Deliv Technol* **2002**; 2:56-62.
- Werner MU, Soholm L, Rotboll-Nielsen P, Kehlet H.** Does an acute pain service improve postoperative outcome? *Anesth Analg* **2002**; 95: 1361–1372

- White PF, Rawal S, Latham P.** Use of a continuous local anesthetic infusion for pain management after median sternotomy. *Anesthesiology* **2003**; 99: 918–923.
- White PF, Schuttler J, Shafer A.** Comparative pharmacology of the ketamine isomers: studies in volunteers. *Br J Anaesth* **1985**; 57: 197–203.
- White PF, Way WL, Trevor AJ.** Ketamine: its pharmacology and therapeutic uses. *Anesthesiology* **1982**; 56: 119–136.
- Woolf CJ, Salter MW.** Neuronal plasticity: increasing the gain in pain. *Science* **2000**; 288(5472):1765–9.
- Woolf CJ.** Windup and central sensitization are not equivalent. *Pain* **1996**; 66(2–3): 105–8.
- Wright D, Paterson C, Scott N et al.** Five-year follow-up of patients undergoing laparoscopic or open groin hernia repair: a randomized controlled trial. *Annals of Surgery* **2002**; 235: 333–337.
- Wu C, Boustany L, Liang H, Brennan TJ.** Nerve growth factor expression after plantar incision in the rat. *Anesthesiology* **2007**; 107: 128–35
- Wu CL, Cohen SR, Richman JM, Cheung K, Lin EE, Liu SS.** Efficacy of postoperative patient-controlled and continuous infusion epidural analgesia versus intravenous patient-controlled analgesia with opioids: A meta-analysis. *Anesthesiology* **2005**; 103:1079–1088.
- Zafar SU, Hamid M, Hoda MQ.** Patient controlled intravenous analgesia (PCIA) in postoperative surgical patients: An audit. *J Pak Med Assoc* **2004**; 54: 353–356.
- Zakine J, Samarcq D, Lorne E.** Postoperative ketamine administration decreases morphine consumption in major abdominal surgery: a prospective, randomized, double-blind, controlled study. *Anesth Analg* **2008**; 106:1856–1861.
- Zisman E, Shenderoy A, Ammar R.** The effects of intrathecal morphine on patients undergoing minimally invasive direct coronary artery bypass surgery. *J Cardiothorac Vasc Anesth* **2005**; 19: 40–43.
- Zohar E, Fredman B, Phillipov A, Jedeikin R, Shapiro A.** The analgesic efficacy of patient-controlled bupivacaine wound instillation after total abdominal hysterectomy with bilateral salpingo-oophorectomy. *Anesth Analg* **2001**; 93:482–487.