

-
- Alper A. and Calhoun D. (2002):** Hypertensive emergencies. In: Antman EM, editor. Cardiovascular therapeutics: a companion to Braunwald's Heart disease. Philadelphia, PA: W.B. Saunders Co; 4(3): 615-27.
- Andersson C., Lithner F., Wikberg A. and Stegmayr B. (2000):** Hypertension and renal impairment in patients with acute intermittent porphyria—a population-based study. *J Hypertens*; 18(4): 98.
- Aronson S., Boisvert D. and Lapp W. (2002):** Isolated systolic hypertension is associated with adverse outcomes from coronary artery bypass grafting surgery. *Anesth Analg*; 94 (5): 1079-84.
- Aronson S., Boisvert D. and Lapp W. (2002):** Isolated systolic hypertension is associated with adverse outcomes from coronary artery bypass grafting surgery. *Anesth Analg*; 94: 1079.
- Attoub S., Gaben A.M., Al-Salam S., et al. (2008):** Captopril as a potential inhibitor of lung tumor growth and metastasis. *Ann N.Y Acad Sci*; 1138: 65–72. doi:10.1196/annals.1414.011.
- Bailey M.A., Paterson J.M., Hadoke P.W., et al. (2008):** A switch in the mechanism of hypertension in the syndrome of apparent mineralocorticoid excess. *Journal of the American Society of Nephrology*; 19(1): 47–58.
- Barba G., Cappuccio F.P., Russ L., et al. (1996):** Renal function and blood pressure response to dietary salt restriction in normotensive men. *Hypertension*; 27: 1160-4.
- Beltran A., McVeigh G., Morgan D., et al. (2001):** Arterial compliance abnormalities in isolated systolic hypertension. *Am J Hypertens*; 14: 1007–1011.
- Berends N., Teunkens A., Vandermeersch E. and Van de Velde M. (2005):** A randomized trial comparing low-dose combined spinal-epidural anesthesia and conventional epidural anesthesia for cesarean section in severe preeclampsia. *Acta Anaesthesiol Belg*; 56: 155-62.
- Berge K.H. and Lanier W.L. (1991):** Myocardial infarction accompanying acute clonidine withdrawal in a patient without a history of ischemic coronary artery disease. *Anesth Analg*; 72: 259–61.
-

-
- Biondi B., Fazio S., Palmieri E.A., et al. (1999):** Left ventricular diastolic dysfunction in patients with subclinical hypothyroidism. *J Clin Endocrinol Metab*; 84: 2064–2067.
- Borzecki A.M., Wong A.T., Hickey E.C., et al. (2003):** Hypertension control: how well are we doing? *PMID: 14662624* .
- Brem A.S. (2001):** Insights into glucocorticoid-associated hypertension. *Am J Kidney Dis*; 37: 1–10.
- Brizio-Molteni L., Molteni A., Cloutier L.C. and Rainey S. (1979):** Incidence of post burn hypertensive crisis in patients admitted to two burn centers and a community hospital in the United States. *Scand J Plast Reconstr Surg*; 13: 21– 28.
- Browne I.M., Birnbach D.J., Stein D.J., et al. (2005):** A comparison of Espocan and Tuohy needles for the combined spinal–epidural technique for labor analgesia. *Anesth Analg*; 101: 535–540.
- Browner W.S., Li J. and Mangano D.T. (1992):** In-hospital and long-term mortality in male veterans following noncardiac surgery. The Study of Perioperative Ischemia Research Group. *JAMA*; 268: 228–32.
- Brunton L., Parker K., Blumenthal D. and Buxton I. (2007):** Therapy of hypertension. *Goodman & Gilman's Manual of Pharmacology and Therapeutics*. New York: McGraw-Hill: 544–6.
- Burnier M. (2001):** Angiotensin II type 1 receptor blockers. *Circulation*; 103: 904-912.
- Cada D.J., Levien T.L. and Baker D.E. (2004):** Clevidipine butyrate injectable emulsion. *Hosp Pharm*; 43(11): 903-12.
- Campbell M. (1970):** Natural history of coarctation of the aorta. *Br Heart J*; 32: 633–640.
- Campese V.M., Tawadrous M., Bigazzi R et al. (1996):** Salt intake
- Cardillo C., Kilcoyne C.M., Cannon R.O. and Panza J.A. (2000):** Inter-actions between nitric oxide and endothelin in the regulation
-

of vascular tone of human resistance vessels in vivo. *Hypertension*; 35: 1237–1241.

Carretero O.A and Oparil S. (2000): "Essential hypertension. Part I: definition and etiology". *Circulation* 101 (3): 329–35. PMID 10645931.

Carretero O.A., Oparil S. (2000): Essential hypertension. Part I: definition and etiology. *Circulation*; 101(3): 329–35.

Casadei B. and Abuzeid H. (2005): Is there a strong rationale for deferring elective surgery in patients with poorly controlled hypertension?. *J Hypertens*; 23: 19.

Chancellor M.B., Erhard M.J., Hirsch I.H. and Stass W.E. (1994): Prospective evaluation of terazosin for the treatment of autonomic dysreflexia. *J Urol*; 151: 111–113.

Chinn S. and Rona R.J. (2001): Prevalence and trends in overweight and obesity in three cross sectional studies of British Children, 1974–94. *BMJ*; 322: 24–26.

Chobanian A.V., Bakris G.L., Black H.R., et al. (2003): Seventh report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. *Hypertension*; 42: 1206.

Clase C.M. and Kiberd B.A. (2000): Prevalence of renal insufficiency in nondiabetic adults: Third National Health and Nutrition Examination Survey (NHANES III). *J Am Soc Nephrol*; 11: 58.

Colao A., Baldelli R., Marzullo P., et al. (2000): Systemic hypertension and impaired glucose tolerance are independently correlated to the severity of the acromegalic cardiomyopathy. *J Clin Endocrinol Metab*; 85: 193–199.

Collins A., Xue J.L., Ma J.Z. and Louis T. (2000): Estimating the number of patients and Medicare cost for end stage renal disease in the U.S. to the year 2010. *J Am Soc Nephrol*; 11: 142.

-
- Coresh J., Wei L., McQuillan G., et al. (2001):** Prevalence of high blood pressure and elevated serum creatinine level in the United States. *Arch Intern Med*; 161: 1207–1216.
- Coriat P, Richer C, Douraki T, et al. (1994):** Influence of chronic angiotensin-converting enzyme inhibition on anesthetic induction. *Anesthesiology*; 81: 299.
- Cosentino F. and Lüscher T.F. (2001):** Effects of blood pressure and glucose on endothelial function. *Curr Hypertens Rep*; 3: 79–88.
- Cowley A.W. (1992) :** Long term control of arterial blood
- Cunningham F.G., Veno K.J., Bloom S.L., et al. (2010):** Pregnancy Hypertension. In: *Williams Obstetrics*. 23e.
- Cygan R and Waitzkin H. (1987):** Stopping and restarting medications in the perioperative period. *J Gen Intern Med*; 2: 270.
- Cygan R. and Waitzkin H. (2006):** Stopping and restarting medications in the Perioperative period. *J Gen Intern Med*; 2: 270-283.
- Dasgupta I., Porter C., Innes A. and Burden R. (2007):** Benign" hypertensive nephrosclerosis". *QJM*; 100(2): 113–9.
- Davies S.J.C., Ghahramani P., Jackson P.R., et al. (1999):** Association of panic disorder and panic attacks with hypertension. *Am J Med*; 107: 310–316.
- Devereux R.B., Case D.B., Alderman M.H., et al. (2000):** Possible role of increased blood viscosity in the hemodynamics of systemic hypertension. *Am J Cardiol*; 85: 1265–1268.
- Duchna H.W., Guilleminault C., Stoohs R.A., et al. (2000):** Vascular reactivity in obstructive sleep apnea syndrome. *Am J Respir Crit Care Med*; 161: 187–191.
- Duley L., Abalos E., Steyn D.W., and Henderson-Smart D.J. (2006):** Antihypertensive drug therapy for mild to moderate hypertension during pregnancy. *Cochrane Database of Systematic*
-

-
- Reviews; Art. No.: CD002252. DOI: 10.1002/14651858.CD002252.
- Duley L., Gulmezoglu A.M. and Henderson-Smart D.J. (2006):** Magnesium sulphate and other anticonvulsants for women with pre-eclampsia. *Cochrane Database Syst. Rev.*; 3: CD000025.
- Dyer R.A., Piercy J.L., Reed A.R., Lombard C.J. et al. (2008):** Hemodynamic changes associated with spinal anesthesia for cesarean delivery in severe preeclampsia. *Anesthesiology*; 108: 802-811.
- Eagle K.M., Berger P.B., Calkins H., et al. (2002):** ACC/AHA Guideline Update for Perioperative Cardiovascular Evaluation for Noncardiac Surgery- Executive Summary. *Circulation*; 105: 1257.
- Ehrenberg H.M., and Mercer B.M. (2006):** Abbreviated postpartum magnesium sulfate therapy for women with mild preeclampsia: a randomized controlled trial. *Obstet Gynecol*; 108: 833-8.
- Eltzchig H.K., Lieberman E.S., Camann W.R. (2003):** Regional anesthesia and analgesia for labor and delivery. *N Engl J Med*; 348-319.
- Ernst A.A., Weiss S.J., Johnson W.D. and Takakuwa K.M. (2000):** Blood pressure in acute vaso-occlusive crises of sickle cell disease. *South Med J*; 93: 590–592.
- Fagot-Campagna A., Narayan K.M.V. and Imperatore G. (2001):** Type 2 diabetes in children. *BMJ*; 322: 377–378.
- Fleisher L.A. (2002):** Preoperative evaluation of the patient with hypertension. *JAMA*; 287: 2043.
- Fleisher L.A. (2002):** Preoperative evaluation of the patient with hypertension. *JAMA*; 287: 2043.
- Fletcher A.K., Weetman A.P. (1998):** Hypertension and hypothyroid-ism. *J Hum Hypertens*; 12: 79–82.
- Foex P., Meloche R., Prys-Roberts C. (1971):** Studies of anaesthesia in relation to hypertension. Pulmonary gas exchange during spontaneous ventilation. *Br J Anaesth*; 43: 644.
-

-
- Fontana F., Bernardi P., Tartuferi L., et al. (2000):** Mechanisms of hypertension in patients with chronic obstructive pulmonary disease and acute respiratory failure. *Am J Med*; 109: 621–627.
- Forrest J.B., Rehder K., Cahalan M.K., et al. (1992):** Multicenter study of general anesthesia. III. Predictors of severe perioperative adverse outcomes. *Anesthesiology*; 76: 3–15.
- Fraser R., Davies D.L., Connel J.M.C. (1989):** Hormones and hypertension. *Clin Endocrinol*; 31: 701–746.
- Frohlich E.D. (1998):** Hypertension: evaluation and treatment. Baltimore, Md. *Williams and Wilkins*; p. 432.
- Fugh-Berman A. (2000): Herb-drug interactions. *Lancet*; 355: 134–138.
- Funck-Brentano C., Pagny J.Y. and Menard J. (1987):** Neurogenic hyper-tension associated with an excessively high excretion rate of catecholamine metabolites. *Br Heart*; 57: 487–489.
- Gajwani P., Muzina D., Gao K. and Calabrese J.R. (2006):** Awareness under anesthesia during electroconvulsive therapy treatment. *J ECT* 22(2): 158–9.
- Ganzevoort W., Rep A., Bonsel G.J., Fetter W.P., et al. (2005):** A randomised controlled trial comparing two temporising management strategies, one with and one without plasma volume expansion, for severe and early onset pre-eclampsia. *BJOG*; 112: 1358-68.
- Giacchetti G., Turchi F., Boscaro M. and Ronconi V. (2009):** Management of primary aldosteronism: its complications and their outcomes after treatment. *Current Vascular Pharmacology*; 7(2): 244–49.
- Gifford R.W., Manger W.M., and Bravo E.L. (2002):** Pheochromocytoma. *Endocrinol Metab Clin North Am*; 23: 387–404.
-

-
- Goldberg M.E. Larijani G.E. (1998):** Perioperative hypertension. *Pharmacotherapy*;18: 911–14.
- Goldman, L. and Caldera D.L. (1979):** Risks of general anesthesia and elective operation in the hypertensive patient. *Anesthesiology*; 50: 285.
- Gratacòs M., Nadal M., Martín-Santos R., et al. (2001):** A polymorphic genomic duplication on human chromosome 15 is a susceptibility factor for panic and phobic disorders. *Cell*; 106(3): 367–379.
- Greenstein R.J., Krakoff L.R. and Felton K. (1987):** Activation of the renin system in acute pancreatitis. *Am J Med*; 82: 401–404.
- Gross P. (2008): Polycystic kidney disease: will it become treatable? *Polskie Archiwum Medycyny Wewntrzej*; 118(5): 298–301.
- Grossman E., Messerli F.H., Grodzicki T. and Kowey P. (1996):** Should a moratorium be placed on sublingual nifedipine capsules given for hypertensive emergencies and pseudoemergencies?. *JAMA*; 276: 1328–31.
- Grote L., Hedner J., Peter J.H. (2000):** Sleep-related breathing disorder is an independent risk factor for uncontrolled hypertension. *J Hypertens*; 18: 679–685.
- Guenthard J. and Wyler F. (1995):** Exercise-induced hypertension in the arms due to impaired arterial reactivity after successful coarctation resection. *Am J Cardiol*; 75: 814–817.
- Hall E., Guyton M., and Arthur C. (2006):** Textbook of medical physiology. *St. Louis, Mo: Elsevier Saunders. P. 228.*
- Halpern N.A., Alicea M., Krakoff L.R., et al. (1990):** Postoperative hypertension: a prospective, placebo-controlled, randomized, double-blind trial, with intravenous nicardipine hydrochloride. *Angiology*; 41: 992–1004.
-

-
- Head B.B., Owen J., Vincent R.D., Shih G., et al (2002):** A randomized trial of intrapartum analgesia in women with severe preeclampsia. *Obstet Gynecol*; 99: 452-7.
- Hoekstra J.W. and Qureshi A. (2008):** Management of Hypertension and Hypertensive Emergencies in the Emergency Department: The EMCREG-International Consensus Panel Recommendations. *Ann Emerg Med*; 51(3): 1-38.
- Horlocker, et al. (2003):** Regional Anesthesia in the Anticoagulated
- Howell S.J., Sear Y.M., Yeates D., et al. (1996):** Hypertension, admission blood pressure and perioperative cardiovascular risk. *Anaesthesia*; 51: 1000.
- Idama T.O. and Lindow S.W. (1998):** Magnesium sulphate: a review of clinical pharmacology applied to obstetrics. *Br J Obstet Gynaecol*; 105: 260-8.
- Isler C.M., Barrilleaux P.S., Rinehart B.K. et al, (2003):** Postpartum seizure prophylaxis: using maternal clinical parameters to guide therapy. *Obstet Gynecol*; 101: 66-9.
- Ito Y., Obara T., Okamoto T., et al. (1998):** Efficacy of single-voided urine metanephrine and normetanephrine assay for diagnosing pheochromocytoma. *World J Surg*; 22: 684–688.
- J.C.P. Kingdom (eds.) (2004):** Pre-eclampsia: Current Perspectives on Management. *London: Parthenon*: 25–39.
- Kang A. and Struben H. (2008):** Pre-eclampsia screening in first and second trimester (in German). *Therapeutische Umschau. Revue Thérapeutique*; 65(11): 663–6.
- Kaplan N.M. (2006):** Hypertensive crises. In: Kaplan's Clinical Hypertension, 9th ed. Neal, W (Ed), Lippincott, Williams and Wilkins, Baltimore: 311.
- Kaplan N.M. (2006):** Treatment of hypertension: Drug therapy. In: Kaplan's Clinical Hypertension, 9th ed. Lippincott, Williams & Wilkins, Baltimore. p.290
-

-
- Karras D.J., Kruus L.K., Cienki J.J., et al. (2006):** Evaluation and treatment of patients with severely elevated blood pressure in academic Emergency Departments: A multicenter study. *Ann Emerg Med*; 47: 230-236.
- Kinney M.A.O., Warner M.E., vanHeerden J.A., et al. (2000):** Perianesthetic risks and outcomes of pheochromocytoma and paraganglioma resection. *Anesth Analg*; 91: 1118–1123.
- Kitaguchi H., Tomimoto H., Miki Y., et al. (2005):** A brainstem variant of reversible posterior leukoencephalopathy syndrome. *Neuroradiol*; 47: 652.
- Kosch M., Hausberg M., Barenbrock M., et al. (2000):** Impaired endothelial function in primary hyperparathyroidism improves after successful parathyroidectomy—a prospective study. *J Hypertens*; 18(2): 144.
- Krum H., Louis W.J., Brow D.J. and Howes L.G. (1992):** Pressor dose responses and baroreflex sensitivity in quadriplegic spinal cord injury patients. *J Hypertens*; 10: 245–250.
- Lavoie J.L. and Sigmund C.D. (2003):** Mini review: overview of the renin-angiotensin system—an endocrine and paracrine system. *Endocrinology*; 144: 2179-2183.
- Lester S.J., Baggott M., Welm S., et al. (2000):** Cardiovascular effects of 3,4-methylenedioxymethamphetamine: a double-blind, placebo-controlled trial. *Ann Intern Med*; 133: 969–973.
- Lew M. and Klonis E. (2003):** Emergency management of eclampsia and severe pre-eclampsia. *Emerg Med (Fremantle)*; 15(4): 361-8.
- Lewis G. (ed.) (2007):** The Confidential Enquiry into Maternal and Child Health (CEMACH). Saving Mothers' Lives: reviewing maternal deaths to make motherhood safer – 2003-2005. *The Seventh Report on Confidential Enquiries into Maternal Deaths in the United Kingdom*, CEMACH, London.
-

-
- Liard J.F. and Spadone J.C. (1985):** Regional circulations in experimental coarctation of the aorta in conscious dogs. *J Hypertens*; 3: 281–291.
- Lindholm J., Juul S., Lorgensen J.O.L., et al. (2001):** Incidence and late prognosis of Cushing's syndrome. *J Clin Endocrinol Metab*; 86: 117–123.
- Lloyd-Mostyn R.H. and Oram S. (1975):** Modification by propranolol of cardiovascular effects of induced hypoglycemia. *Lancet*; 1: 1213–1215.
- Loyola R.N., Dreher H. and Michael R.N. (2003):** Management of Pharmacologically Induced Neuromuscular Blockade Using Peripheral Nerve Stimulation. *Anesth Analg*; 22(4): 157-164.
- Lucas M.J., Sharma S.K., McIntire D.D., et al (2001):** A randomized trial of labor analgesia in women with pregnancy-induced hypertension. *Am J Obstet Gynecol*; 185: 970-5.
- Lumachi F., Behboo R., Scarpa M., et al. (2000):** The effect of serum calcium and serum PTH on arterial blood pressure in patients with primary hyperparathyroidism. Population based study in 194 patients undergoing parathyroidectomy. *Endo Soc*; 421.
- Lyketsos C.G., Garrett E., Liang K.Y. and Anthony J.C. (1999):** Cannabis use and cognitive decline in persons under 65 years of age. *Am J Epidemiol*; 149: 794–800.
- Madejczyk M., Kruszynski G. and Breborowicz G. (2009):** Etiopathology of preeclampsia. *Arch Perinat Med*; 15(3): 144-151.
- Magee L. and Sadeghi S. (2005):** Prevention and treatment of postpartum hypertension. *Cochrane Database of Systematic Reviews* (1); Art. No.: CD004351. DOI: 10.1002/14651858.CD004351.
- Maison P., Démolis P., Young J., et al. (2000):** Vascular reactivity in acromegalic patients: preliminary evidence for regional
-

- endothelial dysfunction and increased sympathetic vasoconstriction. *Clin Endocrinol*; 53: 445–451.
- Malek A.M., Alper S.L. and Izumo S. (1999):** Hemodynamic shear stress and its role in atherosclerosis. *JAMA*; 282: 2035–2042.
- Mangano D.T., Layug E.L., Wallace A. and Tateo I. (1996):** Effect of atenolol on mortality and cardiovascular morbidity after noncardiac surgery. *N Engl J Med*; 335: 1713.
- Mangos G.J., Walker B.R., Kelly J.J., et al. (2000):** Cortisol inhibits cholinergic vasodilatation in the human forearm. *Am J Hypertens*; 13: 1155–1160.
- Mann S.J. and Atlas S.A. (1995):** Hypertensive emergencies. In: Laragh JH, Brenner BM, editors. Hypertension: pathophysiology, diagnosis, and management. *New York: Raven Pr*; 745.
- Mann S.J. and Pickering T.G. (1992):** Detection of renovascular hypertension. State of the art. *Ann Intern Med*; 117: 845–853.
- Marcisz C., Jonderko G. and Jucharz E.J. (2001):** Influence of short-time application of a low sodium diet on blood pressure in patients with hyperthyroidism or hypothyroidism during therapy. *Am J Hypertens*; 14: 995–1002.
- Marik P.E. (2009):** Hypertensive disorders of pregnancy. *Postgraduate Medicine*; 121(2): 69–76.
- McKillion P.C. and Dellinger R.P. (1997):** Hypertensive emergencies and urgencies. In: Civetta JM, Taylor RW, Kirby RR, editors. Critical care. *Philadelphia: Lippincott-Raven*; 245.
- McMurray J.J. (2010):** Systolic heart failure. *N Engl J Med*; 362: 228.
- Metz S., Klein C. and Morton N. (1987):** Rebound hypertension after discontinuation of transdermal clonidine therapy. *Am J Med*; 82: 17.
- Milne F., Redman C., Walker J., et al. (2005):** The pre-eclampsia community guideline (PRECOG): how to screen for and detect onset of pre-eclampsia in the community. *BMJ*; 330: 576–80.

-
- Minami N., Imai Y., Miura Y. and Abe K. (1995):** The mechanism responsible for hypertension in a patient with Guillain-Barré syndrome. *Clin Exp Hypertens*; 17: 607–617.
- Mizuri S., Amagasaki Y., Hosaka H., et al. (1992):** Hypertension in unilateral atrophic kidney secondary to ureteropelvic junction obstruction. *Nephron*; 61: 217–219.
- Montague C.T. and O'Rahilly S. (2000):** The perils of portliness: causes and consequences of visceral adiposity. *Diabetes*; 49: 883–888.
- Morris M.C., Scherr P.A., Herbert L.E., et al. (2000):** The cross-sectional association between blood pressure and Alzheimer's disease in a biracial community population of older persons. *J Gerontol*; 55: 130–136
- Mounier-Vehier C. and Delsart P. (2009):** Pregnancy-related hypertension: a cardiovascular risk situation (in French). *Presse Médicale*; 38(4): 600–8.
- Naftchi N.E., Demeny M., Lowman E.W. and Tuckman J. (1978):** Hypertensive crises in quadriplegic patients. Changes in cardiac output, blood volume, serum dopamine-b-hydroxylase activity, and arterial prostaglandin PGE₂. *Circulation*; 57: 336–41.
- National Task Force on the Prevention and Treatment of Obesity (2000):** Overweight, obesity, and health risk. *Arch Intern Med*; 160: 898–904.
- Newsome L.R., Bramwell R.S. and Curling P.E., (1986):** Severe preeclampsia: hemodynamic effects of lumbar epidural anesthesia. *Anesth Analg*; 65: 31-6.
- Ng T.M., Shurmur S.W., Silver M., et al. (2006):** Comparison of N-acetylcysteine and fenoldopam for preventing contrastinduced nephropathy (CAFCIN). *Int J Cardiol*; 32: 56-58.
- Nørgaard K., Rasmussen E., Jensen T., et al. (1993):** Nature of elevated blood pressure in normoalbuminuric type I diabetic patients. Essential hypertension?. *Am J Hypertens*; 6: 830–836.
-

-
- Norwitz E.R., Hsu C.D. and Repke J.T. (2002):** Acute complications of preeclampsia. *Clin Obstet Gynecol*; 45: 308-29.
- Nzerue C.M., Hewan-Lowe K. and Riley L.J. (2000):** Cocaine and the kidney: a synthesis of pathophysiologic and clinical perspectives. *Am J Kidney Dis*; 35: 783–795.
- O'Brien E., Asmar R., Beilin L., Imai Y., et al. (2003):** on behalf of the European Society of Hypertension Working Group on Blood Pressure Monitoring et al. European Society of Hypertension Recommendations for Conventional, Ambulatory and Home Blood Pressure Measurement. *J Hypertens*; 21: 821–848.
- on Neuraxial Anesthesia and Anticoagulation). *Regional Anesthesia and*
- Oparil S., Zaman M.A. and Calhoun D.A. (2003):** Pathogenesis of hypertension. *Ann. Intern Med*; 139(9): 761–76.
- Pain Medicine*; 28: 172-97.
- Pallini R., Lauretti L. and Fernandez E. (1995):** Chronic arterial hyper-tension as unique symptom of brainstem astrocytoma. *Lancet*; 345: 1573.
- Parati G., Stergiou G.S., Asmar R., Bilo G., et al. (2008):** ESH Working Group on Blood Pressure Monitoring et al. European Society of Hypertension guidelines for blood pressure monitoring at home: a summary report of the Second International Consensus Conference on Home Blood Pressure Monitoring. *J Hypertens*; 26: 1505–1526.
- Paterniti S., Alperovitch A., Ducimetière P., et al. (1999):** Anxiety but not depression is associated with elevated blood pressure in a community group of French elderly. *Psychosom Med*; 61: 77–83.
- Patient: Defining the Risks (The Second ASRA Consensus Conference
- Peppard P.E., Young T., Palta M. and Skatrud J. (2000):** Prospective study of the association between sleep-disordered breathing and hypertension. *N Engl J Med*; 342: 1378–1384.
-

Pharm. 2009; 66(1):1343-52.

Pharm. 2009;66:1448-57.

Plets C. (1989): Arterial hypertension in neurosurgical emergencies. *Am J Cardiol*; 63: 40–42.

Podymow T. and August P. (2008): Update on the use of antihypertensive drugs in pregnancy. *Hypertension*; 51: 960-9.

Poldermans D., Boersma E., Bax J.J., et al. (1999): The effect of bisoprolol on perioperative mortality and myocardial infarction in high-risk patients undergoing vascular surgery. Dutch Echocardiographic Cardiac Risk Evaluation Applying Stress Echocardiography Study Group. *N Engl J Med*; 341:1789.

Portaluppi F., Cortelli P., Avoni P., et al. (1994): Diurnal blood pressure variation and hormonal correlates in fatal familial insomnia. *Hypertension*; 23: 569–576.

pressure. *Physiol Rev*; 72: 231-300.

Psaty B.M., Koepsell T.D., Wagner E.H., et al. (1990): The relative risk of incident coronary heart disease associated with recently stopping the use of beta-blockers. *JAMA*; 263: 1653.

Ram C.V., Holland O.B., Fairchild C. and Gomez-Sanchez C.E. (1979): Withdrawal syndrome following cessation of guanabenz therapy. *J Clin Pharmacol*; 19: 148.

Ram C.V.S. and Fenves A. (2002): Clinical pharmacology of antihypertensive drugs. *Cardiol Clin*; 20(2): 265-8.

Rao P.S. (1995): Coarctation of the aorta. *Semin Nephrol*; 15: 87–105.

Rao P.S. (1995): Coarctation of the aorta. *Semin Nephrol*; 15: 87–105.

Reaven G.M., Lithell H. and Landsberg L. (1996): Hypertension and associated metabolic abnormalities the role of insulin resistance and the sympathoadrenal system. *N Engl J Med*; 334: 374–381.

-
- Rhoney D. and Peacock W.F. (2009):** Intravenous therapy for hypertensive emergencies, part 1. *Am J Health-Syst Pharm*; 66(1): 1343-52.
- Rhoney D., Peacock W.F. (2002):** Intravenous therapy for hypertensive emergencies, part 2. *Am J Health-Syst*; 2: 654:6.
- Roberts J.M., Pearson G., Cutler J. and Lindheimer M. (2003):** Summary of the NHLBI working group on research on hypertension during pregnancy. *Hypertension*; 41: 437– 445.
- Robertson D., Hollister A.S., Biaggioni I., et al. (1993):** The diagnosis and treatment of baroreceptor failure. *N Engl J Med*; 329: 1449–1455.
- Rodriguez-Thompson D. and Lieberman E.S. (2001):** Use of a random urinary protein-to-creatinine ratio for the diagnosis of significant proteinuria during pregnancy. *Am J Obstet Gynecol*; 185(4): 808-11.
- Ross R.D., Clapp S.K., Gunther S., et al. (1992):** Augmented norepinephrine and renin output in response to maximal exercise in hypertensive coarctectomy patients. *Am Heart J*; 123: 1293–1298.
- Ruderman N., Chisholm D., Pi-Sunyer X., Schneider S. (1998):** The metabolically obese, normal-weight individual revisited. *Diabetes*; 47: 699–713.
- Ruggenenti P., Perna A., Lesti M., et al. (2000):** Pretreatment blood pressure reliably predicts progression of chronic nephropathies. *Kidney Int*; 58: 2093–2101.
- Ruggenenti P., Perna A., Lesti M., et al. (2000):** Pretreatment blood pressure reliably predicts progression of chronic nephropathies. *Kidney Int*; 58: 2093–2101.
- Rumantir M.S., Jennings G.L., Lambert G.W., et al. (2000):** The “adrenaline hypothesis” of hypertension revisited. *J Hypertens*; 18: 717–723.
-

-
- Safian R.D. and Textor S.C. (2001):** Renal-artery stenosis. *N Engl J Med*; 344: 431–442.
- Sciscione A., Ivester T., Largoza M., Manley J., et al. (2003):** Acute pulmonary edema in pregnancy. *Obstet Gynecol*; 101: 511-5.
- Seedat Y.K. (2002):** Malignant-accelerated hypertension. In: Mancia G, Chalmers J, Julius S, Saruta T, Weber M, Ferrari A, Wilkinson I, eds. *Manual of Hypertension*. London: Churchill Livingstone; 623–634.
- Shannon J.R., Robertson R.M. and Biaggioni I. (1996):** Treatment of hypertension in autonomic failure. *Circulation*; 94(1): 2676.
- Sharma S.K., Philip J., Whitten C.W., et al. (1999):** Assessment of changes in coagulation in parturients with preeclampsia using thromboelastography. *Anesthesiol*; 90: 385-90.
- Silverberg S.J. (2000):** Cardiovascular disease in primary hyperparathyroidism. *J Clin Endocrinol Metab*; 85: 3513–3514.
- Simard J.M. and Bellefleur M. (1989):** Systemic arterial hypertension in head trauma. *Am J Cardiol*; 63: 32–35.
- Skaer T.L., Robison L.M., Sclar D.A. and Galin R.S. (2000):** Anxiety disorders in the USA, 1990 to 1997: trend in complaint, diagnosis, use of pharmacotherapy and diagnosis of comorbid depression. *Clin Drug Invest*; 20: 255–265.
- Slotwiner D.J., Devereaux R.B., Schwartz J.E., et al. (2001):** Relation of age to left ventricular function and systemic hemodynamics in uncomplicated mild hypertension. *Hypertension*; 37: 1404–1409.
- Solomon C.G., Thiet M.P., Moore F. and Seely E.W. (1996):** Primary hyperaldosteronism in pregnancy. A case report. *J Reprod Med*; 41: 255–258.
- Sriram S. and Robertson M.S. (2008):** Critically ill obstetric patients in Australia: a retrospective audit of 8 years: experience in a tertiary intensive care unit. *Crit Care Resusc*; 10: 124.
-

-
- Stengel B., Tarver-Carr M.E., Powe N.R., et al. (2000):** Lifestyle factors, obesity and the risk of chronic kidney disease (CKD) in U.S. adults. *J Am Soc Nephrol*; 11: 167.
- Stergiou G., Mengden T., Padfield P.L., Parati G. and O'Brien E. (2004):** Working Group on Blood Pressure Monitoring of the European Society of Hypertension. Self monitoring of blood pressure at home. *Br Med J*; 329: 870–871.
- Stewart P.M. (2000):** Dexamethasone-suppressible hypertension. *Lancet*; 356: 697–699.
- Steyn D.W. and Steyn P. (2007):** Low-dose dopamine for women with severe pre-eclampsia. *Cochrane Database of Systematic Reviews* (1); Art. No.: CD003515. DOI: 10.1002/14651858.CD003515.
- Streeten D.H.P, Anderson G.H. Howland T., et al. (1988):** Effects of thyroid function on blood pressure. Recognition of hypothyroid hypertension. *Hypertension*; 11: 78–83.
- Swales J.D. (1990):** Functional disturbance of the cell membrane in hypertension. *J Hypertens*; 8(7): 203–211.
- Swales J.D. (1994):** Overview of essential hypertension. In: *Textbook of hypertension*. Oxford: Blackwell Science; 655–660.
- Textor S.C. and Wilcox C.S. (2000):** Ischemic nephropathy/azotemic renovascular disease. *Semin Nephrol*; 20: 489–502.
- Tisdale J.E., Huang M.B., Borzak S., et al. (2004):** Risk factors for hypertensive crisis: importance of out-patient blood pressure control. *Fam Pract*.
- Toft A.D. (2000):** Surgery for primary hyperparathyroidism—sooner rather than later. *Lancet*; 355: 1478–1479.
- Towne J.B. and Bernhard V.M. (1980):** The relationship of postoperative hypertension to complications following carotid endarterectomy. *Surgery*; 88: 575.
-

-
- Tuffnell D.J., Shennan A.H., Waugh J.J. and Walker J.J. (2006):** The management of severe pre-eclampsia/eclampsia. *Royal College of Obstetricians and Gynaecologists*.
- Utiger R.D. (2000):** Treatment of acromegaly. *N Engl J Med*; 342: 1210–1211.
- Vantyghem M.C., Marcelli-Tourvieille S., Defrance F., Wemeau J.L. (2007):** 11beta-hydroxysteroid dehydrogenases. Recent advances (in French). *Annales D'endocrinologie*; 68(5): 349–56.
- Varon J. and Marik P.E. (2008):** Perioperative hypertension management. *Vasc Health Risk Manag*; 4(3): 615–27.
- Varon J., Marik P. (2000):** The diagnosis and management of hypertensive crises. *Chest*; 118(1): 214–27.
- Vaughan C.J. and Delanty N. (2000):** Hypertensive emergencies. *Lancet*; 356: 411–417.
- Visalyaputra S., Rodanant O., Somboonviboon W., et al. (2005):** Spinal versus epidural anesthesia for cesarean delivery in severe preeclampsia: a prospective randomized, multicenter study. *Anesth Analg*; 101: 862–8.
- Volkl T.M., Degenhardt K., Koch A., Simm D., et al. (2005):** Cardiovascular anomalies in children and young adults with Ullrich-Turner syndrome the Erlangen experience. *Clin Cardiol*; 28(2): 88–92.
- Weiss V., Šonka K., Pretl M., et al. (2000):** Prevalence of the sleep apnea syndrome in acromegaly population. *J Endocrinol Invest*; 23: 515–519.
- Whitaker M.D., McArthur R.G., Corenblum B., et al. (1979):** Idiopathic, sustained, inappropriate secretion of ADH with associated hypertension and thirst. *Am J Med*; 67: 511–515.
- Whitworth J.A., Mangos G.J. and Kelly J.J. (2000):** Cushing, cortisol, and cardiovascular disease. *Hypertension*; 36: 912–916.
- Wilhelm F.H., Gerlach A.L. and Roth W.T. (2001):** Slow recovery from voluntary hyperventilation in panic disorder. *Psychosom Med*; 63: 638–649.
-

-
- Williams S.B., Goldfine A.B., Timimi F.K., et al. (2001):** Acute hyperglycemia attenuates endothelium-dependent vasodilation in humans in vivo. *Circulation*; 97: 1695–1701.
- Wilmshurst P.T., Nuri M., Crowther A. and Webb-Peploe M.M. (1989):** Cold-induced pulmonary oedema in scuba divers and swimmers and subsequent development of hypertension. *Lancet*; 1: 62–65.
- Winchell R.J., Simons R.K. and Hoyt D.B. (1996):** Transient systolic hypertension. *Arch Surg*; 131: 533–539.
- Wolfsthal S.D. (1993):** Is blood pressure control necessary before surgery?. *Med Clin North Am*; 77: 349.
- Wolz M., Cutler J., Roccella E.J., et al. (2000):** Statement from the National High Blood Pressure Education Program: Prevalence of hypertension. *Am J Hypertens*; 13: 103–104.
- Wongprasartsuk P. and Sear J.W. (2003):** Anaesthesia and isolated systolic hypertension: pathophysiology and anaesthesia risk. *Anaesth Intensive Care*; 31: 619–28.
- Zhou Y., Damsky C.H., Fisher S.J. (1997):** Preeclampsia is associated with failure of human cytotrophoblasts to mimic a vascular adhesion phenotype. One cause of defective endovascular invasion in this syndrome?. *J Clin Invest*; 99(9): 2152–64.
- Ziegenhorn A., Roepke S., Schommer N., Merkl A., et al. (2009):** Clonidine improves hyperarousal in borderline personality disorder with or without comorbid posttraumatic stress disorder: a randomized, double-blind, placebo-controlled trial. *J clin psychopharmacol*; 29(2): 170–173.
- Zuccala G., Pahor M., Landi F., et al. (1997):** Use of calcium antagonists and need for perioperative transfusion in older patients with hip fracture: Observational study. *BMJ*; 314: 643.
-