

REFERENCES

- Agro F, Barzoi G, Montechia F**: Tracheal intubation using a Macintosh laryngoscope or a Glidescope in 15 patients with cervical spine immobilization . Br J Anaesth 90:705-6.(2003)
- Angel A, Gratton DA**: The effect of anaesthetic agents on cerebral cortical responses in the rat. Br J Pharmacol:76. 541-549.(2005)
- Angst MS, Koppert W, Pahl I, et al**: Short-term infusion of the mu-opioid agonist remifentanyl in humans causes hyperalgesia during withdrawal. Pain 106:49-57.(2003)
- Aucoin PJ, Kotilainen HR, Gantz NM et al** : Intracranial pressure monitors . Epidemiologic study of risk factors and infections: 80.369-3780. (2006)
- Bake, B., J. Bjure, J. Kasalichy, and A. Nachemson**: Regional pulmonary ventilation and perfusion distribution in patients with untreated idiopathic scoliosis. Thorax. 27: 703-712,(2001).
- Baldus C, Blanke K**. Preoperative nursing care: The Textbook of Spinal Surgery.chapter 1: 3-9. (2002)
- Bergofsky, E.H., G. M. Turino, and A. P. Fishman**: Cardiorespiratory failure in kyphoscoliosis. Medicine (Baltimore). 38:263-317, (2000).
- Bess RS, Lenke LG, Bridwell KH, et al**: Wasting of preoperatively donated autologous blood in surgical treatment of adolescent idiopathic scoliosis.*Spine* 31:2375-80.(2006)
- Black S**: Anaesthesia for spine surgery.The ASA Refresher Courses in Anesthesiology.Rosenblatt F, Butterworth J (Eds): volume 37, chapter 2, pp. 13–23. (2009)
- Blumenthal S, Min K, Marquardt M, Borgeat A**: Postoperative intravenous morphine consumption, pain scores, and side effects with perioperative oral controlled-release oxycodone after lumbar discectomy. Anesth Analg 105:233-7.(2007)
- Boyd SG, Rothwell JC, Cowan JMA, Webb PJ, Morley T, Asselman P, et al**: A method of monitoring function in corticospinal pathways during scoliosis surgery with a note on motor conduction velocities. J Neurol Neurosurg Psychiatry 49.251-7.(2003)

- Branston NM, Symon L:** Cortical EP, blood flow, and potassium changes in experimental ischemia. *Evoked potentials*: 527-530.(2004)
- Calder I, Calder J, Crockard HA:** Difficult direct laryngoscopy in patients with cervical spine disease. *Anaesthesia*: 50:756-63.(2007)
- Carragee E J, Samuels S I, Jaffe R A, Samuels S I:** Spine Surgery. *Anesthesiologists Manual of Surgical Procedures*:23. 791–802.(2006)
- Chen S, Huang T, Lee Y:** Pulmonary function after thoracoplasty in adolescent idiopathic scoliosis. *Clinical orthopedics and related research* 399: 152–161. (2002)
- Cohen AR, Young W, Ransohoff J:** Intraspinal localization of the somatosensory evoked potential. *Neurosurgery*: 9:157-162. (2006)
- Crosby ET:** Airway management in adults after cervical spine trauma: *Anesthesiology*: 104(6).1293-318.(2006)
- Crosby ET:** Considerations for airway management for cervical spine surgery in adults. *Anesthesiol Clin* 25:511-33. (2007)
- Crosby ET:** Perioperative haemotherapy.Risks and complications of blood transfusion: 39.822-40. (2009)
- D. A. Raw, J. K. Beattie and J. M. Hunter:** Anaesthesia for spinal surgery in adults. *British Journal of Anaesthesia*:887-904.(2004)
- Daniel L. Master, MD, George H. Thompson, MD et al:** Spinal Cord Monitoring for Scoliosis Surgery in Rett Syndrome. *J Pediatr Orthop* 28.342-346.(2008)
- Davies, G., and L. Reid., Winter R.B:** Effect of scoliosis on growth of alveoli and pulmonary arteries and on right ventricle. *Arch: Dis. Child.* 46: 623-632.(2002).
- Deletis V, Rodi Z, Amassian VE:** Neurophysiological mechanisms underlying motor evoked potentials in anesthetized humans.*Clin Neurophysiol*: 112.445-52. (2001)
- Deletis V, Sala F :** Intraoperative neurophysiological monitoring of the spinal cord During spinal cord and spine surgery:A review focus on the corticospinal tracts.*Clinical Neurophysiology* 119: 248–264. (2008)

- Dharmavaram S, Jellish WS, Nockels RP, et al:**Effect of prone positioning systems on hemodynamic and cardiac function during lumbar spine surgery.Spine 31.1388-93.(2006)
- Dutton RP:** International Anesthesiology Clinics 40. 103-120.(2002)
- Edward T. Crosby, MD:** Considerations for Airway Management for Cervical Spine Surgery in Adults.elsevier saunders:25.511-533. (2007)
- Elizabeth J. Bridges, PhD:** Pulmonary Artery Pressure Monitoring. AACN Advanced Critical Care17. 286-305.(2006)
- Epstein NE, Hollingsworth R, Nardi D, Singer J:** Can airway complications following multilevel anterior cervical surgery be avoided? .J Neurosurg 94.185-8. (2007)
- Erwin AC, Erwin CW:** Intraoperative evoked potential monitoring-A historical perspective. Am J END Technol:40.98-120.(2005)
- Florentino-Pineda I, Blakemore LC, Adler P, Tripi P:** The Effect of epsilon-aminocaproic acid on perioperative blood loss in patients with idiopathic scoliosis undergoing posterior spinal fusion: A preliminary prospective study. Spine 26:1147-51.(2001)
- Floris E.A. Udink ten Cate, Barend J. van Royen:** Incidence and risk factors of prolonged mechanical ventilation in neuromuscular scoliosis surgery. Journal of Pediatric Orthopaedics 17.204-206.(2008)
- Frances D. Faro, MD, Michelle C. Marks, PT, MA, Peter O. Newton, MD:** Perioperative Changes in Pulmonary Function After Anterior Scoliosis Instrumentation. SPINE 30.1058-1063.(2003)
- Francesco Sala, M.D., Giorgio Palandri, M.D:** motor evoked potential monitoring improves outcome after surgery for intramedullary spinal cord tumors. Neurosurgery:58.1129-1143. (2006)
- Gibson P, et al:** Anaesthesia for spinal surgery in adults. British journal of anaesthesia 92:770- 772. (2004)
- Goodkin R, Mesiwala A.** General principles of operative positioning. Winn HR(eds). Youmans neurological surgery. 5th edition. Saunders, Elsevier Inc. p. 595-621. (2004)
- Goutcher CM, Lochhead V.** Reduction in mouth opening with semi-rigid cervical collars: Br J Anaesth 95.344-8.(2005)

-Grisolia A, Bell RL, Peltier LF: Fractures and dislocations of the spine complicating ankylosing spondylitis: a report of six cases. Clin Orthop 422.129-34. (2004)

-Grottke O, Dietrich PJ, Wiegels S, Wappler F: Intraoperative wake-up test and postoperative emergence in patients undergoing spinal surgery: A comparison of intravenous and inhaled anesthetic techniques using short-acting anesthetics: Anesth Analg 99.1521-7. (2004)

-H. Hermanns, P. Lipfert et al: Cortical somatosensory-evoked potentials during spine surgery in patients with neuromuscular and idiopathic scoliosis under propofol–remifentanyl anaesthesia. British Journal of Anaesthesia:19.363-365.(2007)

-Harper CM, Ambler G, Edge G: The prognostic value of pre-operative predicted forced vital capacity in corrective spinal surgery for Duchenne's muscular dystrophy. Anaesthesia 59:1160-1162.(2004)

-Hetreed M A: Anaesthesia for Major Spinal Surgery. Curr Anaesth Crit Care: 8.264-9. (2004)

-Ian Crabb: Anaesthesia for Spinal Surgery. ANAESTHESIA AND INTENSIVE CARE MEDICINE 45.80-84.(2004)

-Irene Rozet, MD, Monica S. Vavilala, MD: Risks and Benefits of Patient Positioning During Neurosurgical Care. Anesthesiology Clin 25.631–653(2007)

-J. Richard Toleikis, Ph.D, D.ABNM: intraoperative monitoring using somatosensory evoked potentials. Journal of Clinical Monitoring and Computing:24.241-285.(2005)

-J. Richard Toleikis, Ph.D, D.ABNM: intraoperative monitoring using somatosensory evoked potentials. Journal of Clinical Monitoring and Computing:24.241-285.(2005)

-Jameson LC, Popic PM, Harms BA: Hyperkalemic death during use of a high-capacity fluid warmer for massive transfusion. International Anesthesiology Clinics :73.1050-75.(2007)

-Jameson LC, Popic PM, Harms BA: Hyperkalemic death during use of a high-capacity fluid warmer for massive transfusion. International Anesthesiology Clinics :73.1050-75.(2007)

-Jeffrey H. Owen, PhD: The Application of Intraoperative Monitoring During Surgery for Spinal Deformity. *Pediatric Anaesth*:2642-2662.(2003)

-Jeffrey H. Owen, PhD: The Application of Intraoperative Monitoring During Surgery for Spinal Deformity. *Pediatric Anaesth*:2642-2662.(2003)

-Joly V, Richebe P, Guignard B, et al: Remifentanil-induced postoperative hyperalgesia and its prevention with small-dose ketamine.*Anesthesiology* 103:147-55.(2005)

-Joly V, Richebe P, Guignard B, et al: Remifentanil-induced postoperative hyperalgesia and its prevention with small-dose ketamine.*Anesthesiology* 103:147-55.(2005)

-Jones SJ, Edgar MA, Ransford AO:Sensory nerve conduction in the human spinal cord: Epidural recordings made during spinal cord surgery. *J Neurol Neurosurg Psychiatry*: 45. 446-452.(2003)

-Jones SJ, Edgar MA, Ransford AO:Sensory nerve conduction in the human spinal cord: Epidural recordings made during spinal cord surgery. *J Neurol Neurosurg Psychiatry*: 45. 446-452.(2003)

-Jordanoglu, J: Rib movement in health, kyphoscoliosis, and ankylosing spondylitis. *Thorax*. 24: 407-414.(2000).

-Jordanoglu, J: Rib movement in health, kyphoscoliosis, and ankylosing spondylitis. *Thorax* 24: 407-414, (2000).

-K.F. Kothbauer: Intraoperative neurophysiologic monitoring for intramedullary spinal-cord tumor surgery.*Elsavier masson*:37.408 414.(2007)

-Kakiuchi M: Reduction of blood loss during spinal surgery by epidural blockade under normotensive general anaesthesia:*Spine*22.889-94.(2000).

-Kim YJ, Lenke LG, Bridwell KH, et al: Pulmonary function in adolescent idiopathic scoliosis relative to the surgical procedure. *J Bone Joint Surg Am* 87:1534-41.(2005)

-Kokoszka A, Kuflik P, Casden A, et al: Evidence-based review of the role of aprotinin in blood conservation during orthopaedic surgery. *J Bone Joint Surg Am* 87:1129-36.(2005)

- Koolstra JH, van Eijden TMGJ:** Functional significance of the coupling between head and jaw movements. *J Biomech.*37(9).1387-92.(2004)
- Lam AM, Sharar SR, Mayberg TS, et al:**Isoflurane compared with nitrous oxide anaesthesia for intraoperative monitoring of somatosensory evoked potentials. *Can J Anaesth:* 41.295-300.(2003).
- Langen KE, Jellish WS, Ghanayem AJ:**Anesthetic considerations in spine surgery. *Con Spine Surg* 7:1-8.(2006)
- Lars L. Mielke, MD, Elmar K. Entholzner, MD, et al:**Preoperative Acute Hypervolemic Hemodilution with Hydroethylstarch,An Alternative to Acute Normovolemic Hemodilution.*An&h Analg* 84:26-30.(1997)
- Lenke LG, White DK, Kemp JS, et al:** Evaluation of ventilatory efficiency during exercise in patients with idiopathic scoliosis undergoing spinal fusion.*Spine* 27:2041-5.(2002)
- Lennarson PJ, Smith DW, Sawin PD, et al:** Cervical spinal motion during intubation: efficacy of stabilization maneuvers in the setting of complete segmental instability. *J Neurosurg(Spine2)*.94.265-70.(2001)
- Lennarson PJ, Smith DW, Sawin PD, et al.:** Cervical spinal motion during intubation,Efficacy of stabilization maneuvers in the setting of complete segmental instability.*J Neurosurg:* 94.265-70. (2007)
- MacDonald DB:**Safety of intraoperative transcranial electrical stimulation motor evoked potential monitoring. *J Clin Neurophysiol:*19.416-29. (2002)
- Macon JB, Poletti CE, Sweet WH, Ojemann RG, Zervas NT:** Conducted somatosensory evoked potentials during spinal surgery.*J Neurosurg:* 57.354-359 (2005)
- Manninen PH, Jose GB, Lukitto K, Venkatraghavan L, El Beheiry H:** Management of the airway in patients undergoing cervical spine surgery. *J Neurosurg Anesthesiol:* 19:190-4. (2007)
- Mark W.Crawford, MBBS, FRCPC, Chantal Hickey, MD, FRCPC,et al:** development of acute opioid tolerance during infusion of remifentanil for pediatric scoliosis surgery. *Pediatric anaesthesia*102.1662-7.(2006)

- Mary E. McCann, MD, David Hill,et al:** A Comparison of Radial Artery Blood Pressure Determination Between the Vasotrac Device and Invasive Arterial Blood Pressure Monitoring in Adolescents Undergoing Scoliosis Surgery. *Anesth Analg*101:978-85.(2005)
- Mashour GA, Sandberg WS.** Craniocervical extension improves the specificity and predictive value of the Mallampati airway evaluation. *Anesth Analg*103(5):1256-9.(2006)
- Mayhall CG, Archer NH, Lamb V et al:** Ventriculostomy-related infections. A prospective epidemiologic study: 310.553-560. (2008)
- McLeod ADM,Calder I:** Spinal cord injury and direct laryngoscopy: the legend lives on. *Br J Anaesth.*84:705-8.(2000)
- Meert KL, Kannan S, Mooney JF:** Predictors of red cell transfusion in children and adolescents undergoing spinal fusion surgery. *Spine*27:2137-42.(2002)
- MEG A. ROSENBLATT,M.D, JOHN F. BUTTERWORTH IV, M.D:** anaesthesia for spine surgery. *The American Society of Anesthesiologists* 67:13-23.(2009)
- Michael D. Popitz:** Anesthetic Implications of Chronic Disease of the Cervical Spine. *ANESTH ANALG*:672-683.(2005)
- Morris P. Duchenne muscular dystrophy:** a challenge for the anaesthetist. *Pediatric Anaesth* 7. 1-4. (2006)
- Murphy NA, Firth S, Jorgensen T, Young PC:** Spinal surgery in children with idiopathic and neuromuscular scoliosis. What's the difference? *J Pediatr Orthop* 26:216-220.(2006)
- N. Soundararajan , M. Cunliffe:** Anaesthesia for spinal surgery in children. *British Journal of Anaesthesia* 43:86-94.(2007)
- Naik B, Lobato EB, Sulek CA:** Dysphagia, obstructive sleep apnea, and difficult fiberoptic intubation secondary to diffuse idiopathic skeletal hyperostosis. *Anesthesiology*100(5):1311-2.(2004)
- Nielsen CH, Bridwell KH, DeWald RL:** Preoperative and postoperative anesthetic considerations for the spinal surgery patient. In:, eds. *The Textbook of Spinal Surgery*: 31-37.(2002)
- Nuwer, MR:** Evoked potential monitoring in the operating room. *Raven Press.*54-58.(2005).

-Owen JH, Toleikis JR: Nerve root monitoring. The textbook of spinal surgery, 2nd edition: 61-75(2003)

-Padberg AM, Komanetsky RE, Bridwell KH, et al: Neurogenic motor evoked potentials: A prospective comparison of stimulation methods. J Spinal Disord:11:21-4.(2002)

Panjabi MM, Yue JJ, Dvorak J, et al: Cervical spine kinetics and clinical instability. In: Clark CR, editor. The cervical spine. 4th edition. Philadelphia: 55-78. 2005

-Papastefanou SL, Henderson LM, Smith NJ, et al: Surface electrode somatosensory-evoked potentials in spinal surgery. Spine:25:2467-72. (2000).

-Pelosi L, Stevenson M, Hobbs GJ, et al: Intraoperative motor evoked potentials to transcranial electrical stimulation during two anaesthetic regimens. Clin Neurophysiol 112:1076-87.(2001).

-Peter O. Newton, MD, Andrew Perry, MD, Tracey P. Bastrom, MA: Predictors of Change in Postoperative Pulmonary Function in Adolescent Idiopathic Scoliosis. SPINE 32:1875–1882.(2007)

-Powers SK, Bolger CA, Edwards MSB: Spinal cord pathways mediating somatosensory evoked potentials. J Neurosurg:57:472-482. (2003)

-Prior PF: EEG monitoring and evoked potentials in brain ischemia. Br J Anaesth: 57. 63-81.(2003)

-Reid JM, Appleton PJ: A case of ventricular fibrillation in the prone position during back stabilisation surgery in a boy with Duchenne's muscular dystrophy. Anaesthesia: 54. 364-7. 2004

-Richard P. Dutton, MD: Anesthetic Management of Spinal Cord Injury. Clinical Practice and Future Initiatives 103-120.(2006)

-Rosner MJ, Becker DP: ICP monitoring: Complications and associated factors. Clin Neurosurg 23:494-500.(2007)

-Rozet I, Vavilala M: Risks and Benefits of Patient Positioning During Neurosurgical Care, Anesthesiology Clinics 25: 631–653. (2007)

-Rudolph C, Schneider JP, Wallenborn J, et al: Movement of the upper cervical spine during laryngoscopy: a comparison of the Bonfils intubation fiberscope and the Macintosh laryngoscope. *Anaesthesia* 60.668-72. (2005)

-S .w. Suttner,s.n.piper,et al:sodium nitroprosside induced hypotension. *british journal of anaesthesia* 87.699-705.(2001)

-Sala F, Palandri G, Basso E, Lanteri P, Deletis V, Faccioli F, et al: Motor evoked potential monitoring improves outcome during surgery for intramedullary spinal cord tumor. *Neurosurgery*:58.1129-43.(2006)

-Samuel Abraham Joseph, Jr, MD, Ketevan Berekashvili, MD, MPH ,et al: Blood Conservation Techniques in Spinal Deformity Surgery. *SPINE*.33 2310-2315.(2007)

-Schwartz DM, Auerbach JD, Dormans JP, Flynn J, Drummond DS, Bowe JA,Lauffer S, Shah SA, Bowen JR, Pizzutillo PD, Jones KJ: Neurophysiological detection of impending spinal cord injury during scoliosis surgery. *J Bone Joint Surg Am*: 89.2440-9. (2007)

-Scott S. Reuben, MD: Postoperative Analgesic Effects of Celecoxib or Rofecoxib After Spinal Fusion Surgery. *International Anesthesia Research Society*:91.1221-5.(2000)

-Shapiro F, Sethna N, Colan S, et al: Spinal fusion in Duchenne muscular dystrophy: a multidisciplinary approach. *Muscle Nerve*: 15: 604-14.(2005)

-Shapiro F, Zurakowski D, Sethna NF: Tranexamic acid diminishes intraoperative blood loss and transfusion in spinal fusions for Duchenne muscular dystrophy scoliosis. *Spine* 32.2278-83.(2006)

-Shaw, D. B., and J. read: Hypoxia and thoracic scoliosis: *Br. Med. J*.2: 1486-1489.(2001).

-Sheppeard H, Cleak DK, Ward DF, et al: A review of early mortality and morbidity in elderly patients following Charnley total hip replacement. *Arch Orthop Trauma Surg*: 4.243-8.(2002)

Simpson RK Jr, Blackburn JG, Martin HF III, Katz S:Peripheral nerve fibers and spinal cord pathway contribution to the somatosensory evoked potentials. *Exp Neurol*: 73.700-715.(2001)

- Sinha A, Agarwal A, Gaur A, Pandey CK:** Oropharyngeal swelling and macroglossia after cervical spine surgery in the prone position. *J Neurosurg Anesthesiol*: 13.237-9.(2007)
- Sloan TB, Deletis V, Shils J, et al:** Intraoperative neurophysiology and anesthesia management. *Neurophysiology in Neurosurgery*: 451-74.2002.
- Sloan TB:** Anesthetic effects on electrophysiologic recordings. *J Clin Neurophysiol* 15.217-226.(2004)
- Sloan TB.** Evoked potentials. *Textbook of neuroanesthesia with neurosurgical and neuroscience perspectives*: 221-276.(2006)
- Soundararajan N, Cunliffe M:** Anaesthesia for spinal surgery in children. *British journal of anaesthesia* 99: 86–94.(2007)
- Stecker MM, Cheung AT, Pochettino A, Kent G, Patterson T, Weiss SJ, Bavaria JE:** Effects of cooling on electroencephalogram and evoked potentials. *Ann Thorac Surg* 71.22-28.(2001)
- Stiell IG, Clement CM, McKnight RD, et al:** The Canadian C-spine rule versus NEXUS low risk criteria in patients with trauma. *N Eng J Med* 349:2510-20. 2003
- Stocki B, Wimmer C, Innerhofer P, Kofler M, Behensky H:** Delayed anterior spinal artery syndrome following posterior scoliosis correction. *Eur Spine J* 14: 906–9.(2005)
- Supkis DE jr, Varon J Benumof J:** Anesthesia and Uncommon Diseases. 4th Edn. Philadelphia. chapter 5: 545-60. (2005)
- Szelenyi A, Kothbauer KF, Deletis V:** Transcranial electrical stimulation, motor evoked potential monitoring. *J Clin Neurophysiol* 19.416-29. (2002)
- Terao Y, Matsumoto S, Yamashita K, et al:** Increased incidence of emergency airway management after combined anterior-posterior cervical spine surgery. *J Neurosurg Anesthesiology*: 16.282-6.(2004)
- Thompson GH, Poe-Kochert C :** Intraoperative blood cell salvage in obstetrics National Institute for Health and Clinical Excellence. *An&h Analg* 207:23-31.(2005)
- Tobias JD:** Fenoldopam for controlled hypotension during spinal fusion in children and adolescents. *Paediatr Anaesth* 9. 261-6. (2000).

- Tobias JD:** A review of intrathecal and epidural analgesia after spine surgery in children. *Anesth Analg* 98:956-65.(2004)
- Toleikis JR, Carlvin AO, Shapiro DE, Schafer MF:**The use of dermatomal evoked responses during surgical procedures that use intrapedicular fixation of the lumbosacral spine. *Spine*:18. 2401-2407 (2008)
- Turkstra TP, Craen RA, Pelz DM, et al:** Cervical spine motion: a fluoroscopic comparison during intubation with lighted stylet, Glidescope, and Macintosh laryngoscope. *Anesth Analg*.101.910-5.(2005)
- Vernon w.lin, Diana d.cadenas, Nancy c.cutter, Frederick s,frost:**Spinal cord medicine: volume 37, chapter 1, pp. 9-16. (2003)
- Wazeka AN, DiMaio MF, Boachie-Adjei O:** Outcome of pediatric patients with severe restrictive lung disease following reconstructive spine surgery. *Spine* 29:528-534(2004)
- Weber, B, J. P. Smith, W. A. Briscoe, S. A. Friedman, and T. K. C. King:** Pulmonary function in asymptomatic adolescents with idiopathic scoliosis. *Ain. Rev. Respir. Dis* 111. 389-397. (2002)
- WILLIAM. ROSENBLATT, M.D et al:** Awake intubation Made easy. *The American Society of Anesthesiologists* 37.168-174.(2009)
- Wilson-Holden TJ, Padberg AM, Lenke LG, et al:** Efficacy of intraoperative monitoring for pediatric patients with spinal cord pathology undergoing spinal deformity surgery. Presented at the annual meeting of the Scoliosis Research Society: (2002)
- Yew-Long Lo, MD, Yang-Fang Dan,et al:** Intraoperative Motor-evoked Potential Monitoring in Scoliosis Surgery: Comparison of Desflurane/Nitrous Oxide With Propofol Total Intravenous Anesthetic Regimens. *Neurosurg Anesthesiol*:18.211-214.(2006)
- York DH:** Somatosensory evoked potentials in man: Differentiation of spinal pathways responsible for conduction from the forelimb vs hindlimb. *Prog Neurobiol*: 25.1-25. (2002)
- Zhang JG, Wang W, Qiu GX, et al:** The role of preoperative pulmonary function tests in the surgical treatment of scoliosis. *Spine*30:218–21.(2005)