

References

References

1. Apan A, Baydar S, Yilmaz S, Uz A, Tekdemir I, Guney S, Elhan A: Surface landmarks of brachial plexus: ultrasound and magnetic resonance imaging for supraclavicular approach with anatomical correlation. *Eur J Ultrasound* 2001; 13: 191-6
2. Awad IT, Duggan EM: Posterior lumbar plexus block: anatomy, approaches, and techniques. *Reg Anesth Pain Med* 2005; 30: 143-9.
3. Bigeleisen P, Wilson M. A comparison of two techniques for ultrasound guided infraclavicular block. *Br J Anaesth* 2006; 96: 502-507.
4. Bloc S, Garnier T, Komly B, Asfazadourian H, Leclerc P, Mercadal L, Morel B, Dhonneur G. Spread of injectate associated with radial or median nerve-type motor response during infraclavicular brachial-plexus block: an ultrasound evaluation. *Reg Anesth Pain Med* 2007;32:130-135.
5. Carney J, McDonnell JG, Ochana A, et al. The transversus abdominis plane block provides effective postoperative Painalgnesia in patients undergoing total abdominal hysterectomy. *Anesth Analg* 2008;107(6):2056.
6. Carr, J. and J. Brown. *Introduction to Biomedical Equipment Technology*. Prentice-Hall, 1998.
7. Casati A, Danelli G, Baciarello M, Corradi M, Leone S, Di Cianni S, Fanelli G: A prospective, randomized comparison between ultrasound and nerve stimulation guidance for multiple injection axillary brachial plexus block. *Anesthesiology* 2007; 106: 992-6.

References

8. Chan VW, Nova H, Abbas S, McCartney CJ, Perlas A, Quan XD: Ultrasound Examination and Localization of the Sciatic Nerve: A Volunteer Study. *Anesthesiology* 2006; 104: 309-14
9. Chan VW, Perlas A, McCartney CJ, Brull R, Xu D, Abbas S: Ultrasound guidance improves success rate of axillary brachial plexus block. *Can.J.Anaesth.* 2007; 54: 176-82.
10. Chan VW, Perlas A, Rawson R & Odukoya O. Ultrasound-guided supraclavicular brachial plexus block. *Anesth Analg* 2003; 97(5): 1514–1517.
11. Chan VW, Perlas A, Rawson R, Odukoya O: Ultrasound-guided supraclavicular brachial plexus block. *Anesth Analg* 2003; 97: 1514-7
12. Chan VW: Applying ultrasound imaging to interscalene brachial plexus block. *Reg Anesth Pain Med* 2003; 28: 340-3.
13. Chantzi C, Alevizou A, Saranteas T, Zogogiannis J, Iatrou C, Dimitriou V: Usefulness of the two to 5 MHz ultrasound probe in examination and block of the sciatic nerve in orthopedic trauma patients: a preliminary study. *J Clin Anesth* 2007; 19: 486-8
14. Choyce A, Chan V, Middleton W, et al. What is the relationship between paresthesia and nerve stimulation for axillary brachial plexus block? *Reg Anesth. Pain Med* 2001; 26: 100-104.
15. Dalens B. Regional anesthesia in children. In: Miller, ed. *Anesthesia* Philadelphia: Churchill Livingstone, 2000.
16. Demondion X, Herbinet P, Boutry N, Fontaine C, Francke JP, Cotten A: Sonographic mapping of the normal brachial plexus. *Am J Neuroradiol.* 2003; 24: 1303-9.

References

17. Dingemans E, Williams S R, Arcand G, Chouinard P, Harris P, Ruel M, Girard F. Neurostimulation in ultrasound-guided infraclavicular block: a prospective randomized trial. *Anesth Analg* 2007;104:1275-80.
18. Domingo-Triado V, Selfa S, Martinez F, Sanchez-Contreras D, Reche M, Tecles J, Crespo MT, Palanca JM, Moro B: Ultrasound guidance for lateral midfemoral sciatic nerve block: a prospective, comparative, randomized study. *Anesth. Analg.* 2007; 104: 1270-4.
19. Duggan E, Brull R, Lai J, Abbas S: Ultrasound-guided brachial plexus block in a patient with multiple glomangiomatosis. *Reg Anesth. Pain Med.* 2008; 33: 70-3.
20. Ferre RM, Sweeney TW: Emergency physicians can easily obtain ultrasound images of anatomical landmarks relevant to lumbar puncture. *Am.J.Emerg.Med.* 2007; 25: 291-6.
21. Fujiwara Y, Sato Y, Kitayama M, Shibata Y, Komatsu T, Hirota K. Obturator nerve block using ultrasound guidance. *Anesth Analg* 2007;105:888-889.
22. Graif M, Seton A, Nerubai J, Horoszowski H, Itzhak Y: Sciatic nerve: sonographic evaluation and anatomic-pathologic considerations. *Radiology* 1991; 181: 405-8
23. Grau T, Leipold R W, Horter J, Conradi R, Martin E O, Motsch J. Paramedian access to the epidural space: the optimum window for ultrasound imaging. *J Clin Anesth* 2001;13: 213-217.
24. Gray A T, Schafhalter-Zoppoth I. "Bayonet artifact" during ultrasound-guided transarterial axillary block. *Anesthesiology* 2005; 102: 1291-1292.

References

25. Guzeldemir M E, Ustunsoz B. Ultrasonographic guidance in placing a catheter for continuous axillary brachial plexus block. *Anesth Analg* 1995; 81: 882-883.
26. Hebbard P, Royse C. Ultrasound guided posterior approach to the infraclavicular brachial plexus. *Anaesthesia* 2007;62:539.
27. Helayel P E, da Conceicao D B, Pavei P, Knaesel J A, Oliveira Filho G R. Ultrasound-guided obturator nerve block: a preliminary report of a case series. *Reg Anesth Pain Med* 2007;32:221-226.
28. Hurdle MF, Weingarten TN, Crisostomo RA, Psimos C, Smith J: Ultrasound-guided blockade of the lateral femoral cutaneous nerve: technical description and review of 10 cases. *Arch Phys Med Rehabil* 2007; 88: 1362-4.
29. Jan VG, Tielens L, Gielen M: Ultrasound-guided interscalene brachial plexus block in a child with femur fibula ulna syndrome. *Paediatr Anaesth* 2006; 16: 330-2.
30. Kapral S, Krafft P, Eibenberger K, *et al.* Ultrasound-guided supraclavicular approach for regional anesthesia of the brachial plexus. *Anesth Analg* 1994
31. Khabiri B, Arbona F, Norton J: "Gapped supine" position for ultrasound guided lateral popliteal fossa block of the sciatic nerve. *Anesth. Analg.* 2007; 105: 1519.
32. Kirchmair L, Entner T, Kapral S, Mitterschiffthaler G. Ultrasound guidance for the psoas compartment block: an imaging study. *Anesth Analg* 2002; 94: 706-710.

References

33. Kirchmair L, Entner T, Wissel J, Moriggl B, Kapral S, Mitterschiffthaler G. A study of the paravertebral anatomy for ultrasound-guided posterior lumbar plexus block. *Anesth Analg* 2001; 93: 477-81.
34. Kossoff G. Basic physics and imaging characteristics of ultrasound. *World J Surg* 2000; 24: 134-42.
35. La Grange P, Foster P, Pretorius L. Application of Doppler ultrasound blood flow detector in supraclavicular brachial plexus block. *Br J Anaesth* 1978
36. Lang SA, Saito T. Thoracic paravertebral nerve block, nerve stimulator guidance and the endothoracic fascia. *Anaesthesia* 2005;60(9):930.
37. Marhofer P, Greher M, Kapral S. Ultrasound guidance in regional anaesthesia. *Br J Anaesth* 2005; 94:7-17.
38. Marhofer P, Schrogendorfer K, Koinig H, Kapral S, Weinstabl C, Mayer N. Ultrasonographic guidance improves sensory block and onset time of three-in-one blocks. *Anesth Analg* 1997; 85: 854-7
39. Marhofer P, Schrogendorfer K, Wallner T, Koinig H, Mayer N, Kapral S. Ultrasonographic guidance reduces the amount of local anesthetic for 3-in-1 blocks. *Reg Anesth Pain Med* 1998; 23: 584-8
40. Marhofer P, Schrogendorfer K, Wallner T, Koinig H, Mayer N, Kapral S. Ultrasonographic guidance reduces the amount of local anesthetic for 3-in-1 blocks. *Reg Anesth Pain Med* 1998;23:584-588.
41. Martinoli C, Bianchi S, Santacroce E, Pugliese F, Graif M, Derchi LE: Brachial plexus sonography: a technique for assessing the root level. *Am J Roentgenol* 2002; 179: 699-702

References

42. McDonnell JG, Curley G, Carney J, et al. The analgesic efficacy of transversus abdominis plane block after cesarean delivery: a randomized controlled trial. *Anesth Analg* 2008;106(1):186.
43. McDonnell JG, O'Donnell B, Curley G, et al. The analgesic efficacy of transversus abdominis plane block after abdominal surgery: a prospective randomized controlled trial. *Anesth Analg* 2007;104(1):193.
44. Michael KC (1994): *Anesthesia*. Churchill Livingstone; 4:1195-1214.
45. Minville V, Zetlaoui P J, Fessenmeyer C, Benhamou D. Ultrasound guidance for difficult lateral popliteal catheter insertion in a patient with peripheral vascular disease. *Reg Anesth Pain Med* 2004;29: 368-370.
46. Morimoto M, Kim JT, Popovic J, Jain S, Bekker A: Ultrasound guided lumbar plexus block for open reduction and internal fixation of hip fracture. *Pain Pract* 2006; 6: 124-6.
47. Naja MZ, Ziade MF, El Rajab M, et al. Varying anatomical injection points within the thoracic paravertebral space: effect on spread of solution and nerve blockade. *Anaesthesia* 2004;59(5):459.
48. Naja MZ, Ziade MF, Lonnqvist PA. Nerve-stimulator guided paravertebral blockade vs. general anaesthesia for breast surgery: a prospective randomized trial. *Eur J Anaesthesiol* 2003;20(11):897.
49. Oberndorfer U, Marhofer P, Bosenberg A et al. Ultrasonographic guidance for sciatic and femoral nerve blocks in children. *Br J Anaesth* 2007;98:797-801.
50. O'Donnell B D, Mannion S. Ultrasound-guided femoral nerve block, the safest way to proceed? *Reg Anesth Pain Med* 2006;31:387-388.

References

- 51.Ootaki C, Hayashi H, Amano M. Ultrasound-guided infraclavicular brachial plexus block: an alternative technique to anatomical landmark-guided approaches. *Reg Anesth Pain Med* 2000; 25: 600-604
- 52.Paraskeuopoulos T, Saranteas T, Chantzi C, Alevizou A, Zogogiannis J, Dimitriou V, Kostopanagiotou G: Imaging in Anesthesia: The Role of 4 MHz to 7 MHz Sector Array Ultrasound Probe in the Identification of the Sciatic Nerve at Different Anatomic Locations. *Reg Anesth Pain Med* 2007; 32: 537-8
- 53.Peer S, kovacs P, Harpf C, et al. High resolution sonography of lower extremity peripheral nerves: anatomic correlation and spectrum of disease. *J Ultrasound Med* 2002; 21: 315-22.
- 54.Peng P W, Rofaeel A. Using ultrasound in a case of difficult epidural needle placement. *Can J Anaesth* 2006; 53: 325-326.
- 55.Peras A, Chan VW, Simons M: Brachial plexus examination and localization using ultrasound and electrical stimulation: a volunteer study. *Anesthesiology* 2003; 99: 429-35.
- 56.Peras A, Chan VWS. Ultrasound guided interscalene brachial plexus block. *Techniques in Regional Anesthesia and Pain Management* 2004; 8: 143-8.
- 57.Peras A, Niazi A, McCartney C, Chan V, Xu D, Abbas S. The sensitivity of motor response to nerve stimulation and paresthesia for nerve localization as evaluated by ultrasound. *Reg Anesth Pain Med* 2006; 31: 445-450.
- 58.Pham D C, Guilley J, Dernis L et al. Is there any need for expanding the perineural space before catheter placement in continuous femoral nerve blocks? *Reg Anesth Pain Med* 2006;31:393-400.

References

59. Pusch F, Wildling E, Klimscha W, Weinstabl C. Sonographic measurement of needle insertion depth in paravertebral blocks in women. *Br J Anaesth* 2000; 85: 841-843.
60. Rafi AN. Abdominal field block: a new approach via the lumbar triangle. *Anaesthesia* 2001;56(10):1024.
61. Retzl G, Kapral S, Greher M, Mauritz W. Ultrasonographic findings of the axillary part of the brachial plexus. *Anesth Analg* 2001; 92: 1271-1275.
62. Rhodes D W, Bishop P A. A review of diagnostic ultrasound of the spine and soft tissue. *J Manipulative Physiol Ther* 1997; 20:267-273.
63. Rodriguez J, Barcena M, Rodriguez V, Aneiros F, Alvarez J. Infraclavicular brachial plexus block effects on respiratory function and extent of the block. *Reg Anesth Pain Med* 1998; 23:564-8.
64. Saranteas T, Chantzi C, Paraskeuopoulos T, Alevizou A, Zogogiannis J, Dimitriou V, Kostopanagiotou G: Imaging in Anesthesia: The Role of 4 MHz to 7 MHz Sector Array Ultrasound Probe in the Identification of the Sciatic Nerve at Different Anatomic Locations. *Reg Anesth Pain Med* 2007; 32: 537-8
65. Saranteas T, Chantzi C, Zogogiannis J, et al. Ultrasound examination and localization of the sciatic nerve at the mid-femoral. An imaging study with ultrasound. *Acta Anaesthesiol Scand* 2007;51:387-8.
66. Saranteas T, Paraskeuopoulos T, Alevizou A, Kouskouri A, Zogogiannis J, Anagnostopoulou S, Chantzi C. Identification of the obturator nerve divisions and subdivisions in the inguinal region: a study with ultrasound. *Acta Anaesthesiol Scand* 2007;51:1404-1406.

References

67. Schwemmer U, Markus C K, Greim C A, Brederlau J, Roewer N. Ultrasound-guided anaesthesia of the axillary brachial plexus: efficacy of multiple injection approach. *Ultraschall Med* 2005; 26: 114-119.
68. Sheppard DG, Revathy BI, Fenstermacher MJ. Brachial plexus: demonstration at US. *Radiology* 1998
69. Sinha A, Chan V W. Ultrasound imaging for popliteal sciatic nerve block. *Reg Anesth Pain Med* 2004;29: 130-134.
70. Slater M E, Williams S R, Harris P, Brutus J P, Ruel M, Girard F, Boudreault D. Preliminary evaluation of infraclavicular catheters inserted using ultrasound guidance: through-the-catheter anesthesia is not inferior to through-the-needle blocks. *Reg Anesth Pain Med* 2007;32:296-302.
71. Soeding P E, Sha S, Royse C E, Marks P, Hoy G, Royse A G. A randomized trial of ultrasound-guided brachial plexus anaesthesia in upper limb surgery. *Anaesth Intensive Care* 2005; 33: 719-725.
72. Soong J, Schafhalter-Zoppoth I, Gray A T. Sonographic imaging of the obturator nerve for regional block. *Reg Anesth Pain Med* 2007;32:146-151.
73. Soong J, Schafhalter-Zoppoth I, Gray AT: The importance of transducer angle to ultrasound visibility of the femoral nerve. *Reg Anesth Pain Med* 2005; 30: 505
74. Standring S et al. *Gray's Anatomy: The Anatomical Basis of Clinical Practice*, pectoral girdle, shoulder region and axilla, 37th ed. Elsevier: Churchill Livingstone; 2005: 846-849.
75. Ting P L, Sivagnanaratnam V. Ultrasonographic study of the spread of local anaesthetic during axillary brachial plexus block. *Br J Anaesth* 1989; 63: 326-329.

References

76. Tsui B C, Dillane D, Pillay J et al. Cadaveric ultrasound imaging for training in ultrasound-guided peripheral nerve blocks: lower extremity. *Can J Anaesth* 2007;54:475-480.
77. Tsui BC, Finucane BT: The importance of ultrasound landmarks: a "traceback" approach using the popliteal blood vessels for identification of the sciatic nerve. *Reg Anesth. Pain Med.* 2006; 31: 481-2.
78. Urmey W, Stanton J. Inability to consistently elicit a motor response following sensory paresthesia during interscalene block administration. *Anesthesiology* 2002; 96: 552-554.
79. Vallieres E. The costovertebral angle. *Thorac Surg Clin* 2007;17:503-510.
80. Williams SR, Chouinard P, Arcand G, Harris P, Ruel M, Boudreault D, Girard F: Ultrasound guidance speeds execution and improves the quality of supraclavicular block. *Anesth Analg* 2003; 97: 1518-23
81. Winnie AP, Hakansson L. *Plexus Anaesthesia: Perivascular Techniques of Brachial Plexus Blocks*. Revised Edition. Philadelphia: Saunders; 1994,
82. Yang WT, Chui PT, Metreweli C: Anatomy of the normal brachial plexus revealed by sonography and the role of sonographic guidance in anesthesia of the brachial plexus. *Am J Roentgenol* 1998; 171: 1631-6.