

References

Adesanya AO, De Lemos JA, Greilich NB, et al. Management of perioperative myocardial infarction in noncardiac surgical patients. *Chest* **2006**; 130:584–596.

Aitkenhaed AR, Smith G and Rowbotham DJ. The post operative care. text book of Anaesthesia. 5th edition. Chapter 24, 484: 510. Churchill living stone. New York. London. **2006**.

Akhtar S, Barash PG. Perioperative use of beta-blockers: past, present, and future. *Int Anesthesiol Clin* **2002**; 40:133–157.

Albala MZ, Ferrigno M: Short-term noninvasive ventilation in the postanesthesia care unit: A case series. *J Clin Anesth* 2005; 17:636-639.

Albrecht RF, Wagner SR, Leicht CH, et al. Factitious disorder as a cause of failure to awaken after general anesthesia. *Anesthesiology* 1995;83:201–204.

Aldrete J. The post-anesthesia recovery score revisited. *J Clin Anesth* 1995; 7:89.

Aldrete JA. The post-anesthesia recovery score revisited. *J Clin Anesth* 1995; 7:89-91.

American Sleep Apnea Association: Sleep apnea and BMI. The majority of OSA patients are not obese. Available at www.sleepapnea.org 2007.

American Society of Anesthesiologists Task Force on Intraoperative Awareness. Practice advisory for intraoperative awareness and brain function monitoring: a report by the American Society of Anesthesiologists Task Force on Intraoperative Awareness. *Anesthesiology* 2006;104:847–864.

American Society of Anesthesiologists Task Force on Postanesthetic Care. Practice guidelines for postanesthetic care: a report by the American Society of Anesthesiologists Task Force on Postanesthetic Care. *Anesthesiology* 2002; 96:742–752.

Anwari JS. Quality of handover to the postanaesthesia care unit nurse. *Anaesthesia* 2002; 57:488–493.

Apfel CC, Korttila K, Abdalla M, et al. A factorial trial of six interventions for the prevention of postoperative nausea and vomiting. *N Engl J Med*, 10 2004; 350:2441– 2451.

Apfel CC, Kranke P, Eberhart LH. Comparison of surgical site and patient's history with a simplified risk score for the prediction of postoperative nausea and vomiting. *Anaesthesia*, 2004; 59:1078–1082.

Apfelbaum JL, Chen C, Mehta SS, et al.: Postoperative pain experience: results from a national survey suggest postoperative pain continues to be undermanaged. *Anesth Analg* 2003;97:534–540.

Assadian O, Toma CD, Rowley SD: Implications of staffing ratios and workload limitations on healthcare-associated infections and the quality of patient care. *Crit Care Med* 2007; 35:296-298.

Awad IT, Chung F. Factors affecting recovery and discharge following ambulatory surgery. *Can J Anaesth* 2006; 53:858– 872.

Badner NH, Knill RL, Brown JE, et al. Myocardial infarction after noncardiac surgery. *Anesthesiology* 1998;88:572–578.

Barash PG, Cullen PG, Stoelting BF, et al. Handbook of Clinical Anesthesia, 6th Edition. Paul G. Barash; Bruce F. Cullen; Robert K. Stoelting; Michael K. Cahalan; M. Christine Stock (eds.), ippincott Williams & Wilkins, Philadelphia, 530 Walnut St., Philadelphia, PA 19106. 2009.

Baxter BT, Thompson JS, Allison JG, et al.: Temporal patterns of postoperative complications. *Arch Surg* 2003;138:596–603.

Beauregard CL, Friedman WA. Routine use of postoperative ICU care for elective craniotomy: A cost-benefit analysis. *Surg Neurol* 2003; 60:483-489.discussion 489

Benumof JL: Obstructive sleep apnea in the adult obese patient: Implications for airway management. *J Clin Anesth* 2001; 13:144-156.

Borgeat A, Ekatodramis G, Schenker CA. Postoperative nausea and vomiting in regional anesthesia: a review. *Anesthesiology* 2003; 98:530–547.

Brennan TJ, Zahn PK, Pogatzki-Zahn EM: Mechanisms of incisional pain. *Anesthesiol Clin North America* 2005;23:1–20.

Brodsky JB, Lemmens HJ, Brock-Utne JG, et al: Morbid obesity and tracheal intubation. *Anesth Analg* 2002; 94:732-736.

Buggy DJ, Crossley AW: Thermoregulation, mild perioperative hypothermia and postanaesthetic shivering. *Br J Anaesth* 2000; 84:615-628.

Buttner M, Walder B, von Elm E, et al. Is low-dose haloperidol a useful antiemetic?: a meta-analysis of published and unpublished randomized trials. *Anesthesiology* 2004;101:1454–1463.

Caples SM, Gay PC. Noninvasive positive pressure ventilation in the intensive care unit: a concise review. *Crit Care Med* 2005; 33:2651–2658.

Carahan JM: Recovery room for patients. *Am J Nurs* 49:581, 1949.

Carlson K. Night watch. *J Perianesth Nurs* 2005; 20:399–403.

Cashman JN, Dolin SJ. Respiratory and haemodynamic effects of acute postoperative pain management: evidence from published data. *Br J Anaesth* 2004;93:212–223.

Charbit B, Albaladejo P, Funck-Brentano C, et al. Prolongation of QTc interval after postoperative nausea and vomiting treatment by droperidol or ondansetron. *Anesthesiology* 2005;102:1094–1100.

Cullen DJ: Obstructive sleep apnea and postoperative analgesia—a potentially dangerous combination. *J Clin Anesth* 2001; 13:83-85.

Davidson JA, Hosie HE: Limitations of pulse oximetry: respiratory insufficiency—a failure of detection. *BMJ* 1993; 307:372-373.

Debaene B, Plaud B, Dilly MP, et al. Residual paralysis in the PACU after a single intubating dose of nondepolarizing muscle relaxant with an intermediate duration of action. *Anesthesiology* 2003; 98:1042–1048.

Dershwitz M, Michalowski P, Chang Y, et al. Postoperative nausea and vomiting after total intravenous anesthesia with propofol and remifentanyl or alfentanil: how important is the opioid? *J Clin Anesth* 2002; 14:275–278.

Dexter F, Wachtel RE, Epstein RH. Impact of average patient acuity on staffing of the phase I PACU. *J Perianesth Nurs* 2006;21:303–310.

Dhonneur G, Combes X, Leroux B, Duvaldestin P: Postoperative obstructive apnea. *Anesth Analg* 1999; 89:762-767.

Diaconescu D, Grecu L. The post anesthesia care unit. In Dunn PF, ed. *Clinical Anesthesia Procedures of the Massachusetts General Hospital*. 7th ed. Philadelphia: Lippincott, Williams & Wilkins, 2007:623–644.

Dunn PF. Massachusetts General Hospital Beta Blockade Guidelines. Available at: <http://ccmu.massgeneral.org/pathway>. Last accessed 3/15/07.

Eagle KA, Berger PB, Calkins H, et al. ACC/AHA guideline update for perioperative cardiovascular evaluation for noncardiac surgery - executive summary. A report of the American College of Cardiology/ American Heart Association task force on practice guidelines. *Anesth Analg* 2002; 94:1052–1064.

Eagle KA, Berger PB, Calkins H, et al. ACC/AHA guideline update for perioperative cardiovascular evaluation for noncardiac surgery- executive summary. A report of the American College of Cardiology/American Heart Association task force on practice guidelines. *Anesth Analg* 2002; 94:1052–1064.

Edler AA, Mrino ER, Golianu B Kuan C, Pentcheva K. An analysis of factor influencing postanesthesia recovery after pediatric ambulatory tonsillectomy and adenoidectomy. *Anesth Analg*. 2007 Apr; 104 (4): 784-9.

Eikermann M, Blobner M, Groeben H, et al. Postoperative upper airway obstruction after recovery of the train of four ratio of the adductor pollicis muscle from neuromuscular blockade. *Anesth Analg* 2006; 102:937–942.

Finfer S, Bellomo R, Boyce N, et al. A comparison of albumin and saline for fluid resuscitation in the intensive care unit. *N Engl J Med* 27 2004; 350:2247–2256.

Flisberg P, Rudin A, Linner R, et al. Pain relief and safety after major surgery. A prospective study of epidural and intravenous analgesia in 2696 patients. *Acta Anaesthesiol Scand* 2003; 47:457–465.

Frank SM, Kluger MJ, Kunkel SL. Elevated thermostatic setpoint in postoperative patients. *Anesthesiology* 2000;93:1426–1431.

Fu ES, Downs JB, Schweiger JW, et al. Supplemental oxygen impairs detection of hypoventilation by pulse oximetry. *Chest* 2004; 126: 1552–1558.

Gajraj NM, Joshi GP. Role of cyclooxygenase- 2 inhibitors in postoperative pain management. *Anesthesiol Clin North America* 2005; 23:49–72.

Gan TJ, Ing RJ, de L Dear G, et al. How much are patients willing to pay to avoid intraoperative awareness? *J Clin Anesth* 2003; 15:108–112.

Gan TJ, Lubarsky DA, Flood EM, et al. Patient preferences for acute pain treatment. *Br J Anaesth* 2004; 92:681–688.

Gan TJ, Meyer T, Apfel CC, et al. Consensus guidelines for managing postoperative nausea and vomiting. *Anesth Analg* 2003; 97:62–71.

Gan TJ, Soppitt A, Maroof M, et al. Goaldirected intraoperative fluid administration reduces length of hospital stay after major surgery. *Anesthesiology* 2002; 97:820–826.

Gan TJ. Risk factors for postoperative nausea and vomiting. *Anesth Analg* 2006; 102:1884–1898.

Gravenstein D: Transurethral resection of the prostate (TURP) syndrome: a review of pathophysiology and management. *Anesth Analg* 1997; 84:438–446.

Gross JB, Bachenberg KL, Benumof JL, et al.: Practice guidelines for the perioperative management of patients with obstructive sleep apnea: A report by the American Society of Anesthesiologists Task Force on perioperative management of patients with obstructive sleep apnea. *Anesthesiology* 2006; 104:1081-1093.

Haas CE, LeBlanc JM. Acute postoperative hypertension: a review of therapeutic options. *Am J Health-Syst Pharm* 2004; 61: 1661–1673.

Habib AS, Chen YT, Taguchi A, et al. Postoperative nausea and vomiting following inpatient surgeries in a teaching hospital: a retrospective database analysis. *Curr Med Res Opin* 2006; 22:1093–1099.

Hedenstierna G, Rothen HU: Atelectasis formation during anesthesia: causes and measures to prevent it. J Clin Monit Comput 2000;16:329–335.

Hedenstierna G, Rothen HU: Atelectasis formation during anesthesia: causes and measures to prevent it. J Clin Monit Comput 2000; 16:329–335.

Heland M, Retsas A. Establishing a cardiac surgery recovery unit within the post anaesthesia care unit. Collegian 1999; 6:10-13.

Hepner DL, Castells MC: Anaphylaxis during the perioperative period . Anesth Analg 2003 :97:1381- 1395

Hillberg RE, Johnson DC: Noninvasive ventilation. N Engl J Med 1997; 337:1746-1752.

Hillman DR, Platt PR, Eastwood PR: The upper airway during anaesthesia. Br J Anaesth 2003; 91:31–39.

Hines R, Barash PG, Watrous G, et al.: Complications occurring in the postoperative care unit: a survey. Anesth Analg 1992; 74:503–509.

Hodge S. Provision of care for patients. Mechanical ventilation in the PACU. J Periop Pract 2006;16:376.

Hollenberg SM, Dellinger RP: Noncardiac surgery: postoperative arrhythmias. Crit Care Med 2000; 28:N145–N150.

Hollenberg SM, Dellinger RP: Noncardiac surgery: Postoperative arrhythmias. Crit Care Med 2000; 28(10 Suppl): N145-N150.

Huerta S, DeShields S, Shpiner R, et al: Safety and efficacy of postoperative continuous positive airway pressure to prevent pulmonary

complications after Roux-en-Y gastric bypass. J Gastrointest Surg 2002; 6:354-358.

Hugonnet S, Chevrolet JC, Pittet D: The effect of workload on infection risk in critically ill patients. Crit Care Med 2007; 35:76-81.

Hugonnet S, Perneger TV, Pittet D: Alcohol-based handrub improves compliance with hand hygiene in intensive care units. Arch Intern Med 2002; 162:1037-1043.

Irvin SM: Treatment of depression with outpatient electroconvulsive therapy. AORN J 1997; 65:573-578.581-572.

Kabbara A, Goldlust SA, Smith CE, et al.: Randomized prospective comparison of forced air warming using hospital blankets versus commercial blankets in surgical patients. Anesthesiology 2002; 97:338–344.

Keita H, Diouf E, Tubach F, et al.: Predictive factors of early postoperative urinary retention in the postanesthesia care unit. Anesth Analg 2005; 101:592–596.

Keita H, Diouf E, Tubach F, et al.: Predictive factors of early postoperative urinary retention in the postanesthesia care unit. Anesth Analg 2005; 101:592-596.

Kelsaka E, Baris S, Karakaya D, Sarihasan B: Comparison of ondansetron and meperidine for prevention of shivering in patients undergoing spinal anesthesia. Reg Anesth Pain Med 2006; 31:40-45.

Kiekkas P, Pouloupoulou M, Papahatzi A, et al.: Workload of postanaesthesia care unit nurses and intensive care overflow. Br J Nurs 2005;14:434–438.

Kluger MT, Bullock MF: Recovery room incidents: a review of 419 reports from the Anaesthetic Incident Monitoring Study (AIMS). *Anaesthesia* 2002; 57:1060–1066.

Kranke P, Eberhart LH, Roewer N, et al.: Pharmacological treatment of postoperative shivering: a quantitative systematic review of randomized controlled trials. *Anesth Analg* 2002;94:453–460.

Kriner EJ, Shafazand S, Colice GL: The endotracheal tube cuff-leak test as a predictor for postextubation stridor. *Respir Care* 2005; 50:1632–1638.

Lamonerie L, Marret, E, Deleuze A, et al.: Prevalence of postoperative bladder distension and urinary retention detected by ultrasound measurement. *Br J Anaesth* 2004; 92:544–546.

Landry DW, Levin HR, Gallant EM, et al.: Vasopressin deficiency contributes to the vasodilation of septic shock. *Circulation* 1997; 95:1122-1125.

Lepouse C, Lautner CA, Liu L, et al.: Emergence delirium in adults in the postanesthesia care unit. *Br J Anaesth* 2006; 96:747–753.

Lindenauer PK, Pekow P, Wang K, et al.: Perioperative beta-blocker therapy and mortality after major noncardiac surgery. *N Engl J Med* 2005; 353:349–361.

Lindsay M: Is the postanesthesia care unit becoming an intensive care unit?. *J Perianesth Nurs* 1999; 14:73-77.

Lohan D, Walsh S, McLoughlin R, et al.: Imaging of the complications of laparoscopic cholecystectomy. *Eur Radiol* 2005; 15:904–912.

Lovstad RZ, Granhus G, Hetland S: Bradycardia and asystolic cardiac arrest during spinal anaesthesia: A report of five cases. *Acta Anaesthesiol Scand* 2000; 44:48-52.

Macario A, Weinger M, Carney S, et al.: Which clinical anesthesia outcomes are important to avoid? The perspective of patients. *Anesth Analg* 1999; 89:652–658.

Mandell MS, Lezotte D, Kam I, Zamudio S: Reduced use of intensive care after liver transplantation: Influence of early extubation. *Liver Transpl* 2002; 8:676-681.

Mandell MS, Lezotte D, Kam I, Zamudio S: Reduced use of intensive care after liver transplantation: Patient attributes that determine early transfer to surgical wards. *Liver Transpl* 2002; 8:682-687.

Mangano DT, Layug EL, Wallace A, et al.: Effect of atenolol on mortality and cardiovascular morbidity after noncardiac surgery. Multicenter Study of Perioperative Ischemia Research Group. [published erratum appears in *N Engl J Med* 1997;336: 1039]. *N Engl J Med* 1996;335:1713–1720.

Marshall SI, Chung F: Discharge criteria and complications after ambulatory surgery. *Anesth Analg* 1999;88:508–517.

Mathes DD, Conaway MR, Ross WT: Ambulatory surgery: Room air versus nasal cannula oxygen during transport after general anesthesia. *Anesth Analg* 2001; 93:917-921.

Mayson KV, Beestra JE, Choi PT. The incidence of postoperative complications in the PACU. *Can J Anesth* 2005; 52:A62.

Metules TJ, Bauer J: Part 2. JCAHO's patient safety goals: preventing med errors. *RN* 2007; 70:39–43.

Mevorach DL: The management and treatment of recurrent postoperative laryngospasm, *Anesth Analg* 1996; 83:1110–1111.

Mitra S, Sinatra RS: Perioperative management of acute pain in the opioid dependent patient. *Anesthesiology* 2004;101:212–227.

Moss J: Allergic to anesthetics. *Anesthesiology* 2003; 99:521-523.

Myles PS, Williams DL, Hendrata M, et al.: Patient satisfaction after anaesthesia and surgery: results of a prospective survey of 10811 patients. *Br J Anaesth* 2000; 84:6–10.

Nightingale F: Notes on Hospitals, 3rd ed. London, Longman, Green, Longman, Roberts and Green, 1863.

Parikh S, Cohen JR: Perioperative stroke after general surgical procedures. *N Y S J Med* 1993; 93:162–165.

Patel BM, Chittock DR, Russell JA, Walley KR: Beneficial effects of short-term vasopressin infusion during severe septic shock. *Anesthesiology* 2002; 96:576-582.

Petty DS: ECT in the PACU? It's possible. *Nurs Manage* 2000; 31:42-44.

Picheansathian W: A systematic review on the effectiveness of alcohol-based solutions for hand hygiene. *Int J Nurs Pract* 2004; 10: 3-9.

Pittet D, Stephan F, Hugonnet S, et al.: Hand-cleansing during postanesthesia care. *Anesthesiology* 2003; 99:530-535.

Practice Guidelines for Postanesthetic Care: A report by the American Society of Anesthesiologists Task Force on Postanesthetic Care. *Anesthesiology* 2002; 96:742-752.

Rapp SE, Ready LB, Nessly ML: Acute pain management in patients with prior opioid consumption: a case-controlled retrospective review. *Pain* 1995;61:195–201.

Rinfret S, Goldman L, Polanczyk CA, et al.: Value of immediate postoperative electrocardiogram to update risk stratification after major noncardiac surgery. *Am J Cardiol* 2004; 94:1017-1022.

Rock P, Rich PB: Postoperative pulmonary complications. *Curr Opin Anaesthesiol* 2003; 16:123-131.

Rosevear HM, Montgomery JS, Roberts WW, et al.: Characterization and management of postoperative hemorrhage following upper retroperitoneal laparoscopic surgery. *J Urol* 2006;176(4 Pt 1):1458–1462.

Rosseland LA, Stubhaug A, Breivik H: Detecting postoperative urinary retention with an ultrasound scanner. *Acta Anaesthesiol Scand* 2002; 46:279-282.

Rusch D, Eberhart L, Biedler A, et al.: Prospective application of a simplified risk score to prevent postoperative nausea and vomiting. *Can J Anaesth* 2005; 52:478–484.

Saastamoinen P, Piispa M, Niskanen M: Use of postanesthesia care unit for purposes other than postanesthesia observation. *J Peri Anesth Nurs* 2007; 22:102-107.

Sagir O, Gulhas N, Toprak H, et al.: Control of shivering during regional anaesthesia: Prophylactic ketamine and granisetron. *Acta Anaesthesiol Scand* 2007; 51:44-49.

Schweizer A, Khatchatourian G, Hohn L, et al.: Opening of a new postanesthesia care unit: Impact on critical care utilization and complications following major vascular and thoracic surgery. *J Clin Anesth* 2002; 14:486-493.

Sebel PS, Bowdle TA, Ghoneim MM, et al.: The incidence of awareness during anesthesia: a multicenter United States study. *Anesth Analg* 2004; 99:833–839.

Sedrakyan A, Treasure T, Browne J, et al.: Pharmacologic prophylaxis for postoperative atrial tachyarrhythmia in general thoracic surgery: evidence from randomized clinical trials. *J Thorac Cardiovasc Surg* 2005;129:997–1005.

Selim M: Perioperative stroke. *N Engl J Med* 2007; 356: 706–713.

Sessler DI: Complications and treatment of mild hypothermia. *Anesthesiology* 2001; 95:531–543.

Siddiqui N, Teresi J, Arzola C, Fox G: Hypoxemia on arrival to PACU: An observational audit. *Can J Anesth* 2006; 53(Suppl 1):

Siyam MA, Benhamou D: Difficult endotracheal intubation in patients with sleep apnea syndrome. *Anesth Analg* 2002; 95:1098-1102.

Sloan SB, Weitz HH: Postoperative arrhythmias and conduction disorders. *Med Clin North Am* 2001;85:1171–1189.

Squadrone V, Coha M, Cerutti E, et al.: Continuous positive airway pressure for treatment of postoperative hypoxemia: A randomized controlled trial. *JAMA* 2005; 293:589-595.

Sullivan EE: Standards of perianesthesia nursing practice 2002. J Perianesth Nurs 2002;17:275–276.

Swenson JD, Davis JJ, Johnson KB: Postoperative care of the chronic opioid-consuming patient. Anesthesiol Clin North America 2005; 23:37–48.

Thompson JS, Baxter BT, Allison JG, et al.: Temporal patterns of postoperative complications. Arch Surg 2003;138:596–603.

Tobias JD: Noninvasive ventilation using bilevel positive airway pressure to treat impending respiratory failure in the postanesthesia care unit. J Clin Anesth 2000; 12:409-412.

Tomes CS: The administration of ether in America. BMJ 1:297, 1873.

Torn M, Rosenthal FR: Oral anticoagulation in surgical procedures: risks and recommendations. Br J Haematol 2003;123:676–682.

Tornero-Campello G: Letter: randomized clinical trial to evaluate the effects of perioperative supplemental oxygen administration on the colorectal anastomosis (Br J Surg 2006; 93:698–706). Br J Surg 2006;93:1148.

Truong L, Moran JL, Blum P: Post anaesthesia care unit discharge: a clinical scoring system versus traditional time-based criteria. Anaesth Intensive Care 2004; 32:33–42.

Tyler IL, Tantisira B, Winter PM, et al.: Continuous monitoring of arterial oxygen saturation with pulse oximetry during transfer to the recovery room. Anesth Analg 1985;64:1108.

Vaporciyan AA, Correa AM, Rice DC, et al.: Risk factors associated with atrial fibrillation after noncardiac thoracic surgery: Analysis of 2588 patients. *J Thorac Cardiovasc Surg* 2004;127:779-786.

Vidhani K, Langham BT: Obstructive sleep apnoea syndrome: Is this an overlooked cause of desaturation in the immediate postoperative period?. *Br J Anaesth* 1997; 78:442-443.

Vincent JL, Piagnerelli M: Transfusion in the intensive care unit. *Crit Care Med* 2006; 34(5 Suppl):S96–S101.

Walker JR: Anesthesia for cardioversion. *J Perianesth Nurs* 1999; 14:35-38.

Wang K, Asinger RW, Marriott HJ: STsegment elevation in conditions other than acute myocardial infarction. *N Engl J Med* 27 2003; 349:2128–2135.

Warner MA, Warner ME, Weber JG: Clinical significance of pulmonary aspiration during the perioperative period. *Anesthesiology* 1993; 78:56–62.

Weissman C: The enhanced postoperative care system. *J Clin Anesth* 2005; 17:314-322.

White PF, Rawal S, Nguyen J, et al.: PACU fast-tracking: an alternative to “bypassing” the PACU for facilitating the recovery process after ambulatory surgery. *J Perianesth Nurs* 2003; 18:247–253.

Witting MD, Hsu S, Granja CA: The sensitivity of room-air pulse oximetry in the detection of hypercapnia. *Am J Emerg Med* 2005; 23:497-500.

Zeitlin G: Recovery room mishaps in the ASA closed claims study. ASA Newsletter 1989; 53(7):

Ziser A, Alkobi M, Markovits R, Rozenberg B: The postanesthesia care unit as a temporary admission location due to intensive care and ward overflow. Br J Anaesth 2002; 88:577-579.

Zuck D: Anaesthetic and postoperative recovery rooms: Some notes on their early history. Anaesthesia 50:435, 1995.