

REFERENCES

REFERENCES

- Addison, G.M.; Beamish, M.R.; Hales, C.U.; Hodgkins, M.; Jacobs, A.; and Liewellin. (1972) :*
An immunoradiometric assay for ferritin in the serum of normal subjects and patients with iron deficiency and iron overload.
J. Clin. Path . 25. 326 - 329.
- Anoussakis, D. (1977):*
Endocrinological investigation of pituitary gonadal axis in thalassemia major.
Acta. Paediat Scand, 66 : 49.
- Arcasey, A. and Cavdar, A.O. (1975) :*
Changes of trace minerals (serum iron, zinc, copper and magnesium) in thalassemia.
Acta hematol., 53 : 341.
- Aruanitikis, C.; and Cook, A.R. (1978) :*
Diagnostic tests of exocrine pancreatic function and diseases.
Gastroenterology. 74 : 932.
- Asakura A. and Schwartz B. (1990) :*
Human hemoglobin in Miller and Baehner : 6th ed, Blood diseases of infancy and childhood. The C.V. Mosby Company, 358.
- Asbell, M.B. (1964):*
Orthodontic aspects of cooley's anemia.
Ann. NY Acad. Sci., 119 : 6621.
- Atti, G.; Capra, L.; DeSanctis, V.; Vullo, C. and Bagni, B. (1983):*
Beta - cell function assessed by plasma C - peptide evaluation in diabetic thalassemic patients.
Helv - Pediatr - Acta., 38 (2) : 123 - 32.
- Balcerzak, S.P.; Mintz, D.H. and Westerman, M.P. (1968):*
Diabetes mellitus and idiopathic hemochromatosis.
Am. J. Med. Sci., 255 L 53 - 62.

Bank, A. and Marks, P.A. (1966):

Excess α Chain synthesis relative to β Chain synthesis in thalassemia major and minor.
Nature, 212 : 1198.

Bard, H. and Prosmanne, J. (1982):

Postnatal fetal and adult hemoglobin synthesis in preterm infants whose birth weight was less than 1000 grams.
J. Clin. Invest., 70 : 50.

Barry, M. Flynn, D.M.; Ietsky, E.A. and Rison, R.A.; (1974) :

Long - term chelation therapy in thalassemia major : Effect on liver iron concentration, liver histology and clinical progress.
Br Med J. 2 : 16

Barton, B. (1976):

Pancreatic imaging.
Con. J. Radiog. Radiother.
Nucl. Med., 7, 72 - 80.

Baty, J.M.; Blackfan, K.D. and Diamond, L.K. (1932):

Blood studies in infants and in children, I. Erythroblastic anemia : a clinical and pathologic study.
Am. J. Dis. Child., 43 : 665, 1932.

Beard, M.E.J.; (1969) :

Intensive transfusion therapy in thalassemia major.
Pediatrics. 40 : 912

Bell, E.T. (1955):

The relation of portal cirrhosis to hemochromatosis and to diabetes mellitus.
Diabetes. 4 : 485 - 446.

Benz, E.J.Jr and Forget, B.G. (1971):

Defect in messenger RNA for human hemoglobin synthesis in beta thalassemia.
J. Clin invest., 50 : 2755.

Benz, E.J. Jr and Forget, B.G. (1974):

The biosynthesis of hemoglobin.
Semin Hematol., 11 : 463.

Blendis, L.M. (1974) :

Some effects of splenectomy in thalassemia major.
Br. J Haematol 28 : 77

Borgna - Pignatti, C.; Zonata L. and Bongo, I. (1983):

Red blood cell indices in adults and children with heterozygous beta- thalassemia.
Hematologica 68 : 149.

Borgna - Pignatti C.; Destefano, P. and Zonta, M.S. (1985):

Growth and sexual maturation in thalassemia major.
J. pediatr., 106 : 150.

Borgstrom, A.; and Ohlssan, K. (1976) :

Radioimmunological determination and characterization of cathedol trypsin. Like immunoreactivity in normal human plasma.
Scand J. Clin. Lab. Invest. 36 : 809.

Bunn, H.F. and Forget, B.G. (1986):

Hemoglobin : Molecular, Genetics and clinical aspects.
W.B. Saunders Company.

Caffey, J. (1957):

Cooley's anemia : A review of the roentgenographic findings in the skeleton.
AJR., 78 : 381.

Canale, V.C. (1974):

Endocrine function in the thalassemia major.
NY Acad, Sci., 232 : 333.

Capp, G.L.; Riggs, D.A. and Jones, R.T. (1970) :

Evidence for a new hemoglobin chain (ζ chain).
Nature 228 : 278.

Cartwright, G.E. (1948):

Studies on free erythrocyte protoporphyrin, Plasma iron and plasma copper in normal and anemic subjects.
Blood, 3 : 501.

Cavallo - Perin, P.; Pacini, G.; Cerutti, F.; Bessone, A.; Condo, C.; Sacchetti, L.; Piga, A. and Pagano, G. (1995) :
Insulin resistance and hyperinsulinemia in homozygous β -thalassemia.
Metabolism., 44 (3) : 281 - 6.

Cavill, I (1969) :
Erythropoiesis and the effect of transfusion in homozygous γ -thalassemia.
N.Eng. J. Med 298 : 776

Chormis, C. (1959) :
Amino acid tolerance curves and amino - aciduria in cooleys and

sickle - cell anemias.

J. Clin. Pathol., 12 : 245.

Chui, D.H.K.; Wong, S.C.; Chung, S.w.; Patterson, M.; Bhargava, S. and Poon, M.C. (1986):
Embryonic ζ - globin chains in adults : A marker for α - thalassemia -1 haplotype due to a > 17.5 - Kb deletion.
N.Engl. J. Med. 314 : 76.

Clegg, J.B. and Weatherall, D.J. (1972):
Hemoglobin synthesis during erythroid maturation in β -thalassemia.
Nature., 240 : 190.

Clegg, J.B. and Weatherall, D.J. (1974):
Hemoglobin constant spring, an unusual α - chain variant involved in the etiology of hemoglobin H disease.
Ann NY Acad. Sci., 232 : 168.

Clegg, J.B. and Weatherall, D.J. (1976):
Molecular basis of thalassemia.
Br. Med. Bull., 32 : 262.

Cohen, A.; Markenson, A.L. and Schwartz, E. (1980):
Transfusion requirements and splenectomy in thalassemia major.
J. Pediatr., 97 : 100.

Cohen, A.; Martin, M. and Schwartz, E. (1984):

Depletion of excessive liver iron stores with desferroxamine.
Br. J. Hematol., 58 : 369.

Cohen, A.; Mizanin, J. and Schwartz, E. (1985):

Treatment of iron overload.
Semin Liver. Dis., 4 : 228.

Collins, F.S. (1984):

Concordance of a point mutation 5' to the $G\gamma$ - globin gene with
 $G\gamma B$ Hereditary persistence of fetal hemoglobin in the black
population.
Blood, 64 : 1292.

Cook, J.D.; and Finch, C.A. (1979) :

Assessing iron status of a population.
Am. J. Clin. 32 : 2115

Cooley, T.B. and Lee, P. (1925):

Series of cases of splenomegaly in children with anemia and
peculiar bone changes.
Trans, Am. Pediatr. Soc., 47 : 29.

Coren, H.J. (1988):

Insulin action and diabetes.
New York, Raven Press.

Costin, G.; Kogut, M.D.; Hyman, C. and Ortega, J.A. (1977):

Carbohydrate metabolism and pancreatic islet Cell function in
thalassemia major.
Diabetes, 26 : 230 - 40.

Crystal, R.G. (1973):

Initiation of globin chain synthesis in β -Thalassemia.
J. Lab. Clin. Med., 288 : 1096.

Curtis, P.J.; Mantei, N. and Weissman, C. (1977):

Characterization and kinetics of synthesis of 155 B-Globin RNA,
a putative precursor of β -globin mRNA. Quant.
Biol., 42 : 971.

Cutello, S. and Meloni, T. (1974):

Serum concentration of heptoglobin and hemopexin in favism and thalassemia.

Acta. Hematol., 52 : 65.

Czech, M.P. (1984):

New perspectives on the mechanism of insulin action.

Recent prog. Horm. Res. 40 : 347.

Czech, M.P. (1985):

The nature and regulation of the insulin receptors structure and function.

Annu. Rev. Physiol., 47 : 357.

Dagg, J.H.; and Goldbeng, A. (1973) :

Detection and treatment of iron deficiency.

Clin Haematol. 2 : 365

Dandona, P.; Hussain, M.A.M., Varghese, Z.; Politis, D.; Flynn, D.M. and Hoffbrand, A.V. (1983):

Insulin resistance and iron overload.

Ann clin Biochem, 20 : 77 - 79.

DeGraff, C.S.; Taylor, K.J.W.; Simonds, B.D. and Rosarfie, A.J. (1978):

Gray scale echography of the pancreas. Re-evaluation of normal size.

Radiology, 129 : 157 - 161.

Deisseroth, A.; Nienhuis, A.; Turner, P.; Velez, R. and Anderson, W.F. (1977):

Localization of the human α - globin structural gene to chromosome 16 in somatic cell hybrids by molecular hybridization assay.

Cell, 12 : 205.

Deisseroth, A.; Nienhuis, A.; Lawrence, J.; Giles, R.; Turner, P. and Ruddle, F.H. (1978):

Chromosomal localization of human β - globin gene on human chromosome II in somatic cell hybride.

Proc Natl. A cad. Sci. USA., 75 : 1456.

- DeSanctis, V.; Zurlo, M.G.; Senesi, E.; Boffa, C.; Cavallo, L. and DiGregorio, F. (1988) :**
Insulin dependent diabetes in thalassemia.
Arch. Dis. Child., 63 : 58 - 62.
- Dewewy, K.W.; Grossman, H. and Canale, V.C. (1970):**
Cholelithiasis in thalassemia major.
Radiology, 96 : 385.
- Dmochowski, K.; Dianet, T.; Finegood, Francombe, W.; BevTyler and Zinman, B. (1993):**
Factors determining glucose tolerance in patients with thalassemia major.
J. Clin. Endocrinol. Metab., 77 : 478 - 483.
- Doust, B.D. and Pearce, J. D. (1976):**
Grey scale ultrasonic properties of the normal and inflamed pancreas.
Radiology, 120, 693 - 7.
- Dozy, A.M.; Kan, Y.W. and Emberg, S.H. (1979):**
 α Gene organization in blacks precludes the severe form of α thalassemia.
Nature 280 : 605.
- Dymock, I.W. and Williams, R. (1981):**
Hemochromatosis and diabetes.
Postgrad. Med. J. Suppl., 79 - 83.
- Efstratiadis, A.; Posakony, J.W.; Maniatis, T.; Lawn, R.M.; Connell, C.; Spritz, R.A.; DeRiel, J.; Forget, B.; Weissman, S.M.; Slighton, J.L.; Smithies, O.; Barlle, F.E.; Shoulders, C.C. and Prdfoot, N.J. (1980):**
The structure and evaluation of the human β - globin gene family
Cell, 21 : 653.
- Embury, S.H. (1980):**
Two different molecular organization account for the single α - globin gene of the α - thalassemia - 2 genotype.
J. Clin. Invest., 66 : 1319.

Engl, M.A. (1964):

Cardiac involvement in cooley's anemia,
Ann. NY, Acad. Sci., 119 : 694.

Erlandson, M.E. (1964):

Comparison of sixty - six patients with thalassemia major and thirteen patients with thalassemia intermedia : Includingg evaluation of growth, development, maturation and prognosis.
Ann. NY. Acad. Sci., 119 : 727.

Fain, J.N. (1984):

Insulin secretion and action.
Metabolism, 33 : 672.

Fessas, P.; Stamatoyannopoulos, G. and Karaklis, A. (1983):

Hemoglobin "Pylos": study of a hemoglobinopathy resembling thalassemia in the heterozygous homozygous and double heterozygous state.
Blood, 19 : 1.

Fessas, P.; Loukopoulos, D. and Kaltsoya, A. (1966):

Peptide analysis of the inclusions of erythroid cells in β - thalassemia. Biochem, Biophys. Acta., 124 : 439.

Fie, Y.J.; Stoming, T.A. and Kutler, A. (1989):

One form of inclusion body thalassemia is due to a GAA - TAA mutation at codon 121 of the α chain.
Blood. 73 : 1075.

Forget, B.G. (1979):

Molecular genetics of human hemoglobin synthesis.
Ann. Intern. Med., 91 : 605.

Frier, B.M.; Ashby, J.P.; Nairn, I.M.; and Bairs, J.D.; (1981):

Plasma insulin, C-peptide and glucagon concentrations in patients with insulin - independent diabetes treated with chloropamide.
Diab. Metab. 7, 1, 45 - 49.

Gale, R.E.; Clegg, J.B. and Huehns, E.R. (1979):

Human embryonic hemoglobins Gower 1 and Gower 2.
Nature 280 : 162.

Galanello, R. (1983):

A family with segregating triplicated α - globin loci and β -thalassemia.
Blood, 62 : 1035.

Gammeltoft, S. (1984):

Insulin receptors : Binding kinetics and structure function relationship of insulin.
Physiol. Rev., 64 : 1321.

Geiger, R. and Fritzy, H (1984) :

Trypsin in methods of enzymatic analysis 3 rd ed. Vol. 5.H.V. Dargmeyer, J. Bergmeyer, M. Grassl, Eds weinhein, verlag chemie, PP. 119.

Gertner, J.M.; Broadus, A.E. and Anast, C.S.; (1979):

Impaired parathyroid response to induced hypocalcemia in thalassemia major.
J. Pediatr. 95 : 210.

Godet, J. (1977):

β^0 -thalassemia from Algeria : Genetic and molecular characterization.
Blood, 50 : 463.

Goldberg, D.M. (1983) :

Enzymes and isoenzymes in the evaluation of diseases of pancreas. In clinical and analytical concepts in enzymology. H.A. Homburger Ed. Skokie, II, College of American pathology PP. 31.

Gonzalez, Redonado, J.M.; (1988) :

Clinical and genetic heterogeneity in black patients with homozygous β -thalassemia from the southeastern united states.
Blood 72 : 1007.

Grant, G.P. (1986) :

The effect of transfusion on lung capacity, diffusion capacity and arterial oxygen saturation in patients with thalassemia major.
Pediatr. Res., 20 : 20.

Gratwick, G.M. (1978):

Thalassemic osteoarthropathy.
Ann. Intern. Med., 88 : 494.

Grossman, H.; Dischl, M.R.; Winchester, P.H. and Canali, V. (1971):

Renal enlargement in thalassemia major.
Radiology, 100 : 645.

Gullo, L.; Corcioni, E.; Brancati, C.; Bria, M. and Pezzilli, R. (1993):

Morphological and functional evaluation of the exocrine pancreas
in beta - thalassemia major.
Pancreas. Mar., 8 (2) 196 - 80.

**Harton, B.F.; Thompson, R.B.; Dozy, A.M.; Nechtman, C.M.; Nichols,
E. and Huisman, T.H.J. (1984):**

Inhomogeneity of hemoglobin VI. The minor hemoglobin
components of cord blood.
Blood 20 : 302.

Heding, L.G.; (1975) :

Radioimmunological determination of human C-peptide in serum.
Diabetologia. 29 : 321 - 332

Hershko, C. (1978):

Non - specific serum iron in thalassemia. An abnormal serum iron
fraction of potential toxicity.
Br. J. Hematol., 40 : 255.

Higgs, D.R. (1989):

A review of the molecular gene cluster.
Blood. 73 : 1081.

Hilgartner, M.W.; Erlandson, M.E. and Smith, C.H. (1963):

The coagulation mechanism in patients with thalassemia major.
J. Pediatr., 63 : 36.

Hilgartner, M.W. and Smith, C.H. (1964) :

Coagulation studies as a measure of liver function in cooley's
anemia.
Ann. NY, Acad. Sci., 119 : 631.

- Hoekstra, J.B.L.; Van Rijn, H.J.M. and Thijssen, J.H. H. (1982):**
C-peptide reactivity as a measure of insulin dependency in obese diabetic patients treated with insulin.
Diabetes care, 5 : 585 - 591.
- Horwitz, D.L.; Kuzuya, H.; and Rubenstein, A.H. (1976):**
Circulating serum C-peptide : A brief review of diagnostic implications.
N. Engl. J. Med., 295 : 207 - 209.
- Hosea, S.W.; Burch, C.G.; Brown, E.J.; and Berg, R.A. (1981):**
Impaired immune response in splenectomized patients to polyvalent pneumococcal vaccine.
Lancet 1 : 804
- Housman, D. (1973):**
Quantitative deficiency of chain specific globin messenger ribonucleic acids in the thalassemia syndromes.
Proc. Natl. Acad. Sci. USA, 70 : 1809.
- Hoyt, R.W.; Scarpa, N. and Wilmott, R.W. (1988):**
Pulmonary function abnormalities in homozygous thalassemia.
J. Pediatr., 109 : 452.
- Howell, S.L. and Tyhurst, M. (1984):**
Insulin secretion : The effector system.
Experientia, 40 : 1098.
- Huisman, T.H.J. (1971):**
Hereditary persistence of fetal hemoglobin.
N. Engl. J. Med., 285 - 711.
- Huisman, T.H.J.; Schroeder, W.A.; Efremov, G.D.; Duma, H.; Mladenoveski, F.; Human, C.B.; Rachmilewitz, E.A.; bouver, N.; Miller, A.; bordie, A.; Shelton, J.R.; Shelton, J.B. and Apell, G. (1974):**
The present status of the heterogeneity of fetal hemoglobin in β thalassemia : An attempt to unify some observations in thalassemia and related conditions.
Ann. NY, Acad. Sci., 232 : 107.

- Humphries, R.K.; Ley, T.Y.; Anagnou, N.P.; Bouer, A.W. and Nienhuis, A.W. (1984):**
 β^0 - 39 thalassemia gene : A premature termination codon causes β - m RNA without affecting cytoplasmic β - mRNA stability.
Blood 64 : 23.
- Jacobs, S. and Cuatrecasas, P. (1983):**
Insulin receptors.
Annu. Rev. Pharmacol. Toxicol: 23 : 461.
- Johnson, C.S.; Tegos, C. and Beutler, E. (1982):**
Alpha-thalassemia ; Prevalence and hematologic findings in American blacks.
Arch. Intern. Med., 142 : 1280.
- Jones, R.T. (1964):**
Structural studies on aminoethylated hemoglobins by automatic peptide chromatography.
Bio., 24 : 297.
- Junglee, D.; Katrak, A.; Hoffbrand, A.V. and Dandona, P. (1983):**
Pancreatic lipase in serum of patients with beta - thalassemia major. Clin. Chem. 29 : 2003 - 4.
- Kacian, D.L. (1973):**
Decreased globin messenger RNA in thalassemia detected by molecular hybridization.
Proc. Natl. Acad. Sci. USA, 70 : 1886.
- Kan, Y.W. (1975) :**
Demonstration of non functional β - globin mRNA in homozygous β^0 thalassemia.
Proc. Natl. Acad. Sci. USA, 72 : 5140.
- Kan, Y.W.; Dozy, A.M.; Trecartin, R. and Todd, D. (1977):**
Identification of non deletion defect in α - thalassemia.
N. Eng. J. Med. 297 : 1081.
- Karahanyan, E.; Stoyanova, A.; Moumdzhiev, I. and Ivanov, I. (1995):**
Secondary diabetes in children with thalassemia major.
Folia. Med. Plovdiv., 36 (1) : 29 - 35.

Kato, K. and Downey, H. (1933):

Hematology of erythroblastic anemia.
Folia hemtal serum 50 : 55.

Kattamis, C. (1979):

The heterogeneity of normal Hb A2 - β - thalassemia in Greece.
Br. J. Hematol., 42 : 104.

Katz, H.L.; Savage, P.J.; Barclay, - White, B. (1985):

C-peptide measurement in the differentiation of type I (insulin dependet) and type 2 (non - insulin - deperdent) diabetes mellitus.
Diabetologia, 28 : 264 - 268.

Kreimer - Birnabum M, Pinkerton, P.H. and Bonnerman, R.N. (1966):

Urinary dipyrooles, their occurrence and significance in thalassemia and other disorders.
Blood, 28 : 933.

Kreel, L.; Sandin, B. and Slavin, G. (1973):

Pancreatic morphology. A compined radiological and pathological study.
Clin. Radiol., 24 : 154 - 61.

Kuo, B.; Zaino, E. and Roginsky, M.S. (1968):

Endocrine function in thalassemia major.
J. Clin. Endocrinol. Metab., 28 : 805.

Lassman, M.N. and Genel, M. (1974):

Endocrine evaluation in thalassemia major.
Ann. NY, Acad. Sci. 232 : 226.

Lassman, M.N.; Genel, M.; Wise, J.K.; Hindler, R. and Felig, P. (1974):

Carbohydrate homeostasis and pancreatic isiet cell function in thalassemia.
Am. J. Med., 80 : 65.

Lawson, J.P.; Ablow, R.C. and Pearson, H.A. (1981):

The ribs in thalassemia. The relationship of therapy.
Radiology, 140 : 656.

Leopold, G.R. (1975):

Grey scale ultrasonic angiography of the upper abdomen.
Radiology, 117 : 665 - 71.

Lesi, C.; Melzid'Eril, G.V. and Pavesif (1985):

Clinical significance of serum pancreatic enzymes in the quiescent phase of chronic pancreatitis.
Clin Biochem 18 : 235 - 8.

Letsky, E.A. (1976):

A controlled trial of long - term chelation therapy in homozygous β -thalassemia. In iron Metabolism and thalassemia, eds D. Bergsma, A Cerami, CM Peterson, J.H. Graziano.
New York : Alan, R. Liss.

Logothetis, J. Constantoulakis, M. and Economidoud (1972):

Thalassemia major (homozygous beta - thalassemia) : a survey of 138 cases.
Neurology, 22 : 294.

Maioli, M.; Pacifice, A. and Frulio, T. (1983):

Study of A and B - cell function in beta - thalassemia major.
Acta - Diabetolo. Lat., 20 (2) : 143 - 52.

Maniatis, T.; Fritsch, E.F.; Laver, J. and Lawn, R.M. (1980):

The molecular genetics of human hemoglobin.
Annu. Rev. Genet., 14 L 145.

Mann, J.R. (1972) :

δ B-thalassemia in a chinese family.
Br. J. Heamatol., 23 : 393.

Mann Hussan, Dandona P; Fedall, S.S.; Ramdial, L.; Flynn, D.; and

Hoffbrand, A.V (1981) :

Serum immunoreactive trypsin in β -Thalassemia major.
J. clin. Path 34 : 970.

Marks, P.A. (1966):

Thalassemia syndromes : Biochemical, genetic and clinical aspects.
N. Eng. J. Med. 275 : 1363.

Masera, G. (1979):

Role of chronic hepatitis in development of thalassemic liver disease.

Arch. Dis. Child. 51 : 680.

Mauer, H.S.; (1988) :

A prospective evaluation of iron chelation therapy in children with severe B-thalassemia.

Am J. Dis child 142 : 1736.

McDonald. R. (1966) :

Deferoxamine and diethylenetriaminepenta acetic acid (DTPA) in thalassemia major.

Arch Dis child 51 : 195.

McIntosh, N. (1976) :

Endocrinopathy in thalassemia major.

Arch. Dis. Child., 51 : 195 - 201.

Meire, H.B. and Jarront, P. (1979) :

Pancreatic ultrasound - a systematic approach to scanning technique.

Brit. J. Radiol., 52 : 562 - 7.

Merkel, P.A.; Simorson, D.C.; Amiel, S.A.; Plewe, G.; Sherwin, R.S.;

Pearson, W.A. and William, V. (1988):

Insulin resistance and hyperinsulinemia in patients with thalassemia major treated by hypertransfusion.

N.Eng. J. Med. 318 : 809.

Meyer, J.H (1980) :

Pancreatic physiology. In : Gastrointestinal disease : pathophysiology, diagnosis, Management, 4th ed. M. Sleisengev, J.S. fardtran, Eds. philadelphia, W.B. saunders PP. 1777.

Modell, B. (1975) :

Tranfusional hemochromatosis.

In : King H, ed. iron metabolism and its disorders.

New York : Elsevier publishing Co. : 320.

Modell, B. (1976):

Management of thalassemia major.
Br. Med. Bull., 32 : 270.

Modell, B. (1977):

Total management of the thalassemia major.
Arch. Dis. Child., 52 : 484.

Modell, B. (1979) :

Advances in the use of iron - chelating agents for the treatment of iron overload.
Progr Hematol. 11 : 267.

Nakamura, T. (1987):

A δ - globin gene derived from patients with homozygous δ^0 - thalassemia function normally on transient expression in heterologous cells.
Blood, 70 : 804.

Nathan, D.G. (1972):

Thalassemia.
N. Eng. J. Med., 286 : 586.

Nathan, D.G.; Stossel, T.B.; Gunn, R.B.; Zarkowsky, H.S. and Lafort, M.T. (1969):

Influence of hemoglobin precipitation on erythrocyte metabolism in alpha and beta thalassemia.
J. Clin. Invest., 48 : 33.

National, Diabetes, Data Group, (1979):

Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance.
Diabetes 28 : 1039.

Niederau, C.; Berger, M. and Stremmal, W. (1984):

Hyperinsulinemia in non cirrhotic hemochromatosis. Impaired hepatic insulin degradation.

Nienhuis, A.W. and Anderson, W.F. (1973):

Isolation and Translation of hemoglobin messenger RNA from thalassemia, sickle cell anemia and norma human reticulocytes.
J. Clin. Invest., 52 : 1735.
Diabetologia, 26 : 441 - 4.

Nienhuis, A.W. and Benz, E.J. (1977):

Regulation of hemoglobin synthesis during the development of the red cell.

N. Eng. J. Med., 297 : 1318.

Nienhuis, A.W. and Marriettis, T. (1987) :

Structure and experssion of globin genes in erythroid cells. In the molecular Basis of Blood Disease ed.

G. stamatoyannopoulos, A.W. Nienhuisp. Leder, P.W. Majerus.

W.B. Saunders, philadelphia.

O Brien, R.T. (1972) :

Transfusion - induced decrease in spleen size in thalassemia major : Documentation by radioisotopic scan.

J. Pediatr 81 : 105.

Orkin, S.H. and Nathan, D.G. (1976):

Current concepts in gentitics : The thalassemias.

N. Engl. Med., 295 : 710.

Orkin, S.H. and Nathan, D.G. (1981):

The molecular genetics of thalassemia.

Adv. Hum. Genet., 11 : 233.

Perutz, M.F.; Rossman, M.G. and Gullis, A.F. (1960):

Structure of hemoglobin. A three dimentional fourier synthesis at 5.5 A resolution, obtained by X - ray analysis.

Nature, 185 : 416.

Petrsen, O.H. and Findlay, I. (1987):

Electrophysiology of the pancreas.

Physiol. Rev., 67 : 1054.

Petrsen, O.H. and Gallacher, D.V. (1988) :

Electrophysiology of pancreatic and salivary acinar cells.

Annu. Rev. Physiol., 50 : 65.

Polonsky, K.; Frank, B. and Pugh, W. (1986):

The limitations to and valid use of C - peptide as a marker of the secretion of insulin.

Diabetes, 35 : 379 - 386.

Pootrakul, S. (1981):

Thalassemia trait : Hematologic and hemoglobin synthesis studies.
Hemoglobin, 1 : 75.

Propper, R.D. and Nathan, D.G. (1977):

Desferrioxamine and the pump. In Zairo EC, and Roberts R.H, editors : Chelation therapy in chronic iron overload, Miami, Symposia specialists, Inc.

Revel and Mand Govner, Y. (1978):

Post transcriptional and translational controls of gene expression in all - karyocytes.
Ann. Rev. Biochem., 47 : 1079.

Risdon, R. (1975):

Transfusion iron overload : Relationship between Tissue iron concentration and hepatic fibrosis in thalassemia.
J. Pathol., 116 : 83.

Sacks, D.B.; (1994):

Carbohydrate in tietz textbook of clinical chemistry, Burtis C.A and Ashwood E.R. 2nd edition volume 2, 228, WB Saunders, Philadilphia, London, Toronto, Montrial, Sydney, Tokyo

Saenger, P. (1980):

Depressed senum somatomedin activity in β - thalassemia.
J. Pediatr, 96 : 214.

Sample, W.F. (1977):

Techniques for improved delineation of the normal anatomy of the upper abdomen and high retroperitoneum with grey scale ultrasound.
Radiology, 124, 147 - 202.

Schumacker, M.J.; (1970) :

Serum immunoglobulin and transferrin levels after childhood splenectomy.
Arch Dis child 45 : 114.

Schwartz,, M.K. and Fleischer, M. (1970) :

Diagnostic biochemical methods in pancreatic disease.
In advances in clinical chemistry. O.Bodansky and C.P. Stewart,
Eds. New York, Academic press, Vol., 13, P. 114

Smith, C.H. (1948) :

Familial blood studies in cases of Mediterranean (cooley's anemia)
Am J Dis child 65 : 681.

Sonakul, D.; Pacharee, P.; Wasi, P. and Fucharoen, S. (1984):

Cardiac pathology in 47 patients with beta thalassemia
hemoglobin E.
Southeast Asian J. Trop Med. Public Heath 15 : 554 - 63.

Saudek, C.D.; Hemm, R.M. and Peterson, C.M. (1977):

Abnormal glucose Tolerance in β thalassemia major.
Metabolism 26 : 43 - 52.

Schafer, A.I.; Cheron, R.G. and Dluhy, R. (1981) :

Clinical consequences of acquired transfusional iron overload in
adults.
N. Eng. J. med., 304 : 319 - 24.

Schroeder, W.a.; Huisman, T.H.J and Shelton, J.R. (1968) :

Evidence for multiple structural genes for γ chains of human fetal
hemoglobin.
Proc. Natl. Acad. Sci. USA, 60 : 537 - 1.

Shinar, E. (1987):

Erythrocyte membrane skeleton abnormalities in Severe
 β - Thalassemia.
Blood 70 : 158.

Slater, L.M.; Muri, W.A. and Weed, R.I. (1968):

Influence of splenectomy on insoluble hemoglobin inclusion
bodies in β thalassemic erythrocytes.
Blood 31 : 766.

Smith, C.h. (1960):

The role of splenectomy in the management of thalassemia.
Blood, 15 :197.

- Sonakul, D.; Pacharee, P.; Wasi, P. and Fucharoen, S. (1984):***
Cardiac pathology in 47 patients with beta thalassemia /
hemoglobin E.
Southeast Asian, J. Trop. Med. Public health, 15 : 554 - 63.
- Stamatoyannopoulos, G. and Nienhuis, A.W. (1981):***
Organization and expression of globin genes.
New York, Alan, R. Liss.
- Steinberg, M.H. and Adams, J.G. (1983):***
Thalassemic hemoglobinopathies.
Am. J. pathol., 113 : 396.
- Takeshita, K.; Torget, B.G.; Scarpa, A. and Benz, E.J.Jr. (1984):***
Intranuclear defect in β - globin mRNA accumulation due to a
premature translation termination codon.
Blood, 64 : 13.
- Temple, G.F. (1977):***
Authentic, β - globin mRNA sequences in homozygous β^0 -
thalassemia.
Proc. Natt. Acad. Sci. USA., 74 : 3047.
- Thompson, J.C. (1987):***
Gastrointestinal endocrinology.
New York, McGraw - Hill Book Co.
- Tietz, N.W. (1976):***
In fundamentals of clinical chemistry, 2nd ed. NW tietz, Editor,
Saunders, Philadelphia, PP. 924 - 929.
- Todd, D.; Lai, M.C.S.; Beavren, G.M. and Huchns, E.R. (1970):***
The abnormal hemoglobins in homozygous alpha thalassemia.
Br. J. Hematol., 14 : 27.
- Valentine W.N and Neel, J.V. (1944):***
Hematologic and genitic study of transmission of thalassemia.
Arch. Intern. Med., 74 : 185.

- Ventrucci, M.; Pezzilli, R.; Gullo, L.; Plate, L.; Sprovieri, G. and Barbaro, L. (1989) :**
Role of serum pancreatic enzyme assays in diagnosis of pancreatic diseases.
Dig. Dis. Sci., 34 : 39 - 45.
- Walters, G.O.; Miller, F.H. and Worwood, M. (1973):**
Serum ferritin concentration and iron stores in normal subjects.
J. Clini. Path., 26, 770 - 772.
- Wasi, P. (1971) :**
Streptococcal infection leading to cardiac and renal involvement in thalassemia.
Lancet, 1 : 949.
- Watson, D. (1962):**
Enzymatic determination of glucose and easily hydrolyzable glucose esters in blood.
Anal Biochem. 3 : 131.
- Weatherall, D.J. (1976):**
The molecular basis of thalassemia.
Johns, Hopkins, Med. J., 139 : 205.
- Weatherall, D.J. and Clegg, J.B. (1981):**
The thalassemia syndromes.
3rd ed. Oxford, England, Blackwell, Scientific Publications.
- Weatherall, D.J.; (1983) :**
The diagnostic features of the different forms of thalassemia.
In the thalassemias, ed DJ. Weatherall. Edinburgh : Churchill Livingstone.
- Werner, M.; Steinberg, M.W.; Pauley, C. (1989):**
Strategic use of individual and combined enzyme indicators for acute Pancreatitis analyzed by receiver - Operator characteristics.
Clin. Chem., 35 : 967.
- Wintrob, M.M.; Richard, Lee, G.; Bogts, D.R.; Biethell, T.C.; Foerster, J. and Athens, J.W. (1975):**
Production of erythrocytes clinical hematology: Seventh edition
Henry Kimpton on publishers, London, P. 135.

Witzleben, C.L. and Wyatt, J.P. (1961):

The effect of long survival on the pathology of thalassemia major.
J. Pathol., 82 : 1.

Wolf, J.A. and Luke, K.H. (1969) :

Management of thalassemia. A Comparative program.
Ann N Y Acad sci 165 : 423.

Wollman, I.J. and Ortalani, M. (1969):

Some clinical features of cooleys anemia patients as related to
transfusion schedules.
Ann. NY. Acad. Sci. 105 : 407 - 414.

Wood, W.G. (1976):

Hemoglobin synthesis during human fetal development.
Br. Med. Bull, 32 : 282.

Wood, W.G.; Clegg, J.B.; and Weatherall, D.J. (1979) :

Hereditary persistence of fetal hemoglobin (HPFH) and δ B.
Thalassemia.
Br. J. Haematol 43 : 509.

Zago, M.A.; Costa, F.F. and Bottura, C. (1982):

Thalassemia intermedia in a family with β zero thalassemia and
Hb Hasharon.
J Med Genet. 19 : 437.

Zaino, E.C. (1964):

Paleontologic thalassemia. Ann. NY, Acad, Sci., 114 : 402.

Zaino, E.C. (1971):

Gaucher's cells in thalassemia.
Blood, 38 : 457.

Zuppinger, K.; Molinavi, B. and Hirt, A. (1979):

Increased risk of diabetes mellitus in beta - Thalassemia major due
to iron overload.
Helv. Pediatr. Acta., 34 : 197 - 207.