



REFERENCES

- .
- Abdel-Zaher AO. (2000):** The myoneural effects of lithium chloride on the nerve-muscle preparations of rats: role of adenosine triphosphate-sensitive potassium channels. *Pharmacol Res*; 41: 163-78.
- Abrams R. (1997):** Electroconvulsive therapy. 3rd ed. New York: Oxford University Press.
- Adams MR, Brandon EP, Chartoff EH, et al. (1997):** Loss of haloperidol induced gene expression and catalepsy in protein kinase A deficient mice. *Proc Natl Acad Sci U S A*; 94: 12157-61.
- Andersen FA, Arslan D, Holst-Larsen H. (2001):** Effects of combined methohexitone-remifentanyl anaesthesia in electroconvulsive therapy. *Acta Anaesthesiol Scand*; 45 (7): 830-3.
- Andres RL. Social and illicit drug use in pregnancy. (1993):** In: Resnik RR, Cease R editors. *Maternal-fetal medicine*. Philadelphia: WB Saunders; pp.145–164.
- Angelina D., Castro (1998):** Management of anesthesia for specialty procedure. *Introduction to anesthesia*. David E. Longnecker and Frank L. Murphy (eds) 8th edition; 396-409.



- Avramov MN, Husain MM, White PF. (1995):** The comparative effects of methohexital, propofol, and etomidate for electroconvulsive therapy. *Anesth Analg*; 81 (3): 596-602.
- Bader GM, Silk KR, Dequardo JR, et al. (1995):** Electroconvulsive therapy and intracranial aneurysm. *Convuls Ther*; 11 (2): 139-143.
- Baldessarini RJ, Tondo L, Hennen J. (1999):** Effects of lithium treatment and its discontinuation on suicidal behavior in bipolar manic-depressive disorders. *J Clin Psychiatry*; 60: 77-84, discussion 111-116.
- Balon R, Mufti R, Arfken CL. (1999):** A survey of prescribing practices for monoamine oxidase inhibitors. *Psychiatr Serv*; 50: 945-947.
- Berrios G.E. (2004)** Of mania. *History of Psychiatry* 15: 105-124 .
- Bertelsen, A. (2002):** Schizophrenia and Related Disorders: Experience with Current Diagnostic Systems. *Psychopathology*, 35: 89–93
- Bhatia SC, Bhatia SK, Gupta S. (1998):** Concurrent administration of clozapine and ECT: a successful



References

therapeutic strategy for a patient with treatment-resistant schizophrenia. J ECT; 14 (4): 280-283.

Blanch J, Martinez-Palli G, Navines R, et al. (2001): Comparative hemodynamic effects of urapidil and labetalol after electroconvulsive therapy. J ECT 2001; 17 (4): 275-279.

Bland,R.C. (1997): "Epidemiology of Affective Disorders: A Review". Can J Psychiatry 42: 367-377.

Bouckoms AJ, Welch CA, Drop LJ, et al. (1989): Methohexital dosage in electroconvulsive therapy. Convuls Ther; 5 (1): 44-47.

Braga RJ, Petrides G. (2005): The combined use of electroconvulsive therapy and antipsychotics in patients with schizophrenia. J ECT; 21 (2): 75-83.

Carrera MR, Meijler MM, Janda KD.(2004): Cocaine pharmacology and current

pharmacotherapies for its abuse. Bioorg

Med Chem 2004; 12:5019–5030.

Cheam EW, Critchley LA, Chui PT, et al. (1999): Low dose mivacurium is less effective than succinylcholine in electroconvulsive therapy. Can J Anaesth; 46 (1): 49-51.



- Chen G, Zeng WZ, Yuan PX, et al. (1999):** The mood-stabilizing agents lithium and valproate robustly increase the levels of the neuroprotective protein bcl-2 in the CNS. *J Neurochem*; 72: 879-882.
- Clayton A, Keller A, McGarvey EL. (2006):** Burden of phase-specific sexual dysfunction with SSRIs. *J Affect Disord*; 91:27-32.
- Coldecott, Morgan DG, Delon-Jones F, Overstreet DH, Janowsky (1991):** Clinical and biochemical aspects of depression disorders, transmitter, receptor theories. *Synapse*; 9:251.
- Conca A, Germann R, Konig P. (2003):** Etomidate vs thiopentone in electroconvulsive therapy: an interdisciplinary challenge for anesthesiology and psychiatry. *Pharmacopsychiatry*; 36 (3): 94-97.
- Coreyll WH, Tsuang MT (1992):** Epidemiology of schizophrenia. *Arch. Gen. psychiatry*; 39:66.
- Cornelius MD, Taylor PM, Geva D, Day NL.(1995) :** Prenatal tobacco and marijuana use among adolescents: effects on offspring gestational age, growth, and morphology .
- Coulter DM, Bate A, Meyboom RH, et al. (2001):** Antipsychotic drugs and heart muscle disorder in international pharmacovigilance: data mining study. *BMJ*; 322: 1207-1209.



- Cryan JF, Lucki I. (2000):** Antidepressant-like behavioral effects mediated by 5-hydroxytryptamine(2C) receptors. *J Pharmacol Exp Ther*; 295: 1120-1126.
- David Haddox J (1996):** Electroconvulsive therapy. Manual of anesthesia and medically compromised patient. Cheng E. Y. Kay(ed.),1st edition; 606-611.
- Davidson C, Lazarus C, Lee TH, Ellinwood EH. (2004):** Ondansetron, given during the acute cocaine withdrawal, attenuates oral cocaine self-administration. *Eur J Pharmacol*; 503:99–102.
- de Mendoza Asensi D, Rodriguez Jornet A, Carvajal Diaz A, (2004):** Acute renal insufficiency associated to cocaine consumption. *Rev Clin Esp*;204:206–211.
- Ding Z, White PF. (2002):** Anesthesia for electroconvulsive therapy. *Anesth Analg*; 94 (5): 1351-1364.
- Dobkin AB (1980):** Potentiation of thiopental anesthesia by derivatives and analogues of phenothiazine. *Anesthesiology*;21:292.
- Domino E, Sullivan TS, Luby ED (1994):** Barbiturate intoxication in a patient treated with a MAOI. *American journal psychiatry*; 118:941.
- Edwards RP, Miller RD, Roizen MF, et al. (1979):** Cardiac responses to imipramine and pancuronium during



References

anesthesia with halothane and enflurane. *Anesthesiology*; 50:421-425.

El-Ganzouri AR, Ivankovich AD, Braverman B, et al. (1985): monamine oxidase inhibitors: should they discontinued preoperatively? *Anesthesia, analgesia*; 64: 592-596.

Elliot DL, Linz DH, Kane JA. (1982): Electroconvulsive therapy: pre-treatment medical evaluation. *Arch Intern Med*; 142 (5): 979-981.

Evans K., McGrath J., and Milns R. (2003): Searching for schizophrenia in ancient Greek and Roman literature: a systematic review. *Acta Psychiatrica Scandanavica*, 107(5), 323–330.

Fiege M, Wappler F, Weisshorn R.,(2003): Induction of malignant hyperthermia in susceptible swine by 3,4- methylenedioxymethamphetamine (“ecstasy”). *Anesthesiology*; 99:1132–1136.

Figiel GS, DeLeo B, Zorumski CF, et al. (1993): Combined use of labetalol and nifedipine in controlling the cardiovascular response from ECT. *J Geriatr Psychiatry Neurol*; 6 (1): 20-24.

Fink M. (1990): How does convulsive therapy work? *Neuropsychopharmacology*; 3 (2): 73-82.

Fredman B, d’Etienne J, Smith I, et al. (1994): Anesthesia for electroconvulsive therapy: effects of propofol



References

and methohexital on seizure activity and recovery. *Anesth Analg*; 79 (1): 75-79.

Fredman B, Husain MM, White PF. (1994): Anaesthesia for electroconvulsive therapy: use of propofol revisited. *Eur J Anaesthesiol*; 11 (5): 423-425.

Fredman B, Smith I, d'Etienne J, et al. (1994): Use of muscle relaxants for electroconvulsive therapy: how much is enough? *Anesth Analg*; 78 (1): 195-196.

Fu W, Stool LA, White PF, et al. (1997): Acute hemodynamic responses to electroconvulsive therapy are not related to the duration of seizure activity. *J Clin Anesth*; 9 (8): 653-657.

Fu W, Stool LA, White PF, et al. (1998): Is oral clonidine effective in modifying the acute hemodynamic response during electroconvulsive therapy? *Anesth Analg*; 86 (5): 1127-1130.

Gaines III GY, Rees DI. (1986): Electroconvulsive therapy and anesthetic considerations. *Anesth Analg*; 65 (12): 1345-1356.

Gold MI (1984): Profound hypotension associated with preoperative use of phenothiazines. *Anesthesiology*; 53:844.

Opiate withdrawal using clonidine. A safe,



- Goldner EM, Hsu L, Waraich P, Somers JM (2002):** Prevalence and incidence studies of schizophrenic disorders: a systematic review of the literature. *Canadian Journal of Psychiatry*, 47(9), 833–843
- Goodnick PJ, Goldstein BJ. (1998):** Selective serotonin reuptake inhibitors in affective disorders – I. Basic pharmacology. *J Psychopharmacol*; 12 (3 Suppl. B): S5-20.
- Greydanus DE, Patel DR.(2003):** Substance abuse in adolescents: a complex conundrum for the clinician. *Pediatr Clin North Am* 2003; 50:1179–1223.
- Harrison PJ, Owen MJ. (2003):** Genes for schizophrenia? Recent findings and their pathophysiological implications. *Lancet*, 361(9355), 417–419.
- Hatch RC (1984):** Ketamine catalepsy and anesthesia in dogs pretreated with antiserotonergic or antidopaminergic neuroleptics or with anticholinergic agents. *Pharmacology research community*; 6:289.
- Hill GE, Wong KC, Hodges MR (1986):** potentiation of succinylcholine neuromuscular blockade by lithium carbonate. *Anesthesiology*; 44:439.



References

- Hill S, Yau K, Whitwam J. (1992):** MAOIs to RIMAs in anaesthesia – a literature review. *Psychopharmacology (Berl)*; 106 Suppl.: S43-45.
- Hsue PY, Salinas CL, Bolger AF, et al.(2002):** Acute aortic dissection related to crack cocaine. *Circulation* 2002; 105:1592–1595.
- Hughes SC, Dailey PA, Landers D.,(1995):** Parturients infected with human immunodeficiency virus and regional anesthesia. Clinical and immunologic response. *Anesthesiology* 1995; 82:32–37.
- Inouye DS, Navin JJ, Hardman JM.(2004):** Fatal postoperative arrhythmia in a man with a remote history of methamphetamine and cocaine use: a case report. *Hawaii Med J*; 63:82–86.
- Ishikawa T, Kawahara S, Saito T, et al. (2001):** Anesthesia for electroconvulsive therapy during pregnancy: a case report [in Japanese]. *Masui*; 50 (9): 991-997.
- Izard P, Leclercq C, Ducasse JL, et al. (1991):** Use of propofol in 1350 anesthetized patients for electroconvulsive therapy. *Ann Fr Anesth Reanim*; 10 (1): 16-20.
- Jones RM, Knight PR. (1981):** Cardiovascular and hormonal responses to electroconvulsive therapy.



References

Modification of an exaggerated response in an hypertensive patient by beta-receptor blockade. *Anaesthesia*; 36 (8): 795-799.

Kapur S, Remington G. (2001): Atypical antipsychotics: new directions and new challenges in the treatment of schizophrenia. *Annu Rev Med*; 52: 503-1.

Kapur S, Seeman P. (2001): Does fast dissociation from the dopamine D(2) receptor explain the action of atypical antipsychotics? A new hypothesis. *Am J Psychiatry*; 158: 360-369.

Kato, T. (2007). "Molecular genetics of bipolar disorder and depression." *Psychiatry Clin Neurosci* 61(1): 3–19

Kaye AD, Gevirtz C, Bosscher HA,.(2003): Ultrarapid opiate detoxification: a review. *Can J Anaesth*; 50:663–671.

Kellner CH, Nixon DW, Bernstein HJ. (1991): ECT-drug interactions: a review. *Psychopharmacol Bull*; 27 (4): 595-609.

Kellner CH, Pritchett JT, Beale MD, et al. (1997): Handbook of ECT. Washington, DC: American Psychiatric Press Inc.

Kelway B, Simpson KH, Smith RJ, et al. (1986): Effects of atropine and glycopyrrolate on cognitive function



References

following anaesthesia and electroconvulsive therapy (ECT). *Int ClinPsychopharmacol*; 1 (4): 296-302.

Klapheke MM. (1993): Combining ECT and anitpsychotic agents: benefits and risks. *Convuls Ther*; 9 (4): 241-255.

Klein PS, Melton DA. (1996): A molecular mechanism for the effect of lithium on development. *Proc Natl Acad Sci U S A*; 93: 8455-8459.

Klein M, Kramer F.(2004): Rave drugs: pharmacological considerations. *Am Assoc Nurse Anesthetists J*; 72:61–67.

Kober A, Schubert B, Bertalanffy P, et al. (2004): Capnography in nontracheally intubated emergency patients as an additional tool in pulse oximetry for prehospital monitoring of respiration. *Anesth Analg*; 98 (1): 206-210.

Kolano JE, Chhibber A, Calalang CC. (1997): Use of esmolol to control bleeding and heart rate during electroconvulsive therapy in a patient with an intracranial aneurysm. *J Clin Anesth*; 9 (6): 493-495.

Krystal AD, Watts BV, Weiner RD, et al. (1998): The use of flumazenil in the anxious and benzodiazepine-dependent ECT patient. *J ECT*; 14 (1): 5-14.



References

- Kuczkowski KM.(2004):** Marijuana in pregnancy. Ann Acad Med Singapore 2004; 33:336–339.
- Kuczkowski KM, Birnbach DJ, van Zundert A.(2000):** Drug abuse in the parturient.Sem Anesthesiol Periop Med Pain .19:216–224.
- Kuczkowski KM.(2003):** Anesthetic implications of drug abuse in pregnancy. J Clin Anesth ; 15:382–394.
- Kuczkowski KM.(2004):** The cocaine abusing parturient: a review of anesthetic considerations. Can J Anaesth 2004; 51:145–154.
- Kumar , Clark (1998):** Psychological medicine, clinical medicine. Parveen Kumar and Michael Clark (ed.);4th edition: 1117-1121.
- Lange RA, Hillis LD.(2001):** Cardiovascular complications of cocaine use. N Engl J Med 2001; 345:351–358.
- Lee D, Kenny M, Harrop-Griffiths W. (1996):** Anaesthesia for electroconvulsive therapy: personnel and facilities. Anaesthesia; 51 (6): 583-584.
- Lieberman JA, Stroup TS, McEvoy JP, Swartz MS, Rosenheck RA, Perkins DO, Keefe RS, Davis SM, Davis CE, Lebowitz BD, Severe J, Hsiao**



- JK, Clinical Antipsychotic Trials of Intervention Effectiveness (CATIE) Investigators. (2005):** Effectiveness of antipsychotic drugs in patients with chronic schizophrenia. The New England Journal of Medicine, 353 (12). 1209-1223
- Lin JH, Lu AY. (1998):** Inhibition and induction of cytochrome P450 and the clinical implications. Clin Pharmacokinet; 35:361-390.
- Lin JH, Lu AY. (2001):** Interindividual variability in inhibition and induction of cytochrome P450 enzymes. Annu Rev Pharmacol Toxicol; 41: 535-567.
- Liu XS, Chen YC, Li ZH.(2004):** Study on the changes of electrocardiogram and ultrastructural in heroin dependence in rats. Fa Yi Xue Za Zhi;20(129–132):135.
- Lloyd G.G. (2002):** Davidson's principles and practice of medicine: Psychiatric disorders; (17): 978-1019.
- Loimer N, Hofmann P, Chaudhry HR. (1992):** Midazolam shortens seizure duration following electroconvulsive therapy. J psychiatr Res; 26 (2): 97-101.
- Lui PW, Ma JY, Chan KK. (1993):** Modification of tonic-clonic convulsions by atracurium in multiple-



References

monitored electroconvulsive therapy. *J Clin Anesth*; 5 (1): 16-21.

Lyon GD, Bruce DL (1989): Anesthesia for electroconvulsive therapy and cardioversion. *General anesthesia*. Gray IC and Nunn JF (eds.). Butterworths: London, Boston, 5th edition; 985-987.

Malsch E, Gratz I, Mani S, et al. (1994): Efficacy of electroconvulsive therapy after propofol and methohexital anesthesia. *ConvulsTher*; 10 (3): 212-219.

Manji HK, Potter WZ, Lenox RH. (1995): Signal transduction pathways: molecular targets for lithium's actions. *Arch Gen Psychiatry*; 52: 531-543.

McCall WV, Shelp FE, Weiner RD, et al. (1991): Effects of labetalol on hemodynamics and seizure duration during ECT. *Convuls Ther*; 7 (1): 5-14.

McCall WV. (1993): Antihypertensive medications and ECT. *Convuls Ther*; 9 (4): 317-325.

Messick, Machenzie, Nugent (1996): Anesthesia at remote location: Electroconvulsive. Miller RD(ed.); 3rd edition: 2081-2085.

Mills PM, Penfold N.(2003) Cannabis abuse and anaesthesia. *Anaesthesia* 2003;58:1125.



- Milner G, Hills N (1996):** Adynamic ileus and nortriptyline. British medical journal; 1:841.
- Mittmann N, Herrmann N, Einarson TR, et al. (1997):** The efficacy, safety and tolerability of antidepressants in late life depression: a meta-analysis. J Affect Disord; 46 (3): 191-217.
- Modica PA, Tempelhoff R, White PF. (1990):** Pro- and anticonvulsant effects of anesthetics (Part II). Anesth Analg; 70 (4): 433-444.
- Mokriski BK, Nagle SE, Papuchis GC, et al. (1992):** Electroconvulsive therapy-induced cardiac arrhythmias during anesthesia with methohexital, thiamylal, or thiopental sodium. J Clin Anesth; 4 (3): 208-212.
- Mongeau R, Blier P, de Montigny C. (1997):** The serotonergic and noradrenergic systems of the hippocampus: their interactions and the effects of antidepressant treatments. Brain Res Brain Res Rev; 23: 145-195.
- Naguib M, Koorn R. (2002):** Interactions between psychotropics, anaesthetics and electroconvulsive therapy: implications for drug choice and patient management. CNS Drugs; 16 (4): 229-247.
- Musty RE, Reggio P, Consroe P.(1995):** A review of



References

Recent advances in cannabinoid research
and the 1994 International Symposium
Cannabis and the Cannabinoids.

NaguibM, Magboul MM, Jaroudi R. (1997): Clinically significant drug interactions with general anaesthetics: incidence, mechanisms and management. *CNS Drugs*; 8 (1): 51-87.

Nair NP, Ahmed SK, Kin NM, et al. (1995): Reversible and selective inhibitors of monoamine oxidase A in the treatment of depressed elderly patients. *Acta Psychiatr Scand Suppl*; 386: 28-35.

Nguyen TT, Chhibber AK, Lustik SJ, et al. (1997): Effect of methohexitone and propofol with or without alfentanil on seizure duration and recovery in electroconvulsive therapy. *Br J Anaesth*; 79 (6): 801-803.

Oye I (1992): Interaction of the chiral forms of ketamine with the opioid. *Pharmacology & Toxicology*; 77: 355-359.

Panesar SK, Bandiera SM, Abbott FS. (1996): Comparative effects of carbamazepine and carbamazepine-10, 11-epoxide on hepatic cytochromes P450 in the rat. *Drug Metab Dispos*; 24: 619-627.



- Pavy TJG, Kliffer AP, Douglas MJ (1995):** Anesthetic management of labor and delivery in a woman taking long term MAOI. Canadian journal anesthesia; 42:618-620.
- Price LH, Heninger GR. (1994):** Lithium in the treatment of mood disorders. N Engl J Med; 331: 591-598.
- Rasmussen KG, Jarvis MR, Zorumski CF. (1996):** Ketamine anesthesia in electroconvulsive therapy. Convuls Ther; 12 (4): 217-223.
- Reddy RV, Moorthy SS, Dierdorf SF, et al. (1993):** Excitatory effects and electroencephalographic correlation of etomidate, thiopental, methohexital, and propofol. Anesth Analg; 77 (5): 1008-1011.
- Riesenman C. (1995):** Antidepressant drug interactions and the cytochrome P450 system: a critical appraisal. Pharmacotherapy Nov-Dec; 15 (6 Pt 2): 84S-99S.
- Ronald A., Mackenzic, Peter A. Southorn, Paul E. Stensrud (2000):** Anesthesia at remote location. Ronald D. Miller (ed.); fifth edition: 2263-2269.
- Sackeim HA, Devanand DP, Prudic J. (1991):** Stimulus intensity, seizure threshold, and seizure duration: impact on the efficacy and safety of electroconvulsive therapy. Psychiatr Clin North Am; 14 (4): 803-843.



- Sackheim HA. (1994):** Central issues regarding the mechanisms of action of electroconvulsive therapy: directions for future research. *Psychopharmacol Bull*; 30 (3): 281-308.
- Saffer S, Berk M. (1998):** Anesthetic induction for ECT with etomidate is associated with longer seizure duration than thiopentone. *J ECT*; 14 (2): 89-93.
- Saito S, Kadoi Y, Iriuchijima N, et al. (2000):** Reduction of cerebral hyperemia with anti-hypertensive medication after electroconvulsive therapy. *Can J Anaesth*; 47 (8): 767-774.
- Saito S, Kadoi Y, Naro T, et al. (2000):** The comparative effects of propofol versus thiopental on middle cerebral artery blood flow velocity during electroconvulsive therapy. *Anesth Analg*; 91 (6): 1531-1536.
- Saura J, Richards JG, Mahy N. (1994):** Age-related changes on MAO in Bl/C57 mouse tissues: a quantitative radioautographic study. *J Neural Transm Suppl*; 41: 89-94.
- Schenkel, L.S., Spaulding, W.D., Dilillo, D., Silverstein, S.M. (2005):** Histories of childhood maltreatment in schizophrenia: Relationships with premorbid functioning, symptomatology, and cognitive deficits. *Schizophrenia Research*



References

- Schindler CW.(1996):** Cocaine and cardiovascular toxicity. *Addict Biol* 1996; 1:31–47.
- Selvin BL (1995):** Eelectroconvulsive Therapy. *Anesthesiology*;67:267.
- Shih JC. (1991):** Molecular basis of human MAO A and B. *Neuropsychopharmacology*; 4: 1-7.
- SpacekA, Neiger FX, SpissCK, et al. (1997):** Atracurium-induced neuromuscular block is not affected by chronic anticonvulsant therapy with carbamazepine. *Acta Anaesthesiol Scand*; 41: 1308-1311.
- Spencer KM, Nestor PG, Perlmutter R, Niznikiewicz MA, Klump MC, Frumin M, Shenton ME, McCarley (2004):** Neural synchrony indexes disordered perception and cognition in schizophrenia. *Proceedings of the National Academy of Sciences*, 101, 17288-17293.
- Stewart JT. (2000):** Lithium and maintenance ECT. *J ECT*; 16: 300-301.
- Stoelting R. K. (1999):** Pharmacology and physiology in anesthetic practice;3rd edition :357-376.
- Stoelting R.K., Dierdorf S.F. (2002):** Anesthesia and co-existing disease;4th edition:629-639.



- Stoudemire A, Knos G, Gladson M, et al. (1990):** Labetalol in the control of cardiovascular responses to electroconvulsive therapy in high-risk depressed medical patients. *J Clin Psychiatry*; 51 (12): 508-512.
- Swartz CM. (1993):** Anesthesia for ECT. *Convuls Ther*; 9 (4): 301-316.
- Symons IE.(2002)** Cannabis smoking and anaesthesia. *Anaesthesia* 2002;57:1142–1143.
- Tanaka N, Saito Y, Hikawa Y, et al. (1997):** Effects of thiopental and sevoflurane on hemodynamics during anesthetic management of electroconvulsive therapy [in Japanese]. *Masui*; 46 (12): 1575-1579.
- Tang WK, Ungvari GS. (2003):** Efficacy of electroconvulsive therapy in treatment-resistant schizophrenia: a prospective open trial. *Prog Neuropsychopharmacol Biol Psychiatry*; 27 (3): 373-379.
- Taylor D. (2003):** Ziprasidone in the management of schizophrenia: the QT interval issue in context. *CNS Drugs*; 17 (6): 423-430.
- Terra Filho M, Yen CC, Santos Ude P. (2004):** Pulmonary alterations in cocaine users. *Sao Paulo Med*; 122:26–31.



- Tharyan P, Adams CE. (2002):** Electroconvulsive therapy for schizophrenia. *Cochrane Database Syst Rev*; (2): CD000076.
- Vagts DA, Boklage C, Galli C.(2004):** Intraoperative ventricular fibrillation in a patient with chronic cocaine abuse – a case report [in German].*Anaesthesiol Reanim*; 29:19–24.
- Van Den Berg AA, Honjol NM. (1998):** Electroconvulsive therapy and intraocular pressure. *Middle East J Anesthesiol*; 14 (4): 249-258.
- van Os J, Burns T, Cavallaro R, Leucht S, Peuskens J, Helldin L, Bernardo M, Arango C, Fleischhacker W, Lachaux B, Kane JM. (2006):** Standardized remission criteria in schizophrenia. *Acta Psychiatrica Scandinavica*, 113(2), 91-95.
- Van Praag, HM (2005):** *World J Biol Psychiatry* 6 Suppl: 5-22
- Vizi E, (1982):** The effect of lithium on acetylcholine release and synthesis. *Neuropharmacology*; 11:521.
- Wajima Z, Yoshikawa T, Ogura A, et al. (2002):** Intravenous verapamil blunts hyperdynamic responses during electroconvulsive therapy



without altering seizure activity. *Anesth Analg*; 95(2): 400-402.

Warltier DC, Mitra S, Sinatra RS.(2004): Perioperative management of acute pain in the opioid-dependent patient. *Anesthesiology*; 101:212–227.

Webb MC, Coffey CE, Saunders WR, et al. (1990): Cardiovascular response to unilateral electroconvulsive therapy. *Biol Psychiatry*; 28 (9): 758-766.

Weiner RD. (2001): American Psychiatric Association committee on electroconvulsive therapy. The practice of electroconvulsive therapy: recommendations for treatment, training, and privileging. 2nd ed. Washington, DC: American Psychiatric Press.

Weinger MB, Partridge BL, Hauger R,et al. (1991): Prevention of the cardiovascular and neuroendocrine response to electroconvulsive therapy I. hemodynamics. *Anesth. Analg*; 73 (5): 556-562

Wells DG, Bjorksten AR (1994): Monoamine oxidase inhibitors revisited. *Canadian journal anesthesia*; 36:64-74.



- Wells DG, Davies GG, Rosewarne F. (1989):** Attenuation of electroconvulsive therapy induced hypertension with sublingual nifedipine. *Anaesth Intensive Care*; 17 (1): 31-33.
- Wells DG, Davies GG. (1987):** Hemodynamic changes associated with electroconvulsive therapy. *Anesth Analg*; 66 (11): 1193-1195.
- Whittington RA, Virag L, Vulliemoz (2002):** Dexmedetomidine increases the cocaine seizure threshold in rats. *Anesthesiology*; 97:693–700.
- Woo D, Haverbusch M, Sekar P, et al. (2004):** Effect of untreated hypertension on hemorrhagic stroke. *Stroke*; 35 (7): 1703-1708.
- Wulfson HD, Askanazi J, Finck AD. (1984):** Propranolol prior to ECT associated with asystole. *Anesthesiology*; 60 (3): 255-256.
- WWW.mayoclinic.com/health/bipolar-disorder/DS00356/DSECTION=2.**
- WWW.mayoclinic.com/health/bipolar-disorder/DS00356/DSECTION=7**
- Xia Z, Ying G, Hansson AL, et al. (2000):** Antidepressant-induced lipidosis with special reference to tricyclic compounds. *Prog Neurobiol* 2000; 60: 501-512.



References

- Zaidi NA, Khan FA. (2000):** Comparison of thiopentone sodium and propofol for electro convulsive therapy (ECT). J Pak Med Assoc; 50 (2): 60-63.
- Zielinski RJ, Roose SP, Devanand DP, et al. (1993):** Cardiovascular complications of ECT in depressed patients with cardiac disease. Am J Psychiatry; 150: 904-909.
- Zimmer R, Gieschke R, Fischbach R, et al. (1990):** Interaction studies with moclobemide. Acta Psychiatr Scand Suppl; 360: 84-86.