

References

- Absalom AR, Lal A 2009.** Accuracy of the ‘Paedfusor’ in children undergoing cardiac surgery or catheterization. *Br J Anaesth*; 91: 468–75.
- A Rodgers, N Walker and S Schug et al.2000.** Reduction of postoperative mortality and morbidity with epidural or spinal anaesthesia: results from an overview of randomised trials, *BMJ* 321, pp. 1493–1504.
- A Taguchi, N Sharma and RM Saleem et al. 2001.** Selective postoperative inhibition of gastrointestinal opioid receptors, *N Engl J Med* 345, pp. 935–940.
- Ackerman, M.S 2002.** Outpatient Anaesthesia. In: *Clinical Anaesthesiology*, Morgan, G.E.jr.; Mikhail, M.S.; Murray, M.J. and Larson, C.P.jr.(Eds.), Mc Graw-Hill professional, 3rd edition; (2);46: 882-888.
- Adams JP, Murphy PG. 2000.** Obesity in anaesthesia and intensive care. *Br J Anaesth*; 85: 91–108.
- J Andre’s, E Monzo’ 2000.** Regional Techniques for Day Surgery: Intraarticular Anesthesia and Analgesia. *Techniques in Regional Anesthesia and Pain Management* ;4 (1): 54-61.
- Aldrete JA 1995.** The post-anesthesia recovery score revisited (Letter). *J Clin Anesth*; 7: 89–91.
- Allen GS, Cox CS Jr, White N, Khalil S, Rabb M, Lally KP. 1998.** Postoperative respiratory complications in ex-premature infants after inguinal herniorrhaphy. *J Pediatr Surg*; 33: 1095–8.

Alijani A, Hanna GB, Cuschieri A 2004. Abdominal wall lift versus positive- pressure capnoperitoneum for laparoscopic cholecystectomy: randomized controlled trial. *Ann Surg*;239(3):388 - 94.

American Society of Anesthesiologists Task Force on Postanesthetic Care. Practice guidelines for postanesthetic care: a report by the American Society of Anesthesiologists Task Force on Postanesthetic Care. *Anesthesiology* 2002; 96: 742–52.

Andersen FH, Nielsen K, Kehlet H 2005. Combined ilioinguinal blockade and local infiltration anaesthesia for groin hernia repair--a double-blind randomized study *Anesthesia; Br J Anaesth*; 94: 520–3.

Apfel CC, Kranke P, Eberhart LH, Roos A, Roewer N 2002. Comparison of predictive models for postoperative nausea and vomiting. *Br J Anaesth*; 88: 234–40.

Apfel CC, Laara E, Koivuranta M, Greim CA, Roewer N 1999. A simplified risk score for predicting postoperative nausea and vomiting: conclusions from cross-validations between two centers. *Anesthesiology*; 91: 693–700.

Apfel CC, Korttila K, Abdalla M, Kerger H, Turan A, Vedder I, et al.2004. A factorial trial of six interventions for the prevention of postoperative nausea and vomiting. *N Engl J Med*; 350(24):2441–51.

Apfelbaum JL, Lichtor JL, Lane BS, Coalson DW, Korttila KT 1996. Awakening, clinical recovery, and psychomotor effects after desflurane and propofol anesthesia. *Anesth Analg*; 83: 721–5.

Apfelbaum JL, Walawander CA, Grasela TH, et al 2002. Eliminating intensive postoperative care in same-day surgery patients using short acting anesthetics. *Anesthesiology*; 97: 66–74.

Arain SR, Ebert TJ 2002. The efficacy, side effects, and recovery characteristics of dexmedetomidine versus propofol when used for intraoperative sedation. *Anesth Analg*; 95: 461–6.

Atkins M, White J, Ahmed K.2002. Day surgery and body mass index: results of a national survey. *Anaesthesia* 2002; 57: 180–2.

AW Majeed, S Brown, N Williams, DR Hannay and AG Johnson, 1995. Variations in medical attitudes to postoperative recovery period, *BMJ* 311, p. 296.

Awad IT, Moore M, Rushe C, Elburki A, O'Brien K, Warde D 2004. Unplanned hospital admission in children undergoing day-case surgery. *Eur J Anaesthesiol*; 21: 379–83.

AV, Vaghadi H, Collins L, Mitchell GW.2001 Small-dose selective spinal anaesthesia for short-duration outpatient gynaecological laparoscopy: recovery characteristics compared with propofol anaesthesia. *Br J Anaesth*; 86(4):570 - 2.

Axelsson K, Mollefors K, Olsson JO, Lingardh G, Widman B 1985. Bladder function in spinal anaesthesia. *Acta Anaesthesiol Scand*; 29: 315–21.

Azurin DJ, Cwik JC, Schuricht AL 2001. The efficacy of epidural anesthesia for endoscopic preperitoneal herniorrhaphy: a prospective study. *J Laparoendosc Surg*;6(6):374 - 81.

Bailey CJ, Turner RC. Metformin. N Engl J Med 1996; 334: 574–9.

Bain J, Kelly H, Snadden D, Staines H. 1999. Day surgery in Scotland: patient satisfaction and outcomes. *Qual Health Care* 1999;8:86-91.

Barker JP, Robinson PN, Vafidis GC, Burrin JM, Sapsed-Byrne S, Hall GM.1995 Metabolic control of non-insulin-dependent diabetic patients undergoing cataract surgery: comparison of local and general anaesthesia. Br J Anaesth 1995; 74: 500–5

Barker JP, Robinson PN, Vafidis GC, Burrin JM, Sapsed-Byrne S, Hall GM. 1995 Metabolic control of non-insulin-dependent diabetic patients undergoing cataract surgery: comparison of local and general anaesthesia. Br J Anaesth; 74: 500–5.

Ben-David B, Levin H, Solomon E, Admoni H, Vaida S 1996. Spinal bupivacaine in ambulatory surgery: the effect of saline dilution. Anesth Analg; 83: 716–20.

Bonnet F, Marret E.2005. Influence of anaesthetic and analgesic techniques on outcome after surgery. Br J Anaesth; 95(1):52–8.

Bouchut JC, Dubois R, Foussat C, et al.2001. Evaluation of caudal anaesthesia performed in conscious ex-premature infants for inguinal herniotomies. Paediatr Anaesth; 11: 55–8.

British Association of Day Surgery 2004. Spinal anaesthesia apractical guide, London; Available from /www.badsco.ukS.

Brodsky JB, Lemmens HJ, Brock-Utne JG, Vierra M, Saidman LJ. 2002.Morbid obesity and tracheal intubation. Anesth Analg; 94: 732–6.

Brown AR,Weiss R, Greenberg C, et al. 1993. Interscalene block for shoulder arthroscopy: Comparison with general anesthesia. Arthroscopy; 9: 295–300.

Bryson GL, Chung F, Cox RG, et al.2004. Canadian Ambulatory Anesthesia Research Education Group. Patient selection in ambulatory

anesthesia - an evidence-based review: part II. *Can J Anesth* 2004; 51: 782–94.

Buckenmaier 3rd. CC, Nielsen KC, Pietrobon R Klein SM, Martin AH, Greengrass RA, et al 2002. Small-dose intrathecal lidocaine versus ropivacaine for anorectal surgery in an ambulatory setting. *Anesth Analg*; 95(5): 1253–7.

Buell JF, Thomas MJ 2009. An initial experience and evolution of laparoscopic hepatic resectional surgery. *Surgery* ;136(4):812 - 16
Cahill, C.J 1999. Basket cases and trolleys. *Day surgery Proposals for the Millennium. J One-day Surgery*; 9: 11-12.

Capogna G, Celleno D, Fusco P, Lyons G, Columb M 1999. Relative potencies of bupivacaine and ropivacaine for analgesia in analgesia in labour *Br J Anaesth*:82(3):371-3.

Cappelleri G, Aldegheri G, Danelli G, et al 2005. Spinal anesthesia with hyperbaric levobupivacaine and ropivacaine for outpatient knee arthroscopy: a prospective, randomized, double-blind study. *Anesth Analg*; 101: 77–82.

Carr DB, Goudas LC. Acute pain. Lancet 1999.353(9169): 2051–8.

Carlisle JB, Stevenson CA.2006. Drugs for preventing postoperative nausea and vomiting. *Cochrane Database Syst Rev*3: CD00125.

Cartwright DP, Kvalsvik O, Cassuto J, Jansen JP, Wall C, Remy B, et al 1997. A randomized, blind comparison of remifentanyl and alfentanil during anesthesia for outpatient surgery. *Anesth Analg*;85(5):1014 9.

Cheng CR, Sessler DI, Apfel CC 2005. Does neostigmine administration produce a clinically important increase in postoperative nausea and vomiting? *Anesth Analg*;101(5): 1349–55.

Chew LD, Bradley KA, Flum DR, Cornia PB, Koepsell TD 2004. The impact of low health literacy on surgical practice. *Am J Surg*; 188: 250–3.

Christiansen CL, Schurizek BA, Malling B, Knudsen L, Alberti KG, Hermansen K.1988. Insulin treatment of the insulin-dependent diabetic patient undergoing minor surgery. Continuous intravenous infusion compared with subcutaneous administration. *Anaesthesia* 43: 533–7.

Chung F, Chan VW, Ong D 1995. A post-anesthetic discharge scoring system for home readiness after ambulatory surgery. *J Clin Anesth*; 7: 500–6.

Chung F, Imasogie N, Ho J, Ning X, Prabhu A, Curti B 2005. Frequency and implications of ambulatory surgery without a patient escort. *Can J Anesth*; 52: 1022–6.

Chung F, Kayumov L, Sinclair DR, Edward R, Moller HJ, Shapiro CM 2005 . What Is the Driving Performance of Ambulatory Surgical Patients after General Anesthesia? *Anesthesiology*; 103: 951–6.

Chung F, Lane R, Spraggs C, et al 1999. Ondansetron is more effective than metoclopramide for the treatment of opioid-induced emesis in post-surgical adult patients. Ondansetron OIE Post-Surgical Study Group. *Eur J Anaesthesiol*; 16: 669–77.

Chung F, Mezei G 1999 . Factors contributing to a prolonged stay after ambulatory surgery. *Anesth Analg*; 89: 1352–9.

Chung F, Ritchie E, Su J 1997. Postoperative pain in ambulatory surgery. *Anesth Analg*; 85: 808–16.

Chung F.1995. Recovery pattern and home-readiness after ambulatory surgery. *Anesth Analg* 1995; 80: 896–902.

Chung, F, Mezei, R. and Tong, D.1999. Adverse events in ambulatory surgery. A comparison between elderly and younger patients. *Can J Anaesth*, 46: 309-312.

Chung, F, Mezei, G. and Tong, D.1999. Pre-existing medical conditions as predictors of adverse events in day-case surgery. *Br J Anaesth*, 83: 262-70.

Claxton AR, McGuire G, Chung F, Cruise C 1997. Evaluation of morphine versus fentanyl for postoperative analgesia after ambulatory surgical procedures. *Anesth Analg*; 84: 509–14.

Claxton AR, McGuire G, Chung F, Cruise C.1997. Evaluation of morphine versus fentanyl for postoperative analgesia after ambulatory surgical procedures. *Anesth Analg*,84(3): 509–14.

Clergue F, Seebacher J, Lefebvre G, Viars P 1997. Ventilatory effects of laparoscopy under epidural anesthesia. *Anesth Analg*;70(4):357- 61.

Cohen MS, Finkelstein SE, Brunt LM, et al 2005. Outpatient minimally invasive parathyroidectomy using local/regional anesthesia: a safe and effective operative approach for selected patients. *Surgery*; 138: 681–7.

Cohen MM, Cameron CB.1991. Should you cancel the operation when a child has an upper respiratory tract infection? *Anesth Analg* 1991; 72: 282–8.

Coley KC, Williams BA, DaPos SV, Chen C, Smith RB 2002. Retrospective evaluation of unanticipated admissions and readmissions after same day surgery and associated costs. *J Clin Anesth*; 14: 349–53.

Collins LM, Vaghadia H 2001. Regional anesthesia for laparoscopy. *Anesthesiol Clin North America*; 19(1):43-55.

Coloma M, Chiu JW, White PF, Armbruster SC 2001. The use of esmolol as an alternative to remifentanyl during desflurane anesthesia for fast-track outpatient gynecologic laparoscopic surgery. *Anesth Analg*;92(2):352–7.

Conlay, L.A 2005. Ambulatory Anaesthesia then and now. In: *Ambulatory Anesthesia and Perioperative Analgesia*, Steele, S.M.; Nielsen, K.C and Klein, S.M. (Eds.), Mc Graw-Hill professional. Ch.1, Pp.3-9.

Cox RG, Goresky GV.1990. Life-threatening apnea following spinal anesthesia in former premature infants. *Anesthesiology* 1990; 73: 345–7.

Craig J, Cooper GM, Sear JW 1982. Recovery from day case anaesthesia. Comparison between methohexitone, althesin and etomidate. *Br J Anaesth*; 54: 447–51.

D'Alessio JG, Rosenblum M, Shea KP et al.1995. A retrospective comparison of interscalene block and general anesthesia for ambulatory surgery shoulder arthroscopy. *Regional Anesthesia*; 20: 62–68.

Dahl V, Gierløff C, Omland E et al 1997. Spinal, epidural or propofol anaesthesia for outpatient knee arthroscopy? *Acta Anaesthesiologica Scandinavica*; 41: 1341–1345.

Davies KE, Houghton K, Montgomery JE. 2001. Obesity and day-case surgery. *Anaesthesia* 2001; 56: 1112–5.

De Kock M, Gautier P, Fanard L, Hody JL, Lavand'homme P 2001. Intrathecal ropivacaine and clonidine for ambulatory knee arthroscopy: a dose-response study. *Anesthesiology*; 94(4):574–8.

De Lathouwer C, Poullier JP 2000. How much ambulatory surgery in the World in 1996-1997 and trends? *Ambul Surg*;8:191-210.

Department of Health 2000. The NHS plan. A plan for investment A plan for reform.

Descottes B, Glineur D, Lachachi F, et al.2003. Laparoscopic liver resection of benign liver tumors. Surg Endosc 2003;17(1):23 - 30.

DeSoto H, Patel RI, Soliman IE, Hannallah RS. 1988. Changes in oxygen saturation following general anesthesia in children with upper respiratory infection signs and symptoms undergoing otolaryngological procedures. Anesthesiology, 68: 276–9.

Dexter F, Macario A, Manberg PJ, Lubarsky DA 1999. Computer simulation to determine how rapid anesthetic recovery protocols to decrease the time for emergence or increase the phase I postanesthesia care unit bypass rate affect staffing of an ambulatory surgery center. Anesth Analg; 88: 1053–63.

Dexter F, Tinker J 1995. Analysis of strategies to decrease postanesthesia care unit costs. Anesthesiology; 82: 94–101.

Diemunsch PA, Torp KD, Van Dorsselaer T, et al 2000. Nitrous oxide fraction in the carbon dioxide pneumoperitoneum during laparoscopy under general inhaled anesthesia in pigs. Anesth Analg; 90(4): 951- 3.

DI Sessler,1997. Mild perioperative hypothermia, N Engl J Med 336 (1997), pp. 1730–1737.

Ding Y, Fredrich, White PF 2000. Use of mivacurium during laparoscopic surgery: effect of reversal drugs on postoperative recovery. Anesth Analg;78(3):445 - 50.

DN Lobo, KA Bostock and KR Neal et al.2002. Effect of salt and water balance on recovery of gastrointestinal function after elective colonic resection: a randomised controlled trial, Lancet 359, pp. 1812–1818.

Dobrydnjov I, Axelsson K, Thorn SE, Matthiesen P, Klockhoff H, Holmstrom B, et al. 2003. Clonidine combined with small-

dosebupivacaine during spinal anesthesia for inguinal herniorrhaphy: a randomized double-blinded study. *Anesth Analg*;96(5):1496–503.

Dornhoffer J, Manning L 2000. Unplanned admissions following outpatient otologic surgery: the University of Arkansas experience. *Ear Nose Throat J*; 79: 710, 713–7.

Duarte JB, Kux P, Costa JR 2010. Fast track endoscopic thoracic sympathectomy. *Clin Auton Res*;13(Suppl 1): I66 - 72.

Duncan PG, Cohen MM, Tweed WA, et al.1992. The Canadian four-centre study of anaesthetic outcomes: III. Are anaesthetic complications predictable in day surgical practice? *Can J Anaesth* 1992; 39(5 Pt 1): 440–8.

Duncan PG, Shandro J, Bachand R, Ainsworth L 2001. Pilot study of recovery room bypass (“fast-track protocol”) in a community hospital. *Can J Anesth*; 48: 630–6.

DW Wilmore, 2002. From Cuthbertson to fast-track surgery:70 years of progress in reducing stress in surgical patients , *Ann Surg* 236, pp. 643–648.

EJ Norris, C Beattie and BA Perler et al.2001. Double-masked randomised trial comparing alternate combinations of intraoperative anesthesia and postoperative analgesia in abdominal aortic surgery, *Anesthesiology* 95, pp. 1054–1067.

Enneking FK, Chan V, Greger J, Hadzic A, Lang SA, Horlocker TT 2005. Lower-extremity peripheral nerve blockade: essentials of our current understanding. *Reg Anesth Pain Med*; 30: 4–35.

Eriksson H, Korttila K 2000. Recovery profile after desflurane with or without ondansetron compared with propofol in patients undergoing outpatient gynecological laparoscopy. *Anesth Analg*;82(3): 533-8.

Fisher DM. 1995. When is the ex-premature infant no longer at risk for apnea?(Editorial). *Anesthesiology*, 82: 807–8.

Fisher QA, McComiskey CM, Hill JL, et al. 2001. Postoperative voiding interval and duration of analgesia following peripheral or caudal nerve blocks in children. *Anesth Analg*; 72: 173–7.

Fleisher LA, Pasternak LR, Herbert R, Anderson GF 2004. Inpatient hospital admission and death after outpatient surgery in elderly patients: importance of patient and system characteristics and location of care. *Arch Surg*; 139: 67–72.

Fleisher LA, Pasternak R, Lyles A. 2002. A novel index of elevated risk for hospital admission or death immediately following outpatient surgery. *Anesthesiology*, 96: A38 (abstract).

Fortier J, Chung F and Su J 1998. Unanticipated admission after ambulatory surgery-a prospective study. *Can J Anaesth*; 45: 612-619.

Friedman Z, Wong DT, Chung F.2003. What are the ambulatory surgical patient selection criteria in Canada? *Can J Anesth*, 50(Suppl): A16 (abstract).

Gan T, Sloan F, Dear Gde L, El-Moalem HE, Lubarsky DA 2001. How much are patients willing to pay to avoid postoperative nausea and vomiting? *Anesth Analg*; 92: 393–400.

Gan TJ.2002. Postoperative nausea and vomiting: can it be eliminated? *JAMA* 2002; 287: 1233–6.

Gan TJ, Meyer T, Apfel CC, Chung F, Davis PJ, Eubanks S et al. 20003. Consensus guidelines for managing postoperative, nausea and vomiting. *Anesth Analg*, 97(1):62–71.

Ganesan S, Prior AJ, Rubin JS 2000. Unexpected overnight admissions following day-case surgery: an analysis of a dedicated ENT day care unit. *Ann R Coll Surg Engl*; 82: 327–30.

Ghatge S, Lee J, Smith I 2003. Sevoflurane: an ideal agent for adult day-case anesthesia? *Acta Anaesthesiol Scand*; 47(8):917–31.

GJ Rubin and M Hotopf, 2002. Systematic review and meta-analysis of interventions for postoperative fatigue, *Br J Surg* 89 (2002), pp. 971–984

GM Hall and P Salmon, 2002. Physiological and psychological influences on postoperative fatigue, *Anesth Analg* 95, pp. 1446–1450

Goel S, Bhardwaj N, Grover VK 2003. Intrathecal fentanyl added to intrathecal bupivacaine for day case surgery: a randomized study. *Eur J Anaesthesiol*;20(4): 294–7.

Gold BS, Kitz DS, Lecky JH, Neuhaus JM.1989. Unanticipated admission to the hospital following ambulatory surgery. *JAMA* 1989; 262: 3008–142 Fisher DM. When is the ex-premature infant no longer at risk for apnea?(Editorial). *Anesthesiology* 1995; 82: 807–8.

Golembiewski J, Chernin E, Chopra T.2005. Prevention and treatment of postoperative nausea and vomiting. *Am J Health Syst Pharm*; 62: 1247–60.

Gonzalez R, Smith CD, McClusky III DA, et al 2004. Laparoscopic approach reduces likelihood of perioperative complications in patients undergoing adrenalectomy. *Am Surg*;70(8):668 - 74.

Greenberg CP 1995. Practical, cost-effective regional anesthesia for ambulatory surgery. *J Clin Anaesth*; 7:614-21.

Grewal H, Sweat J, Vazquez WD 2004. Laparoscopic appendectomy in children can be done as a fast-track or same-day surgery.*JSLS*;8(2):1514.

Guignard B, Bossard AE, Coste C, Sessler DI, Lebrault C, Alfonsi P, et al.2000. Acute opioid tolerance: intraoperative remifentanyl increases

postoperative pain and morphine requirement. *Anesthesiology*;93(2):409–17.

Gunter JB, Watcha MF, Forestner JE, et al. 1991. Caudal epidural anesthesia in conscious premature and high-risk infants. *J Pediatr Surg*, 26: 9–14.

Gupta A, Bodin L, Holmström B et al. 2001. A systematic review of the peripheral analgesic effects of intraarticular morphine. *Anesthesia and Analgesia*; 93: 761–770.

Gupta A, Axelsson K, Thorn SE, Matthiessen P, Larsson LG, Holmstrom B, et al 2003. Low-dose bupivacaine plus fentanyl for spinal anesthesia during ambulatory inguinal herniorrhaphy: a comparison between 6 and 7.5 mg of bupivacaine. *Acta Anaesthesiol Scand*;47(1):13–9.

Gupta A, Stierer T, Zuckerman R, Sakima N, Parker SD, Fleisher LA 2004 . Comparison of recovery profile after ambulatory anesthesia with propofol, isoflurane, sevoflurane and desflurane: a systematic review. *Anesth Analg*;98(3): 632–41.

Gutt CN, Oniu T, Mehrabi A, et al 2004. Circulatory and respiratory complications of carbon dioxide insufflation. *Dig Surg*;21(2): 95- 105.

H Kehlet and DW Wilmore, 2002. Multimodal strategies to improve surgical outcome, *Am J Surg* 183, pp. 630–644

H Kehlet and K Holte, 2001. Effect of postoperative analgesia on surgical outcome, *Br J Anaesth* 87 (2001), pp. 62–72.

H Kehlet, 1998. Modification of responses to surgery by neural blockade: clinical implications. In: MJ Cousins and PO Bridenbaugh, Editors, *Neural blockade in clinical anesthesia and management of pain*, Lippincott, Philadelphia , pp. 129–175.

Hadzic A, Arliss J, Kerimoglu B, et al 2004. A comparison of infraclavicular nerve block versus general anesthesia for hand and wrist day-case surgeries. *Anesthesiology*; 101: 127–32.

Hadzic A, Karaca PE, Hobeika P, et al 2005. Peripheral nerve blocks result in superior recovery profile compared with general anesthesia in outpatient knee arthroscopy. *Anesth Analg*; 100: 976–81.

Hadzic A, Williams BA, Karaca PE, et al.2005. For outpatient rotator cuff surgery, nerve block anesthesia provides superior same-day recovery over general anesthesia. *Anesthesiology* 2005; 102: 1001–7.

Haldane G, Stott S, McMenemin I 2000. Pouch of Douglas block for laparoscopic sterilisation. *Anaesthesia*; 53(6):605- 617.

Hamad MA, El-Khattary OA 2003. Laparoscopic cholecystectomy under spinal anesthesia with nitrous oxide pneumoperitoneum: a feasibility study. *Surg Endosc*; 17(9):1426- 8.

Hannington-Kiff JG 1970. Measurement of recovery from outpatient general anaesthesia with a simple ocular test. *Br Med J*; 3: 132–5.

Hasaniya NW, Zayed FF, Faiz H, Severino R 2001. Preinsertion local anesthesia at the trocar site improves perioperative pain anddecreases costs of laparoscopic cholecystectomy. *Surg Endosc*; 15(9):962 - 4.

Haydon GH, Dillon J, Thomas H, Hayes PC 2001. Hypoxemia during diagnostic laparoscopy: a prospective study. *Gastrointest Endosc*;44(2):144 -8.

Hedayati B, Fear S.1999. Hospital admission after day-case gynaecological laparoscopy. *Br J Anaesth*, 83(5):776–9.

Hemal AK, Singh I 2004. Hand assisted laparoscopic radical cystectomy for cancer bladder. *Int Urol Nephrol*;36(2):191 - 5.

Henderson RC, Campion ER, DeMasi RA et al 1990. Postarthroscopy analgesia with bupivacaine. A prospective, randomized, blinded evaluation. *The American Journal of Sports Medicine*; 18: 614–617.

Henderson-Smart DJ, Steer P.2002. Prophylactic caffeine to prevent postoperative apnea following general anesthesia in preterm infants. *Cochrane Database Syst Rev*, 4: CD000048.

Henzi I, Walder B, Tramer MR.1999. Metoclopramide in the prevention of postoperative nausea and vomiting: a quantitative systematic review of randomized, placebo-controlled studies. *Br J Anaesth*, 83(5):761–71.

Higgins PP, Chung F, Mezei G 2002. Postoperative sore throat after ambulatory surgery. *Br J Anaesth*; 88: 582–4.

Hirschberg T, Olthoff D, Borner P 2002. Comparative studies of total extraperitoneal hernioplasty in combined spinal epidural anesthesia versus balanced general anesthesia. *Anaesthesiol Reanim*; 27(6):144- 51.

Hopkins PM 2005. Nitrous oxide: a unique drug of continuing importance for anaesthesia. *Best Pract Res Clin Anaesthesiol*;19(3):381–9.

Hovorka J, Korttila K, Nelskyla K, et al 1997. Reversal of neuromuscular blockade with neostigmine has no effect on the incidence or severity of postoperative nausea and vomiting. *Anesth Analg*;85(6):1359-61.

Hsieh CH 2003. Laparoscopic cholecystectomy for patients with chronic obstructive pulmonary disease. *J Laparoendosc Adv Surg Tech A*;13(1):5-9.

Ian Jackson 2007. Day surgery overview: Where are we now, how did we get here and where are we going? *Current Anaesthesia & Critical Care*; 18: 176–80.

Imarengiaye CO, Song D, Prabhu AJ, Chung F 2003. Spinal anesthesia: functional balance is impaired after clinical recovery. *Anesthesiology*;98(2):511–5.

Imasogie N, 2001. Effect of return hospital visits on economics of ambulatory surgery. *Curr Opin Anaesthesiol*; 14: 573–8.

Jain NB, Pietrobon R, Guller U, Ahluwalia AS, Higgins LD 2005. Influence of provider volume on length of stay, operating room time, and discharge status for rotator cuff repair. *J Shoulder Elbow Surg*; 14: 407–13.

Jackson I, Gandhimani P.2006. Adequacy of postoperative pain relief after discharge. In: *Raising the standard: a compendium of audit recipes for continuous quality improvement in anaesthesia. Section 5, day surgery services*, 2nd ed. London: The Royal College of Anaesthetists, p. 108–9.

Jarrett PE, Staniszewski A 2006. The development of ambulatory surgery and future challenges. In: Lemos P, Jarrett P, Philip B, editors. *Day surgery: development and practice*. Porto, Portugal: Clássica Artes Gráficas; 21–34.

Javerliat I, Coggia M, Bourriez A, Goeau- Brissonniere OA 2009. Total laparoscopic aortomesenteric bypass. *Vascular*;12(2):126– 9.

Jenkins K, Grady D, Wong J, Correa R, Armanious S, Chung F.2001. Post-operative recovery: day surgery patients’ preferences. *Br J Anaesth* 2001; 86: 272–4.

Jenkins K, Grady D, Wong J, Correa Armanio, Chung F 2001 . Postoperative recovery: day surgery patients preferences. *Br J Anaesth*; 86:272-4.

Jense HG, Dubin SA, Silverstein PI, O’Leary-Escolas U.1991. Effect of obesity on safe duration of apnea in anesthetized humans. *Anesth Analg* 1991; 72: 89–93.

JH.1999. Comparison of vital capacity induction with sevoflurane to intravenous induction with propofol for adult ambulatory anesthesia. *Anesth Analg*, 89(3):623–7.

Jin FL, Norris A, Chung F, Ganeshram T.1998. Should adult patients drink fluids before discharge from ambulatory surgery? *Anesth Analg*, 87: 306–11.

Jones P, Yate P.2003. Contraindications to use of metformin. Blanket banning of metformin two days before surgery may not be a good idea (Letter). *BMJ* 2003; 326: 762.

Joshi GP 2005. The Society for Ambulatory Anesthesia: 19th Annual Meeting Report. *Anesth Analg*; 100: 982–6.

JR Rigg, K Jamrozik and PS Myles et al.2002. Epidural anaesthesia and analgesia and outcome of major surgery: a randomised trial, *Lancet*, 359:1276–82.

Junger A, Klasen J, Benson M, et al. 2001. Factors determining length of stay of surgical day-case patients. *Eur J Anaesthesiol*; 18: 314–21.

Kallio H, Snall EV, Kero MP, Rosenberg PH 2004. A comparison of intrathecal plain solutions containing ropivacaine 20 or 15 mg versus bupivacaine 10 mg. *Anesth Analg*; 99(3): 713–7.

Kalso E, Smith L, McQuay et al 2002. No pain, no gain: clinical excellence and scientific rigour – lessons learned from IA morphine. *Pain*; 98: 269–275.

Kalso E, Tramer MR, Carroll D, McQuay HJ, Moore RA.1997. Pain relief from intra-articular morphine after knee surgery: a qualitative systematic review. *Pain*, 71(2):127–34 21.

KHolte and H Kehlet,2002. Compensatory fluid administration for preoperative dehydration - does it improve outcome?, *Acta Anaesthesiol Scand* ;46:1089–93.

K Holte and H Kehlet, Postoperative ileus.2002. Progress towards effective management, *Drugs* 62, pp. 2603–2615

K Holte, NE Sharrock and H Kehlet,2002. Pathophysiology and clinical implications of perioperative fluid excess, *Br J Anaesth* 89;82: 622–32.

Kearney R, Mack C, Entwistle L.1998. Withholding oral fluids from children undergoing day surgery reduces vomiting. *Paediatr Anaesth* 1998; 8: 331–6.

Kehlet H.1997. Multimodal approach to control postoperative pathophysiology and rehabilitation. *Br J Anaesth*;78(5): 606–17.

H Kehlet and DW Wilmore.2002. Multimodal strategies to improve surgical outcome, *Am J Surg* 183, pp. 630–644

Klein SM, Nielsen KC, Greengrass RA, Warner DS, Martin A, Steele SM 2002. Ambulatory discharge after long-acting peripheral nerve blockade: 2382 blocks with ropivacaine. *Anesth Analg*; 94: 65–70.

Kohli M, Yuan L, Escobar M, et al.2003. A risk index for sternal surgical wound infection after cardiovascular surgery. *Infect Control Hosp Epidemiol*; 24: 17–25.

Koivuranta M, Laara E, Snare L, Alahuhta S 1997. A survey of postoperative nausea and vomiting. *Anaesthesia*; 52: 443–9.

Korhonen AM, Valanne JV, Jokela RM, Ravaska P, Korttila K 2003. Intrathecal hyperbaric bupivacaine 3 mg+fentanyl 10 microg for outpatient knee arthroscopy with tourniquet. *Acta Anaesthesiol Scand*;47(3):342-6.

Kooger Infante NE, Van Gessel E, Forster A, Gamulin Z 2000. Extent of hyperbaric spinal anesthesia influences the duration of spinal block. *Anesthesiology*;92(5):1319-23.

Kopman AF, Yee PS, Neuman GG 1997. Relationship of the train-of-four fade ratio to clinical signs and symptoms of residual paralysis in awake volunteers. *Anesthesiology*;86(4):765 - 71.

Korttila K, Linnoila M, Ertama P, Hakkinen S 1975. Recovery and simulated driving after intravenous anesthesia with thiopental, methohexital, propanidid, or alphadione. *Anesthesiology*; 43: 291–9.

Korttila K, Tammisto T, Ertama P, Pfaffli P, Blomgren E, Hakkinen S 1977. Recovery, psychomotor skills, and simulated driving after brief inhalational anesthesia with halothane or enflurane combined with nitrous oxide and oxygen. *Anesthesiology*; 46: 20–7.

Korttila KT 1995. Post-anaesthesia psychomotor and cognitive function. *Eur J Anaesthesiol Suppl*; 10: 43–6.

Kouri ME, Kopacz DJ 2004. Spinal 2-chloroprocaine: a comparison with lidocaine in volunteers. *Anesth Analg*; 98: 75–80.

Krane EJ, Haberkern CM, Jacobson LE.1995. Postoperative apnea, bradycardia, and oxygen desaturation in formerly premature infants: prospective comparison of spinal and general anesthesia. *Anesth Analg*; 80: 7–13.

Kuusniemi KS, Pihlajamaki KK, Pitkanen MT, Korkeila JE 1999. Low-dose bupivacaine: a comparison of hypobaric and near isobaric solutions for arthroscopic surgery of the knee. *Anaesthesia*;54(6):540–

Labaille T, Mazoit JX, Paqueron X, Franco D, Benhamou D 2002. The clinical efficacy and pharmacokinetics of intraperitoneal ropivacaine for laparoscopic cholecystectomy. *Anesth Analg*; 94(1):100 - 5.

Laffey JG, Carroll M, Donnelly N, Boylan JF 1998. Instructions for ambulatory surgery-patient comprehension and compliance. *Ir J Med Sci*; 167: 160–3.

Larsen JF, Svendsen FM, Pedersen V 2004. Randomized clinical trial of the effect of pneumoperitoneum on cardiac function and haemodynamics during laparoscopic cholecystectomy. *Br J Surg*;91(7):848- 54.

Lattermann R, Carli F, Wykes L, Schricker T. 2002. Epidural blockade modifies perioperative glucose production without affecting protein catabolism. *Anesthesiology*; 97: 374–81.

Lau H, Brooks DC 2001. Predictive factors for unanticipated admissions after ambulatory laparoscopic cholecystectomy. *Arch Surg*; 136: 1150–3.

Lean ME. 1999. Obesity and cardiovascular disease: the wasted years. *Br J Cardiol*, 6: 269–73.

Ledin T, Gupta A, Tytor M 1995. Postural control after propofol anaesthesia in minor surgery. *Acta Otolaryngol Suppl*; 520(Pt 2): 313–6.

Lee TH, Marcantonio ER, Mangione CM, et al. Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation* 1999; 100: 1043–9.

Leeder PC, Matthews T, Krzeminska K, Dehn TC 2004. Routine day-case laparoscopic cholecystectomy. *Br J Surg*; 91: 312–6.

Lemo P 2005. Development of clinical indicators for ambulatory surgery. *Euroanaesthesia refresher course Vienna*;28-31

Lennox PH, Vaghadia H, Henderson C, Martin L, Mitchell GW.2001. Small-dose selective spinal anesthesia for short-duration outpatient laparoscopy: recovery characteristics compared with propofol anaesthesia. *Br J Anaesth*;86(4):570 – 2

Lennox PH, Vaghadia H, Henderson C, Martin L, Mitchell GW 2002. Small-dose selective spinal anesthesia for short-duration outpatient laparoscopy: recovery characteristics compared with desflurane anesthesia. *Anesth Analg* ;94(2): 346–50.

Levy L, Pandit UA, Randel GI, Lewis IH, Tait AR.1992. Upper respiratory tract infections and general anaesthesia in children. Perioperative complications and oxygen saturation. *Anaesthesia*; 47: 678–82.

Lichtor JL, Alessi R, Lane BS 2002. Sleep tendency as a measure of recovery after drugs used for ambulatory surgery. *Anesthesiology*; 96: 878–83.

Liu SS, Strodtbeck WM, Richman JM et al 2005. A comparison of regional versus general anesthesia for ambulatory anesthesia: A meta-

analysis of randomized controlled trials. *Anesthesia and Analgesia*; 101: 1634–1642.

Liu SS, Strodtbeck WM, Richman JM, Wu CL 2005. *A Br J Anaesth*; 94: 520–3.

Lubarsky DA 1996. Fast-track in the postanesthesia unit: unlimited possibilities? *J Clin Anesth*; 8(Suppl): 70–2.

Lu PP, Brimacombe J, Yang C, Shyr M 2002. ProSeal versus the classic laryngeal mask airway for positive pressure ventilation during laparoscopic cholecystectomy. *Br J Anaesth*; 88(6): 824–7.

Macario A, Weinger M, Carney S, Kim A.1999. Which clinical anesthesia outcomes are important to avoid? The perspective of patients. *Anesth Analg*; 89: 652–8

Macario A, Weinger M, Carney S, Kim A.1999. Which clinical anesthesia outcomes are important to avoid? The perspective of patients. *Anesth Analg* 1999;89(3):652–8.

Magner JJ, McCaul C, Carton E, Gardiner J, Buggy D 2004. Effect of intraoperative intravenous crystalloid infusion on postoperative nausea and vomiting after gynaecological laparoscopy: comparison of 30 and 10 ml kg⁻¹. *Br J Anaesth*; 93: 381–5.

Malone DL, Genuit T, Tracy JK, Gannon C, Napolitano LM.2002. Surgical site infections: reanalysis of risk factors. *J Surg Res*; 103: 89–95.

Maltby JR, Beriault MT, Watson NC, Liepert D, Fick GH. The LMA-2002. ProSeal is an effective alternative to tracheal intubation for laparoscopic cholecystectomy. *Can J Anaesth*; 49(8): 857–62.

Maltby JR, Beriault MT, Watson NC, Liepert DJ, Fick GH 2002. LMA Classic and LMA-ProSeal are effective alternatives to endotracheal intubation for gynecologic laparoscopy. *Can J Anaesth*; 50(1): 71–7.

Marshall SI, Chung F 1999. Discharge criteria and complications after ambulatory surgery. *Anesth Analg*;88:508-517.

Marshall S, Chung F 1997. Assessment of 'home-readiness' -- discharge criteria and postdischarge complications. *Curr Opin Anesthesiol*; 10: 445–50.

Mattila K, Toivonen J, Janhunen L, Rosenberg PH, Hynynen M.2005. Postdischarge symptoms after ambulatory surgery: firstweek incidence, intensity, and risk factors. *Anesth Analg*;101(6):1643–50.

Mattioli G, Repetto P, Carlini C, et al 2002. Laparoscopic vs open approach for the treatment of gastroesophageal reflux in children. *Surg Endosc*;16(5):750 - 2.

MaxPherson GH, de Jonge M, Griffiths JM 1998. Rectus sheath and mesosalpinx block for laparoscopic sterilization. *Anaesthesia*;46(10):875-7.

Mazdisnian F, Palmieri A, Hakakha B, Hakakha M, Cambridge C, Lauria BJ 2002 . Office microlaparoscopy for female sterilization under local anesthesia. A cost and clinical analysis *Reprod Med*;47(2):97-100.

McAnulty GR, Robertshaw HJ, Hall GM.2000. Anaesthetic management of patients with diabetes mellitus. *Br J Anaesth*; 85: 80–90.

McCartney CJ, Brull R, Chan VW, et al. 2004. Early but no long-term benefit of regional compared with general anesthesia for ambulatory hand surgery. *Anesthesiology*; 101: 461–7.

McGrath B, Elgendy H, Chung F, Kamming D, Curti B, King S.2004. Thirty percent of patients have moderate to severe pain 24 h after ambulatory surgery: a survey of 5703 patients. *Can J Anaesth*;51(9):886–91.

McHugh GA, Thoms GM.2002. The management of pain following day-case surgery. *Anaesthesia* 2002;57(3):270–5.

McLeod GA 2004. Density of spinal anaesthetic solutions of bupivacaine, levobupivacaine, and ropivacaine with and without dextrose. *Br J Anaesth*;92(4):547–51 Erratum in: *Br J Anaesth*;93(4):602.

McLeod DH, Mitchell GW, Merrick PM, Chilvers CR 2000. Small-dose hypobaric lidocaine- fentanyl spinal anesthesia for short duration outpatient laparoscopy. I. A randomized comparison with conventional dose hyperbaric lidocaine. *Anesth Analg*;84(1):59-64.

Menes T, Spivak H 2000. Laparoscopy: searching for the proper insufflation gas. *Surg Endosc*; 14(11):1050 - 6.

Mezei G, Chung F.,1999. Return hospital visits and hospital readmissions after ambulatory surgery. *Ann Surg*; 230: 721–7.

Millar J.2004. Fast-tracking in day surgery. Is your journey to the recovery room really necessary? (Editorial) *Br J Anaesth* 2004; 93: 756–8.

Millar,J.M 2000. Patients selection, assessment, and preparation. In: *Day Care Anaesthesia*, Smith, 1. (Ed.), BMJ Books, London, Ch.1, Pp.1-30.

Miller LS, Rohling ML.2001. A statistical interpretative method for neuropsychological test data. *Neuropsychol Rev*; 11: 143–69. 1. Hedayati B, Fear S. Hospital admission after day-case

Milligan KA, Mowbray MJ, Mulrooney L et al.1988. Intra-articular bupivacaine for pain relief after arthroscopic surgery of the knee joint in daycase patients. *Anaesthesia*; 43: 563–564.

Moemen M E 2004. Recovery Characteristics after Ambulatory Anaesthesia and Surgery. *Eg J Anaesth*;20:449-457

Moiniche S, Mikkelsen S, Wetterslev J, Dahl JB.1999. A systematic review of intra-articular local anesthesia for postoperative pain relief after arthroscopic knee surgery. *Reg Anesth Pain Med*;24(5):430–7.

Moizo E, Marchetti C, Bergonzi PC, Putzu M, Lemi K, De LM, et al 2004. Unilateral spinal anesthesia for inguinal hernia repair: a prospective, randomized, double-blind comparison of bupivacaine, levobupivacaine, or ropivacaine. *Minerva Anesthesiol*;70:542–3.

Mulroy MF, Salinas FV, Larkin KL, Polissar NL 2002. Ambulatory surgery patients may be discharged before voiding after short-acting spinal and epidural anesthesia. *Anesthesiology*; 97: 315–9.

Muzi M, Ebert TJ, O'Brien TJ 2001. Induction of anesthesia and tracheal intubation with sevoflurane in adults. *Anesthesiology*;85(3):536-43.

Nadal JL, Fernandez BG, Escobar IC, Black M, Rosenblatt WH.1998. The palm print as a sensitive predictor of difficult laryngoscopy in diabetics. *Acta Anaesthesiol Scand*; 42: 199–203.

Nagasaka H, Awad H & Yaksh T 1996. Peripheral and spinal actions of opioids in the blockade of the autonomic response evoked by compression of the inflamed knee joint. *Anesthesiology* 1996; 85: 808 816.

Naja MZ, Ziade MF, Lonnqvist PA 2004. General anaesthesia combined with bilateral paravertebral blockade(T5-6) vs. general anaesthesia for laparoscopic cholecystectomy: a prospective, randomized clinical trial. *Eur J Anaesthesiol*;21(6):489- 95.

Neudecker J, Sauerland S, Neugebauer E, et al. 2002. The European Association for Endoscopic Surgery clinical practice guideline on the pneumoperitoneum for laparoscopic surgery. *Surg Endosc*;16(7):1121-43.

Newman MG, Trieger N, Miller JC.1969. Measuring recovery from anesthesia: a simple test. *Anesth Analg*; 48: 136–40.

Nielsen KC, Tucker MS, Steele SM 2005. Outcomes after regional anesthesia. *Int Anesthesiol Clin*; 43: 91–110.

Nordin P, Zetterstrom H, Gunnarsson U, Nilsson E 2003. Local, regional, or general anaesthesia in groin hernia repair: multicentre randomised trial. *Lancet*; 362: 853–8.

Osborne GA, Rudkin GE 1993. Outcome after day-care surgery in a major teaching hospital. *Anaesth Intensive Care*; 21: 822–7.

Palter SF.1999. Office microlaparoscopy under local anesthesia. *Obstet Gynecol Clin North Am*; 26(1): 109- 20.

Parnis SJ, Barker DS, Van Der Walt JH.2001. Clinical predictors of anaesthetic complications in children with respiratory tract infections. *Paediatr Anaesth*; 11: 29–400

Paulo Lemos, Ana Pinto, Gustavo Morais, José Pereira, et al.2009. Patient satisfaction following day surgery. *Journal of Clinical Anesthesia*; 21: 200–5.

Pavlin DJ, Chen C, Penaloza DA, Buckley FP.2004. A survey of pain and other symptoms that affect the recovery process after discharge from an ambulatory surgery unit. *J Clin Anesth*;16(3):200–6.

Pavlin DJ, Chen C, Penaloza DA, Polissar NL, Buckley FP 2002. Pain as a factor complicating recovery and discharge after ambulatory surgery. *Anesth Analg*; 95: 627–34.

Pavlin DJ, Pavlin EG, Gunn HC, Taraday JK, Koerschgen ME 1999. Voiding in patients managed with or without ultrasound monitoring of bladder volume after outpatient surgery. *Anesth Analg*; 89: 90–7.

Pavlin DJ, Rapp SE, Polissar NL, Malmgren JA, Koerschgen M, Keyes H 1998. Factors affecting discharge time in adult outpatients. *Anesth Analg*; 87: 816–26.

Pezzarossa A, Taddei F, Cimicchi MC, et al.1988. Perioperative management of diabetic subjects. Subcutaneous versus intravenous insulin administration during glucose-potassium infusion. *Diabetes Care*; 11: 52–8.

Pflug AE, Aasheim GM, Foster C 1978. Sequence of return of neurological function and criteria for safe ambulation following subarachnoid block (spinal anaesthetic). *Can Anaesth Soc J*; 25: 133–9.

Philip BK, Lombard LL, Roaf ER, Drager LR, Calalang I, Philip JH 1999. Comparison of vital capacity induction with sevoflurane to intravenous induction with propofol for adult ambulatory anesthesia. *Anesth Analg* ;89(3):623–7.

Philip BK, Scuderi PE, Chung F, et al 2000. Remifentanyl compared with alfentanil for ambulatory surgery using total intravenous anesthesia. *Anesth Analg*;84(3):521- 32.

Pittoni G, Toffoletto F, Calcarella G, Zanette G, Giron GP 1995. Spinal anesthesia in outpatient knee surgery: 22-gauge versus 25-gauge Sprotte needle. *Anesth Analg*; 81: 73–9.

Pomposelli JJ, Baxter JK III, Babineau TJ, et al.1998. Early postoperative glucose control predicts nosocomial infection rate in diabetic patients. *JPEN J Parenter Enteral Nutr* 1998; 22: 77–81.

Power I, Barratt S 2000 . Analgesic agents for the postoperative period. Nonopioids. *Surg Clin North Am*; 79(2):275- 95.

R Greif, O Akca and EP Horn et al.2000. Supplemental perioperative oxygen to reduce the incidence of surgical-wound infection, *N Engl J Med* 342, pp. 161–167.

Ræder J 2000. Epidural blocks for day surgery. *Techniques in Regional Anesthesia and Pain Management*; 4: 10–2.

Rolf N, Cote CJ.1992. Frequency and severity of desaturation events during general anesthesia in children with and without upper respiratory infections. *J Clin Anesth*; 4: 200–3

Rasmussen LS, Larsen K, Houx P, Skovgaard LT, Hanning CD, Moller JT 2001. The International Study of Postoperative Cognitive Dysfunction. The assessment of postoperative cognitive function. *Acta Anaesthesiol Scand*; 45: 275–89.

Rawal N 2001. Analgesia for day-case surgery. *British Journal of Anaesthesia* 87(1):73-87.

Rawal N, Allvin R, Amilon A, Ohlsson T, Hallen J 2001. Postoperative analgesia at home after ambulatory hand surgery: a controlled comparison of tramadol, metamizol, and paracetamol. *Anesth Analg*;92(2):347–51.

Rawal N 2005. Ambulatory wound and intraarticular infusions. In **Steele S, Nielsen K & Klein SM (eds.)**. Ambulatory anesthesia, perioperative analgesia. The McGraw Hill Companies;503–18.

Recart A, Gasanova I, White PF, et al. 2003. The effect of cerebral monitoring on recovery after general anesthesia: a comparison of the auditory evoked potential and Bispectral Index devices with standard clinical practice. *Anesth Analg*;97(6):1667- 74.

Reitan JA, Porter W, Braunstein M 1986. Comparison of psychomotor skills and amnesia after induction of anesthesia with midazolam or thiopental. *Anesth Analg*; 65: 933–7.

Reuben SS & Sklar J 2000. Pain management in patients who undergo outpatient arthroscopic surgery of the knee. *The Journal of Bone and Joint Surgery*; 82-A: 1754–1766.

Reuben SS, Pristas R, Dixon D, Faruqi S, Madabhushi L, Wenner S 2006. The incidence of complex regional pain syndrome after fasciectomy for Dupuytren's contracture: a prospective observational study of four anesthetic techniques. *Anesth Analg*; 102: 499–503.

Ritchie ED, Tong D, Chung F, Norris AM, Miniaci A, Vairavanathan SD 1997. Suprascapular nerve block for pain relief after arthroscopic shoulder surgery: a new modality? *Anesth Analg*; 84: 1306–12.

Robinson TN, Biffl WL, Moore EE, Heimbach JK, Calkins CM, Burch JM 2002. Predicting failure of outpatient laparoscopic cholecystectomy. *Am J Surg*; 184: 515–9.

Rodriguez-Chinchilla R, Rodriguez-Pont A, Pintanel T, Vidal-Lopez F 1996. Bilateral severe pain at L3-4 after spinal anaesthesia with hyperbaric 5% lignocaine. *Br J Anaesth*; 76: 328–9.

Rose DK, Cohen MM, Wigglesworth DF, DeBoer DP.1994. Critical respiratory events in the postanesthesia care unit: patient, surgical, and anesthetic factors. *Anesthesiology*; 81: 410–8.

Rose DK, Cohen MM. The airway: problems and predictions in 18,500 patients. *Can J Anaesth* 1994; 41(5 Pt 1): 372–83.

Rosseland LA, Stubhaug A, Breivik H 2002. Detecting postoperative urinary retention with an ultrasound scanner. *Acta Anaesthesiol Scand*; 46: 279–82.

Rudkin GE1997. Local and regional anaesthesia in the adult day surgery patient. In: *Practical anaesthesia and analgesia for day surgery*. UK: Bios Scientific Publishers Oxford; 207–10.

Russon K, Sardesai AM, Ridgway S, Whitear J, Sildown D, Boswell S, et al.2006. Postoperative shoulder surgery initiative (POSSI): an interim report of major shoulder surgery as a day case procedure. *Br J Anaesth* 2006;97(6):869–73.

Salihoglu Z, Demiroglu S, Dikmen Y.2003. Respiratory mechanics in morbid obese patients with chronic obstructive pulmonary disease and hypertension during pneumoperitoneum. *Eur J Anaesthesiol*;20(8):658-61.

Salpeter S, Greyber E, Pasternak G, Salpeter E.2002. Risk of fatal and nonfatal lactic acidosis with metformin use in type 2 diabetes mellitus. *Cochrane Database Syst Rev*; 2: CD002967.

Santanen U, Rautoma P, Luurila H, Erkola O, Pere P 2004. Comparison of 27-gauge (0.41-mm) Whitacre and Quincke spinal needles with respect to post-dural puncture headache and non-dural puncture headache. *Acta Anaesthesiol Scand*; 48: 474–9.

Sartorelli KH, Abajian JC, Kreutz JM, Vane DW.1992. Improved outcome utilizing spinal anesthesia in high-risk infants. *J Pediatr Surg* 1992; 27: 1022–5.

Schmidt J, Carbajo MA, Lampert R, Zirngibl H 2001. Laparoscopic intraperitoneal onlay polytetrafluoroethylene mesh repair(IPOM) for inguinal hernia during spinal anesthesia in patients with severe medical conditions. *Surg Laparosc Endosc Percutan Tech*; 11(1):34 - 7.

Schneider M, Ettlin T, Kaufmann M, et al.1993. Transient neurologic toxicity after hyperbaric subarachnoid anesthesia with 5% lidocaine. *Anesth Analg*; 76: 1154–7.

Schreiner MS, Nicolson SC 2000. Should children drink before discharge from day surgery? *Anesthesiology*; 76: 538–45.

Schreiner MS, Nicolson SC, Martin T, Whitney L.1992. Should children drink before discharge from day surgery? *Anesthesiology*; 76: 528–33.

Schreiner MS, O'Hara I, Markakis DA, Politis GD.1996. Do children who experience laryngospasm have an increased risk of upper respiratory tract infection? *Anesthesiology*; 85: 475–80.

Schricker T, Wykes L, Carli F.2000. Epidural blockade improves substrate utilization after surgery. *Am J Physiol Endocrinol Metab*; 279: E646–53.

Scott NB, Hodson M.1997. Public perceptions of postoperative pain and its relief. *Anaesthesia*;52(5):438–42.

Sener Mpektas, Z.O. Yilmaz, I Turkoz, A, Uckan, S, Donmez, A. and Arslan, G 2005. Comparison of Preemptive Analgesic Effects of a Single Dose of Nonopioid Analgesics for Pain Management After Ambulatory Surgery: A Prospective, Randomized, Single-Blind Study in Turkish Patients. *Curr Ther Res Clin Exp*; 66 (6): 541-551.

Shaikh S, Chung F, Imarengiaye C, Yung D, Bernstein M 2003. Pain, nausea, vomiting and ocular complications delay discharge following ambulatory microdiscectomy. *Can J Anesth*; 50: 514–8.

Shakir AAK, Ramachandra V, Hasan MA.1997. Day surgery postoperative nausea and vomiting at home related to peroperative fentanyl. *J One-day Surg*;6(3):10–1.

Shakir AAK, Ramachandra V, Hasan MA 1997. Day surgery postoperative nausea and vomiting at home related to peroperative fentanyl. *J One-day Surg*;6(3):10–1.

Shenkman Z, Hoppenstein D, Litmanowitz I, et al.2002. Spinal anesthesia in 62 premature, former-premature or young infants—technical aspects and pitfalls. *Can J Anesth*; 49: 262–9.

Shimizu S, Tanaka M, Konomi H, Tamura T, Mizumoto K, Yamaguchi K 2004. Spleen-preserving laparoscopic distal pancreatectomy after division of the splenic vessels. *J Laparoendosc Adv Surg Tech A*;14(3):173 - 7.

Sims C, Johnson CM.1994. Postoperative apnoea in infants. *Anaesth Intensive Care*; 22: 40–5.

Sinclair DR, Chung F, Smiley A 2003. General anesthesia does not impair simulator driving skills in volunteers in the immediate recovery period - a pilot study. *Can J Anesth*; 50: 238–45.

SJ Lewis, M Egger, PA Sylvester and S Thomas,2001. Early enteral feeding versus “nil by mouth” after gastrointestinal surgery: systematic review and meta-analysis of controlled trials, *BMJ* 323, pp. 773–776.

Smith AF, Pittaway AJ 2000. Premedication for anxiety in adult day surgery. *Cochrane Database Syst Rev*; (3):CD002192

Smith I 2007. Avoidance of fentanyl reduces nausea and vomiting associated with volatile induction and maintenance of anaesthesia with sevoflurane (abstract). *Anaesthesia*;62:309.

Smith I, Ding Y, White PF 1999. Muscle pain after outpatient laparoscopy: influence of propofol versus thiopental and enflurane. *Anesth Analg*;76(6): 1181- 4.

Smith I, Van Hemelrijck JV, White PF et al.1991. Effects of local anaesthesia on recovery after outpatient arthroscopy. *Anesthesia and Analgesia*; 73: 536–539.

Smith I.2007.Avoidance of fentanyl reduces nausea and vomiting associated with volatile induction and maintenance of anaesthesia with sevoflurane (abstract). *Anaesthesia*; 62:309.

Smith I.2007. Avoidance of fentanyl reduces nausea and vomiting associated with volatile induction and maintenance of anaesthesia with sevoflurane (abstract). *Anaesthesia* 2007;62:309.

Smith I 2001. Cost considerations in the use of anaesthetic drugs. *Pharmacoeconomics*;19(5 Pt 1):469–81.

Sneyd JR, Carr A, Byrom WD, Bilski AJ 1998 . A meta-analysis of nausea and vomiting following maintenance of anaesthesia with propofol or inhalational agents. *Eur J Anaesthesiol*;15(4):433–45.

Somri M, Gaitini L, Vaida S, Collins G, Sabo E, Mogilner G.1998. Postoperative outcome in high-risk infants undergoing herniorrhaphy: comparison between spinal and general anaesthesia. *Anaesthesia*; 53: 762–6.

Song D, Chung F, Ronayne M, Ward B, Yogendran S, Sibbick C 2004. Fast-tracking (bypassing the PACU) t 400 not reduce nursing workload after ambulatory surgery. *Br J Anaesth*; 93: 768–74.

Song D, Greilich NB, White PF, Watcha MF, Tongier WK 2000. Recovery profiles and costs of anesthesia for outpatient unilateral inguinal herniorrhaphy. *Anesth Analg*; 91: 876–81.

Song D, Joshi GP, White PF 1998. Fast-track eligibility after ambulatory anesthesia: a comparison of desflurane, sevoflurane, and propofol. *Anesth Analg*;86(2):267–73.

Song D, van Vlymen J, White PF 1998 . Is the bispectral index useful in predicting fast-track eligibility after ambulatory anesthesia with propofol and desflurane? *Anesth Analg*; 87: 1245–8.

Song D, White PF 1999. Remifentanyl as an adjuvant during desflurane anesthesia facilitates early recovery after ambulatory surgery. *J Clin Anesth*;11(5):364 – 7.

Song D, White PF 2000. Fast-track eligibility after ambulatory anesthesia: a comparison of desflurane, sevoflurane, and propofol . *Anesth Analg*;86(3):278- 88.

Spaliviero M, Gill IS 2010. Laparoscopic nephrectomy for large renal arteriovenous malformation. *Urology*;66(6):1176- 7.

Spivak H, Nudelman I, Fuco V, et al 2001. Laparoscopic extraperitoneal inguinal hernia repair with spinal anesthesia and nitrous oxide insufflation. *Surg Endosc*; 13(10):1026 -9.

Stevens JB, Walker SC, Hickey R 2000. Tracheal intubation using alfentanil and no muscle relaxant: is the choice of hypnotic important? *Anesth Analg*;84(6):1232- 6.

Sukhani R, Vazquez J, Pappas AL, Frey K, Aasen M, Slogoff S.1996. Recovery after propofol with and without intraoperative fentanyl in patients undergoing ambulatory gynecologic laparoscopy. *Anesth Analg*;83(5):975–81.

Sukhani R, Vazquez J, Pappas AL, Frey K, Aasen M, Slogoff S.1996. Recovery after propofol with and without intraoperative fentanyl in patients undergoing ambulatory gynecologic laparoscopy. *Anesth Analg*;83(5):975–81.

T Bisgaard, B Klarskov, J Rosenberg and H Kehlet,2001. Factors determining convalescence after uncomplicated laparoscopic cholecystectomy, *Arch Surg* 136, pp. 917–921.

Tait AR, Knight PR. The effects of general anesthesia on upper respiratory tract infections in children. *Anesthesiology* 1987; 67: 930–5.

Tait AR, Malviya S, Voepel-Lewis T, Munro HM, Siewert M, Pandit UA.2001. Risk factors for perioperative adverse respiratory events in children with upper respiratory tract infections. *Anesthesiology*; 95: 299–306.

Tait AR, Pandit UA, Voepel-Lewis T, Munro HM, Malviya S.1998. Use of the laryngeal mask airway in children with upper respiratory tract infections: a comparison with endotracheal intubation. *Anesth Analg*; 86: 706–11.

Tait AR, Reynolds PI, Gutstein HB.1995. Factors that influence an anesthesiologist's decision to cancel elective surgery for the child with an upper respiratory tract infection. *J Clin Anesth*; 7: 491–9.

Taitelman U, Reece EA, Bessman AN.1977. Insulin in the management of the diabetic surgical patient. Continuous intravenous infusion vs subcutaneous administration. *JAMA*; 237: 658–60.

Tang J, Wang B, White PF, Watcha MF, Qi J, Wender RH.1998. The effect of timing of ondansetron administration on its efficacy, cost-effectiveness, and cost-benefit as a prophylactic antiemetic in the ambulatory setting. *Anesth Analg*;86(2):274–82.

Taniguchi M, Bollen AW, Drasner K 2004. Sodium bisulfite: scapegoat for chloroprocaine neurotoxicity? *Anesthesiology*;100(1):85–91.

Tarkilla P, Huhtala J, Tuominen M 1995. Transient radicular irritation after spinal anaesthesia with hyperbaric 5% lignocaine. *Br J Anaesth*; 74: 328–9.

Thomas DJ, Platt HS, Alberti KG.1984. Insulin-dependent diabetes during the perioperative period. An assessment of continuous glucose-insulin-potassium infusion, and traditional treatment. *Anaesthesia*; 39: 629–37.

Thornes R. 2000.Just for the day. London: NAWCH.

Tiras MB, Gokce O, Noyan V, et al 2001. Comparison of microlaparoscopy and conventional laparoscopy for tubal sterilization under local anesthesia with mild sedation. *J Am Assoc Gynecol Laparosc*; 8(3):385 - 8.

Tobias JD, Lowe S, Hersey S, Brock III JW 1999. Caudal epidural block for analgesia following herniorrhaphy with laparoscopy in children. *J Laparoendosc Surg*; 4(2):117- 20.

Tobias JD, Holcomb III GW, Brock III JW, et al. 2001. Analgesia after inguinal herniorrhaphy with laparoscopic inspection of the peritoneum in children. Caudal block versus ilioinguinal/iliohypogastric block. *Am J Anesthesiol*; 22(4):167- 77.

Toivonen J, Permi J, Rosenberg PH 2001. Effect of preincisional ilioinguinal and iliohypogastric nerve block on postoperative analgesic requirement in day-surgery patients undergoing herniorrhaphy under spinal anaesthesia. *Acta Anaesthesiol Scand*; 45: 603–7.

Tramer M, Moore A, McQuay H 1997. Meta-analytic comparison of prophylactic antiemetic efficacy for postoperative nausea and vomiting: propofol anaesthesia vs omitting nitrous oxide vs total i.v. anaesthesia with propofol. *Br J Anaesth*; 78(3):256–9.

Tramer M, Moore A, McQuay H 2001. Omitting nitrous oxide in general anaesthesia: meta-analysis of intraoperative awareness and postoperative emesis in randomized controlled trials. *Br J Anaesth*;76(3):194- 99.

Tucker DH, Sieker HO.1960. The effect of change in body position on lung volumes and intrapulmonary gas mixing in patients with obesity, heart failure, and emphysema. *Am Rev Respir Dis*; 82: 787–91.

Twersky R, Fishman D, Homel P 1997. What happens after discharge? Return hospital visits after ambulatory surgery. *Anesth Analg*; 84: 319–24.

Tzabar Y, Asbury AJ, Millar K 1996. Cognitive failures after general anaesthesia for day-case surgery. *Br J Anaesth*; 76: 194–7.

Uemura N, Nomura M, Inoue S, et al 2002. Changes in hemodynamics and autonomic nervous activity in patients undergoing laparoscopic cholecystectomy: differences between the pneumoperitoneum and abdominal wall-lifting method. *Endoscopy*;34(8):643- 50.

Vaghadia H, McLeod DH, Mitchell GW, Merrick PM, Chilvers CR 1997. Small-dose hypobaric lidocaine-fentanyl spinal anesthesia for short duration outpatient laparoscopy. I. A randomized comparison with conventional dose hypobaric lidocaine. *Anesth Analg*; 84: 59–64.

Vaghadia H, Scheepers L, Merrick PM 1998. Readmission for bleeding after outpatient surgery. *Can J Anaesth*; 45: 1079–83.

Vaghadia H 2000. Spinal anesthesia for outpatients: controversies and new techniques. *Can J Anesth*;45(5 Pt 2):77-91.

Vaghadia H, Viskari D, Mitchell GW, Berrill A 2001. Selective spinal anesthesia for outpatient laparoscopy. I: characteristics of three hypobaric solutions. *Can J Anaesth*;48(3):256- 60.

van den Berg AA, Chitty DA, Jones RD, Sohel MS, Shahan A 2005. Intravenous or inhaled induction of anesthesia in adults? An audit of preoperative patient preferences. *Anesth Analg*;100(5):1422–4.

Vath JS, Kopacz DJ 2004. Spinal 2-chloroprocaine: the effect of added fentanyl. *Anesth Analg* ;98(1):89–94.

Verghese C, Brimacombe JR 1999. Survey of laryngeal mask airway usage in 11,910 patients: safety and efficacy for conventional and nonconventional usage. *Anesth Analg*;82(1): 133 - 38.

Veverka TJ, Henry DN, Milroy MJ, et al.1991. Spinal anesthesia reduces the hazard of apnea in high-risk infants. *Am Surg*; 57: 531–4

Vickers MD 1965. The measurement of recovery from anaesthesia. *Br J Anaesth*; 37: 296–302.

Vofsi O, Barak M, Moscovici R, Bustan M, Katz Y 2004. Cardiorespiratory parameters during conventional or gasless gynecological laparoscopy under general or regional anesthesia. *Med Sci Monit*; 10(4):CR152- 5.

Wang JJ, Ho ST, Tzeng JI, Tang CS.2000. The effect of timing of dexamethasone administration on its efficacy as a prophylactic antiemetic for postoperative nausea and vomiting. *Anesth Analg*;91(1):136–9.

Ward B, Imarengiaye C, Peirovy J, Chung F 2005. Cognitive function is minimally impaired after ambulatory surgery. *Can J Anesth*; 52: 1017–21.

Warner MA, Sheilds SE, Chute CG.1993. Major morbidity and mortality within 1 month of ambulatory surgery and anaesthesia. *JAMA*; 270: 1437–41.

Warner ME, Contreras MG, Warner MA, Schroeder DR, Munn SR, Maxson PM.1998. Diabetes mellitus and difficult laryngoscopy in renal and pancreatic transplant patients. *Anesth Analg* 1998; 86: 516–9.

Warner,M.A, Shields, S.E. and Chute, C.G 1993. Major morbidity and mortality within 1 month of ambulatory surgery and anaesthesia.*JAMA*;; 270-1437.

Watcha MF, Thach BT, Gunter JB.1989. Postoperative apnea after caudal anesthesia in an ex-premature infant. *Anesthesiology*; 71: 613–5.

Watson BJ, Allen JG 1999. Spinal anaesthesia in day surgery—an audit of the first cases. *J One-day Surg* 2003;12(4): 1 Marshall SI, Chung F. Discharge criteria and complications after ambulatory surgery. *Anesth Analg*; 88: 508–17.

Watson B, Howell V 2007. Spinal anaesthesia: The saviour of day surgery?. *Current Anaesthesia & Critical Care*; 18: 193–99.

Welborn LG, Hannallah RS, Fink R, Ruttimann UE, Hicks JM.1989. High-dose caffeine suppresses postoperative apnea in former preterm infants. *Anesthesiology*; 71: 347–9.

Welborn LG, Hannallah RS, Luban NL, Fink R, Ruttimann UE.1991. Anemia and postoperative apnea in former preterm infants. *Anesthesiology*; 74: 1003–6.

Welborn LG, Rice LJ, Hannallah RS, Broadman LM, Ruttimann UE, Fink R.1990. Postoperative apnea in former preterm infants: prospective comparison of spinal and general anesthesia. *Anesthesiology*; 72: 838–42.

Welge JA, Myer CM 2007. Steroids for improving recovery following paediatric tonsillectomy. *Cochrane Database Syst Rev*; 1: CD003997.

White H, Black RJ, Jones M, Mar Fan GC.2007. Randomized comparison of two anti-emetic strategies in high-risk patients undergoing day-case gynaecological surgery. *Br J Anaesth*;98(4):470–6.

White PF, Ma H, Tang J, Wender RH, Sloninsky A, Kariger R 2004. Does the use of electroencephalographic bispectral index or auditory evoked potential index monitoring facilitate recovery after desflurane anesthesia in the ambulatory setting? *Anesthesiology*; 100: 811–7.

White PF, Song D 1999. New criteria for fast-tracking after outpatient anesthesia: a comparison with the modified Aldrete's scoring system. *Anesth Analg*; 88: 1069–72.

White PF 1998. Bypassing (fast-tracking) of the recovery room after ambulatory surgery. *Acta Anaesthesiol Scand*; 42: 189–91.

Whiteside JB, Burke D, Wildsmith JA 2003. Comparison of ropivacaine 0.5% (in glucose 5%) with bupivacaine 0.5% (in glucose 8%) for spinal anaesthesia for elective surgery. *Br J Anaesth*;90(3):304–8.

Williams BA, Kentor ML, Williams JP, et al 2002. PACU bypass after outpatient knee surgery is associated with fewer unplanned hospital admissions but more phase II nursing interventions. *Anesthesiology*; 97: 981–8.

Williams JM, Stoddart PA, Williams SA, Wolf AR.2001. Post-operative recovery after inguinal herniotomy in ex-premature infants: comparison between sevoflurane and spinal anaesthesia. *Br J Anaesth* 2001; 86: 366–71.

Wilmore DW, Kehlet H 2001. Management of patients in fast track surgery. *BMJ*;322(7284):473 - 6.

Wilson AT, Nicholson E, Burton L, Wild C 2004. Analgesia for day-case shoulder surgery. *Br J Anaesth*; 92: 414–5.

Wu CL, Berenholtz SM, Pronovost PJ, Fleisher LA 2002. Systematic review and analysis of postdischarge symptoms after outpatient surgery. *Anesthesiology*; 96: 994–1003.

Wu CL, Hsu W, Richman JM, Raja SN 2004. Postoperative cognitive function as an outcome of regional anesthesia and analgesia. *Reg Anesth Pain Med*; 29: 257–68.

Wu CL, Rouse LM, Chen JM, Miller RJ.2002. Comparison of postoperative pain in patients receiving interscalene block or general anesthesia for shoulder surgery. *Orthopedics*; 25: 45–8.

WY Park, JS Thomson and KK Lee,2001. Effect of epidural anesthesia and analgesia on perioperative outcome: a randomised controlled Veterans Affairs cooperative study, *Ann Surg* 234 (2001), pp. 560–569.

Yang H, Choi PT, McChesney J, Buckley N 2004. Induction with sevoflurane-remifentanil is comparable to propofol- fentanyl-rocuronium in PONV after laparoscopic surgery. *Can J Anaesth*; 51(7):660- 7.

Yao FS, Savarese JJ.1998. Morbid obesity in anesthesiology, problem-oriented patient management. *In:* Yao FS, Artusio JF Jr (Eds). *Anesthesiology. Problem-Oriented Patient Management*, 3rd ed. PA: Lippincott-Raven;; 1001–19.

Yogendran S, Asokumar B, Cheng DC, Chung F.1995. A prospective randomized double-blinded study of the effect of intravenous fluid

therapy on adverse outcomes on outpatient surgery. *Anesth Analg*;80(4):682–6.

Yogendran S, Asokumar B, Cheng DC, Chung F 1995. A prospective randomized double-blinded study of the effect of intravenous fluid therapy on adverse outcomes on outpatient surgery. *Anesth Analg*; 80: 682–6.

Yoos JR, Kopacz DJ 2005. Spinal 2-chloroprocaine: a comparison with small-dose bupivacaine in volunteers. *Anesth Analg*; 100: 566–72.

Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S.1993. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med*; 328: 1230–5.

Zullo F, Pellicano M, Zupi E, Guida M, Mastrantonio P, Nappi C 2000. Minilaparoscopic ovarian drilling under local anesthesia in patients with polycystic ovary syndrome. *Fertil Steril*; 74(2):376- 9.

Zupi E, Marconi D, Sbracia M, et al. 2000. Is local anesthesia an affordable alternative to general anesthesia for minilaparoscopy? *J Am Assoc Gynecol Laparosc*; 7(1):111 - 4.

Zvara DA, Mathes DD, Brooker RF, McKinley CA 1996. Video as a patient teaching tool: does it add to the preoperative anesthetic visit? *Anesth Analg*; 82: 1065–8.