References



- Adams M, Freeman B, Nelson I and Dolan P (2000): Mechanical initiation of intervertebral disc degeneration. Spine; 25: 1625-36.
- Advanced Trauma Life Support for Doctors (2004): Student Course Manual 7th ed. Chicago, Ill: American College of Surgeons Committee on Trauma.
- Albin MS, Ritter RR, Pruett CE and Kalff K (1991): Venous air embolism during lumbar laminectomy in the prone position: Report of three cases. Anesth Angl; 73: 346-349.
- Albrecht RM, Kingsley D and Schermer CR (2001): Evaluation of cervical spine in intensive care patients following blunt trauma. World J Surg; 25(8): 1089-1096.
- **Aldrich TK and Rochester DF (2002):** The lungs and neuromuscular disease. In: Murray JR, Nadell J, eds. Textbook of Respiratory Medicine. Philadelphia: WB Saunders; 2329-2356.
- **Amar AP and Levy ML (1999):** Pathogenesis and pharmacological strategies for mitigating secondary damage in acute spinal cord injury. Neurosurgery; 44(5): 1027-1040.
- American Association of Neurological Surgeons and the Congress of Neurological Surgeons (2002): Cervical spine immobilization before admission to the hospital: Chapter 1. Neurosurgery; 50(Suppl): S7-17.
- American Association of Neurological Surgeons and the Congress of Neurological Surgeons (2002): Blood pressure

management after acute spinal cord injury: Chapter 8. Neurosurgery; 50: S58-62.

- American Association of Neurological Surgeons and the Congress of Neurological Surgeons (2002): Pharmacological therapy after acute cervical spinal cord injury: Chapter 9. Neurosurgery; 50: S63-72.
- American College of Surgeons Committee on Trauma (2004):

 Advanced trauma Life Support for Doctors Student Course

 Manual. Chicago: American College of Surgeons.
- American Society of Anaesthesiologists (2003): Practice guidelines for pulmonary artery catheterization: An updated report task force on pulmonary artery catheterization. Anaesthesiology; 99(4): 988-1014.
- American Society of Anaesthesiologists (2004): Standards for basic anaesthetic monitoring. www.asahq.org.
- American Society of Anaesthesiologists (2005): Practice guidelines for management of the difficult airway: An updated report on the task force on management of the difficult airway. Anaesthesiology; 98(5): 1269-1277.
- American Spinal Injury Association (2005): Standard neurological classification of SCI. www.asia-spinalinjury.org.
- **Amzallag M (1993):** Autonomic hyperrefexia. International Anaesthesiology Clinics; 31: 87-102.
- Anatomy and pathophysiology of traumatic spinal cord injury. Fractures of Cervical, Thoracic, and Lumbar Spine; 2: 9-21.

Anderton JM. (1991): The prone position for the surgical patient. Br J Anaesth; 32-37.

- Andreas Erdmann (2001): Anatomy of the spinal cord. Concise Anatomy for Anaesthesia; 44-47.
- Anke A, Aksnes AK, Stanghelle Jk and Hjltnes N (1993): Lung volumes in tetraplegic patients according to cervical spinal cord injury level. Scandinavian Journal of Rehabilitation medicine; 25: 73-77.
- Arrica M and Bissonnette B. (2007): Therapeutic hypothermia.

 Semin Cardiothorac Vasc Anaesth; 11: 6-15.
- Association of Anaesthetists of Great Britain and Ireland (2000):

 Recommendations for standards of monitoring (3rd ed).

 London: Association of Anaesthetists of Great Britain and Ireland.
- **Attygale D and Rodrigo N. (1999):** Magnesium sulphate for the control of spasms in severe tetanus. Anaesthesia; 54: 297-310.



- **Ball PA (2001):** Critical care of spinal cord injury. Spine; 26(suppl24): S27-S30.
- Baptiste and Fehlings (2008): Emerging drugs for spinal cord injury. Expert Opin. Emerg. Drugs; 13: 63-80.
- **Barker and Saulino (2002):** First ever guidelines for spinal cord injuries. RN; 65(10): 32-37.

Basso, DM (2004): Behavioral testing after spinal cord injury: Congruities, complexities, and controversies. J. Neurotrauma; 21: 395-404.

- **Belanger E and Levi AD (2000):** The acute and chronic management of spinal cord injury. J Am Colleg Surg; 190: 603-18.
- **Bendo AA, Kass IS and Hartung J (2006):** Anaesthesia for neurosurgery. In: Barash PG, et al., eds. Clinical anaesthesia, 5th ed. Philadelphia, PA: Lippincott Williams and Wilkins; 746-789.
- Bensmail D, Ecoffey C, Ventura M and Albert T (2009): Chronic neuropathic pain in patients with spinal cord injury. What is the efficacy of regional interventions? Sympathetic blocks, nerve blocks and intrathecal drugs. Annals of Physical and Rehabilitation Medicine; 52: 42-148.
- **Benumof JL** (1998): Neurologic diseases in anaesthesia and uncommon diseases, 4th ed, Saunders; 16-21.
- Benzel EC and Doezema D (1996): Prehospital management of the spinally injured patient. In: Narayan RK, Wilberger JE, Povlishock JT (eds). Neurotrauma. McGraw Hill.
- Bernhard JM, Passuti N and Pinaud M (1992): Long term hypotensive technique with nicardipine and nitroprusside during isoflurane anaesthesia for spinal surgery. Anaesth. Analg; 75: 179-185.
- Bernhard M, Helm M, Aul A and Gries A (2004): Preclinic management of multiple trauma. Anaesthesist; 53: 887-904.

Bethea JR (2000): Spinal cord injury-induced inflammation: a dualedged sword. Prog Brain Res; 128: 33-42.

- Bhasker KR, Brown R, O'Sulliyan DD, Melia S,Duggan M and Reid L (1991): Bronchial mucus hypersecretion in acute quadriplegia. Macromolecular yields and glycocorjugate composition. American Review of respiratory Disease; 143: 640-648.
- **Biering-Sorensen F (2002):** Urinary tract infection in individuals with spinal cord lesion. Curr Opin Urol; 12(1): 45-49.
- **Blackmer J (2003):** Autonomic dysreflexia: Rehabilitation medicine: CMAJ; 169(9): 931–935.
- Bledsoe BE, Wesley AK and Salomone JP (2004): High-dose steroids for acute spinal cord injury in emergency medical services. The National Association of EMS Physicians and Clinical Practice Committee Prehosp Emerg Care; 8: 313-6.
- **Bob Winter and Dave Knight (2008):** Cord injuries how to prevent secondary injury. Anaesthesia and Intensive Care Medicine; 9(9): 401-403.
- Bolinger B, Shartz M and Marion D (2004): Bedside fluoroscopic flexion and extension cervical spine radiographs for clearance of the cervical spine in comatose trauma patients. J Trauma; 56(1): 132-136.
- **Bonny JB and Thomas MS (2007):** Spinal cord injuries: Emergency medicine: A comprehensive study guide, 6th ed. Chapter 256.
- Bracken MB and Holford TR (2002): Neurological and functional status one year after acute spinal cord injury: estimates of


functional recovery in National Acute Spinal Cord Injury Study II from results modeled in National Acute Spinal Cord Injury Study III. J Neurosurg; 96(3): 259-266.

- Brambilla S, Ruosi C and La Maida GA (2004): Prevention of venous thromboembolism in spinal surgery. Eur Spine J; 13(1): 1-8.
- Brian KK, Cody M, Hong MS and Alan SH (2004): Hypothermia for spinal cord injury. The Spine Journal; 8: 859-874.
- Brian KK, Wolfram Tetzlaff and Jonathan NG (2004):

 Pathophysiology and pharmacologic treatment of acute spinal cord injury. The Spine Journal; 4: 451-464.
- Brian T, King BS, Gupta R and Narayan RK (2000): The early assessment and intensive care unit management of patients with severe traumatic brain and spinal cord injuries. Surgical Clinics of North America; 72: 80.
- **Briggs ED and Kirsch JK (2003):** Anaesthetic implications of neuromuscular disease: Spinal cord injury. Journal of Anaesthesia; 17: 177-185.
- **British Trauma Society (2002):** Guidelines for the initial management and assessment of spinal injury. Injury; 34: 405-425.
- Brock M and Romsbacher J (1998): Lumbar spinal stenosis. Crit Rev Neurosug; 8(6): 333-337.
- Brohi K, Healy M and Fotheringham T (2005): Helical computed tomographic scanning for the evaluation of the cervical spine in the unconscious intubated trauma patient. J Trauma; 58: 897-901.

Buhre W, Weland A, Buhre K and Kazmaier S (2000): Effects of the sitting position on the distribution of blood volume in patients undergoing neurosurgical procedures. Br J Anaesth; 84(3): 354-357.

Burns AS, Lee BS, Ditunno JF Jr and Tessler A (2003): Patient selection for clinical trials: The reliability of the early spinal cord injury. J Neurotrauma; 20: 477-82.



- Cao Q, Zhang YP, Iannotti, and DeVries (2005): Functional and electrophysiological changes after graded traumatic spinal cord injury in adult rat. Exp. Neurol.; 191: S3-S16.
- Catharyn T, Liverman Bruce and Altevogt M (2005): Spinal cord injury: Progress, Promise and Priorities.
- Christensen MD and Hulseboseh CE (1997): Chronic Central pain after spinal cord injury. J Neurotrauma; 14: 517-537.
- Christie SD, Comeau B, Myers T, Sadi D, Purdy M and Mendez I (2008): Duration of lipid peroxidation after acute spinal cord injury in rats and the effect of methylprednisolone. Neurosurg Focus; 25: E5.
- Claire EH (2002): Recent advances in pathophysiology and treatment of spinal cord injury.
- Colachis SC (1992): Autonomic hyperreflexia with spine cord injury. Journal of the American Paraplegia Society; 15: 171-186.

Coleman WP, Benzel D and Cahill DW (2000): A critical appraisal of the reporting of the national acute spinal cord injury studies (II and III) of methylprednisolone in acute spinal cord injury. J Spinal Disord; 13(3): 185-99.

Cooper DJ, Myles PS and McDermott FT (2004):

- Crosby E (2002): Airway management after upper cervical spine injury: what have we learned? Can J Anaesth; 49: 733-744.
- Curt A, Schwab ME and Dietz V (2004): Providing the clinical basis for new interventional therapies: Refined diagnosis and assessment of recovery after spinal cord injury. Spinal Cord; 42(1): 1-6.
- Cuzzocrea S, Riley DP, Caputi AP and Salvemini D (2001):

 Antioxidant therapy: a new pharmacological approach in shock, inflammation, and ischaemia/ reperfusion injury.

 Pharmacol Rev; 53(1): 135-59.



- **Dangor A and Lam AM (1999):** Perioperative management of patients with head and spinal cord trauma. Anaesthesiolgy Clinics of North America; 17: 155-170.
- **Danon J, Druz WS and Goldberg NB (2002):** Function of the isolated paced diaphragm and the cervical accessory muscles in C₁ quadriplegics. Am Rev Respir Dis; 119: 909.
- **Davis BD, Fowler R, Kupas DF and Ropollo LP (2002):** Role of rapid sequence induction in the prehospital setting: helpful or harmful? Curr Opin Crit Care; 8: 571-7.

Davis JW, Kaups KL and Cunningham MA (2001): Routine evaluation of the cervical spine in head-injured patients with dynamic fluoroscopy: A reappraisal. J Trauma; 50(6): 1044-1047.

- De Latorre F, Nolan J, Robertson C, Chamberlain D and Baskett P (2001): European resuscitation council guidelines for adult advanced life support. A statement from the advanced life support working group and approved by the executive committee of the European resuscitation council. Resuscitation; 48: 211-21.
- **Deletis V (2006):** Intraoperative neurophysiology of the corticospinal tract in the spinal cord. Functional Neuroscience; and related techniques. Supplements to Clinical Neurophysiology; 59: 107-112.
- **Demetriades D, Charalambides K and Chahwan S (2000):** Non-skeletal cervical spine injuries: Epidemiology and diagnostic pitfalls. J Trauma; 48(4): 724-727.
- **Denise IC (2009):** Autonomic dysreflexia in spinal cord injury. Spinal Cord; 132-136.
- **Diaz JJ, Aulino JM and Collier B (2005):** The early work-up for isolated ligamentous injury of the cervical spine: Does computed tomography scan have a role? J Trauma; 59(4): 897-904.
- **Dietrich WD, Chatzipanteli K and Vitarbo E (2004):** The role of inflammatory processes in the pathophysiology and treatment of brain and spinal cord trauma. Acta Neurochir Suppl; 89: 69-74.

Ditunno JF and Formal CS (2004): Chronic spinal cord injury. Spinal Cord; 132-136.

- **Ditunno JF, Little JW and Tessler A (2004):** Spinal shock revisited: A four-phase model. Spinal Cord; 42(7): 383-395.
- **Dobkin BH and Havton LA (2004):** Basic advances and new avenues in therapy of spinal cord injury. Annu RevMed; 55: 255-282.
- **Dunham CM, Barraco RD, Clark DE and Daley BJ (2003):**Guidelines for emergency tracheal intubation immediately after traumatic injury. J Trauma; 55: 162-179.




- Ebert TJ, Kanitz DD and Kampine JP (1990): Inhibition of Sympathetic neural outflow during Thiopental anaesthesia in human. Anaesth. Analg.; 71: 319-326.
- Edlich RF, Wish JR, Britt LD and Long WB (2004): An organized approach to trauma care: Legacy of Adams Cowley.

 J Long Term Eff Med Implants; 14(6): 481-511.
- Edward Morgan G, Maged SM and Michael JM (2006): Clinical anesthesiology. 3rd edition; 16: 253-282.
- Ellis H and Feldman S (1997): Anatomy for anaesthetists, 7th ed. Blackwell Science, Oxford.
- Eltorai IM, Wong DH and Lacerna M. (1997): Surgical aspects of autonomic dysreflexia. J Spinal Cord Med; 20(3): 361-4.
- Esmail Z, Shalasky KF, Sunderji R, Anton H, Chambers K and Fish (2002): Evaluation of captopril for the management of

hypertension in autonomic dysreflexia: A pilot study. Arch Phys Med Rehabil; 83: 604-608.

Estenne M, Van Muylem A, Gorini M and Kinnear W (1994):

Evidence of dynamic airway compression during cough in tetraplegic patients. American Journal of Respiratory and Critical Care Medicine; 150: 1081-1085.



- **Fawcett JW, Curt A and Steeves JD (2007):** Guidelines for the conduct of clinical trials for spinal cord injury as developed by the ICCP panel: Spontaneous recovery after spinal cord injury and statistical power needed for therapeutic clinical trials. Spinal Cord; 45: 190-205.
- Fawcett JW, Haxby EJ and Mala DA (1999): Magnesium: Physiology and pharmacology. Br J Anaesth; 83: 302-20.
- **Fehlings MG (Editorial) (2001):** Recommendations regarding the use of methylprednisolone in acute spinal injury: making sense out of the controversy. Spine; 26(24S): S56-S57.
- **Fehlings MG and Perrin RG (2006):** The timing of surgical intervention in the treatment of spinal cord injury: a systematic review of recent clinical evidence. Spine; 31: S28-35.
- **Fehlings M and Sekhon L (2001):** Acute interventions in SCI: What do we know, what should we do? Clin Neurosurg; 48: 226-242.
- **Fife C, Otto G and Capsuto EG (2001):** Incidence of pressure ulcers in a neurologic intensive care unit. Crit Care Med; 29(2): 283-290.

Fife D and Kraus J (1986): Anatomic location of spinal cord injury :Relationship to the cause of injury. Spine; 11: 2-5.

- **Fisher CG, Noonan VK, Smith DE and Wing PC (2005):** Motor recovery, functional status, and health-related quality of life in patients with complete spinal cord injuries. Spine; 30: 2200-7.
- **Fox R and Watling (2001):** Focus on spinal injury: Anaesthesia for patients with chronic spinal cord injury. Current Anaesthesia and Critical Care; 12: 154-158.
- **Fox RSR and Mulnier (2001):** Focus on spinal injury: The anaesthetic management of acute spinal injury patients: Current Anaesthesia and Critical Care; 12: 147-153.
- Francisco de Assis Aquino Gondim (2005): Spinal cord, topographical and functional anatomy. Spine; 177-185.
- **Frederic SB and Darryl YS (2003):** Cervical spinal cord injuries: Neurosurgical critical care. Current Critical Care Diagnosis and Treatment; 31: 741-747.
- Freedman I, van Gelderen D and Cooper DJ (2005): Cervical spine assessment in the unconscious trauma patient: a major trauma service's experience with passive flexion-extension radiography. J Trauma; 58(6): 1183-1188.



- **Gary RS, Randall MS and Daniel JC (2001):** Spinal cord injury. Anesthesia and Neurosungery; 33: 715-747.
- **Gavelli G, Canini R and Bertaccini P (2002):** Traumatic injuries: imaging of thoracic injuries. European Radiology; 12(6): 1273-1294.

Geerts WH, Pineo GF and Heit J (2004): Prevention of venous thromboembolism: the seventh ACCP conference on antithrombotic and thrombolytic therapy. Chest; 126: 338S-400S.

- **Geron Press Release (2009):** Geron receives FDA clearance to begin world's first human clinical trial of embryonic stem cell based therapy.
- **Gibson P, Lum Hee W and Vadodaria B (2004):** Anaesthesia for spinal surgery in adults. Br. J. Anaesth.; 92(5): 770-772.
- Goldberg W, Mueller C and Panacek E (2001): Distribution and patterns of blunt traumatic cervical spine injury. Ann Emerg Med; 38: 17-21.
- Gondim FA, Lopes AC and Oliveira GR (2004): Cardiovascular control after spinal cord injury. Curr Vasc Pharmacol; 2: 71-79.
- **Goy J (1994):** Spinal injuries. In: Loach A., ed. Orthopedic Anaesthesia. Edward Arnold: London; 145-157.
- Gray H, Williams PL and Bannister LH (1995): Nervous system. In: Williams PL, ed. Gray's Anatomy. The anatomical basis of medicine and surgery, 38th ed. New York, NY: Churchill Livingstone; 901-1398.
- **Gregory D and Susan A (2005):** Basic and clinical anatomy of the spine: Spinal cord, and ANS. Spine; 177-185.
- **Grundy D and Swain A (2002):** ABC of spinal cord injury, fourth ed. Early management and complications. BMJ Publishing Group, London; 17-20.


Gunn AJ and Thoresen M (2006): Hypothermic neuroprotection. NeuroRx; 3: 154-69.



- Hadley MN, Walters BC and Grabb PA (2002): Guidelines for the management of acute cervical spine and spinal cord injuries. Clin Neurosurg; 49: 407-498.
- **Hall ED (2003):** Drug development in spinal cord injury: What is the FDA looking for? J. Rehabil. Res. Dev.; 40: 81-91.
- Hambly PR and Martin B (1998): Anaesthesia for chronic spinal cord lesions. Anaesthesia; 53: 273-289.
- Hampton T (2005): Spinal cord network. JAMA; 293: 2463.
- **Harrison P (2000):** Systemic effects of spinal cord injury: Respiratory system. Managing spinal injury: Critical care. The Spinal Injuries Association, London; 66-74.
- Hauser CJ, Visvikis G and Hinrichs C (2003): Prospective validation of computed tomographic screening of the thoracolumbar spine in trauma. J Trauma; 55(2): 228-234.
- Heller J and Pedlow F (1998): Anatomy of the cervical spine. Clark C, ed. The cervical spine. Philadelphia: Lippincott-Raven; 12-14.
- Hendey GW, Wolfson AB, Mower WR and Hoffman JR (2002): Spinal cord injury without radiographic abnormality: Results of the national emergency X-radiography utilization study in blunt cervical trauma. J Trauma; 53: 1-4.

Hess D. (2000): Detection an monitoring of hypoxia and oxygen therapy. Resp. Care; 45(10): 65-80.

- Hill L and Zanaboni P (2004): Vasodilators. In: Evers AS, Maze M, eds. Anaesthetic pharmacology: Physiologic principles and cinical practice. A companion to anaesthesia. Philadelphia, PA: Churchill Livingstone; 691-702.
- Holly LT, Kelly DF, Counelis GJ and Blinman T (2002): Cervical spine trauma associated with moderate and severe head injury: Incidence, risk factors, and injury characteristics. J Neurosurg; 96: 285-91.
- Holmes JF, Miller PQ, Panacek EA and Lin S (2001): Epidemiology of thoracolumbar spine injury in blunt trauma. Acad Emerg Med; 8: 866-872.
- **Hulbert RJ and Moulton RJ (2002):** Why do you prescribe methylprednisolone for acute spinal cord injury? A Canadian perspective and position statement. Can J Neurol Sci; 29: 236-9.
- **Hurlbert RJ (2006):** Strategies of medical intervention in the management of acute spinal cord injury. Spine; 31: S16-S21.



- **Intensive Care Society (2002):** Guidelines for the transport of the critically ill adult. Intensive Care Society. www.ics.ac.uk.
- International Liaison Committee on Resuscitation (ILCOR)


 (2005): International Consensus on Cardiopulmonary

Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Resuscitation; 67: 157-341.

Inufusa A, An HS, Lim TH and Hasegawa T (1996): Anatomic changes of the spinal canal and intervertebral foramen associated with flexion-extension movement. Spine; 21(21):



- **James L. and Larson Jr (2007):** Injuries to the spine, emergency medicine: A comprehensive study guide, 6th ed.; 272.
- José EP, Luis MC and Antonio MC (2009): Methylprednisolone fails to improve functional and histological outcome following spinal cord injury in rats: Experimental Neurology; 220: 71-81.
- Joshi S, Young WL and Ornstein E (2001): Cerebral and spinal cord blood flow. In: Cottrell JE, Smith DS, eds. Anaesthesia and neurosurgery, 4th ed. St. Louis, MO: Mosby; 19-68.
- **Julio CF and Michael GF (2009):** Cardiovascular complications after acute spinal cord injury. Pathophysiology, Diagnosis and Management; 25(5): E13.




- **Kalinowski CP and Kirsch JR (2004):** Strategies for prophylaxis and treatment for aspiration. Best Pract Res Clin Anaesthesiol; 18(4): 719-737.
- **Karen JH and Richard EE (2000):** Textbook of Neuroanaesthesia and Critical Care; 17: 241-252.

Karim Brohi (1999): Spinal stabilization and management of trauma. Org; 7: 14.

- **Karlsson AK (1999):** Autonomic dysreflexia. Spinal Cord; 37(6): 383-91.
- Kersten JR, Pagel PS and Warltier DC (2004): Cardiovascular pharmacology of positive inotropic drugs. In: Vers AS, Maze M, eds. Anaesthetic pharmacology: Physiologic principles and clinical practice. A companion to anaesthesia. Philadelphia, PA: Churchill Livingstone: 655-670.
- Killian JT, Mayberry S and Wilkinson L (1999): Current concepts in adolescent idiopathic. Pediatr Ann; 28(12): 755-61.
- Kim YT, Caldwell JM and Bellamkonda RV (2009):

 Nanoparticle-mediated local delivery of methylprednisolone after spinal cord injury. Biomaterials; 30: 2582-2590.
- **Kiran S,Gombar S,Chhabra BK and Gombar (1997):** Another hazard of the prone position. Anaesth. Analg.; 85: 949.
- Kirshblum SC, Groah SL and McKinaley WO (2002): Spinal cord injury medicine: Etiology, classification, and acute medical management. Arch Phys Med Rehabil; 83(Suppl 1): S50-8.
- **Kreimeier U and Messmer K (2002):** Small-volume resuscitation from experimental evidence to clinical routine: Advantages and disadvantages of hypertonic solutions. Acta Anaesthesiol Scand; 46: 625-38.
- **Kwan I, Bunn F, Roberts I (2001):** Spinal immobilization for trauma patients.

Kwon BK, Tetzlaff W and Grauer JN (2004): Pathophysiology and pharmacologic treatment of acute spinal cord injury. Spine J; 4(4): 451-464.



- Lali HS, Sehkon MB and Fehlings MG (2001): Epidemiology, demographics, and pathophysiology of acute spinal cord injury. Spine; 26: S2-S12.
- Lee JM, Yan P, Xiao Q, Chen S, Lee KY, and Xu J (2008): Methylprednisolone protects oligodendrocytes but not neurons after spinal cord injury. J. Neurosci.; 28: 3141-3149.
- Li S and Stys PK (2000): Mechanisms of ionotropic glutamate receptor- mediated excitotoxicity in isolated spinal cord white matter. J Neurosci 20: 1190-1198.
- **Linsenmeyer TA and Culkin D (1999):** APS recommendations for the urological evaluation of patients with spinal cord injury. J Spinal Cord Med; 22(2): 139-142.
- Liu L and Yenari MA (2007): Therapeutic hypothermia: Neuroprotective mechanisms. Front Biosci; 12: 816-25.
- Lyeette CA, Doberstein C, Rodts GE and MeBride DQ (2003):

 Neurosurgical critical care: Current. Critical Care Diagnosis and Treatment; 31: 741-747.




MacDonald DB and Deletis V (2007): Safety issues during surgical monitoring. In: Nuwer M, editor. Monitoring neural function during surgery. Handbook of Clinical Neuophysiology, 3rd ed. Elsevier.


- Maehama T, Izena H and Kanazawa K (2000): Management of autonomic hyperreflexia with magnesium sulfate during labor in a woman with spinal cord injury. Am J Obstet Gyne: 183: 492-3.
- Matta, B.F.; Menon, D.K. and Turner, J.M. (2008): The anaesthetic management of spinal injuries. Textbook of Neuroanaesthesia and Critical Care; 17: 241-252.
- Maxwell RA, Chavarria Aguilar M and Cockerham WT (2002):
 Routine prophylactic vena cava filtration is not indicated after acute spinal cord injury. J Trauma; 52(5): 902-906.
- McDonald JW and Sadowsky C (2002): Spinal cord injury. Lancet; 359(9304): 417-25.
- McKinley WO, Gittler MS and Kirshblum SC (2002): Spinal cord injury medicine: II. Medical complication after SCI: Identification and management. Arch Phys Med Rehabil; 83(Suppl 1): S58-98.
- McSwain NE, Salomone JP and Fraunl SB (2003): Basic and advanced prehospital trauma life support, 5th ed. St. Louis, MO: Mosby.
- Michael Bernhard, Andre Gries, Paul Kremer and Bernd WB (2005): Spinal Cord Injury (SCI): Prehospital management. Resuscitation; 66: 127-139.
- Michael GF and Richard GP (2005): The role and timing of early decompression for cervical spinal cord injury: Update with a

review of recent clinical evidence: Injury. Int. J. Care Injured; 36: 13-26.

- Michelle Denton and Justin McKinlay (2009): Cervical cord injury and critical care. Cont Edu Anaesth Crit Care and Pain; 9(3): 82-86.
- **Miller SM (2008):** Methylprednisolone in acute spinal cord injury: A tarnished standard. J. Neurosurg. Anaesthesiol.; 20: 140-142.
- Mitcho K and Yanko JR (1999): Acute care management of spinal cord injuries. Critical Care; 22(2): 60-79.
- **Moore T (2000):** Supporting respiration. In: Bassett C, Makin L (Eds). Caring for the seriously III patient. Arnold Publishers, London; 51-79.
- Mutlu GM, Mutlu EA and Factor P (2001): GI complications in patients receiving mechanical ventilation. Chest; 119(4): 1222-



- **Nockels RP (2001):** Non-operative management of acute spinal cord injury. Spine; 26(suppl 24): S31-S37.
- Norenberg MD, Smith J and Marcillo A (2004): The pathology of human spinal cord injury: Defining the problems. J. Neurotrauma; 21: 429-440.



O'Connor MI, Currier BL (1992): Metastatic disease of the spine.
Orthopaedies; 15(5): 611-620.


Oren GB and Douglas ME (2003): Autonomic hyperreflexia in a pregnant paraplegic patient. Regional Anaesthesia; 19: 415-417.



- Palmon SC, Kirsh JR, Depper JA and Tounge TJ (1996): The effect of the prone position on pulmonary mechanics is frame-dependent Anaesth Analg; 87: 1175-1180.
- Patel RV, DeLong W and Vresilovic EJ (2004): Evaluation and treatment of spinal injuries in the patient with polytrauma. Clinical Orthopaedics and Related Research; 422: 43-54.
- **Patterson H (2004):** Emergency department intubation of trauma patients with undiagnosed cervical spine injury. Emerg Med Journal; 21(3): 302-305.
- **Paul Licina and Adrian MN (2005):** Approach and considerations regarding the patient with spinal injury: Injury. Int. J. Care Injured; 36: SB2-SB12.
- Pelosi P, Croci M, Calappi E and Cerisara M (1995): The prone positioning during general anesthesia minimally affects respiratory mechanics while improving functional residual capacity and increasing oxygen tension. Anaesth Analg; 80: 955-960.
- **Penny L, Kimball Jones and Randall MS (1999):** The use of remifentanyl infusion to allow intraoperative awakening for international Fracturing of the anterior cervical spine. Anaesth. Analge.; 89: 1095.

Petrozza PH and Prough DS (2001): Postoperative and intensive care. In: Cottrell JE, Smith DS, eds. Anaesthesia and Neurosurgery. St. Louis, MO: Mosby: 623-662.

Phillis JW and Goshgarian HG (2001): Adenosine and neurotrauma): therapeutic perspectives. Neurol Res; 23: 183-189.



- Rabinowitz RS, Eck JC and Harper JR (2008): Urgent surgical decompression compared to methylprednisolone for the treatment of acute spinal cord injury: A randomized prospective study in beagle dogs. Spine; 33: 2260-2268.
- **Ralph JM (2003):** Spinal injury: Etiology, demographics, and outcomes. Fractures of Cervical, Thoracic and Lumbar Spine; 1: 1-8.
- Raw DA, Beattie JK and Hunter JM (2003): Anaesthesia for spinal surgery in adults. Br J Anaesth; 91: 886-904.
- Raw DA, Beattie JK and Hunter JM (2003): Anaesthesia for spinal surgery in adults. Br J Anaesth; 91: 886-904.
- **Richards CF (2004):** Initial management of the trauma patient. Crit Care Clin; 20(1): 1-11.
- **Robert KS and Stephen FD (2002):** Spinal cord transaction: Disease of the nervous system. Anaesthesia and Co-Existing Disease; 278-282.

Robert WT, Jane TH and Jo Anne LA (2009): Venous thromboembolism after spinal cord injury. Archives of Physical Medicine and Rehabilitation; 90: 232-245.

- **Rogers LF (2006):** Fractures and dislocations of the spine. Garfin's Spine Trauma? Jefferson's Series.
- Rose JC and Mayer SA (2004): Optimizing blood pressure in neurological emergencies. Neurocrit Care; 4: 287-299.
- Rosenblatt WH, PJ Wagner, A Ovassapian and ZN Kain (1998):

 Practice patterns in managing the difficult airway by anaesthesiologists. Anaesth. Analge.; 87: 153-157.
- **Rovlias A and Kotsou S (2000):** The influence of hyperglycemia on neurological outcome in patients with severe head injury. Neurosurgery; 46: 335-42.
- Roy F, Cucchiara and Susan Black John DM (1998): Vertebral column and spinal cord surgery. Clinical Neuroanaesthesia; 13: 403-448.



- Sara Palmer; Kay Harris Kriegsman and Jeffrey BP (2000): Spinal cord injury: A guide for living.
- Sayer FT, Kronvall E and Nilsson OG (2006): Methylprednisolone treatment in acute spinal cord injury: The myth challenged through a structured analysis of published literature. Spine J.; 6: 335-343.
- **Schwab ME (2002):** Repairing the injured spinal cord. Science; 295: 1029-1031.

Segal JL and Milne Brunnemann SR (1995): Gastric emptying is impaired in patients with spinal cord injury. American journal of Gastroenterology; 90: 466-470.

- Sidhu, VS, Whitchead EM, Ainsworth QP, Smith M and Calde I (1993): A technique of awake fiberoptic intubation. Anaesthesia; 48: 910-913.
- **Silver JR (2000):** Early autonomic dysreflexia. Spinal Cord; 38: 229-33.
- Skellett S, Tibby SM, Durward A and Murdoch IA (2002): Immobilization of the cervical spine in children. BMJ; 324(7337): 591-3.
- **Snell RS (1997):** Clinical neuroanatomy for medical students, 4th edn. Chapter 12, Lippincott-Raven, Philadelphia.
- **St John RE and Thomson PD (1999):** Noninvasive respiratory monitoring. Crit. Care Nurs. Clin North Am.; 4: 423-35.
- Stassen NA, Williams, Valerie A and Gestring ML (2006): Magnetic resonance imaging in combination with helical computed tomography provides a safe and efficient method of cervical spine clearance in the obtunded trauma patient. J Trauma; 60(1): 171-177.
- Stevens RD, Bhardwaj A, Kirsch JR and Mirski MA (2003): Critical care and perioperative management in traumatic spinal cord injury. Journal of Neurosurgical Anaesthesiology; 15(3): 215-229.
- **StoeIting RK and Dierdorf SF (1996):** Diseases of the central nervous system: Anaesthesia and coexisting disease, 3rd ed. New York. Churchill Livingstone; 226-230.

Takami T, Oudega M, Bethea JR and Wood PM (2002): Methylprednisolone and interleukin-10 reduce gray matter damage in the contused fisher rat thoracic spinal cord but do not improve functional outcome J Neurotrauma; 19: 653-666.

- **Taneichi H and Kaneda K (1997):** Risk factors and probability of vertebral body collapse in metastases of the thoracic and lumbar spine. Spine; 22(3): 239-245.
- **Tator CH (1996):** Management of associated spine injuries in head injured patients. Neurotrauma. Mcgraw Hill.
- **Teasell RW, Arnold JM, Krassioukov A and Delaney GA** (2000): Cardiovascular consequences of loss of supraspinal control of the sympathetic nervous system after spinal cord injury. Arch Phys Med Rehabil; 81(4): 506-16.
- **Tins BJ and Cassar Pullicino VN (2004):** Imaging of acute cervical spine injuries): review and outlook. Clin Rad; 59: 865-880.
- **Tommasino** C (2002): Fluids and the neurosurgical patient. Anaesthesiol Clin North America; 20: 329-346.



Vaccaro AR and Blam OG (2002): Adult spine trauma. In: Koval KJ, ed. Orthopaedic knowledge update 7. Rosemont, IL: American Academy of Orthopaedic Surgeons; 693-607.

Vaccarro A, Ahmad S, Rausching W and Garfin S (1998):

Anatomy and pathophysiology of spinal cord injury with

APSCI. Levine A, ed. Spine Trauma. Philadelphia: WB

Saunders; 75-86.

- Van Elstraete AC, JC Mami and Hmehdaoui (1998):

 Nasotracheal intubation in patients with immobilized cervical spine a comparison of tracheal tube cuff inflation and fiberoptic bronchoscopy. Anesth Analg; 87: 400-402.
- **Vedran Deletis and Francesco Sala (2008):** Intraoperative neurophysiological monitoring of the spinal cord during spinal cord and spine surgery: A review focus on the corticospinal tracts. Clinical Neurophysiology; 119: 248-264.
- Velmahos GC, Toutouzas K and Chan L (2003): Intubation after cervical spinal cord injury: To be done selectively or routinely? Am Surg; 69: 891-4.
- Vera-Portocarrero LP and Mills CD (2002): Rapid changes in expression of glutamate transporters after spinal cord injury. Brain Res; 927: 104-110.
- VK Grover, Tewari MK, Gupta SK and Kumar (2001):

 Anaesthetic and intensive care aspects of spinal cord injury.

 Neurol India; 49: 11-8.



Warren J, Fromm RE and Orr RA (2004): Guidelines for the inter-and intrahospital transport of critically ill patients. Critical Care Medicine; 32: 256-262.

Watt J (1998): Anaesthesia for chronic spinal cord lesions and multiple sclerosis. Anaesthesia; 53: 825-826.

White DC (1996): The maintenance of anaesthesia: Clinical Anaesthesia and intensive care aspects. Churchill Livingstone; 7: 173-188.



Ziser A, RJ Friedhoff and SH Rose (1996): Prone position visceral hypoperfusion and rhabdomyolysis. Anaesth Analg; 82: 412-415.