

REFERENCES

- **Albert N, Hail M, Li J, Young JB.(2004):** Equivalence of bioimpedance and thermodilution methods in measuring cardiac output in patients with advanced, decompensated chronic heart failure hospitalized in critical care. *J Am J Crit Care* 2004; 13:469-79.
- **Albrecht K, Nave H, Breitmeier D, et al.(2004):** Applied anatomy of the superior vena cava—the carina as a landmark to guide central venous catheter placement. *Br J Anaesth* ; 92:75-77.
- **Amar D, Melendez JA, Zhang H, et al.(2001):** Correlation of peripheral venous pressure and central venous pressure in surgical patients. *J Cardiothorac Vasc Anesth* ; 15:40-43.
- **Auler JO, Galas F, Hajjar L, et a.(2008)l:** Online Monitoring of Pulse Pressure Variation to Guide Fluid Therapy After Cardiac Surgery. *Anesth Analg*; 106:1201–6
- **Auler Jr JO, Galas F, Hajjar L, et al.(2008):** Online monitoring of pulse pressure variation to guide fluid therapy after cardiac surgery. *Anesth Analg* ; 106:1031-1033.
- **Bailey PL, Glance LG, Eaton MP, et al. (2007):** A survey of the use of ultrasound during central venous catheterization. *Anesth Analg* ; 104:491-497.
- **Bein B, Worthmann F, Tonner PH, Paris A, Steinfath M, Hedderich J, et al.(2004):** Comparison of esophageal Doppler, pulse contour analysis and real- time pulmonary artery thermodilution for continuous measurement of cardiac output. *J Cardiothorac Vasc Anaesth*; 18:185-9.
- **Berton C, Cholley B.(2002):** Equipment review: New techniques for cardiac output measurement-oesophageal Doppler, Fick principle

using carbon-dioxide and pulse contour analysis. Crit Care; 6:216-21.

- **Berwick DM, Calkins DR, McCannon CJ, Hackbarth AD.(2006):** The 100,000 Lives Campaign. Setting a goal and a deadline for improving health care quality. JAMA; 295(3):324-7.
- **Binanay C, Califf RM, Hasselblad V, et al.(2005):** Evaluation study of congestive heart failure and pulmonary artery catheterization effectiveness: The ESCAPE trial. JAMA ; 294:1625-1633.
- **Boldt J.(2002):**Clinical review: Hemodynamic monitoring in the intensive care unit. Crit Care; 6:52-9.
- **Botero M, Kirby D, Lobato EB, Staples ED, Gravenstein N.(2004):** Measurement of cardiac output before and after cardiopulmonary bypass: Comparison among aortic transit-time ultrasound, thermodilution and noninvasive partial CO₂ rebreathing. J Cardiothorac Vasc Anaesth; 18; 563-72.
- **Botero M, Labato EB.(2001):** Advances in noninvasive cardiac output monitoring: An update. J Cardiothorac Vasc Anaesth; 15:631-40.
- **Boulnois JL, Pechoux T.(2000):** Non invasive cardiac output monitoring by aortic blood flow measurement with Dynemo 3000. J Clin Monit Comput; 16:127-40.
- **Brooks AJ, Alfredson M, Pettigrew B, MorrisDL.(2005):** Ultrasound-guided insertion of subclavian venous access ports. Ann R Coll Surg Engl ; 87:25-27.
- **Caruso LJ, Gravenstein N, Layon AJ, et al.(2002):** A better landmark for positioning a central venous catheter. J Clin Monit Comput ; 17:331-334.
- **Chaney JC, Derdak S. (2002):** Minimally invasive hemodynamic monitoring for the intensivists. Current and emerging technology. Crit Care Med; 30:2338-2345.

- **Chew MS, Poelaert J.(2003):** Accuracy and repeatability of pediatric cardiac output measurement using Doppler: 20 year review of literature. *Intensive Care Med*; 29:1889-94.
- **Cholley BP, Payen D.(2005):** Noninvasive technique for measurements of cardiac output. *Curr Opin Crit Care*; 11:424-9.
- **Cholley BP, Singer M.(2003):** Esophagel Doppler: Noninvasive cardiac output monitor. *Echocardiography*; 20:763-9.
- **Cohen MG, Kelly RV, Kong DF, et al.(2005):** Pulmonary artery catheterization in acute coronary syndromes: Insights from the GUSTO IIb and GUSTO III trials. *Am J Med* ; 118:482-488.
- **Conway DH, Mayall R, Abdul-Latif MS, Gillingan S, Tackaberry C.(2002) :** Randomised controlled trial investigating the influence of intravenous fluid titration using esophageal Doppler monitoring during bowel surgery. *Anesthesia*; 57:845-9.
- **Dalen JE.(2001):** The pulmonary artery catheter—friend, foe, or accomplice? *JAMA* ; 286:348-350.
- **Dark PM, Singer M.(2004):** The validity of trans-esophageal Doppler ultrasonography as a measure of cardiac output in critically ill adults. *Intensive Care Med*; 30:2060-6.
- **De Waal EE, Kalkman CJ, Rex S, et al.(2007):** Validation of a new arterial pulse contour-based cardiac output device. *Crit Care Med*; 35:1904–1909
- **Dhingra VK, Fenwick JC, Walley KR, et al.(2002):** Lack of agreement between thermodilution and Fick cardiac output in critically ill patients. *Chest* ; 122:990-997.
- **Domino KB, Bowdle TA, Posner KL, Spitellie PH, Lee LA, Cheney FW.(2004):** Injuries and liability related to central vascular catheters. A Closed Claims analysis. *Anesthesiology*; 100(6):1411-8.

-
- **Donati A, Loggi S, Preiser JC, et al.(2007):** Goal-directed intraoperative therapy reduces morbidity and length of hospital stay in high-risk surgical patients. *Chest* ; 132:1817-1824.
 - **Durairaj L, Schmidt GA.(2008):** Fluid Therapy in Resuscitated Sepsis: Less is more. *Chest*; 133:252–263
 - **Ezri T, Szmuk P, Cohen Y, et al.(2001):** Carotid artery–internal jugular vein fistula: A complication of internal jugular vein catheterization. *J Cardiothorac Vasc Anesth* ; 15:231-232.
 - **Fakler U, Pauli Ch, Balling G, et al.(2007):** Cardiac Index monitoring by pulse contour analysis and thermodilution after pediatric cardiac surgery. *J Thorac Cardiovasc Surg* 2007; 133:224–228
 - **Fangio P, Mourgeon E, Romelaer A, et al.(2002):** Aortic injury and cardiac tamponade as a complication of subclavian venous catheterization. *Anesthesiology* ; 96:1520-1522.
 - **Felbinger TW, Reuter DA, Eltzschig HK, et al.(2002):** Comparison of pulmonary arterial thermodilution and arterial pulse contour analysis: Evaluation of a new algorithm. *J Clin Anesth*; 14:296–301
 - **Feller-Kopman D.(2007):** Ultrasound-guided internal jugular access: A proposed standardized approach and implications for training and practice. *Chest* ; 132:302-309.
 - **Friese RS, Shafi S, Gentilello LM.(2006):** Pulmonary artery catheter use is associated with reduced mortality in severely injured patients: A National Trauma Data Bank analysis of 53,312 patients. *Crit Care Med* ; 34:1597-1601.
 - **Gan TJ, Soppit A, Maroof M, el-Moalem H, Robertson KM, Moretti E, et al.(2002) :** Goal-directed intraoperative fluid administration reduces length of hospital stay after major surgery. *Anesthesiology*; 97:820-6.
-

-
- **Garcia-Rodriguez C, Pittman J, Cassell CH, et al.(2002):** Lithium dilution cardiac output measurement: A clinical assessment of central venous and peripheral venous indicator injection. Crit Care Med ; 30:2199-2204.
 - **Garland JS, Alex CP, Mueller CD, et al.(2001):** A randomized trial comparing povidone-iodine to a chlorhexidine gluconate–impregnated dressing for prevention of central venous catheter infections in neonates. Pediatrics ; 107:1431-1436.
 - **Gelman S.(2008):** Venous function and central venous pressure. A physiologic story. Anesthesiology; 108:735-48.
 - **Ghani MK, Boccalandro F, Denktas AE, Barasch E.(2003):** Right atrial thrombus formation associated with central venous catheters utilization in hemodialysis patients. Intensive Care Med ; 29:1829-1882.
 - **Girard TD, Schectman JM.(2005):** Ultrasound guidance during central venous catheterization: A survey of use by house staff physicians. J Crit Care ; 20:224-229.
 - **Godje O, Hoeke K, Goetz AE, Felbinger TW, Reuter DA, Reichart B, et al.(2002):** Reliability of a new algorithm for continuous cardiac output determination by pulse-contour analysis during hemodynamic instability. Crit Care Med; 30:52-8.
 - **Graham AS, Ozment C, Tegtmeyer K, et al. (2007):** Videos in clinical medicine. Central venous catheterization. N Engl J Med ; 356(21):e21.
 - **Halvorsen PS, Espinoza A, Lundblad R, et al.(2006):** Agreement between PiCCO pulsecontour analysis, pulmonic artery thermodilution and transthoracic thermodilution during off-pump coronary artery bypass surgery. Acta Anaesthesiol Scand; 50:1050 – 1057
-

- **Harvey S, Harrison DA, Singer M, et al.(2005):**Assessment of the clinical effectiveness of pulmonary artery catheters in management of patients in intensive care (PAC-Man): a randomised controlled trial. *Lancet*; 366:472-477
- **Harvey S, Stevens K, Harrison D, et al.(2006):** An evaluation of the clinical and cost-effectiveness of pulmonary artery catheters in patient management in intensive care: A systematic review and a randomised controlled trial. *Health Technol Assess* ; 10(29):iii-iv.ix-xi, 1-133
- **Heerdt PM, Pleimann BE, Hogue CW.(2001):** Flow dependency of error in thermodilution measurement of cardiac output during acute tricuspid regurgitation. *J Cardiothorac Vasc Anesth* ; 15:183- 186.
- **Hind D, Calvert N, McWilliams R, et al.(2003):** Ultrasonic locating devices for central venous cannulation: Meta-analysis. *BMJ* ; 327:361.
- **Huang G-S Wang H-J, Chen C-H, et a.(2002)l:** Pulmonary artery ruptures after attempted removal of a pulmonary artery catheter. *Anesth Analg* ; 95:299-301.
- **Jacka MJ, Cohen MM, To T, et a.(2002)l:** Pulmonary artery occlusion pressure estimation: How confident are anesthesiologists? *Crit Care Med* ; 30:1197-1203.
- **Jain M, Canham M, Upadhyay D, Corbridge T.(2003):** Variability in interventions with pulmonary artery catheter data. *Intensive Care Med* ; 29:2059-2062.
- **Jansen JR, Schreuder JJ, Mulier JP, Smith NT, Settes KK, Wesseomg KH(2001).** A comparison of cardiac output derived from the arterial pressure wave against thermodilution in cardiac surgery patients. *Br J Anaesth*;87:212-22

-
- **Laupland KB, Bands CJ.(2002):** Utility of esophageal Doppler as a minimally invasive hemodynamic monitor: A review. *Can J Anaesth*; 49:393-401.
 - **Levy I, Katz J, Salter E, et al.(2005):** Chlorhexidine-impregnated dressing for prevention of colonization of central venous catheters in infants and children: A randomized controlled study. *Pediatr Infect Dis J* ; 24:676-679.
 - **Levy RJ, Chivacci RM, Nicolson SC, Rome JJ, Lin RJ, Helfaer MA, et al.(2004):** An evaluation of a noninvasive cardiac output measurement using partial carbon dioxide rebreathing in children. *Anesth Analg*; 99:1642-7.
 - **Lieshout JJ, Wesseling KH.(2001):** Continuous cardiac output by pulse contour analysis? *Br J Anaesth* ; 86:467-469.
 - **Linton NWF, Linton RAF.(2001):** Estimation of changes in cardiac output from the arterial blood pressure waveform in the upper limb. *Br J Anaesth* ; 86:486-496.
 - **London MJ, Moritz TE, Henderson WG, et al.(2002):** Standard versus fiberoptic pulmonary artery catheterization for cardiac surgery in the Department of Veterans Affairs. *Anesthesiology* ; 96:860-870.
 - **Mahajan A, Shabanie A, Turner J, et al. (2003):** Pulse contour analysis for cardiac output monitoring in cardiac surgery for congenital heart disease. *Anesth Analg*; 97:1283–1288
 - **Manecke JRJ, Brown JC, Landau AA, et al.(2002):** An unusual case of pulmonary artery catheter malfunction. *Anesth Analg* ; 95:302-304.
 - **Manning TS, Shykoff BE, Izzo JL Jr.(2002):** Validity and reliability of diastolic pulse contour analysis (Windkessel model) in humans. *Hypertension*; 39:963-8.
 - **Mayer J, Bold J, Wolf MW.(2008):** Cardiac output derived from arterial pressure waveform analysis in patients undergoing cardiac
-

-
- surgery: Validity of a second generation device. *Anesth Analg*; 106: 867-872.
- **McGee DC, Gould MK.(2003):** Preventing complications of central venous catheterization. *N Engl J Med*; 348(12):1123-33.
 - **Merrer J, De Jonghe B, Golliot R, et al.(2001):** Complications of femoral and subclavian venous catheterization in critically ill patients. A randomized controlled trial. *JAMA* ; 286:700-707.
 - **Michard F, Schachtrupp A, Toens C.(2005):** Factors influencing the estimation of extravascular lung water by transpulmonary thermodilution in critically ill patients. *Crit Care Med* ; 33:1243-1247.
 - **Michard F. (2000):** Pulse contour cardiac analysis:Fairy tale or new reality. *Crit Care Med*;35:1791–1792
 - **Milling Jr TJ, Rose J, Briggs WM, et al.(2005):** Randomized, controlled clinical trial of point-of-care limited ultrasonography assistance of central venous cannulation: The Third Sonography Outcomes Assessment Program (SOAP-3). *Trial. Crit Care Med* ; 33:1764-1769.
 - **Nilsson LB, Eldrup N, Berthelsen PG.(2001):** Lack of agreement between thermodilution and carbon dioxide-rebreathing cardiac output. *Acta Anaesthesiol Scand*; 45:680-5.
 - **Osman D, Ridel C, Ray P, et al.(2007):** Cardiac filling pressures are not appropriate to predict hemodynamic response to volume challenge. *Crit Care Med*; 35:64–68
 - **Ostergaard M, Nielsen J, Rasmussen JP, et al.(2006) :** Cardiac output-pulse contour analysis vs. pulmonary artery thermodilution. *Acta Anaesthesiol Scand*; 50:1044–1049
 - **Parmley CL, Pousman RM.(2002):** Noninvasive cardiac output monitoring. *Curr Opin Anaesthesiol*; 15:675-80.
-

-
- **Parry G.(2004):** Trendelenburg position, head elevation and a midline position optimize right internal jugular vein diameter. Can J Anaesth ; 51:379-381.
 - **Pearse R, Dawson D, Fawcett J, et al. (2005):** Early goal-directed therapy after major surgery reduces complications and duration of hospital stay. A randomized controlled trial Crit Care 9: R687
 - **Pearse RM, Ikram K, Barry J. (2004):** Equipment review: An appraisal of the LiDCO plus method of measuring cardiac output. Crit Care 8 190-195
 - **Pirotte T, Veyckemans F.(2007):** Ultrasound-guided subclavian vein cannulation in infants and children: A novel approach. Br J Anaesth ; 98:509-514.
 - **Pittman J, Bar-Yosef S, SumPing J, et a.(2005)l:** Continuous cardiac output monitoring with pulse contour analysis: A comparison with lithium indicator dilution cardiac output measurement. Crit Care Med ; 33:2015-2021.
 - **Polanczyk CA, Rohde LE, Goldman L, et al.(2001):** Right heart catheterization and cardiac complications in patients undergoing noncardiac surgery: an observational study. JAMA; 286:309-314
 - **Rauch H, Muller M, Fleisher F, et al.(2002):** Pulse contour analysis versus thermodilution in cardiac surgery patients. Acta Anaesthesiol Scand; 46:424–429
 - **Reisner A, Shaltis PA, McCombie D, Asada HH.(2008):** Utility of the photoplethysmogram in circulatory monitoring. Anesthesiology ; 108:950-958.
 - **Rothschild JM.(2001):** Ultrasound guidance of central vein catheterization. Evidence Report/Technology Assessment No. 43. Making Health Care Safer. A Critical Analysis of Patient Safety Practices. Agency for Healthcare Research and Quality, , pp 245-253.
-

-
- **Rubinfeld GD, McNamara-Aslin E, Robinson L.(2007):** The pulmonary artery catheter, 1967-2007: Rest in peace?. JAMA ; 298:458-461.
 - **Sandham JD, Hull RD, Brant RF, et al.(2003):** A randomized, controlled trial of the use of pulmonary-artery catheters in high-risk surgical patients. N Engl J Med ; 348:5-14.
 - **Seo JH, Jung CW, Bahk JH.(2007):** Uppermost blood levels of the right and left atria in the supine position: Implication for measuring central venous pressure and pulmonary artery wedge Pressure. Anesthesiology ; 107:260-263.
 - **Seoudi HM, Perkal MF, Hanrahan A, Angood PB.(2003):** The esophageal Doppler monitor in mechanically ventilated surgical patients: Does it work? J Trauma; 55:720-5.
 - **Shah MR, Hasselblad V, Stevenson LW, et al.(2005):**Impact of the pulmonary artery catheter in critically ill patients. Meta-analysis of randomized clinical trials. JAMA;294(13):1664-70.
 - **Sloth E, Pedersen J, Olsen KH, Wanscher M, Hansen OK, Sorensen KE.(2001):** Transoesophageal echocardiographic monitoring during pediatric cardiac surgery: Obtainable information and feasibility in 532 children. Pediatr Anesth; 11:657-62.
 - **Squara P, Bennett D, Perret C.(2002):** Pulmonary artery catheter. Does the problem lie in the users? Chest ; 121:2009-2015.
 - **Stouffer G. (2008):** Arterial pressure. In: Stouffer G, ed. Cardiovascular Hemodynamics for the Clinician, Malden, MA: Blackwell Futura; :57-66.
 - **Su NY, Huang CJ, Tsai P, Hsu YW, Hung YC, Cheng CR.(2002):** Cardiac output measurement during cardiac surgery: Oesophageal Doppler versus pulmonary artery catheter. Acta Anaesthesiol Scan; 40:127-33.
-

- **Tibby SM, Hatherill M, Murdoch IA.(2000):** Use of transoesophageal Doppler ultrasonography in ventilated pediatric patients: Derivation of cardiac output. *Crit Care Med*; 28:2045-50.
- **Turner MA.(2003):** Doppler-based hemodynamic monitoring: A minimally invasive alternative. *AACN Clin Issues*; 14:220-31.
- **Van De Water JM, Miller TW:** Impedance cardiography: The next vital sign technology. *Chest*; 123:2028-33.
- **Watson MO, Sanderson P.(2007):** Designing for attention with sound: Challenges and extensions to ecological interface design. *Hum Factors* ; 49:331-346.
- **Wiesenack C, Prasser C, Keyl C, Rödig G.(2001):** Assessment of intrathoracic blood volume as an indicator of cardiac preload: Single transpulmonary thermodilution technique versus assessment of pressure preload parameters derived from a pulmonary artery catheter. *J Cardiothorac Vasc Anesth* ; 15:584-588.
- **Wigmore TJ, Smythe JF, Hacking MB, et al.(2007):** Effect of the implementation of NICE guidelines for ultrasound guidance on the complication rates associated with central venous catheter placement in patients presenting for routine surgery in a tertiary referral centre. *Br J Anaesth* ; 99:662-665.