




REFERENCES



REFERENCES

- Abbas, A.K.; Lichtman, A.H. and Pober, J.S. (1991) : Effector mechanisms of immune response. *Cell Mol. Immunol.*; 1: 229.
- Akerman, P.; cote, P.; Yang, S.; McClain, C.; Nelson, S.; Bagby, G. and Diehl, A.M. (1992) : Antibodies to tumor necrosis factor-alpha inhibit liver regeneration after partial hepatectomy. *Am. J. Physiol.*; 263: G579-585.
- Ambrus, J.L. Jr.; Chesky, L. and Chused, T. et al. (1991) : Intracellular signaling events associated with induction of proliferation of normal human B lymphocytes by two different antigenically related human B cell growth factors (high molecular weight B cell growth factor [HMW-BCGF] and the complement factor 3b). *J. Biol. Chem.*; 266: 3702-2708.
- Ambrus, J.L. Jr.; Pippin, J. and Joseph, A. et al. (1993) : Identification of a cDNA for human high-molecular-weight B-cell growth factor. *Proc. Natl. Acad. Sci., USA*; 90: 6330-6334.
- Ameglio, F.; Capobianchi, M.R. and Castilletti, C. et al. (1994) : Recombinant gp 120 induces IL-10 in resting peripheral blood mononuclear cells: Correlation with the induction of other cytokines. *Clin. Exp. Immunol.*; 95: 455.

References

- Asadullah, K.; Sterry, W. and Leupold, M. (1998) : IL-10 is a key cytokine in psoriasis. *J. Clin. Invest.*, Vol. 101, No. 4, 783-794.
- Asadullah, K.H.; Schmid, M.; Friedrich, F.; Rando, H.D.; Voik, W. Sterry and W.D. Döcke (1997) : Influence of monomethylfumarate on monocytic cytokine formation: Explanation for adverse and therapeutic effects in psoriasis? *Arch. Dermatol. Res.*; 289: 623-630.
- Aukrust, P.; Liabakk, N.B. and Muller, F. (1994) : Serum levels of TNF α and soluble TNF receptors in human immunodeficiency virus type 1 infection correlation to clinical immunologic, and virologic parameters. *J. Inf. Dis.*; 169: 420-424.
- Baker, C.J. and Barrett, F.F. (1973) : Transmission of group B streptococcus among parturient women and their neonates. *J. Pediatr.*; 83: 919-925.
- Baker, C.J. and Edwards, M.S. (1994) : Group B streptococcal infection. In: Remington, J.S.; Klein, O., eds. *Infectious Diseases of the Fetus and Newborn Infant*. Philadelphia: W.B. Saunders; 980-1208.
- Bale, J.F. (1994) : Meningitis and encephalitis. In: McKendall, R.R.; Stroop, W.G., eds. *Handbook of Neurovirology*. New York: Marcel Dekker; 141-158.

References

- Berger, J.R. and Levy, R.M. (1993) : The neurologic complications of human immunodeficiency virus infection. Med. Clin. North Am.; 77: 1-23.**
- Beutler, B. (editor) (1992) : Tumor necrosis factors: The molecules and their emerging role in medicine. Raven Press.**
- Beutler, B. and Van Huffer, C. (1994) : Unraveling function in the TNF ligand and receptor families. Science; 264: 667.**
- Beutler, B.A.; Krochin, N.; Milsark, I.W. and Cerami, A. (1986) : Control of cachectin (tumour necrosis factor) synthesis: Mechanisms of endotoxin resistance. Science; 232: 977.**
- Beutler, B.A.; Mahoney, J.; Pakala, P. and Cerami, A. (1985b): Purification of cachectin, a lipoprotein-lipase suppressing hormone secreted by endotoxin included RAW 2647 Cells. J. Exp. Med.; 161: 984.**
- Bogdan, C.; Paik, J.; Vodovotz, Y. and Nathan, C. (1992) : Contrasting mechanisms for suppression of macrophage cytokine release by transforming growth factor- β and interleukin-10. J. Biol. Chem.; 267: 23301 - 23308.**
- Bonadio, W.A. (1987) : Incidence of serious infections in afebrile neonates with history of fever. Pediatr. Inf. Dis. J.; 6: 911-914.**

References

- Bonadio, W.A. (1992) : The cerebrospinal fluid: Physiologic aspects and alterations associated with bacterial meningitis. *Pediatr. Inf. Dis. J.*; 11: 423-432.
- Borish, L.; Aarons, A. and Wenzel, S. (1996) : Interleukin-10 regulation in normal subjects and patients with asthma. *J. Allergy Clin. Immunol.*; 97 (6): 1288-9.
- Bot, F.J.; Dorssers, L. and Wagemaker, G. et al. (1988) : Stimulating spectrum of human recombinant multi-CSF (IL-3) on human marrow precursors: Importance of accessory cells. *Blood*; 71: 1609.
- Bree, A.; Schierman, F. and Timony, G. et al. (1991) : Pharmacokinetics and thrombopoietic effects of recombinant human interleukin-11 (rhIL-11) in non-human primates and rodents. *Blood*; 78 (Suppl.): 132a.
- Brown, Z.A.; Benedetti, J. and Ashley, R. et al. (1991) : Neonatal herpes simplex virus infection in relation to asymptomatic maternal infection at the time of labor. *N. Engl. J. Med.*; 324: 1247-1252.
- Burdach, S.E. and Levitt, L.J. (1987) : Receptor specific inhibition of bone marrow erythropoiesis by recombinant DNA-derived IL-2. *Blood*; 69: 1368.

References

- Burke, J.M.; Roberts, C.W.; Hunter, C.A.; Murray, M. and Alexander, J. (1994) : Temporal differences in the expression of mRNA for IL-10 and IFN- γ in the brains and spleens of C57BL/10 mice infected with *Toxoplasma gondii*. *Parasite Immunol.*; 16: 305-314.
- Butt, A.M. and H.G. Jenkins (1994) : Morphological changes in oligodendrocytes in the intact mouse optic nerve following intravitreal injection of tumor necrosis factor. *J. Neuroimmunol.*; 51: 27-33.
- Campbell, A.G.M. and McIntosh, N. (1998) (eds): Forfar and Arneil's Textbook of Pediatrics (5th edn.). Churchill Livingstone, New York, Edinburgh, London, Madrid, Melbourne, San Francisco, and Tokyo; 1288-1292.
- Cannella, B. and C.S. Raine (1995) : The adhesion molecule and cytokine profile of multiple sclerosis lesions. *Ann. Neurol.*; 37: 424-435.
- Carter, A.; Huhad, N.; Droxler, I. and Rowe, J.M. (1996) : Expression and role in growth regulation of tumour necrosis factor receptors P55 and P75 in acute myeloblastic leukaemia cells. *Br. J. Haematol.*; 92: 116.
- Cassatella, M.A.; Meda, L. and Bonora, S. (1993) : Interleukin-10 inhibits the release of proinflammatory cytokines from human polymorphonuclear leukocytes. Evidence for and autocrine role of tumor necrosis factor and IL-1 β in mediating the production of IL-8 triggered by lipopolysaccharide. *J. Exp. Med.*; 178: 2207-11.

References

- Cassatella, M.A.; Meda, L.; Gasperini, S.; Calzetti, F. and Bonora, S. (1994) :** Interleukin 10 (IL-10) upregulates IL-1 receptor antagonist production from lipopolysaccharide-stimulated human polymorphonuclear leukocytes by delaying mRNA degradation. *J. Exp. Med.*; 179: 1695-1699.
- Chees, Brough, M. (1984) (Ed.) :** Medical Laboratory Manual for Tropical Countries. Vol. 11: Microbiology. El BS, P. 160-165.
- Cohen, M.C. and Cohen, S. (1996) :** Cytokine function: A study in biologic diversity. *Am. J. Clin. Pathol.*; 105: 589-98.
- Committee on Infectious Diseases American Academy of Pediatrics (1991) :** Haemophilus influenzae type b conjugate vaccines: Recommendations for immunization of infants and children 2 months of age and older. *Pediatrics*; 88: 169-172.
- Connolly, K.J. and Hammer, S.M. (1990) :** The acute aseptic meningitis syndrome. *Infect. Dis. Clin. North Am.*; 4: 599-622.
- Cruikshank, W.W.; Long, A. and Tarpy, R.E. et al. (1995) :** Early identification of interleukin-16 (lymphocyte chemoattractant factor) and macrophage inflammatory protein 1 alpha (MIP-1 α) in bronchoalveolar lavage fluid of antigen-challenged asthmatics. *Am. J. Resp. Cell Mol. Biol.*; 13: 738-747.

References

- Cuevas, L.E. and Hart, C.A. (1993) : Chemoprophylaxis of bacterial meningitis. *J. Antimicrob. Chemother.*; 31: (Suppl. B): 79-91.
- D'Andrea, A.; M. Aste-Amezaga; N.M. Valiante; X. Ma; M. Kubin; and G. Trinchieri (1993) : Interleukin 10 (IL-10) inhibits human lymphocyte interferon γ -production by suppressing natural killer cell stimulatory factor/IL-12 synthesis in accessory cells. *J. Exp. Med.*; 178: 1041-1048.
- D'Andrea, A.; Rengaraju, M. and Valiante, N.M. et al. (1992) : Production of natural killer cell stimulatory factor (NKSF/IL-12) by peripheral blood mononuclear cells. *J. Exp. Med.*; 176: 1387-1398.
- D'Souza, S.; K. Alinauskas; E. McCrea; C. Goodyer and J.P. Antel (1995) : Differential susceptibility of human CNS-derived cell populations to TNF-dependent and independent-immune-mediated injury. *J. Neurosci.*; 15: 7293-7300.
- Dacie S.J. and Lewis, S.M. (1984) (eds): *Practical haematology*. Sixth edition, Churchill Livingstone, Edinburgh, London, Melbourne and New York, p. 38 - 50.
- Dawson, M.M. (1991) : Tumour necrosis factor. In: *Lymphokines and interleukins*. CRC Pres Inc. Boca Raton. Florida 1st edn., Ch. 10, pp. 210-225.
- De Serio, A.; Graziani, R.; Laufer, R.; Ciliberto, G. and Paonessa, G. (1995) : In vitro binding of ciliary neurotrophic factor to its receptors: Evidence for the formation of an IL-6-type hexameric complex. *J. Mol. Biol.*; 254: 795-800.

References

- de Waal Malefyt, R.; Abrams, J. and Bennett, B. (1991) : Interleukin-10 inhibits cytokine synthesis by human monocytes: An autoregulatory role of IL-10 produced by monocytes. *J. Exp. Med.*; 174: 1209-20.
- DeBries, J.E. and Zurawski, G. (1995) : Immunoregulatory properties of IL-13: Its potential role in atopic disease. *Int. Arch. Allergy Immunol.*; 106: 175-179.
- Defrance, T.; Vanbervliet, B.; Briere, F.; Duran, I.; Rousset, F. and Banchereau, J. (1992) : Interleukin-10 and transforming growth factor beta cooperate to induce anti-CD40-activated naive human B cells to secrete immunoglobulin A. *J. Exp. Med.*; 175: 671-682.
- Del Prete, G.; De Carli, M.; Almerigogna, F.; Giudizi, M.G.; Biagiotti, R. and Romagnani, S. (1993) : Human IL-10 is produced by both type 1 helper (Th1) and type 2 helper (Th2) T cell clones and inhibits their antigen-specific proliferation and cytokine production. *J. Immunol.*; 150: 353-360.
- Detmers, P.A.; Los, K. and Cohn, Z.A. (1990) : Neutrophil-activating protein: IL-8 stimulates the binding activity of

References

- the leucocyte adhesion receptor CD11, CD18 on human neutrophils. *J. Exp. Med.*; 171: 1155.
- Dillon, H.C.; Gray, E.; Pass, M.A. and Gray, B.M. (1982) : Anorectal and vaginal carriage of group B streptococcus in pregnancy. *J. Inf. Dis.*; 145: 794-799.
- Dinarello, C. (1991) : Interleukin-1 and interleukin-1 antagonism. *Blood*; 77: 1627.
- Dinarello, C.A. (1984) : Interleukin-1. *Rev. Infect. Dis.*; 6: 51-95.
- Dougherty, J.M. and Roth, R.M. (1986) : Cerebral spina fluid. *Emerg. Med. Clin. North. Am.*; 4: 281-297.
- Durack, D.T. and Spanos, A. (1982) : End-of-treatment spinal tap in bacterial meningitis. Is it worthwhile? *JAMA*; 248: 75-78.
- Durand, M.L.; Calderwood, S.B.; Weber, D.J. and Miller, S.I. et al. (1993) : Acute bacterial meningitis in adults: A review of 493 episodes. *N. Engl. J. MEd.*; 328: 21-28.

References

- Durandy, A.; Schiff, C. and Bonnefoy, J.Y. et al. (1993) : Induction by anti-CD40 antibody or soluble CD40 ligand and cytokines of IgG, IgA syndrome. *Eur. J. Immunol.*; 23: 2294.
- Emilie, D.; Touitou, R.; Raphael, M. et al. (1992) : In vivo production of IL-10 by malignant cells in AIDS lymphomas. *Eur. J. Immunol.*; 22: 2937.
- Enk, A.H.; Angeloni, V.L.; Udey, M.C. and Katz, S.I. (1993) : Inhibition of Langerhans cell antigen-presenting function by IL-10: A role for IL-10 in induction of tolerance. *J. Immunol.*; 151: 2390-8.
- Enk, C.D.; D. Sredni; A. Blauvelt; and S.I. Katz (1995) : Induction of IL-10 gene expression in human keratinocytes by UVB exposure in vivo and in vitro. *J. Immunol.*; 154: 4851-4856.
- Faix, R.G. (1992) : Invasive neonatal meningitis: Comparison of albicans and parapsilosis infection. *Pediatr. Inf. Dis. J.*; 11: 88-93.
- Farrar, W.L.; Vinovour, M. and Hill, J.M. (1989) : In situ hybridization histochemistry localization of IL-3 mRNA in mouse brain. *Blood*; 73: 137.
- Feigin, R.D.; McCracken, G.H. and Klein, J.O. (1992) : Diagnosis and management of meningitis. *Pediatr. Inf. Dis. J.*; 11: 785-814.

References

- Feigin, R.D.; McCracken, G.H. and Klein, J.O. (1992) : Diagnosis and management of meningitis. *Pediatr. Infect. Dis. J.*; 11: 785-814.
- Feingold, K.; Soued, M. and Grunfeld, C. (1988) : Tumor necrosis factor stimulates DNA synthesis in the liver of intact rats. *Biochem. Biophys. Res. Commun.*, 153: 576-582.
- Feldman, M. (1996) : Cytokines. *Immunol. Today*, June (3) 11.8: 11.10.
- Fiorentino, D.; Zlotnik, A. and Viera, P. et al. (1991) : IL-10 acts on the antigen presenting cells to inhibit cytokine production by Th1 cells. *J. Immunol.*; 146: 3444.
- Fishman, R.A. (1992) : Composition of the cerebrospinal fluid. In: Fishman, R.A., ed. *Cerebrospinal Fluid in Diseases of the Nervous System*. Philadelphia: W.B. Saunders Company; 183-252.
- Ford, R.; Tamayo, A. and Martin, B. et al. (1995) : Identification of B-cell growth factors (interleukin-14; high molecular weight-B-cell growth factors) in effusion fluids from patients with aggressive B-cell lymphomas. *Blood*; 86: 283-293.
- Fortunato, S.J.; Menon, R. and Lombardi, S.J. (1997) : Interleukin-10 and transforming growth factor- β inhibit amniochorion tumor necrosis factor alpha production by contrasting mechanisms of action: Therapeutic

References

- implications in prematurity. *Am. J. Obstet. Gynecol.*; 177 (4): 803-9.
- Frei, K.; Nadal, D.; Pfister, H.W. and Fontana, A. (1993) : *Listeria meningitis: Identification of a cerebrospinal fluid inhibitor of macrophage listericidal function as interleukin 10.* *J. Exp. Med.*; 178: 1255-1261.
- Freichman, R.B.; Anderson, R.E. and Entire, S.M. (1987) : *Clin. Chem*; 26: 1D.
- Friede, R.L. (1989) : *Purulent leptomeningitis in newborns and infants.* In: Friede, R.L., ed. *Developmental Neuropathology.* Berlin: Springer-Verlag; 169-179, 180-181.
- Fukushima, K.; Ishiguro, A. and Shimbo, T. (1995) : *Transient elevation of granulocyte colony-stimulating factor levels in cerebrospinal fluid at the initial stage of aseptic meningitis in children.* *Pediatr. Res.*; 37: 160-164.
- Fukushima, K.; Ishiguro, A.; Nakamura, T.; Suzuki, Y.; Nagayama, S.; Abe, M.; Umezawa, T.; Nakahata, T.; Komiyama, A.; Shimbo, T. (1993) : *Elevated levels of interleukin-6 in the cerebrospinal fluid in childhood aseptic meningitis. (in Japanese with English Summary).* *Jpn. J. Inflam.*; 13: 263-268.
- Gamero, A.M.; Ussery, D. and Reintgen, D.S. et al. (1995) : *Interleukin-15 induction of lymphokine-activated killer cell function against autologous tumor cells in*

References

- melanoma patient lymphocytes by a CD18-dependent, perforin-related mechanism. *Cancer Res.*; 55: 4988-4994.
- Gazzinelli, R.T.; Hieny, S. and Wynn, T.A. et al. (1993) : Interleukin-12 is required for the T-lymphocyte independent induction of interferon- γ by an intracellular parasite and induces resistance in T-deficiency hosts. *Proc. Natl. Acad. Sci., USA*; 90: 6115-6119.
- Gerard, C.; Bruyns, C. and Marchant, A. (1993) : Interleukin-10 reduces the release of tumor necrosis factor and prevents lethality in experimental endotoxemia. *J. Exp. Med.*; 177: 547-50.
- Gilhar, A.; N. Grossman, S.; Kahanovicz; H. Reuveni; S. cohen; and A. Eitan (1996) : Antiproliferative effect of pentoxifylline on psoriatic and normal epidermis. *Acta. Derm. Venereol. (Stockh.)*; 76: 437-441.
- Gilmaker, M.; Kragstbjerg, P.; Forsgren, M. and Olcen, P. (1993) : Tumor necrosis factor-alpha (TNF- α) in cerebrospinal fluid from patients with meningitis of different etiologies: High levels of TNF α indicate bacterial meningitis. *J. Infect. Dis.*; 167: 882-889.
- Giri, J.G.; Anderson, D.M. and Kumaki, S. et al. (1995) : IL-15, a novel T cell growth factor that shares activities and receptor components with IL-2. *J. Leuk. Biol.*; 57: 763-766.

References

- Goh, C.R. (1990) : Tumour necrosis factor in clinical practice. *Am. Acad. Med. Singapore*; 19: 235.
- Gomez, R.; Ghezi, F.; Romero, R.; Munoz, H.; Tolosa, J.E. and Rojas, I. (1995) : Premature labor and intra-amniotic infection: Clinical aspects and role of cytokines in diagnosis and pathophysiology. *Clin. Perinatol.*; 22: 281-342.
- Gotoff, S.D. and Kliegman, R.M. (1992) : The fetus and the neonatal infant. In: Berhman, R.E.; Kleigman, R.M.; Nelson, W.E.; Vaughan, V.C., eds. *Nelson Textbook of Pediatrics*. Philadelphia: W.B. Saunders; 509-512.
- Grabstein, K.H.; Eisenman, J. and Shanebeck, K. et al. (1994) : Cloning of a T-cell growth factor that interacts with the beta chain of the interleukin-2 receptor. *Science*; 264: 965-968.
- Greenlee, J.E. (1990) : Approach to diagnosis of meningitis: Cerebrospinal fluid evaluation. *Infect. Dis. Clin. North Am.*; 4: 583-598.
- Grove, R.I. et al. (1993) : Oncostatin M is a mitogen for rabbit vascular smooth muscle cells. *Proc. Natl. Acad. Sci., USA*; 90: 823.
- Hardonk, M.J.; Dijkhuis, F.W.; Hulstaert, C.; Koudstaal, J. (1992) : Heterogeneity of rat liver and spleen macrophages in gadolinium chloride-induced elimination and repopulation. *J. Leukoc. Biol.*; 52: 296-302.

References

- Hayashi, T.; Ichiyama, T. and Fucukaura, S. (1997) : Levels of transforming growth factor β 1, tumor necrosis factor alpha and IL-6 in cerebrospinal fluid: Association with clinical outcome for children with bacterial meningitis. *Clinical Infect. Dis.*; 25: 11-12.**
- Henny, C.S. (1989) : Interleukin-7: Effects on early events in lymphopoiesis. *Immunol. Today*; 10: 170.**
- Herbst, H.; Foss, H.D. and Samol, J. et al. (1996) : Frequent expression of IL-10 by EBV harboring tumor cells of HD. *Blood*; 87 (7): April.**
- Ho, A.S.Y.; Liu, Y.; Khan, T.A.; Hsu, D.H.; Bazan, J.F. and Moore, K.W. (1993) : A receptor for interleukin-10 is related to interferon receptors. *Proc. Natl. Acad. Sci., USA*; 90: 11267-11271.**
- Houssiau, F.A.; Schandené, L. and Stevens, M. et al. (1995) : A cascade of cytokines is responsible for IL-9 expression in human T cells. *J. Immunol.*; 154: 2624-2630.**
- Ishiguro, A.; Suzuki, Y. and Inaba, Y. (1996) : Production of IL-10 in the cerebrospinal fluid in aseptic meningitis of children. *Pediatr. Res.*; 40 (4): 610-4.**
- Jakubowski, A.A. and Golde, D.W. (1996) : Therapeutic use of cytokines. In: William's, J.; Beutler, E.; Erslev, A.; Lichtman, M.A. (eds.). *Hematology*. McGraw-Hill Book Company, P. 155.**

References

- Johnson, R.A. and White, M. (1992) :** Syphilis in the 1990s: Cutaneous and neurologic manifestations. *Semin. Neurol.*; 12: 287-298.
- Kernig, V.M. (1882) :** Ueber ein Krankheitssymptom der acuten Meningitis. *St. Petersburg Med. Wochenschr*; 7: 398.
- Kessler, S.L. and Dajani, A.S. (1990) :** *Listeria meningitis in infants and children. Pediatr. Inf. Dis. J.*; 9: 61-63.
- Kimberly, J.V.Z.; fischer, E. and Hawes, A.S. et al. (1992) :** Effect of intravenous IL-8 administration in non-human primate. *J. Immunol.*; 148: 1746.
- Klein, J.O. (1994) :** Neonatal sepsis. *Semin. Pediatr. Inf. Dis.*; 5: 3-14.
- Klein, J.O. and Marcy, S.M. (1994) :** Bacterial sepsis and meningitis. In: Remington, J.S.; Klein, J.O. eds. *Infectious Diseases of the Fetus and Newborn Infant. Philadelphia: W.B. Saunders*; 835-890.
- Kornelisse, R.F.; Savelkoul, H.F.J. and Mulder, P.H.G. (1996) :** Interleukin-10 and soluble tumor necrosis factor receptors in cerebrospinal fluid of children with bacterial meningitis. *J. of Infect. Dis.*; 173: 1498-502.
- Kurzrock, R.; Redman, J. and Cabanillas, F. et al. (1993) :** Serum IL-6 levels are elevated in lymphoma patients and correlate with survival in advanced Hb and with B symptoms. *Cancer Res.*; 54: 782.

References

- Kusmo, J.W. (1984) : Basic statistics for health science. Mayfield Publishing Co., California.**
- Laberge, S.; Cruikshank, W.W. and Beer, D.J. et al. (1996) : Secretion of IL-16 (lymphocyte chemoattractant factor) from serotonin-stimulated CD8+ T cells in vitro. J. Immunol.; 156: 310-315.**
- Laberge, S.; Cruikshank, W.W. and Kornfeld, H. et al. (1995) : Histamine-induced secretion of lymphocyte chemoattractant factor from CD8+ T cells is independent of transcription and translation: Evidence for constitutive protein synthesis and storage. J. Immunol.; 155: 2902-2910.**
- Lambert, H.P. (1994) : Meningitis. J. Neurol. Neurosurg. Psych.; 57: 405-415.**
- LeBeau, M.M.; Lemons, R.S. and Espinosa, R. (1989) : IL-4 and IL-5 map to human chromosome 5 in a region encoding growth factors and receptors and are deleted in myeloid leukemias with a del (5q). Blood; 73: 647.**
- Lehmann, A.K.; Halstensen, A. and Rokke, O. (1995) : High levels of interleukin-10 in serum are associated with fatality in meningococcal disease. Infect. Immun.; 63: 2109-12.**
- Leist, M.; Gantner, F.; Jilg, S. and Wendel, A. (1995) : Activation of the 55 kDa TNF receptor is necessary and sufficient for TNF-induced liver failure, hepatocyte apoptosis, and nitrite release. J. Immunol.; 154: 1307-1316.**

References

- Leonardy, J.M. and Des Prez, R.M. (1990) : Tuberculous meningitis. *Infect. Dis. Clin. North Am.*; 4: 769-787.
- Leopard, K.A.; Reisbach, L.M. and Plauda, C. et al. (1992) : Gene expression for IL-1 receptor. *J. Exp. Med.*; 85: 121.
- Lindsten, T.; June, C.H. and Ledbetter, J.A. (1989) : Regulation of lymphokine messenger RNA stability as a surface-mediated T-cell activation pathway. *Science*, 244: 339.
- Liu, Y.; Wei, S.H.Y.; Ho, A.S.Y.; Malefy, R.W. and Moore, K.W. (1994) : Expression cloning and characterization of a human IL-10 receptor. *J. Immunol.*; 152: 1821-1829.
- Loetscher, H. (1992) : Purification and structural properties of two distinct TNF receptors on human cells. In: Aggarwal, B.B.; Vilcek, J., eds. *Tumor necrosis factor: Structure, function and mechanism of action*. New York: Marcel Dekker, pp. 161-188.
- Longermans, J.A.M.; Van Furth, A.M. and Van Fusth, R. (1995) : High levels of interleukin-10 and TNF α in cerebrospinal fluid during the onset of bacterial meningitis. *Clinical Infect. Dis.*; 21: 220-2.
- Louis, J-C; E. Magal; S. Takayama and S. Varon (1993) : CNTF protection of oligodendrocytes against natural and tumor necrosis factor-induced death. *Science (Wash. DC)*; 259: 689-692.

References

- Louvois, J.; Blackbourn, J.; Hurley, R. and Harvey, D. (1991) : Infantile meningitis in England and Wales: A two year study. Arch. Dis. Child.; 66: 630-607.
- Marchant, A.; Devire, J.; Byl, B.; De Groote, D.; Vincent, J-L and Goldman, M. (1994) : Interleukin-10 production during septicaemia. Lancet; 343: 707-708.
- Marsters, S.A.; Pennica, D.; Bach, E.; Schreiber, R.D. and Ashkenazi, A. (1995) : Interferon γ signals via a high-affinity multisubunit receptor complex that contains two types of polypeptide chain. Proc. Natl. Acad. Sci., USA; 92: 5401-5405.
- Masood, R.; Lunardi, Y. and Modgil, T. et al. (1994) : IL-10 inhibits HIV-1 replication. Biochem. Biophys. Res. Commun.; 202: 374.
- Masood, R.; Zhang, Y. and Bond, M.W. et al. (1995) : IL-10 is an autocrine growth factor for AIDS-related B-cell lymphoma. Blood; 85 (12), (June 15).
- Matsushima, K.; Baldwin, E. and Mukaida, N. et al. (1992) : Interleukin-8 and MCAF: Novel leukocyte recruitment and activating cytokines. In: Kishimoto, T., editor, Interleukins: Molecular Biology and Immunology. Chem. Immunol., Basel, Karger; 51: 236.

References

- McNiece, I.K.; Bertoncello, I. and Kriegler, A.B. et al. (1990) :**
Colony forming cells with high proliferative potential.
Int. J. Cell Cloning; 8: 146.
- Metcalf, D. (1989) :** The molecular control of cell division,
differentiation, commitment and maturation in
hemopoietic cells. Nature (Lond.): 339: 27.
- Metcalf, D. (1992) :** Leukemia inhibitory factor: A puzzling
polyfunctional regulator. Growth Factors; 7: 169.
- Michie, H.R.; Manogue, K.R. and Spriggs, D.R. (1988) :** Detection
of circulating TNF after endotoxin administration. N.
Engl. J. Med.; 318: 1481.
- Mizuno, T.; Sawada, M.; Marunouchi, T. and Suzumura, A. (1994)**
: Production of interleukin-10 by mouse glial cells in
culture. Biochem. Biophys. Res. Commun.; 205:
1907-1915.
- Modlin, J.F. (1986) :** Perinatal echovirus infections: Insights from a
literature review of 61 cases of serious infection and 16
reported nursery outbreaks. Rev. Inf. Dis.; 9: 918-926.
- Modlin, J.F. (1994) :** Neonatal enterovirus infections. Semin. Inf.
Dis.; 5: 70-77.

References

- Moore, K.W.; O'Garra, A.; de Waal Malefyt, R.; Vieira, P. and Mostmann, T.R. (1993) : Interleukin-10. *Annu. Rev. Immunol.*; 11: 165-190.
- Mosmann, T. (1996) : Cytokines and immune regulation. In *clinical immunology principles and practice*, Rich, R.R.; Fleisher, T.A. and Shearer, W.T. (eds.) Mosby St. Louis, Baltimore, Boston, Catsbad, Chicagon Naples, New York; pp. 217-229.
- Mosmann, T.R. (1994) : Properties and functions of interleukin-10. *Adv. Immunol.*; 56: 1-26.
- Mozes, T. (1991) : Sequential release of TNF, platelet activating factor and eicosanoides during endotoxin shock in anaesthetized pigs, protective effects of indomethacin. *Br. J. Pharmacol.*; 104: 691.
- Nagata, Y.; Tanaka, K. and Orita, K. (1994) : Endotoxin-induced liver injury after extended hepatectomy and the role of Kupffer cells in the rat. *Surg. Today*; 24: 441-448.
- Naldini, A.; Fleischmann, W.R. and Ballas, Z.K. et al. (1987) : IL-2 inhibits in vitro GM-CSF formation. *J. Immunol.*; 139: 1880.

References

- Nelson, J.D. (1995) : Antifungal therapy. In: Pine, J.W. Jr., ed.: *Pocketbook of Pediatric Antimicrobial Therapy*. Baltimore: Williams & Wilkins; 54-55.
- Nelson, J.S. (1990) : Pathology of the central nervous system. In: Kissone, J.M., ed. *Anderson's Pathology*. St. Louis: Mosby Company; 2154-2155.
- Nickoloff, B.J. (1991) : The cytokine network in psoriasis. *Arch. Dermatol.*; 127: 871-884.
- Nicord, L.P.; Habre, F.; Dayer, J.M. and Boehringer, N. (1995) : Interleukin-10 decreases tumor necrosis factor alpha and beta in alloreactions induced by human lung dendritic cells and macrophages. *Am. J. Respir. Cell Mol. Biol.*; 13: 83-90.
- Niemeyer, C.M.; Sieff, C.A.; Mathey-Prevot, B. et al. (1989) : Expression of human IL-3 is restricted to human lymphocytes and T cell tumor line. *Blood*; 73: 945.
- Ockenfels, H.M.; S.N. Wagner; C. Keim-Maas; R. Funk; G. Nussbaum and M. Goos (1996) : Lithium and psoriasis: Cytokine modulation of cultured lymphocytes and psoriatic keratinocytes by lithium. *Arch. Dermatol. Res.*; 288: 173-178.
- Oksanen, V.; Fyhrquist, F.; Somer, H. and Gronhagen-Riska, C. (1985) : Angiotensin converting enzyme in cerebrospinal fluid: A new assay. *Neurology*; 35: 1220-1223.

References

- Oppenheim, J.J.; Ruscetti, F.W. and Flatynek, C. (1994) :**
Cytokines in basic and clinical immunology (8th edn.),
Stites, D.P. and Terr, A.I. (eds.): Appleton and Lange,
Northwalk, Connecticut/San mateo, California; pp.
105-123.
- Overall, J.C. (1993) :** Is it bacterial or viral; laboratory
differentiation. *Pediatr. Rev.*; 14: 251-261.
- Paul, S. and Schendel, P. (1992) :** The cloning and biological
characterization of recombinant human interleukin-11.
Int. Cell Cloning; 10: 135.
- Paul, S.; Bennett, F. and Calvetti, J. (1990) :** Molecular cloning of a
cDNA encoding interleukin-11, a stromal cell derived
lymphopoietic and hematopoietic cytokine. *Proc. Natl.*
Acad. Sci., USA; 87: 7512.
- Paul, W.E. (1991) :** A proteolytic immunoregulatory lymphokine.
Blood; 77: 1859.
- Pawankar, R.U.; Okuda, M. and Hasegawa, S. et al. (1995) :**
Interleukin-13 expression in the nasal mucosa of
perennial allergic rhinitis. *Am. J. Resp. Crit. Care Med.*;
152: 2059-2067.
- Petit-Frere, C.; Ongas, B. and Mencia-Huerta, J. (1993) :** IL-9
potentiates the IL-4 induced IgE and IgG release from
murine B-lymphocytes. *Immunol.*; 79: 146.

References

- Pfister, H.W.; Fontana, A.; Tauber, M.G.; Tomasz, A. and Scheld, W.M. (1994) :** Mechanisms of brain injury in bacterial meningitis: Workshop summary. *Clin. Infect. Dis.*; 19: 463-479.
- Philip, A.G.S. (1982) :** Detection of neonatal sepsis of late onset. *JAMA*; 247: 489-492.
- Philip, A.G.S. and Hewitt, J.R. (1980) :** Early diagnosis of neonatal sepsis. *Pediatrics*; 65: 1036-1041.
- Plata-Salamán, C.R. (1991) :** Immunoregulators in the nervous system. *Neurosci. Biobehav. Rev.*; 15: 185-215.
- Platzer, C.; C. Meisel; K. Vogt; M. Platzer; and H.D. Volk (1995) :** Upregulation of monocytic IL-10 by tumor necrosis factor-alpha and cAMP elevating drugs. *Int. Immunol.*; 7: 517-523.
- Powell, K.R. and March, S.M. (1994) :** Laboratory aids for diagnosis of neonatal sepsis. In: Remington, J.S.; Klein, J.O. eds. *Infectious diseases of the fetus and newborn infant*. Philadelphia: W.B. Saunders; 1223-1239.
- Punnonen, J.; De Waal Malefyt, R.; van Vlasselaer, P.; Gauchat J-F; de Vries, J.E. (1993) :** IL-10 and viral IL-10 prevent IL-4-induced IgE synthesis by inhibiting the accessory cell function of monocytes. *J. Immunol.*; 151: 1280-9.
- Puren, A.J.; Fantuzzi, G. and Dinarello, C.A. (1998) :** Interleukin-18 (IFN γ -inducing factor induces IL-18 and

References

- IL-1 β via TNF α production from non-CD14+ human blood mononuclear cells. *J. Clin. Invest.*, Vol. 101, No. 3, 711-721.
- Quagliarello, V. and Scheld, W.M. (1992) : Bacterial meningitis: Pathogenesis, pathophysiology, and progress. *N. Engl. J. Med.*; 327: 864-872.
- Quesenberry, P.J. (1995) : Hemopoietic stem cells, progenitor cells and cytokines. In: William's J.; Beutler, E.; Erslev, A.J.; Lichtman, M.A. (eds.). *Hematology McGraw-Hill Book Company*; P. 211.
- Rai, R.; Yang, S.Q.; McClain, C.; Karp, C.L.; Klein, A.S. and Diehl, A.M. (1996) : Kupffer cell depletion by gadolinium chloride enhances liver regeneration after partial hepatectomy in rats. *Am. J. Physiol.*; 270: G909-G918.
- Raine, C.S. (1994) : The immunology of the multiple sclerosis lesions. *Ann. Neurol.*; 36: 561-572.
- Reichlin, S. (1993) : Neuroendocrine-immune interactions. *N. Engl. J. Med.*; 329: 1246-1253.
- Reickman, N.P.; M. Albrecht; B. Kitze; T. Weber; H. Tumani; A. Broocks; W. Luer; A. Helwig and S. Poser (1995) : Tumor necrosis factor-alpha messenger RNA expression in patients with relapsing-remitting multiple sclerosis is associated with disease activity. *Ann. Neurol.*; 37: 82-88.

References

- Renauld, J-C; Kermouni, A. and Vink, A. et al. (1995) :**
Interleukin-9 and its receptor: Involvement in mast cell
differentiation and T-cell oncogenesis. *J. Leuk. Biol.*;
57: 353-360.
- Rench, M.A. and Baker, C.J. (1993) :** Neonatal sepsis caused by a
new group B streptococcal serotype. *J. Pediatr.*; 122:
638-640.
- Robbins, J.B.; McCracken, G.H. Jr.; Gotshlich, E.C.; Orskov, F.
and Orskov, I. (1974) :** Escherichia coli K1 capsular
polysaccharide associated with neonatal meningitis. *N.
Engl. J. Med.*; 290: 1216-1220.
- Rodriguez, A.F.; Kaplan, S.L. and Mason, E.O. (1990) :**
Cerebrospinal fluid values in the very low birth weight
infant. *J. Pediatr.*; 116: 971-974.
- Roitt, I.; Brostoff, J. and Male, D. (1998) (Eds) :** Immunology (th
edn.). Mosby, London, Philadelphia, St. Louis, Sydney,
Tokyo; P. 121-137.
- Romero, R. and Mazor, M. (1988) :** Infection and preterm labor.
Clin. Obstet. Gynecol.; 31: 553-84.
- Roos, K.L. (1997) (editor):** Central nervous system infections.
Diseases and therap. Marcel Dekker Inc., New York, p.
99-139.

References

- Rousset, F.; Garcia, E.; Defrance, T.; Peronne, C.; Vezzio, N.; Hsu, D.H.; Kastelein, R.; Moore, K.W. and Banchereau, J. (1992) : Interleukin-10 is a potent growth and differentiation factor for activated human B lymphocytes. *Proc. Natl. Acad. Sci., USA*; 89: 1890-1893.
- Rubeiz, H. and Roos, R.P. (1992) : Viral meningitis and encephalitis. *Semin. Neurol.*; 12: 165-177.
- Saez-Llorens, X.O.; Ramilo, M.M. and Mertsola, M.J. (1990) : Molecular pathophysiology of bacterial meningitis: Current concepts and therapeutic implications. *J. Ped.*; 116: 671.
- Sanderson, C. (1992) : Interleukin-5, eosinophils, and disease. *Blood*; 79: 3101.
- Schaafsma, M.; Falkenburg, J. and Duinkerken, N. (1993) : Interleukin-9 stimulates the proliferation of enriched human erythroid progenitor cells: Additive effect with GM-CSF. *Ann. Hematol.*; 66: 45.
- Schlaak, J.F.; M. Buslau, W.; Jochum, E.; Hermann, M.; Girndt, H. Gallati; K.H. Meyer zum Büschenfelde and B. Fleischer (1994) : T cells involved in psoriasis vulgaris belong to the Th1 subset. *J. Invest. Dermatol.*; 102: 145-149.

References

- Schlesinger, Y.; Sawyer, M.H. and Storch, G.A. (1994) :
Enteroviral meningitis in infancy: Potential role for
polymerase chain reaction in patient management.
Pediatrics; 94: 157-162.
- Scissions, C.F. (1991) : Bacterial infections of the newborn. In:
Taeusch, H.W.; Ballard, R.A.; Avery, M.E., eds.
Schaffer and Avery's Diseases of the Newborn.
Philadelphia: W.B. Saunders; 350-369.
- Seigel, J.; Harper, M.E. and Wong-Staal, F. (1984) : Gene for
T-cell growth factor location on human chromosome 4q.
Science; 223: 175.
- Selmaj, K.W. and C.S. Raine (1988) : Tumor necrosis factor
mediates myelin and oligodendrocyte damage in vitro.
Ann. Neurol.; 23: 339-346.
- Shalaby, M.R.; Sundan, A.; Loetscher, H. and Espevik, T. (1990) :
Binding and regulation of cellular functions by
monoclonal antibodies against human TNF receptors. *J.*
Exp. Med.; 172: 1517.
- Sieff, C.A.; Neimyer, C.M. and Nathan, D.G. (1987) : Stimulation
of human hematopoietic colony formation by
recombinant gibbon multi-colony stimulating factor or
interleukin-3. *J. Clin. Invest.*; 80: 818.
- Smith, K.A. (1988) : IL-2 inception, impact and implications.
Science; 240: 1169.

References

- Smith, R.R. (1992) : Neuroradiology of intracranial infections. *Pediatr. Neurosurg.*; 8: 92-104.**
- Springer, T.A. (1990) : Adhesion receptors of the immune system. *Nature (Lond.)*; 346: 425-434.**
- Squire, E.; Favara, B. and Todd, T. (1979) : Diagnosis of neonatal bacterial infections: Hematologic and pathologic findings in fatal and non-fatal cases. *Pediatrics*; 64: 60-64.**
- Stern, B.J.; Krumholz, A.; Johns, C.; Scott, P. and Nissim, J. (1985) : Sarcoidosis and its neurological manifestations. *Arch. Neurol.*; 42: 909-917.**
- Stroop, W.G. (1994) : Viral pathogenesis. In: McKendall, R.R.; Stroop, W.G., eds. *Handbook of Neurovirology*. New York: Marcel Dekker; 27-54.**
- Sutherland, G.R.; Baker, E. and Callen, D.F. (1988) : IL-5 is at 5q31 and is detected in the Sq-syndrome. *Blood*; 71: 1150.**
- Svare, J.; Andersen, L.F.; Langhoff-Roos, J.; Madsen, H. and Bruun, B. (1991) : Maternal-fetal listeriosis: Two case reports. *Gyn. Obstet. Invest.*; 3: 179-181.**
- Taga, K.; Mostowski, H. and Tosato, G. (1993) : Human interleukin-10 can directly inhibit T-cell growth. *Blood*; 81: 2964.**

References

- Takanaski, S.; Nonaka, R.; Xing, Z.; O'Bryne, P.; Dolovich, J. and Kordana, M. (1994) : Interleukin-10 inhibits lipopolysaccharide-induced survival and cytokine production by human peripheral blood eosinophils. *J. Exp. Med.*; 180: 711-5.
- Tan, J.C.; Braun, S.; Rong, H.; DiGiacomo, R.; Dolphin, E.; Baldwin, S.; Narula, S.K.; Zavodny, P.J. and Chou, C.C. (1995) : Characterization of recombinant extracellular domain of human interleukin-10 receptor. *J. Biol. Chem.*; 270: 12906-12911.
- Tan, J.C.; Indelicato, S.R.; Narula, S.K.; Zavodny, P.J. and Chou, C.C. (1993) : Characterization of interleukin-10 receptors on human and mouse cells. *J. Biol. Chem.*; 268: 21053-21059.
- Tartaglia, L.A.; Goeddel, D.V.; Reynolds, C. and Palladino, P.M. (1993) : Stimulation of human T cell proliferation by specific activation of the 75KDa tumour necrosis factor receptors. *J. Immunol.*; 151: 4637.
- Tartaglia, L.A.; Waber, R.F.; Figari, I.S. and Goeddel, D. (1991) : The two different receptors for tumour necrosis factor mediate distinct cellular responses. *Proc. Natl. Acad. Sci., USA*; 88: 9292.
- Tracy, K.J. and Cerami, A. (1988) : Cachectin/Tumour necrosis factor and other cytokines in infectious disease. *Curr. Opin. Immunol.*; 1: 454.

References

- Traugott, U.; E.L. Reinberz; and C.S. Raine (1982) : Multiple sclerosis: Distribution of T cell subsets within active chronic lesions. *Science (Wash. DC)*; 219: 308-310.**
- Trentin, L.; Zambello, R.; Agostini, C.; Pizzolo, G. and Semenzato, G. (1993) : Expression and function role of tumour necrosis factor receptors on leudaemic cells from patients with type B chronic lymphoproliferative disorders. *Blood*; 81: 752.**
- Treseler, C.B. and Sugar, A.M. (1990) : Fungal meningitis. *Infect. Dis. Clin. North Am.*; 4: 789-808.**
- Trinchieri, G. (1995) : Interleukin-12: A proinflammatory cytokine with immunoregulatory functions that bridge innate resistance and antigen-specific adaptive immunity. *Ann. Rev. Immunol.*; 13: 251-276.**
- Tripp, C.S.; Gately, M.K. and Hakimi, J. et al. (1994) : Neutralization of IL-12 decreases resistance to *Listeria* in SCID and CB-17 mice. *J. Immunol.*; 152: 1883-1887.**
- Trujillo, M. and McCracken, G.H. (1997) : Neonatal meningitis in central nervous system infectious diseases and therapy. Edited by Karan L, Roos, Marcel Dekker, Inc., New York, p. 25 - 44.**
- Uren, E.C.; Johnson, P.D.; Montanoro, J. and Gilbert, G.L. (1993) : Herpes simplex virus encephalitis in pediatrics: Diagnosis by detection of antibodies and DNA in CSF. *Pediatr. Inf. Dis. J.*; 12: 1001-1006.**

References

- Wanidworanun, C.; Strober, W. (1993) : Predominant role of tumor necrosis factor-alpha in human monocyte IL-10 synthesis. *J. Immunol.*; 151: 6853-6861.
- Ward, L.D.; Howlett, G.J.; Discolo, G.; Yasukawa, K.; Hammacher, A.; Moritz, R.L. and Simpson, R.J. (1994) : High affinity interleukin-6 receptor is a hexameric complex consisting of two molecules each of interleukin-6, interleukin-6 receptor, and gp-130. *J. Biol. Chem.*; 269: 23286-23289.
- Wheat, J.J.; Batteiger, B.E. and Sathapatayavongs, B. (1990) : Histoplasma capsulatum infections of the central nervous system: A clinical review. *Medicine*; 69: 244-260.
- Whitley, J.D.; Arvin, A.M. and Prober, C.G. et al. (1991) : Predictors of morbidity and mortality in neonatal herpes simplex infection. *N. Engl. J. Med.*; 324: 450-454.
- Whitley, R.J. and Arvin, A.M. (1994) : The natural history, pathogenesis and treatment of neonatal herpes simplex virus infections. *Semin. Pediatr. Inf. Dis.*; 5: 56-63.
- Whitley, R.J. and Gnann, J.W. (1992) : Acyclovir: A decade later. *N. Engl. J. Med.*; 11: 782-789.
- Whitley, R.J. and Lakeman, F. (1995) : Herpes simplex virus infections of the central nervous system: Therapeutic and diagnostic considerations. *Clin. Infect. Dis.*; 20: 414-420.

References

- Wigglesworth, J.S. (1991) : Miscellaneous conditions affecting the fetus and neonate. In: Wigglesworth, J.S.; Singer, G.B., eds. Textbook of Fetal and Perinatal Pathology, Boston: Blackwell Scientific; 838-841.
- Wodner-Filipowicz, A.; Heusser, C. and Handoroni, C. (1989) : Production of the hemopoietic growth factor GM-CSF and IL-3 by mast cells in response to IgE receptor-mediated activation. Nature (Lond.), 339: 150.
- Wölfer, L.U.; S. Goerdts; K. Schröder; C.C. Zouboulis; and C.E. Orfanos (1996) : Interferon- α induced psoriasis. Hautarzt; 47: 124-128.
- Wynn, T.A.; Eltoum, I. and Oswald, I.P. et al. (1994) : Endogenous interleukin-12 (IL-12) regulates granuloma formation induced by eggs of *Schistosoma mansoni* and endogenous IL-12 both inhibits and prophylactically immunizes against egg pathology. J. Exp. Med.; 179: 1551-1561.
- Yang, Y.; Yin, T. and McKinley, D. et al. (1991) : The molecular cloning and characterization of human interleukin-11. J. Cell Biochem.; 15F: 42.
- Yao, Z.; Painter, S.L. and Fanslow, W.C. et al. (1995) : Human IL-17: A novel cytokine derived from T cells. J. Immunol.; 155: 5483-5486.

References

- Zdanov, A.; Menon, S. and Mookerjee, K.W. (1997) : Crystal structure of Epstein-Barr virus protein BCRF1, a homology of cellular interleukin-10. *J. Med. Biol.*; 268 (2): 460-7.
- Zhu, X.; Zellweger, R.; Zhu, X.; Ayala, A. and Chaudry, I.H. (1995) : Cytokine gene expression in splenic macrophages and Kupffer cells following haemorrhage. *Cytokine*; 7: 8-14.
- Zwahlen, A.; Nydegger, U.E.; Vaudaux, P.; Lambert, P.H. and Waldvogel, F.A. (1982) : Complement-mediated opsonic activity in normal and infected human cerebrospinal fluid: Early response during bacterial meningitis. *J. Infect. Dis.*; 145: 635-646.