
REFERENCES

- Abbadie C, Brown JL, Mantyh PW, et al. (1996):** Spinal cord substance P receptor immunoreactivity increases in both inflammatory and nerve injury models of persistent pain. *Neurosci*; 70(1): 201-209.
- Abdulatif M, El-Sanabary (2002):** Caudal neostigmine,bupivacaine and their combination for postoperative pain management after hypospadias surgery in children. *Anesth Analg* 95:1215-8
- Adewale L, Dearlove O, Wilson B, et al. (2000):** The caudal canal in children: a study using magnetic resonance imaging. *Pediatr Anesth*; 10(2): 137-41.
- Agarwal R, Gutlove DP, and Lockhart CH (1992):** Seizures occurring in pediatric patients receiving continuous infusion of bupivacaine. *Anesth Analg*; 75: 284.
- Agur AM, and Lee MJ (1991):** Sacrum. In: Agur AM, and Lee MJ (eds) Grant's Atlas of Anatomy, 9th edition. Williams and Wilkins, Baltimore, p. 219.
- Al-Chaer ED, Traub RJ (2002):** Biological basis of visceral pain: recent developments. *Pain* 96(3): 221-225.
- Anand KJS (2003):** Neurophysiological and Neurobiological correlates of supraspinal pain processing: measurement techniques. In: Finley GA, McGrath PJ (Eds). *Measurement of Pain in Infants and Children*. Seattle: IASP Press: 21–46.

- Ansermino M, Basu R, Vandebek C, et al. (2003):** Nonopioid additives to local anesthetics for caudal blockade in children: a systematic review. *Paediatr Anaesth*; 13: 561-573.
- Arendt-Nielsen L, Petersen-Felix S, Fisher M, et al. (1995):** The effect of N-methyl D-aspartate antagonist (ketamine) on single and repeated nociceptive stimuli: A placebo-controlled experimental human study. *Anesth Analg*; 81: 63-68.
- Baris S, Karakaya D, Kelsaka E, et al. (2003):** Comparison of fentanyl-bupivacaine or midazolam-bupivacaine mixtures with plain bupivacaine for caudal anesthesia in children. *Pediatr Anaesth*; 13:126-131.
- Batra YK, Arya VK, Mahajan R, Chari P (2003):** Dose Response study of caudal neostigmine in pediatric patients Undergoing genitourinary surgery. *Pediatric Anaesth*; 13: 515-21.
- Basbaum AI, and Fields HL (1984):** Endogenous pain control systems: Brainstem spinal pathways and endorphin circuitry. *Annu Rev Neurosci*; 7:309.
- Beyer JE, and Wells N (1989):** The assessment of pain in children. *Ped Clin North Am*; 36: 837-854.
- Beyer JE, McGrath PJ, and Berde CB (1990):** Discordance between self-report and behavioral pain measures in children aged 3-7 years after surgery. *J Pain Symptom Manag*; 5(6): 350.
- Borgbjerg FM, Svensson BA, Frigast C, et al. (1994):** Histopathology after repeated intrathecal injections of preservative free ketamine in rabbit: a light and electron microscopic examination. *Anesth Analg*; 79: 98-104.

- Brau M, Sander F, Vogel W, et al. (1997):** Blocking mechanisms of ketamine and its enantiomers in enzymatically demyelinated peripheral nerve as revealed by single-channel experiments. *Anesthesiology*; 86: 394-404.
- Brown AM, and Birnbaumer L (1990):** Ionic channels and their regulation by G protein subunits. *Annu Rev Physiol* 52:197.
- Clark, and Gilbert (2001):** Regional analgesia in the intensive care unit: Principles and practice. *Critical Care Clinics*; 17 (4).
- Crighton IM, Barry BP, and Hobbs GJ (1997):** A study of the anatomy of the caudal space using magnetic resonance imaging. *Br J Anaesth*; 78: 391.
- D'Angelo R, Dean LS, Meitner GC, Nelson KE (2005):** Neostigmine combined with bupivacaine, clonidine, and sufentanyl for spinal labor analgesia. *Anesth Analg*; 93: 1560-4.
- Dalens B, and Hasnaoui A (1989):** Caudal anesthesia in pediatric surgery: success rate and adverse effects in 750 consecutive patients. *Anesth Analg*; 68: 83.
- Dalens BJ (2002):** Regional anesthetic techniques. In: Bissonnette B, and Dalens BJ (eds): *Pediatric Anesthesia: Principles and Practice*. McGraw-Hill, pp.528-575.
- Dalens BJ, and Mazoit JX (1998):** Adverse effects of regional anesthesia in children. *Drug Saf*; 19: 251-68.
- De Beer DAH, Thomas ML (2003):** Caudal additives in children: solutions or problems? *Br J Anaesth*; 90: 487-98.

- De Negri P, Ivani G, Visconti C, et al. (2001):** How to prolong postoperative analgesia after caudal anesthesia with ropivacaine in children: S-ketamine versus clonidine. *Pediatr Anesth*; 11: 679-683.
- Derbyshire SWG (2006):** Can fetuses feel pain? *Britis Medical Journal*, 332 (7546), 909-12.
- Desparmet JF (1993):** Control blocks in children and adolescents. In: Schoechter N, Berde Ch B, and Yasterm (eds): *Pain in infants, children and adolescents*. Williams and Wilkins, Baltimore, p.245.
- Dinesh Kaushal, Vinita S, Haider A and Anita M (2009):** Caudal bupivacaine-neostigmine for perioperative analgesia in pediatric patients undergoing infraumbilical surgeries: A prospective, randomized, double blind, controlled study. *The Internet Journal of Anesthesiology*. 2009 Volume 21 No.1:1092-406.
- Donevan SD, and Rogaski MA (1993):** GYKI-52466, a 2, 3-benzodiazepine, is a highly selective non-competitive antagonist of AMPA/kainate-receptor responses. *Neuron*; 10: 51-59.
- Duggan AW, Hope PJ, Jarrott B, et al. (1990):** Release, spread and persistence of immunoreactivity neurokinin A in the dorsal horn of the cat following noxious cutaneous stimulation: Studies with antibody microprobes. *Neurosci*; 35: 195-202.
- Ellis JL, Harman D, Gonzalez J, et al. (1999):** Development of muscarinic analgesics derived from epibatidine: role of the M4 receptor subtype. *J Pharmacol Exp Ther*; 288: 1143–50.

- Eranardini N, Sauer SK, Haberberger, Tallarida RJ (2001):** Drug Synergism and Dose-Effect Data Analysis Chapman Hall/CRC.
- Evan EE, Friedrichsdorf S, Vesel T, Zeltzer LK (2007):** Pediatric Pain and Palliative Care Programs: Developing U.S. Models of Clinical Service and Education. Symposium presented at the American Pain Society, Washington DC.
- Ferrante FM, and Timothy RV (1993):** Alpha-2 antagonists. *Postoperative pain management*; 1st edition. Churchill Livingstone.
- Fischer RA (2004):** Excitatory nicotinic and desensitizing muscarinic (M2) effects on C-nociceptors in isolated ratskin. *J. Neurosci.* 21: 3295–3302.
- Frank AJ, Moll JM, and Hort JF (1982):** A comparison of three ways of measuring pain. *Rheumatology and Rehabilitation*; 24: 211-217.
- Giaufre' E, Dalens B, and Gombert A (1996):** Epidemiology and morbidity of regional anesthesia in children: A one-year prospective survey of the French-Language Society of Pediatric Anesthesiologists (ADARPEF). *Anesth Analg*; 83: 904
- Gregory GA (2002):** Pediatric regional anesthesia. In: Gregory GA (ed): *Pediatric Anesthesia*, 4th edition. Churchill Livingstone.
- Grubb BD (1998):** Peripheral and central mechanisms of pain. *Br J Anaesth*; 81:811.
- Grunau RV, and Craig KD (1987):** Pain expression in neonates: facial action and cry. *Pain*; 28(3): 395-410.
- Gunter JB (1991):** Caudal anesthesia in children: a survey. *Anesthesiology*; 75:936.

- Gunter JB, and Eng C (1992):** Thoracic epidural anesthesia via the caudal approach in children. *Anesthesiology*; 75: A936.
- Hammond DL (1986):** Control systems for nociceptive afferent processing: the descending inhibitory pathways. *Spinal Afferent Processing*. Plenum Press, New York.
- Hanesch U, Heppelmann B, and Schmidt RF (1992):** Neurokinin A-like immunoreactivity in articular afferents of the cat. *Brain Research*; 586:332-335.
- Hanesch U, Pfrommer U, Grubb BD, et al. (1993):** Acute and chronic phases of unilateral inflammation in rat's ankle are associated with an increase in the proportion of calcitonin gene-related peptide-immunoreactive dorsal root ganglion cells. *European Journal of Neurosciences*; 5:154-161.
- Hannallah RS, Broadman LM, and Rice LJ (1987):** Testing the validity of an objective pain scale for infants and children. *Anesthesiology*; 69(3A): 770.
- Hirota K, and Lambert DG (1996):** Ketamine; its mechanisms of action and unusual clinical uses. *Br J Anaesth*; 77: 441-44.
- Hurstveit O, Maurset A, and Oye I (1995):** Interaction of the chiral forms of ketamine with opioid, phencyclidine, and muscarinic receptors. *Pharmacology and Toxicology*; 77: 355-59.
- Imai M, and Kemmotsu O (1992):** Direct observation of the epidural space by the superfine fiberscope. *Mar*; 41(3): 474-9.
- Jansen KL, Faull RL, and Dragunow M (1989):** Excitatory amino acid receptors in the human cerebral cortex: a quantitative autoradiographic study

comparing the distributions of [^3H] TCP, [^3h] glycine, [^3H] AMPA, [^3H] kainic acid binding sites. *Neuroscience*; 32: 587-607.

Johnson, Gordon E (2002): PDQ Pharmacology. 2nd edition. BC Decker. 2002. ISBN 1-55009-109-3.

Kangarga I, Randic M (1991): Outflow of endogenous aspartate and glutamate from the rat spinal dorsal horn in vitro by low activation of low- and high-threshold primary afferent fibres: Modulation by mu-opioids. *Brain Research*; 553:347-352.

Kast-Byerclly S, Tabin JR, and Green RS (1993): Colonization and infection rates of continuous pediatric epidural catheters: A prospective study. *Anesthesiology*; 79: A1132.

Katz N and Ferrante FM (1993): Nociception. *Postoperative pain management*; 1st edition. Churchill Livingstone.

Khalil S, Campos C, Farag AM, et al. (1999): Caudal block in children: Ropivacaine compared with bupivacaine. *Anesthesiology*; 91: 1279-84.

Kissin I (2002): Study design to demonstrate clinical value of preemptive analgesia: is the common used approach valid? *Reg Anesth Pain Med*; 27: 242–4.

Kumar P, Rudra A, Pan AK, et al. (2005): Caudal additives in pediatrics: A comparison among Midazolam, ketamine, and neostigmine coadministered with bupivacaine. *Anesth Analg*; 101: 69-73.

- Lauretti GR, de Oliveira R, Reis MP, Juliao MC, Pereira NL (1999):** Study of three different doses of epidural neostigmine co administered with lidocaine for postoperative analgesia. *Anesthesiology*; 90: 1534-8.
- Lee Susan J, Ralston Henry J Peter, Drey Eleanor A, Partridge John Colin, Rosen Mark A (2005):** Fetal Pain: A Systematic Multidisciplinary Review of the Evidence. *Journal of the American Medical Association*, 294 (8), 947- 954.
- Lee HM, and Sanders GM (2000):** Caudal ropivacaine and ketamine for postoperative analgesia in children. *Anesthesia*; 55: 798-810.
- Lei SZ, Pan ZH, Aggarwal SK, et al. (1992):** Effect of nitric oxide production on the redox modulatory site on the NMDA receptor-channel complex. *Neuron*; 8: 1087-1099.
- Levine JD, Fields HL, and Basbaum AI (1993):** Peptides and the primary afferent nociceptors. *J Neurosci*; 13: 2273-2286.
- Lidow MS (2002):** Long-term effects of neonatal pain on nociceptive systems. *Pain*; 90: 377-383.
- Liu SS, Hodgson PS, Moore JM, et al (1999):** Dose-response effects of spinal neostigmine added to bupivacaine spinal anesthesia in volunteers. *Anesthesiology* .90:710–7.
- Marchant WA, and Black AE (2002):** Caudal injections for postoperative analgesia in children. *Pediatr Anesth*; 12: 192-3.
- Maze M, and Tranquilli W (1991):** Alpha-2-adrenergic agonists: defining the role in clinical anesthesia. *Anesthesiology*; 74: 581.

McCarthy SN, and Lawson PW (1989): Cell type and conduction velocity of rat primary sensory neurones with substance P-like immunoreactivity. *Neurosciences*; 28:745-753.

McCarthy SN, and Lawson PW (1990): Cell type and conduction velocity of rat primary sensory neurones with calcitonin gene-related peptide-like immunoreactivity. *Neurosciences*; 34:623-632.

McClain BC, and Redd SA (1993): Barrier flaps for continuous caudal anesthesia in pediatric patients. *Anesthesiology*; 79: 400.

McGrath PJ, Johnston G, Goodman JT, Unruh (1985): CHEOPS: a behavioral scale for rating postoperative pain In children. In: Fields HL, Ed *Advances in Pain Research And Therapy*. New York: Raven Press: 395-402.

McGrath PJ, Seiferi CE, and Speechly KN (1996): A new analogue scale for assessing children's pain: an initial validation study. *Pain*; 64: 435-443.

McGrath PJ, and Unruh AM (1994): Measurement and assessment of pediatric pain. In: Wall PD, and Melzack R (eds): *Textbook of Pain*, 3rd edition. Churchill Livingstone, New York, pp. 303-313.

McGrath PJ, Hsu E, and Capelli M (1990): Pain from pediatric cancer: A survey of an outpatient oncology clinic. *Journal of Psychosocial Oncology*; 8: 109-124.

Meistelman, Claude (2001): Claude; Update of neuromuscular pharmacology. *Curr Opin*; 14(4): 399-404.

Melzack R (1987): The Short-Form McGill Pain Questionnaire. *Pain*; 30(2):191-7.

- Melzack R, and Wall PD (1996):** The challenge of pain, 2nd edition. Penguin Books, London.
- Mendelson WB (1992):** Neuropharmacology of sleep induction by benzodiazepines. *Neurobiology*; 16:221.
- Messer, William S (2000):** Acetylcholine. 20 Jan 2000.University of Toledo. Accessed 11 September 2005.
- Mody I, and McDonald JF (1995):** NMDA receptor-dependent excitotoxicity: The role of intracellular Ca^{++} release. *Trends in Pharmacological Science*; 16: 356-59.
- Morgan M (1989):** The rational use of intrathecal and extradural opioids. *Br J Anaesth*; 63: 165.
- Morgan GE, Mikhail MS, and Murray MJ (2002):** Nonvolatile anesthetic agents. In: Morgan GE, Mikhail MS, and Murray MJ (eds): *Clinical Anesthesiology*, 3rd edition. Appleton and Lange, Stamford, p.151-177.
- Morton CR, and Hutchison WD (1989):** Release of sensory neuropeptides in the spinal cord: studies with calcitonin gene-related peptide and galanin. *Neurosciences*; 31:807-815.
- Mycek MJ, Harvey RA, and Pamela CC (1997):** Opioid analgesics and antagonists. In: Lippincott's Illustrated Review: Pharmacology, 2nd edition. Lippincott-Raven Publishers, Philadelphia, PA, chapter 14.
- Naguib M, Sharif AM, Seraj M, et al (1991):** Ketamine for caudal analgesia in children: comparison with caudal bupivacaine. *Br J Anaesth*. 67:559–64.
- Naguib M, Yaksh TL.(1997):** Characterization of muscarinic receptor

subtypes that mediate antinociception in the rat spinal cord. *Anesth Analg*.85:847–53.

Nicoletti F, Bruni V, Copani A, et al. (1996): Metabotropic glutamate receptors; a new target for therapy of neurodegenerative disorders. *Trends in Neurosciences*; 19: 267-271.

Olubukola O, Nafiu and Kolawole(2007): Comparison of Caudal Ketamine with or without Bupivacaine in Pediatric SubumbilicalSurgery. JOURNAL OF THE NATIONAL MEDICAL ASSOCIATION. VOL. 99, NO. 6

Omote K, Kitahata CM, and Collins JG (1991): Interactions between opiate subtype and alpha-2 adrenergic agonists in suppression of noxiously evoked activity of neurons in the spinal dorsal horn. *Anesthesiology*; 74: 737.

Pin JP, and Duvoisin R (1995): Neurotransmitter receptors 1. The metabotropic glutamate receptors: structure and functions (a review). *Neuropharmacology*; 34: 1-26.

Porter F, and Porges S (1991): Vagal tone: An index of stress and pain in high risk newborn infants. *J Pain Symptom Management*; 6: 206.

Pradhan B, Bajracharya GR (2008): Midazolam for caudal analgesia in children: Comparison with caudal bupivacaine. *Katmandu Univ. Med J (KUMJ)*; 6(2): 166.

- Rajesh Mahajan, MD, Vinod K. Grover, MD MNAMS and Pramila Chari, MD FAMS (2004):** Caudal neostigmine with bupivacaine produces a dose-independent analgesic effect in children. *Canadian Journal of Anesthesia* 51:702-706.
- Reich DL, and Silvay G (1989):** Ketamine; An update on the first twenty five years of clinical experience. *Can J Anesth*; 77: 441-44.
- Rice LJ, and Britton (1994):** Pediatric regional anesthesia. In: Lake CL, Barash PG, and Sperry RJ (eds): *Advances in Anesthesia*; vol.11. Mosby-Year Book, St. Louis.
- Robertson B, Hao JX, Wiesenfeld-Halin Z, et al. (1997):** Interferon-gamma receptors in nociceptive pathways: role in neuropathic pain-related behavior. *Neurosci*; 8(5): 1311-1316.
- Ross DM, and Ross SA (1988):** Childhood Pain: Current issues, research and management. Urban and Schwarzenberg, Baltimore.
- Rowney DA, and Doyle E (1998):** Epidural and subarachnoid block in children. *Anaesthesia*; 53: 980.
- Rusin KI, Jiang MC, Cerne R, et al. (1993):** Interactions between excitatory amino acids and tachykinins in the rat spinal dorsal horn. *Brain Research Bulletin*; 30:329-338.
- Schott GD, Loh L (2004):** Anticholinesterase drugs in Treatment of chronic pain. *Pain* 2004, 20:201-206.
- Seeburg BH (1993):** The molecular biology of glutamate receptor channels. *Trends in Pharmacol Sci*; 14: 297-303.

- Somasundrana S. and Madhu G (2008):** A comparative Study of Ketamine And Tramadol As Additives To Plain Bupivacaine In Caudal Anaesthesia In Chilren. *The internet journal of anesthesiology*. Volume 17 number 2.
- Sonada H, and Omote K (1996):** Analgesic mechanism of ketamine. *Masui*; 45(6): 689-97.
- Sorkin LS, Westlund KN, Sluka KA, et al. (1992):** Neural changes in acute arthritis in monkeys. IV. Time-course of amino acid release into the lumbar dorsal horn. *Brain Research Reviews*; 17:39-50.
- Stoetling RK (1991):** Pharmacology and physiology in anesthetic practice. JB Lippincott.
- Sweet SD, McGrath PJ (2005):** Physiological measures of Pain. In: Finley GA, McGrath PJ (Eds). *Measurement of Pain in Infants and Children*. Seattle: IASP Press: 59–81.
- Takahashi H, Miyazaki M, Nanbu T, et al. (1998):** The NMDA receptor antagonist ketamine abolishes neuropathic pain after epidural administration in a clinical case. *Pain*; 75:391-394.
- Tallarida RJ (2001):** J. Pharmacol. Exp. Ther. 298: 865–872. | PubMed | ISI | ChemPort |.
- Taylor P (2001):** Anticholinesterase agents. In: Hardman JG, Limbird LE, eds. *Goodman and Gilman's The Pharmacological basis of Therapeutics*, 10th edn. New York: McGraw Hill,: 175-192.
- Tsao JCI (2006):** CAM for pediatric pain: What is state of the Research? *Evidence-based Complementary and Alternative Medicine*, 3, 143-144.

- Vicker MD, Morgan M, and Spencer PS (1991):** *Drugs in anesthetic practice*, 7th edition. Butter-Worth Heinemann, London, p.200.
- Wall PD (1984):** Introduction. In: Wall PD, and Melzack R (eds): *Textbook of pain*. Churchill Livingstone, Edinburgh, pp. 1-16.
- White MC (2004):** Pain and stress in the human fetus. *Best Practice and Research Clinical Anesthesiology* 18(2): 205- 220.
- Whitwam (1987):** Benzodiazepines. *Anesthesia*; 42: 1255.
- Wilkie J, Holzema WL, and Tester MD (1990):** Measuring pain quality: validity and reliability of children's and adolescents' pain language. *Pain*; 44: 151.
- Williams Pl, Warwick R, and Dyson M (1989):** Spinal medulla or cord. In: Williams PL, and Bannister LH (eds): *Grays Anatomy*, 37th edition. Churchill Livingstone, Edinburgh, p.922.
- Willis RJ (1998):** Caudal epidural blockade. In: Causins MJ, Bridenbaugh PO (eds): *Neural Blockade in Clinical Anesthesia and Management of Pain*. Philadelphia, Lippincott-Raven, p.323.
- Willis WD (1984):** The origin and destination of pathways involved in pain transmission. In: Wall PD, and Melzack R (eds): *Textbook of Pain*. Churchill Livingstone, Edinburgh, pp.88-99.
- Willis WD, Westlund KN (1997):** Neuroanatomy of the Pain System and of the Pathways that Modulate Pain. A comprehensive and recent review, not for the fainthearted. *J Clin Neurophys*; 14: 2-31.

Yang CY, Wong CS, Chang JY, et al. (1996): Intrathecal ketamine reduces morphine requirements in patients with terminal cancer pain. *Can J Anesth*; 43(4): 379-383.

Yaster M, Tobin JR, and Billett C (1994): Epidural analgesia in the management of severe vaso-occlusive sickle cell crisis. *Pediatrics*; 93: 310.

Zadra N, and Glusti F(2001): Caudal block in pediatrics. *Minerva Anesthesiol*; 67: 126-31.

Zorumski CF, and Isenberg KE (1991): Insights into the structure and function of GABA-benzodiazepine receptors: Ion channels and psychiatry. *An J Psychiatry*; 148:162.