

## REFERENCES

- Abdel-Razek, S.; El-Tabbakh, H. and Francis, M. (1989):**  
Detection of hepatitis B virus persistence and carcinogenesis. A histopathological study.  
Egyptian J. Lab. Med., 1:129.
- Abruzzo, L.V.; Thornton, A.J.; Liebert, M. and Kunkel, S.L. (1992):**  
Cytokine-induced gene expression of interleukin-8 in human transitional cell carcinomas and renal cell carcinomas.  
Am. J. Path., 140: 365.
- Aloisi, F.; Care, A. and Borsettino, G. (1992):**  
Production of hamolymphopoietic cytokines (IL-6, IL-8, Colony- stimulating factors) by normal human astrocytes in response to IL-1B and tumor necrosis factor- $\alpha$ .  
J. Immunol., 149: 2358.
- Al-Wabel, A.; Al-Janadi, M. and Raziuddin, S. (1994):**  
Cytokine profile of viral and autoimmune chronic active hepatitis.  
J. Allergy Clin. Immunol., 92: 902.
- Al-Wabel, A.; Al-Knawy, B. and Raziuddin, S. (1995):**  
IL-8 and granulocyte macrophage colony-stimulating factor secretion in hepatocellular carcinoma and viral chronic active hepatitis.  
Cl. Immunol. Immunopath., 74 (3): 2315.
- Anastassakos, C.; Alexander, G.J.; Wolstencroft, R.A. and Williams, R. (1988):**  
Interleukin-1 and interleukin-2 activity in chronic hepatitis B virus infection.  
Gastroenterology, 94: 999.
- Andreeff, M. and Welte, K. (1989):**  
Haemaopoietic colony stimulating factor.  
Semin. Oncol., 16: 211.
- Andus, T.; Baver, J. and Gerok, W. (1991):**  
Effects of cytokines on the liver.  
Hepatology, 13: 364-375.

- Anthony, P.B. (1987):**  
Liver cell dysplasia: What is its significance?  
Hepatology; 7: 394.
- Antman, K.S.; Griffin, J.D. and Elias, A. (1988):**  
Effects of recombinant human granulocyte macrophage colony stimulating factor on chemotherapy induced myelosuppression.  
N. Engl. J. Med., 79: 593.
- Aoki, K. (1978):**  
Cancer of the liver: International mortality trends, World Health Stat.  
Rep. 31: 28.
- Arakawa, M.; Kage, M. and Matsumoto, S. (1986):**  
Frequency and significance of tumor thrombi in oesophageal varices in hepatocellular carcinoma associated with cirrhosis.  
Hepatology, 6: 419.
- Baggiolini, M.; Dewald, B. and Moser, B. (199):**  
Interleukin-8 and related chemotactic cytokines: CXC and CC chemokines.  
Adv. Immunol., 55: 96.
- Baggiolini, M.; Walz, A. and Kunkel, S.L. (1989):**  
Neutrophil-activating peptide-1/interleukin-8, a novel cytokine that activates neutrophils.  
J. Clin. Invest., 84: 1045.
- Baldwin, G.C.; Gasson, J.C.; Kaufman, S.E.; Quan, S.G. and Dpersio, J.F. (1989a):**  
Non heamatopoietic tumor cells express functional GM-CSF receptors.  
Blood 73: 1033.
- Baldwin, G.C.; Benveniste, E.N.; Chung, G.Y.; Gasson, J.C. and Golde, D.W. (1993):**  
Identification and characterization of a high affinity granulocyte-macrophage colony-stimulating factor receptor on primary rat oligodendrocytes.  
Blood, 82: 3279.

- Ballesta, A.M.; Cartel, X.; Filella, X. and Brui, J. (1989):**  
Alpha-fetoprotein in the early diagnosis of HCC.  
J.Nucl. Med. Allied. Sci. 33 (3 suppl.) 39.
- Barlogie, B.; Jagannath, S.; Dixon, D.; Cheson, B.; Smallwood, L. and Alexanian, R. (1990):**  
High dose melpahlan and granulocyte macrophage colony stimulating factor for refractory multiple myeloma.  
Blood, 76:680.
- Bassendine, M.F.; Seta, L. and Salmeron, J. (1983):**  
Incidence of hepatitis B virus infection in alcoholic liver disease. HBsAg negative chronic active liver disease and primary liver cell cancer in Britain.  
Liver, 3: 65.
- Beasley, R.P. (1982):**  
Hepatitis B virus as the etiologic agent in hepatocellular carcinoma-epidemiologic considerations.  
Hepatology, 2: 215.
- Beasley, R.P.; Hwang, L.Y.; Lin, C.C. and Chien, C.S. (1981):**  
Hepatocellular carcinoma and hepatitis B virus.  
Lancet, ii: 1129.
- Belfield, A. and Goldberg, D. (1971):**  
Colorimetric determination of alkaline phosphatase activity.  
Enzyme, 12: 561.
- Berdel, W.E.; Danhauser-Riedle, S.; Steinhauser, G. and Winton, E.F. (1989):**  
Various human haematopoietic growth factors (IL-3, G-CSF, GM-CSF) stimulate clonal growth of non haematopoietic tumor cells.  
Blood, 73: 80.
- Bertagolli, M.; Lin, B. and Young, D. (1992):**  
IL-12 augment antigen dependent proliferation of activated T-lymphocytes.  
J. Immunol., 140: 3778.

- Bignold, L.P.; Rogers, S.D.; Siawa, T.M. and Bahnish, J. (1992):**  
Inhibition of chemotaxis of neutrophil leukocytes to interleukin-8 by endotoxins of various bacteria. *Infect. Immunol.*, 29: 4255.
- Bird, G.L.; Sheron, N.; Goka, J.; Alexander, G. and William, R.S. (1990):**  
Increased plasma tumor necrosis factor in severe alcoholic hepatitis. *Ann. Int. Med.*, 113: 197.
- Blanchard, D.K.; Norris, M.B. and Dejeu, J. (1991):**  
Production of granulocyte-macrophage colony stimulating factor by large granular lymphocytes stimulated with candida albicans: Role in activation of human neutrophil function. *Blood*, 77: 2259.
- Borcerick, T.W.; Gosink, B. and Menuck, L. (1980):**  
Echographic and radionuclide detection of hepatoma. *Radiology*, 135: 149.
- Borzio, M.; Bruno, S. and Rocalli, M. (1991):**  
Liver cell dysplasia and risk of hepatocellular carcinoma in cirrhosis: a preliminary report. *Br. Med. J.*, 302: 1312.
- Brechet, C. (1987):**  
Hepatitis B virus (HBV) and hepatocellular carcinoma. *J. Hepatol.*, 4: 269.
- Brechet, C.; Pourcel, C. and Louise, A. (1980):**  
Presence of integrated hepatitis B virus DNA sequence in cellular DNA of human hepatocellular carcinoma. *Nature*, 286: 533.
- Brown, Z.; Strieter, R.M.; Chensue, S.W. and Westwich, J. (1991):**  
Cytokine-activated human mesangial cells generate the neutrophil chemoattractant, interleukin-8. *Kidney. Int.* 40: 86.
- Bruix, J.; Barrera, J.M. and Calvet, X. (1989):**  
Prevalence of antibodies to hepatitis C virus in Spanish patients with hepatocellular carcinoma and hepatic cirrhosis. *Lancet*; ii: 1004.

- Burgess, A.W.; Camakaris, J. and Metcalf (1977):**  
Purification and properties of colony stimulating factor from mouse lung-conditioned medium.  
J. Biol. Chem., 252: 1998.
- Burgess, A.W.; Begely, C.G.; Johnson, G.R.; Lopez, A.F.; Mermod, J.J. and DeLamarier, J.F. (1987):**  
Purification and properties of bacterially synthesized human granulocyte macrophage colony stimulating factor.  
Blood, 69: 43.
- Bussolino, F.; Wang, J.M.; Defilippi, P.; Turrini, F.; Sanavio, F.; Edgell, C.L. and Mantovani, A. (1989):**  
Granulocyte and granulocyte-macrophage colony stimulating factors induce human endothelial cells to migrate and proliferate.  
Nature, 337: 471.
- Campbell, H.D.; Tucker, W.Q.J. and Hort, Y. (1987):**  
Molecular cloning and expression of the gene encoding human eosinophil differentiation factor.  
Proc. Natl. Acad. Sci., U.S.A., 84: 6629.
- Cannistran, S.A. and Griffin, J.D. (1988):**  
Regulation of the production and function of granulocytes and monocytes.  
Semin. Hematol., 25: 173.
- Cannistran, S.; Groshek, P.; Garlick, R. and Miller, J. (1990):**  
Regulation of surface expression of the granulocyte macrophage colony stimulating factor in normal human myeloid cells.  
Proc. Natl. Acad. Sci., USA; 87: 93.
- Case, J.P.; Lafyatis, R.; Remmers, E.F.; Kumkumian, G.K. and Wilder, R.L. (1990):**  
IL-1 regulation of transin/stromelysin transcription in Rheumatoid synovial fibroblasts appears to involve two antagonistic transduction pathways, an inhibitory prostaglandin dependent pathway mediated by CAMP, and a stimulatory protein kinase C-dependent pathway.  
J. Immunol., 145: 3755.

- Cebon, J.; Nicola, N.; Ward, M. and Maher, D. (1990):**  
Granulocyte-macrophage colony stimulating factor from human lymphocytes. The effect of glycosylation on receptor binding and biological activity.  
J. Biol. Chem., 265: 4483.
- Champlin, R.E.; Nimer, S.D.; Ireland, d. and Golde, D.W. (1989):**  
Treatment of refractory aplastic anaemia with recombinant human granulocyte-macrophage colony stimulating factor.  
Blood, 73: 694.
- Chen, W.F. and Zlotnick, A. (1991):**  
A novel cytotoxic T-cell differentiation factor.  
J. Immunol., 147: 528.
- Cheng, W.K. and Lightdale, C.J. (1987):**  
Primary liver cancer: diagnosis and laboratory functions. In: Hepatic and Biliary Cancer. Edited by Wanebo, H.J. Marcel Dekker New York, P. 45.
- Chiba, S.; Shibuya, K.; Piao, Y.F.; Tojo, A.; Sasaki, N.; Matsuki, S.; Miyagawa, K.; Miyazono, K. and Takaku, F. (1990):**  
Identification and cellular distribution of distinct proteins forming human GM-CSF receptor.  
Cell Regul, 1:327.
- Choo, Q.L.; Kuo, G. and Weiner, A.J. (1989):**  
Isolation of c-DNA clone derived from a blood-borne non-A, non-B viral hepatitis genome.  
Science, 244: 359.
- Cline, M.J. and Golde, D.W. (1974):**  
Production of colony stimulating activity by human lymphocytes.  
Nature, 248:703.
- Colditz, I.; Znahlen, R.; Dewald, B. and Baggiolini, M. (1989):**  
In vivo inflammatory activity of neutrophil activity factor, a novel chemotactic peptide derived from human monocytes.  
Am. J. Pathol., 134: 755.
- Colombo, M.; DeFranchis, R. and Del Ninno, E. (1991):**  
Hepatocellular carcinoma in Italian patients with cirrhosis.  
N. Engl. J. Med., 325: 675.

- Conrade, D.H.; Ben-Sasson, S.Z.; Legros, G.; Finkelman, F.D. and Paul, W.E. (1990):**  
Infection with *Nippostrongylus brasiliensis* or infection of anti-IgD antibodies markedly enhance FC-receptor mediated interleukin-4 production by non B, non T-cells.  
*J. Exp. Med.*, 171: 1497.
- Crabtree, J.E.; Wyatt, J.I.; Nichols, P.H.; Ramsay, N. and Primrose, J.N. (1994):**  
Interleukin-8 expression in helicobacter pylori infected, normal and neoplastic gastroduodenal mucosa.  
*J. Clin. Pathol.*, 47: 61.
- Cybulsky, M.I.; Chan, M.K. and Movat, H.Z. (1988):**  
Acute inflammation and microthrombosis induced by endotoxin, interleukin-1 and tumor necrosis factor and their implication in gram negative infection.  
*Lab. Invest.*, 58: 365.
- Dannelly, S.; Strieter, R.; Kunkel, S.L. and Carter, D.C. (1993):**  
Interleukin-8 and development of adult respiratory distress syndrome in risk patient group.  
*Lancet*, 344: 643.
- Darwish, M.A.; Issa, S.A.; Aziz, A.M.; Darwish, N.M.; Soliman, A.H. (1993):**  
Hepatitis C and B viruses, and their association with hepatocellular carcinoma in Egypt.  
*J. Egypt Public Health Assoc.*, 68: 1.
- Dedhar, S.; Gaboury, L.; Galloway, P. and Eaves, C. (1988):**  
Human granulocyte macrophage colony stimulating factor is a growth factor active on a variety of cell types of non haematopoietic origin.  
*Proc. Natl Acad. Sci. USA.*, 85: 9253.
- Demetri, G.; Zenzic, B.; Rheinwald, J. and Griffin, J. (1989):**  
Expression of colony stimulating factor genes by normal human mesothelial cells and malignant mesothelioma cell lines in vitro.  
*Blood*, 74: 940.

- Denzlinger, W.; Gerhartz, H.T.; Sagebiel, S.P.; Haberl, C. and Wilmanns, W. (1993):**  
Differential activation of the endogenous leukotriene biosynthesis by two different preparation of granulocyte-macrophage colony stimulating factor in healthy volunteers. *Blood*, 81: 2007.
- De-Oliveira, E.; Silva, A.; Roldan-Molina, F.; Dos-Santos, T.E.; Santo, G.C.; Miranda, M.P.; Genzini, T.; Neves, L.B. and Albuquerque, L.A. (1990):**  
Diagnosis of hepatocellular carcinoma performed by searching for serologic tumour markers. *Arch. Gastroenterol.*, 27 (2): 83.
- Devereux, S.; Linch, D.C.; Campos, C.D.; Spitte, M.F. and Jelliffe, A.M. (1987):**  
Transient leukopenia induced by granulocyte macrophage colony stimulating factor. *Lancet*, 2: 1523.
- Dienstag, J.L. and Alter, H.J. (1986):**  
Non-A, non-B hepatitis: evolving epidemiologic and clinical perspective. *Sem. Liver Dis.*, 6: 67.
- Dinareello, C.A. (1988):**  
Biology of interleukin-1. *FASEB J.* 2: 108.
- Dipersio, J.; Billing, P.; Kaufman, S.; Williams, R.E. and Gasson, J.C. (1988a):**  
Characterization of the human granulocyte macrophage colony stimulating factor receptor. *J. Biol. Chem.*, 263:1834.
- Dipersio, J.; Hedvat, C.; Ford, C.F.; Golde, D.W. and Gasson, J.C. (1991):**  
Characterization of the soluble human granulocyte-macrophage colony stimulating factor receptor complex. *J. Biol. Chem.*, 266: 279.

- Donahue, R.E.; Yang, Y.C. and Clark, S.C. (1990):**  
Human P40 T-cell growth factor (IL-9) supports erythroid colony formation.  
Blood, 75: 2271.
- Douglas, S.R. and Sabiston, D.C. (1987):**  
The liver. In: Essentials of Surgery. Edited by Sabiston, D.C. Jr. W.B. Saunders Company, Philadelphia, London, Toronto, Sydney, Tokyo. P. 513.
- Du, X.X.; Neben, T.; Goldman, S. and Williams, D.A. (1993):**  
Effects of recombinant human interleukin-11 on haematopoietic reconstitution in transplant mice, acceleration of recovery of peripheral blood neutrophils and platelets.  
Blood, 81: 27.
- Dunk, A.A.; Spiliadis, H.; Sherlock, S.; Fowler, M.; Monjardino, J. and Thomas, H.C. (1988):**  
Hepatocellular carcinoma: clinical, aetiological and pathological features in british patients.  
Int. J. Cancer, 41: 17.
- Eddleston, A.L.; Donaldson, P.T.; Hegarty, J.E. and Reed, B.D. (1991):**  
Immunological aspects of liver disease.  
Gut, S40.
- Edmondson, H.A. and Craig, J.R. (1987):**  
Neoplasms of the liver. In: Diseases Of The Liver. Edited by Schiff, L. and Eugene, R. J.B. Lippincott Company Philadelphia., P. 1109.
- El-Zayadi, A. (1989):**  
Primary hepatic malignancies: high lighting the Egyptian situation. Medical Forum, No. 33.
- Emerson, S.G.; Yang, Y.G.; Clark, S.C. and Long, M.W. (1988):**  
Recombinant granulocyte- macrophage colony stimulating factor and interleukin-3 have overlapping but distinct haematopoietic activities.  
J. Clin. Invest., 82: 1282.

- Engstrom, P.F.; McGlynn, K. and Weese, J.L. (1993):**  
Primary neoplasms of the liver. In: Cancer Medicine. Edited by Holland, J.F., Frei III, E.; Bast, R.C.Jr.; Kufe, D.W.; Morton, D.L. and Weichselbaum, R.R. Lea and Febiger, Philadelphia. London, P. 1430.
- Enwonwu, C.O. (1984):**  
The role of dietary aflatoxin in the genesis of hepatocellular cancer in developing countries.  
Lancet; ii: 956.
- Estey, E.H.; Kurzrock, R. and Talpaz, E. (1991):**  
Effects of low doses of recombinant human GM-CSF in patients with myelodysplastic syndromes.  
Br. J. Haematol., 77: 291.
- Eto, H.; Toriyama, K. and Itakura, H. (1994):**  
A clinicopathological study of hepatocellular carcinoma in Nagasaki, South-Wester Japan: The association of hepatitis B and C viruses. Southeast-Asian.  
J. Trop. Med. Public Health, 25 (1): 88.
- Farhi, D.C.; Shikes, R.H.; Muxai, P.J. and Silverberg, S. (1983):**  
Hepatocellular carcinoma in young people.  
Cancer, 52: 1516.
- Feldman, M. (1996):**  
Cytokines.  
Immunol. Today, June (3): 11.8.
- Fibbe, W.E.; Kuck, P.M.C.; Duinkerken, N.; Voogt, P.J. and Willemze, R. (1988b):**  
Factors influencing release of granulocyte macrophage colony stimulating factor activity from human mononuclear phagocytes.  
Eur. J. Haematol., 41: 352.
- Fielder, H.G.; Suciu, E.; Wittlief, C.; Ostertag, W. and Hossfeld, D.K. (1990):**  
Mechanisms of growth factor expression in acute myeloid leukemia.  
Leukemia, 4: 459.

- Fiehn, C.; Wermann, M.; Pezzutto, A.; Hufner, M. and Heilig, C. (1992):**  
Plasma granulocyte- macrophage colony stimulating factor concentrations in rheumatoid arthritis, systemic lupus erythematosus and spondyloarthritis.  
J. Rheumatol., 51 (3): 121.
- Fleischmann, J.; Golde, D.W.; Weisbart, R.H. and Gasson, J.C. (1986):**  
GM-CSF enhances phagocytosis of bacteria by human neutrophils.  
Blood 68: 708.
- Fortson, W.C.; Tedesco, F.T.; Starnes, E.C. (1985):**  
Marked elevation of serum transaminase activity associated with extrahepatic biliary tract disease.  
J. Clin. Gastroenterol., 7: 502.
- Foss, H.D.; Herbst, H.; Gottstein, S. and Demel, G. (1996):**  
Interleukin-8 in Hodgkin's disease.  
Am. J. Pathol., 148: 1229-1236.
- Gasson, J. (1991):**  
Molecular physiology of GM-CSF.  
Blood, 77: 1131.
- Gasson, J.C.; Weisbart, R.H.; Kaufman, S.E.; Clark, S.C.; Wong, G.G. and Golde, D.W. (1984):**  
Purified human granulocyte macrophage colony stimulating factor: Direct action on neutrophils.  
Science, 226: 1339.
- Gasson, J.C.; Kaufman, S.E.; Weisbart, R.H.; Tomonaga, M. and Golde, D.W. (1986):**  
High affinity binding of GM-CSF to normal and leukemic human myeloid cells.  
Proc. Natl. Acad. Sci. USA., 83: 669.
- Gately, M.; Desai, B. and Quinn, R. (1991):**  
Regulation of human lymphocyte proliferation by a heterodimeric cytokine IL-12 (CLME).  
J. Immunol., 147: 874.

- Gause, A.; Kejmis, S.; Schalz, R.; Schobert, T.; Jong, W. and Pohl, C. (1992):**  
Increased levels of circulating cytokines in patients with untreated Hodgkin's disease.  
Lympho cytokine Res. 11 (2): 109.
- Gerhartz, H.H.; Marcus, R. and Delmer, A. (1990):**  
Randomized phase II study with GM-CSF and low dose ARAC in patients with high risk myelodysplastic syndrome.  
Blood, 76 (Supplement 1): 337a.
- Gimbrone, M.; Obin, M.; Luis, E. and Hass, R. (1989):**  
Endothelial IL-8, a novel inhibitor of leukocyte endothelial interactions.  
Science, 246: 1601.
- Goitzuka, R.; Ohashi, T.; Ono, K.; Yasukawa, Y.; Koishibara, Y.; Fukui, H.; Ohsugi, Y. and Hasegawa, A. (1990):**  
IL-6 activity in infectious peritonitis.  
J. Immunol., 144: 2599.
- Gough, N.M.; Gough, J.; Metcalf, D.; Kelso, A. and Dunn, A.R. (1984):**  
Molecular cloning of cDNA encoding a murine haematopoietic growth regulator granulocyte/macrophage colony stimulating factor.  
Nature, 309: 763.
- Gralnik, H.R.; Givelber, H. and Abrams, E. (1978):**  
Dysfibrinogenemia associated with hepatoma.  
N. Engl. J. Med., 299: 221.
- Gulati, S.C. and Bennett, C.L. (1992):**  
Granulocyte macrophage colony stimulating factor as adjunct therapy in relapsed Hodgkin disease.  
Ann. Inter. Med., 116:1777.
- Holstensen, A.; Ceska, M.; Brandtzaeg, P. and Wage, A. (1993):**  
Interleukin-8 in serum and cerebrospinal fluid from patients with meningococcal disease.  
J. Infect. Dis., 167: 471.

- Hardell, L.; Bengtsson, N.O. and Jonsson, H. (1984):**  
Aetiological aspects of primary liver cancer with special regard to alcohol, organic solvents and acute intermittent porphyria. An epidemiological investigation.  
Br. J. Cancer, 50: 389.
- Harper, H.A. and Rodwell, V. (1979):**  
Review of physiological chemistry, 11<sup>th</sup> edition, Lebanon, Large medical publication. P. 145.
- Hebert, C.A.; Luscinskas, F.W.; Kiely, J.M.; Luis, C.C.; Obin, M.S.; Gimbrone, M.A. and Baker, J.B. (1990):**  
Endothelial and leukocyte forms of IL-8, conversion by thrombin and interactions with neutrophils.  
J. Immunol., 145: 3033.
- Hiraoka, K.; Iwai, K. and Yamashita, R. (1986):**  
Metastasis from hepatocellular carcinoma in sclerosed oesophageal varices in cirrhotic patients.  
Br. J. Surg., 73: 932.
- Hober, D.; Ajana, F.; Petit, M.; Sartiaux, C.; Boniface, M.; Caillaux, M. and Mouton, Y. (1993):**  
Granulocyte-macrophage colony stimulating factor and tumor necrosis factor-alpha in patients with human immunodeficiency virus type I infection.  
Microbiol Immunol., 37: 785.
- Holbrook, S.T.; Ohls, R.K. and Yang, Y.C. (1991):**  
Effect of IL-9 on clonogenic maturation and cell cycle status of fetal and adult haematopoietic progenitors.  
Blood, 77: 2129.
- Holmes, W.E.; Lee, J.; Kuang, W.J.; Rice, G.C. and Wood, W.I. (1991):**  
Structure and functional expression of a human interleukin-8 receptor.  
Science, 253: 1278.
- Ihde, D.C.; Sherlock, P. and Winawer, S. (1974):**  
Clinical manifestations of hepatoma. A review of 6 years experience at a cancer hospital.  
Am. J. Med., 56: 83.

- Itch, Y.; Okanou, T.; Enjo, F.; Sakamoto, S.; Ohmoto, Y.; Hirai, Y.; Kagawa, K. and Kashima, K. (1994):**  
Serum levels of macrophage colony stimulating factor in liver disease.  
J. Hepatol., 21 (4): 527.
- Jacobsen, G.K.; Gammelgard, J. and Fuglo, M. (1983):**  
Coarse needle biopsy versus fine needle aspiration biopsy in the diagnosis of focal lesions of the liver.  
Acta. Cytol., 27: 152.
- Jakubowski, A.A. and Golde, D.W. (1996):**  
Therapeutic use of cytokines. In: Haematology. Edited by Williams, J.; Beutler, E.; Erslev, A. and Lichtman, M.A. McGraw-Hill Book Company, P. 155.
- Jendrassik, L. and Groft, P. (1938):**  
Quantitative methods for measuring urobilinogen.  
Biochemistry, 297: 81.
- Johnson, P.J. and Williams, R. (1987):**  
Cirrhosis and the aetiology of hepatocellular carcinoma.  
J. Hepatol., 5: 140.
- Kaklamani, E.; Trichopoulos, D. and Tzonou, A. (1991):**  
Hepatitis B and C viruses and their interaction in the origin of hepatocellular carcinoma.  
JAMA, 265: 1974.
- Kaneko, S.; Miller, R.H.; Feinstone, S.M.; Unoura, M.; Kobayashi, K.; Hollori, N. and Purcell, R.H. (1989):**  
Detection of serum hepatitis B virus DNA in patients with chronic hepatitis using the polymerase chain reaction assay.  
Proc. North. Acad. Sci., USA., 86: 312.
- Kaplan, M.M. (1972):**  
Alkaline phosphatase.  
Gastroenterology; 62: 452.
- Kashala, L.O.; Conne, B.; Kapanci, Y.; Frei, P.C.; Lambert, P.H.; Kalengayi, M.R. and Essex, M. (1992):**  
Hepatitis B virus, AFP synthesis and hepatocellular carcinoma in Zaire.  
Liver, 12 (5): 330.

- Kato, K.; Yokoi, T.; Takano, N.; Kanegane, H.; Yachie, A.; Miyawaki, T. and Taniguchi, N. (1990):**  
Detection by in situ hybridization and phenotypic characterization of cell expressing IL-6 mRNA in human stimulated blood.  
J. Immunol., 144: 1317.
- Kaushansky, K.; O'Hara, P.J.; Berkener, K.; Segal, G.M.; Hagen, F.S. and Adamson, J.W. (1986):**  
Genomic cloning, characterization, multilineage growth-promoting activity of human granulocyte-macrophage colony stimulating factor.  
Proc. Natl. Acad. Sci. USA., 83: 3101.
- Keeling, J.W. (1971):**  
Liver tumours in infancy and childhood.  
J. Pathol., 101: 245.
- Kew, M.C. (1981):**  
Clinical pathological and etiological heterogenicity in HCC. Evidence from Southern Africa.  
Hepatology, 1:366.
- Kew, M.C. and Geddes, E.W. (1982):**  
Hepatocellular carcinoma in rural South African blacks.  
Medicine (Baltimore), 61: 98.
- Kew, M.C.; Houghton, M. and Choo, Q.L. (1990):**  
Hepatitis C virus antibodies in Southern African blacks with hepatocellular carcinoma.  
Lancet, 335: 873.
- Kimberly, J.V.Z.; Fischer, E. and Hawes, A.S. (1992):**  
Effect of intravenous IL-8 administration in non-human primate.  
J. Immunol., 148: 1746.
- Kitamura, T.; Salo, N.; Arai, K. and Miyajima, A. (1991):**  
Expression cloning of the human IL-3 receptor cDNA reveals a shared Beta subunit for human IL-3 and GM-CSF receptors.  
Cell, 66: 1165.

- Koch, A.E.; Polverini, P.J.; Harlow, L.A. and Strieter, R.M. (1992):**  
Interleukin-8 as a macrophage-derived mediator of angiogenesis.  
Science, 258: 1798.
- Koch, A.E.; Evunoff, H.L.; Haines, G.K. and Strieter, R.M. (1993):**  
Enhanced production of the chemotactic cytokines interleukin-8 and monocytes chemoattractant protein-1 in human abdominal aortic aneurysms.  
Am. J. Path., 142: 1423.
- Koff, W.C. and Fann, A.V. (1986):**  
Human tumor necrosis factor-alpha kills herpes virus-infected but not normal cells.  
Lymphokine. Res., 5: 215.
- Kupper, T.; Food, P.; Coleman, D. and Horowitz, M. (1987):**  
Growth of an interleukin-2 interleukin-4 dependent T-cell line induced by granulocyte macrophage colony stimulating factor (GM-CSF).  
J. Immunol., 138: 4288.
- Kurzrock, R.; Talpaz, M. and Estrov, Z. (1991):**  
Phase I study of recombinant human IL-3 in patients with bone marrow failure.  
J. Clin. Oncol., 9: 1241.
- Larsen, C.G.; Anderson, A.O.; Appella, E. and Oppenheim, J.J. (1989):**  
The neutrophil-activating protein (NAP-1) is also chemotactic for T-lymphocytes. Science, 243: 1464.
- Le Beau, M.M.; Westbrook, C.A.; Diaz, M.O.; Larsen, R.A. and Sherr, C.J. (1986):**  
Evidence for the involvement of GM-CSF and FMS in the deletion (5q) in myeloid disorders.  
Science 231: 984.
- Lee, H.S.; Chung, Y.H. and Kim, C.Y. (1991):**  
Specificities of serum alpha-fetoprotein in HbsAg +ve and HBsAg -ve patients in the diagnosis of hepatocellular carcinoma.  
Hepatology, 14: 68.

- Leopard, K.A.; Reisbach, L.M. and Plauda, C. (1992):**  
Gene expression for IL-1 receptor.  
J. Exp. Med., 85: 121.
- Lieschke, G.J. and Burgess, A.W. (1992):**  
Granulocyte colony stimulating factor and granulocyte-macrophage colony stimulating factor (part I).  
N. Engl. J. Med., 327: 28.
- Lin, D.Y.; Liaw, Y.F.; Chu, C.M.; Chang-Chein, W.U.; Chen, P.C. and Sheen, I.S. (1984):**  
Hepatocellular carcinoma. A laparoscopic study of 92 cases in Taiwan.  
Cancer, 54: 1466.
- Lindsten, T.; June, C.H. and Ledbetter, J.A. (1989):**  
Regulation of lymphokine messenger RNA stability as a surface-mediated T-cell activation pathway.  
Science, 244: 339.
- Lingao, A.L.; Domingo, E.O. and Nishioka, K. (1981):**  
Hepatitis B virus profile of hepatocellular carcinoma in the Philippines.  
Cancer, 48: 1590.
- Lok, A.S.F. and Lai, C.L. (1989):**  
Alpha-fetoprotein monitoring in Chinese patients with chronic hepatitis B virus infection: role in the early detection of hepatocellular carcinoma.  
Hepatology, 9: 110.
- Lopez, A.F.; Nicola, N.A.; Burgess, A.W. and Metcalf, D. (1983):**  
Activation of granulocyte cytotoxic function by purified mouse colony-stimulating factors.  
J. Immunol., 131: 2983.
- Lopez, J.B.; Thambyrajah, V.; Balasengaram, M. and Satguasingam, N. (1994):**  
Hepatitis markers in Malaysians with hepatocellular carcinoma.  
Br. J. Biom. Sci., 51(2): 177.

- Mahida, Y.R.; Ceska, M.; Effenberg, F. and Hawkey, C.J. (1992):**  
Enhanced synthesis of neutrophil-activating peptide-1,  
interleukin-8 in active ulcerative colitis.  
Clin. Sci., 82: 273.
- Mano, H.; Nishida, J.; Usuki, K.; Maru, Y. and Takaku (1987):**  
Constitutive expression of the granulocyte macrophage colony  
stimulating factor gene in human solid tumors.  
J. Cancer Res., 78: 1041.
- Markusse, H.M.; Breddveld, F.C. and Fibbe, W.E. (1990):**  
Granulocyte macrophage colony stimulating factor treatment  
of neutropenia associated with Felty's syndrome.  
Arthritis and rheumatism, 33: 1865.
- Martin, J.; Quiroga, J.A.; Bosch, O. and Carreno, V. (1994):**  
Changes in cytokine production during therapy with  
granulocyte macrophage colony stimulating factor in patients  
with chronic hepatitis B.  
Hepatology, 20 (5): 1156.
- Marucha, P.T.; Zeff, R.A. and Kreutzer, D.L. (1990):**  
Cytokine regulation of IL-1B gene expression in the human  
polymorph nuclear leucocyte.  
J. Immunol., 145: 2932.
- Matsushima, K.; Morishita, K.; Yoshimor, T. and Oppenheim, J.J.  
(1988):**  
Molecular cloning of human monocyte derived neutrophil  
chemotactic factor (MDNCF) and induction of MDNCF by  
interleukin-1 and tumor necrosis factor.  
J. Exp. Med., 167: 1883.
- Matsushima, K. and Oppenheim, J.J. (1989):**  
Interleukin-8 and MCAF: novel inflammatory cytokines  
inducible by IL-1 and TNF.  
Cytokine, 1: 2.

- Matsushima, K.T.A.; Yamada, M.; Fuutani, Y. and Oppenheim, J.J. (1992):**  
Properties of a specific IL-1 receptor on human Epstein Barr virus transformed B lymphocytes, identity of IL-1 alpha and IL-1 beta.  
J. Immunol., 136: 4496.
- McIntire, K.R.; Vogel, C.L. and Primack, A. (1976):**  
Effect of surgical and chemotherapeutic treatment on alpha-fetoprotein levels in patients with HCC.  
Cancer, 37: 677.
- Metcalf, D. (1983):**  
Clonal analysis of the response of HL60 human myeloid leukemia cells to biological regulators.  
Leuk. Res., 7: 117.
- Metcalf, D. (1986):**  
The molecular biology and functions of the granulocyte-macrophage colony stimulating factors. Blood, 67: 257.
- Metcalf, D. (1988):**  
The colony stimulating factors and their receptors. In: The molecular control of Blood cells. Edited by Metcalf, D. Cambridge, M.A.: Harverd University, Press, P. 26.
- Metcalf, D. (1993):**  
Haematopoietic regulation, redundancy or subtlety.  
Blood, 82: 3515.
- Miller, W.J.; Federle, M.P. and Campbell, W.L. (1991):**  
Diagnosis and staging of hepatocellular carcinoma: comparison of CT and sonography in 36 liver transplantation patients.  
Am. J. Reontgenol., 157: 303.
- Mitsuyasu, R.; Levine, J. and Miles, S.A. (1988):**  
Effect of long term subcutaneous administration of recombinant human granulocyte macrophage colony stimulating factor in patients with HIV-related neutropenia.  
Blood, 72: Suppl. 1: 356 abstract.

- Miyajima, A.; Kitamura, T.; Harada, N.; Yokota, T. and Aria, K. (1992):**  
Cytokine receptors and signal transduction.  
Annu Rev. Immunol., 10: 295.
- Miyatake, S.; Otsuka, T.; Yokota, T.; Lee, F. and Aria, K. (1985):**  
Structure of the chromosomal gene for granulocyte  
macrophage colony stimulating factor: Comparison of the  
mouse and human genes.  
EMBO, J. 4: 2561.
- Mokhtar, N. (1990):**  
Cancer pathology. Registry, 1985-1988.  
National Cancer Institute. Department of Pathology.
- Moore, K.W. (1993):**  
Annu. Rev.  
Immunol., 11: 165.
- Nagato, Y.; Kondo, F. and Kondo, Y. (1991):**  
Histological and morphometrical indicators for a biopsy  
diagnosis of well-differentiated hepatocellular carcinoma.  
Hepatology, 14: 473.
- Nakao, K.; Kimura, K. and Miura, Y. (1966):**  
Erythrocytosis associated with carcinoma of the liver (with  
erythropoietin assay of tumour extract).  
Am. J. Med. Sci., 251: 161.
- Nakao, A.; Suzuki, Y. and Isshiki, K. (1991):**  
Clinical evaluation of plasma abnormal prothrombin  
(desgammacarboxyprothrombin) in hepatobiliary  
malignancies and other diseases.  
Am. J. Gastroenterol., 86: 62.
- Nakashima, T.; Okuda, K.; Kojiro, M. and Jimi, A. (1983):**  
Pathology of hepatocellular carcinoma in Japan. 232  
consecutive cases autopsied in ten years.  
Cancer, 51: 863.

**Nemunaitis, J.; Singer, J.W. and Bukner, C.D. (1988):**

Use of recombinant human granulocyte macrophage colony stimulating factor in autologous marrow for lymphoid malignancies.  
Blood, 72: 834.

**Nimer, S.D.; Golde, D.W.; Kwan, K.; Lee, K. and Champlin, R. (1991):**

In vitro production of granulocyte macrophage colony stimulating factor in aplastic anemia: possible action of anti-thymocyte globulin.  
Blood, 78: 1634.

**Nishioka, K.; Watanabe, J. and Furuta, S. (1991):**

A high prevalence of antibody to the hepatitis C virus in patients with hepatocellular carcinoma in Japan.  
Cancer, 67: 429.

**Nolan, J.P. (1989):**

Intestinal endotoxins as mediators of hepatic injury. An idea whose time has come again.  
Hepatology, 10: 887.

**Ohnishi, S.; Murakami, T. and Moriyama, T. (1986):**

Androgen and estrogen receptors in hepatocellular carcinoma and in the surrounding non-cancerous liver tissue.  
Hepatology, 6: 440.

**Okuda, K.; Kotoba, K. and Obata, H. (1975):**

Clinical observations during a relatively early stage of HCC with special reference to serum AFP.  
Gastroenterology, 69: 226.

**Okuda, K. and Okuda, H. (1991):**

Primary liver carcinoma. In Oxford Textbook of Clinical Hepatology. Edited by Neil, M. Jean-Pierre, B.; Johannes, B.; Mario, R. and Jaun, R. Oxford, University Press, Oxford, New York, Tokyo, P. 1019.

**Omori, F.; Okamura, S.; Shimoda, K.; Otsuka, Harada, M. and Niho, Y. (1992):**

Levels of human granulocyte colony stimulating factor and granulocyte macrophage colony stimulating factor under pathological condition.  
Biotherapy, 24: 147.

- Oppenheim, J.J.; Ruscetti, F.W. and Connie, F. (1991):**  
Cytokines. In: Basic and Clinical Immunology, Edited by Stites, D.P. and Terr, A.I.; Appleton and Lange, London, Sydney, Toronto, Mexico, New Delhi, P. 78.
- Oppenheim, J.J. and Ruscetti, F.W. (1997):**  
Cytokines. In: Medical Immunology. Edited by Stites, D.P.; Terr, A.I. and Parslow, T.G. Lange medical book, London, Sydney, Toronto, Mexico, New Delhi, Tokyo, Singapore, Rio de Janeiro, New Jersey, P. 146.
- Oshimi, K. and Hoshino, S. (1992):**  
Function and molecular structure of IL-12.  
Nippon. Rinsho, 50 (8): 1840.
- Park, L.S.; Friend, D.; Gillis, S. and Urdal, D.I. (1986):**  
Characterization of the cell surface receptor for human granulocyte/macrophage colony stimulating factor.  
J. Exp. med., 164: 251.
- Park, L.S.; Martin, U.; Sorensen, R.; Luhr, S.; Morrissey, J.J.; Cosman, D. and Larsen, A. (1992):**  
Cloning of low affinity murine granulocyte macrophage colony stimulating factor receptor and reconstitution of a high affinity receptor complex.  
Proc. Natl. Acad. Sci., USA. 89: 4295.
- Parker, Jr.; Dahlin, D.C. and Stuffer, M.H. (1970):**  
Malignant hepatoma: Evaluation of surgical (including needle biopsy) material from 69 cases.  
Mayo Clin. Proc., 45: 25.
- Patil, R. and Borch, R. (1995):**  
Granulocyte-macrophage colony stimulating factor expression by human fibroblasts upregulated and downregulated by interleukin-1.  
Blood, 85: 80.
- Paul, S.R.; Barut, B.A.; Bennett, F.; Cochran, M.A. and Anderson, K.C. (1990):**  
Lack of a role of interleukin-11 in the growth of multiple myeloma.  
Leuk. Res., 16 (3): 247.

- Paul, S.R. and Schendel, P. (1992):**  
The cloning and biological characterization of recombinant human interleukin-11.  
*Int. J. Cell cloning*, 10 (3): 135.
- Pavesi, F.; Lotzniker, M. and Scarabelli, M. (1989):**  
Efficiency of composite laboratory tests in the diagnosis of liver malignancies.  
*Int. J. Biol. Markers*, 4(3): 163.
- Perri, P.; Waltz, A.; Dewald, B. and Baggiolini, M. (1988):**  
A novel neutrophil activating factor produced by human mononuclear phagocytes.  
*J. Exp. Med.*, 167: 1547.
- Peschel, C.; Green, I. and Ohara, I. (1987):**  
Role of B-cell stimulatory factor 1. IL-4 in clonal proliferation of B-cells.  
*J. Immunol.*, 139: 3338.
- Peter, M.; Wellek, S.; Dienes, H.P.; Junginger, T.; Meyer, J.; Meyer, Zum, Buschenfelde, K.H. and Gerken, G. (1994):**  
Epidemiology of HCC. Evaluation of viral and other risk factors in a low-endemic area for hepatitis B and C.  
*J-Gastroenterol.*, 32 (3): 146.
- Petit-Frere, C.; Ongas, B. and Mencia-Huerta, J. (1993):**  
IL-9 potentiates the IL-4 induced IgE and IgG release from murine B-lymphocytes.  
*Immunol.*, 79: 146.
- Plaut, M.; Pierce, J.H.; Watson, C.J.; Hanely-Hyde, J.; Nordan, R.P. and Paul, W.E. (1989):**  
Mast cell lines produce lymphokines in response to cross linkage Fc R1 or to calcium ionophores.  
*Nature*, 339: 64.
- Podlaski, F.J.; Nanduri, V.B.; Hulmes, J.D.; Pan, Y.C.; Levin, W.; Danho, W.; Chizzonite, R.; Gately, M.K. and Stern, A.S. (1992):**  
Molecular characterization of interleukin-12.  
*Arch Biochem. Biophys.*, 294 (1): 230.

- Punnonen, J.G.; Aversa, B.G.; Cocks, A.N.J.; McKenzie, S.; Menon, G.; Zurawski, R.D.; Malefyt and Vries, J.E.de. (1993):**  
Interleukin-13 induces interleukin-4 independent IgG4 and IgE synthesis and CD23 expression by human B-cells.  
Proc. Natl. Acad. Sci. USA., 90: 3730.
- Puren, A.J.; Fantuzzi, G. and Dinarello, C.A. (1998):**  
Interleukin-18 (IFN- $\gamma$  inducing factor) induces IL-8 and IL-1 $\beta$  via TNF- $\alpha$  producing from non-CD14+ human blood mononuclear cells.  
J. Clin. Invest., 101 (3): 711.
- Quesenberry, P.J. (1995):**  
Haemopoietic stem cells, progenitor cells and cytokines. In: Haematology. Edited by William's, J.; Beutler, E.; Erslev, A.J. and Lichtman, M.A. McGraw-Hill Book Company, P. 211.
- Ramadori, G. and Bushenfeld, H.H. (1993):**  
Liver cells and cytokines.  
Curr. Opinions Gastroenterol., 9: 359.
- Reitman, S. and Frankel, S. (1957):**  
Colorimetric determination of GOT and GPT activity.  
Am. J. Clin. Path., 28: 56.
- Rifaat, A.M. (1988):**  
A manual of surgery.  
The Scientific Book Center, Cairo, Egypt.
- Rook, G.; Roitt, I. and Brostoff, J. (1996):**  
Cytokine network: cytokines form a complex communication network between cells.  
Immun., 96: 212.
- Rosalki, S.B. (1975):**  
Gamma-glutamyl transpeptidase.  
Adv. Clin. Chem., 17:53.
- Rosellini, S.R.; Arienti, V.; Nanni, O.; Uenti, F.; Tassinari, M.; Camporesi, C.; Boriani, L.; Versari, G.; Osta, P.L.; Amadori, D.; Miglio, F. and Gasbarrini, G. (1992):**  
Hepatocellular carcinoma. Prognostic factors and survival analysis in 135 Italian patients.  
J. Hepatol., 16: 66.

- Ruff, M.R.; Farrar, W.L. and Bert, C.B. (1986):**  
Interferon gamma and granulocyte macrophage colony stimulating factor inhibit growth and induction antigens characteristic of myeloid differentiation in small cell lung cancer cell lines.  
Proc. Natl. Acad. Sci. USA., 13: 6613.
- Rummeny, E.; Weissleder, R. and Stark, D.D. (1989):**  
Primary liver tumours: diagnosis by MR imaging.  
Am. J. Roentgenol., 63: 72.
- Sanderson, C.F. (1992):**  
Interleukin-5, eosinophils and disease.  
Blood, 79: 3101.
- Scarffe, J.H. (1991):**  
Emerging clinical uses for GM-CSF.  
Eur. J. Cancer, 27: 1493.
- Schadendorf, D.; Moller, A.; Worm, M. and Czarmetzki, B.M. (1993):**  
IL-8 production by human malignant melanoma cells in vitro is an essential autocrine growth factor.  
J. Immunol., 152: 2667.
- Scheuer, P.J. (1991):**  
Classification of chronic viral hepatitis. A need for reassessment.  
J. Hepatol., 13: 372.
- Schmouder, R.L.; Strieter, R.M.; Wiggins, R.C.; Chensue, S.W. and Kunkel, S.L. (1992):**  
In vitro and in vivo interleukin-8 production in human renal cortical epithelia.  
Kidney Int., 41: 191.
- Schrezenmeier, H.; Raghavachar, A. and Heimpel, H. (1993):**  
Granulocyte-macrophage colony stimulating factor in the sera of patients with aplastic anaemia.  
Clin. Invest., 71: 102.
- Seigel, J.; Harper, M.E. and Wong-Staal, F. (1984):**  
Gene for T-cell growth factor location on human chromosome 4q.  
Science, 223: 175.

- Seitz, M.; Dewald, B.; Gerber, N. and Baggialini, M. (1991):**  
Enhanced production of neutrophil-activating peptide -  
1/interleukin-8 in rheumatoid arthritis.  
J. Clin. Invest., 87: 463.
- Shamma, S.; El-Bedawy, A.F.; Azzam, F.; Hamed, M.; Shehab, El-Din,  
M. and Aita, M. (1992):**  
Hepatocellular carcinoma in Egypt: Analysis of 192 cases.  
Egypt Nati Cancer Inst., 5 (4): 761.
- Sherlock, S. (1989):**  
Virus hepatitis. In: Diseases of the Liver and Biliary System.  
Edited by Sherlock, S. Blackwell Scientific Publication,  
Oxford, London, Edimburgh, 301-338.
- Sherlock, S. (1993):**  
Hepatic tumors. In: Diseases of the liver and biliary system.  
Ed. by Sherlock, S. and Dooley, J. Blackwell Scientific  
Publications, Oxford, London, Edimburgh, USA, Australia, P.  
503.
- Sheron, N.; Bird, G.; Koskinas, J. and Williams, R.S. (1993):**  
Circulating and tissue levels of neutrophil chemotaxin  
interleukin-8 are elevated in severe alcoholic hepatitis and  
tissue levels correlate with neutrophil infiltration.  
Hepatology, 18: 41.
- Shirafuji, N.; Mastuda, S.; Ogura, H. and Takaku, F. (1990):**  
Granulocyte colony stimulating factor stimulates human  
mature neutrophilic granulocytes to produce interferon- $\alpha$ .  
Blood, 75: 17.
- Simonetti, R.G.; Camma, C.; Fiarella, F.; Politi, f.; D'Amico, G. and  
Pagliaro, L. (1991):**  
Hepatocellular carcinoma: A world-wide problem. The major  
riskfactors.  
Dig. Dis. Sci., 36: 962.
- Sisson, S.D. and Dinarello, C.A. (1988):**  
Production of interleukin-1 alpha, interleukin-1 beta and  
tumor necrosis factor by human mononuclear cells stimulated  
by granulocyte macrophage colony stimulating factor.  
Blood, 72: 1368.

- Skjomsberg, C.; Erikstein, B.K.; Smeland, E.B.; Lie, S.O.; Funderud, S.; Beiske, K. and Blomhoff, H.K. (1991):**  
Interleukin-7 differentiates a subgroup of acute lymphoblastic leukemias.  
Blood, 77 (11): 2445.
- Soini, I. and Kojola, H. (1983):**  
Time-resolved fluorometer for lanthanide chelates-a new generation of non isotopic immunoassays.  
Clin. Chem., 29: 65.
- Sparrow, L.G.; Metcalf, D.; Hunkapiller, M.W.; Hood, L.E. and Burgess, A.W. (1985):**  
Purification and partial amino acid sequence of asialo murine granulocyte macrophage colony stimulating factor.  
Proc. Natl. Acad. Sci. USA, 82: 292.
- Steward, W.P. (1993):**  
Granulocyte and granulocyte macrophage colony stimulating factors.  
Lancet, 342: 153.
- Strieter, R.M.; Kunkel, H.J. and Showell, H.J. (1989):**  
Endothelial cell gene expression of a neutrophil chemotactic factor by TNF, IL-1b or ILPs.  
Science, 243: 1467.
- Strieter, R.M.; Kasahara, K.; Allen, R.M.; Standiford, T.J.; Rolfe, M.W.; Becker, F.S.; Chensue, S.W. and Kunkel, S.L. (1992):**  
Cytokine-induced neutrophil-derived interleukin-8.  
Am. J. Pathol., 141: 397.
- Strieter, R.M.; Kunkel, S.L.; Elner, V.M.; Martonyi, C.L.; Koch, A.E.; Polverini, P.J. and Elner, S.G. (1992):**  
Interleukin-8, corneal factor that induces neovascularization.  
Am. J. Pathol., 141: 1279.
- Suga, M.; Senota, A.; Arima, K.; Kodama, T.; Ikuta, S.; Sakamoto, H.; Ohe, Y.; Sugai, S. and Yachi, A. (1994):**  
Prevalence of HBV and HCV infection in Japanese patients with hepatocellular carcinoma.  
Hepatogastro-enterol., 41 (5): 436.

- Sumida, M.; Ohto, M. and Ebara, M. (1986):**  
Accuracy of arteriography in the diagnosis of small hepatocellular carcinoma.  
Am. J. Roentgnol., 147: 531.
- Sun, Y.; Katsutushi, K.; Isono, E. and Obata, H. (1992):**  
Elevated serum interleukin-6 levels in patients with acute hepatitis.  
J. Clin. Immunol., 12, 197.
- Swain, S.L. and Dutton, R.W. (1982):**  
Production of a B-cell growth promoting activity from a cloned T-cell line and its assay.  
J. Exp. Med., 156: 1821.
- Symann, M. (1992):**  
Interleukin-3 and peripheral blood stem cell harvesting. In: Perspectives on interleukin-3 symposium. New connaught Rooms, grand hall.  
London, 1:18.
- Takatsu, K.; Tominaga, A. and Mamaoka, T. (1980):**  
Antigen induced T-cell replacing factor (TRF).  
J. Immunol., 124: 2414.
- Takeda, S.; Gillis, S. and Palacios, R. (1989):**  
In vitro effects of recombinant IL-7 on growth and differentiation of bone marrow pro-B and pro-T-lymphocytes clones and fetal thymocyte clones.  
Proc. Natl. Acad. Sci., U.S.A., 86: 1634.
- Taketa, K. (1990):**  
Alpha-fetoprotein: Re-evaluation in hepatology.  
Hepatology, 12: 1420.
- Tavernier, J.; Devos, R.; Cornelis, S.; Tuypens, T.; Van der Heyden, j.; Fiers, W. and Plaetinck, G. (1991):**  
A human high affinity interleukin-5 receptor (IL-5R) is composed of an IL-5 specific alpha chain and a beta chain shared with the receptor for GM-CSF.  
Cell 66: 1175.

**Teates, C.D. (1987):**

Radiological techniques in the diagnosis and treatment of liver tumours. In: Hepatic and Biliary Cancer. Edited by Wanebo, H.J. Marcel Dekker, New York, P. 57.

**Thornton, A.; Lindley, I. and Stierter, M. (1990):**

Cytokine induced gene expression of neutrophil chemotactic factor (IL-8) in human hepatocytes.  
J. Immunol., 144: 2609.

**Tilg, H.; Wilmer, A.; Vogel, W. and Huber, C. (1992):**

Serum levels of cytokines in chronic liver diseases.  
Gastroenterology, 103: 264.

**Tsai, J.F.; Chang, W.Y.; Jeng, J.E.; Ho, M.S.; Lin, Z.Y. and Tsai, J.H. (1994):**

Effects of hepatitis C and B viruses infection on the development of HCC.  
J. Med. Virol., 44 (1): 92.

**Vadas, M.A.; Nicola, N.A.; Metcalf, D. (1983):**

Activation of antibody dependent cell mediated cytotoxicity of the human neutrophils and eosinophils by separate colony-stimulating factors.  
J. Immunol., 130: 795.

**Van Damme, t. and Opdenakker, G. (1991):**

Interaction of interferons with skin reactive cytokins, from interleukin-1 to interleukin-8.  
J. Invest. Dermatol., 95 (6): 90.

**Van Leeuwen, B.H.; Martinson, M.E.; Webb, G.C. and Young, J.G. (1989):**

Molecular organization of the cytokines gene cluster involving the human IL-3, IL-4, IL-5 and GM-CSF genes, on human chromosome 5.  
Blood, 73: 1142.

**Van Meir, E.; Ceska, M.; Effenberger, F. (1992):**

Interleukin-8 is produced in neoplastic and infectious diseases of the human central nervous system.  
Cancer Res. 52: 4297.

- Van Zee, K.J.; Deforge, L.E.; Fischer, E. and Remick, D.G. (1991):**  
IL-8 in septic shock. endotoxaemia and after IL-1 administration.  
J. Immunology, 146: 3478.
- Wanebo, H.J.; Falkson, G. and Stanley, E.O. (1989):**  
Cancer of the hepatobiliary system. In Cancer Principles and Practice of Oncology. Edited by: Devita, V.T.; Hellman, S. and Rosenberg, S.A. J.B. Lippincott Company Philadelphia. P. 320.
- Weisbart, R.H.; Golde, D.W.; Clark, S.C. and Gasson, I.C. (1985):**  
Human granulocyte macrophage colony stimulating factor is a neutrophil activator.  
Nature, 314: 361.
- Wickenhauser, C.; Lorenzen, J.; Thiele, H.; Hillienhof, A. and Fisher, R. (1995):**  
Secretion of cytokines (IL-1 alpha, IL-3, IL-6 and GM-CSF) by normal human megakaryocytes.  
Blood, 85: 685.
- Wimperis, J.Z.; Niemeyer, C.M.; Sieff, C.A.; Nathan, D.G. and Arceci, R.J. (1989):**  
Granulocyte macrophage colony stimulating factor and interleukin-3 mRNA are produced by a small fraction of blood mononuclear cells.  
Blood, 74: 1525.
- Wognum, A.; Westerm, Y.; Visser, T. and Wagemaker, G. (1994):**  
Distribution of receptors for granulocyte-macrophage colony stimulating factor on immature CD34 positive bone marrow cells, differentiating non myeloid progenitors and mature blood cells subset.  
Blood, 84: 764.
- Wolters, G.; Kuikbers, L.; Kacaki, K. and Schurs, S. (1976):**  
Solid phase enzyme immunoassay for detection of HBsAg.  
J. Virol., 67: 4365-4368.
- Wood, J.W.; Rawlings, M.; Evans, H. and Lim, C.N. (1988):**  
HCC: Importance of histologic classification as a prognostic factor.  
Am. J. Surg., 155: 663.

- Wright, D.; Kenney, R.; Oette, D.; Boxer, L. and Malech, H. (1994):**  
Contrasting effects of recombinant human granulocyte macrophage colony stimulating factor and granulocyte colony stimulating factor treatment on the cycling of blood elements in childhood-onset cyclic neutropenia.  
Blood, 84: 1257.
- Xu, W.D.; Firestein, G.S.; Taetle, R.; Kaushansky, K.; Zvaifler, N.J. (1989):**  
Granulocyte macrophage colony stimulating factor in rheumatoid synovial effusions.  
J. Clin. Invest., 83: 876.
- Yao, Z.; Painter, S.L. and Fanslow, W.C. (1995):**  
Human IL-17: A novel cytokine derived from T-cells.  
J. Immunology, 155: 5483.
- Zaman, S.N.; Melia, W.M. and Johnson, R.D. (1985):**  
Risk factors in development of hepatocellular carcinoma in cirrhosis: Prospective study of 613 patients.  
Lancet I: 1357.
- Zsebo, K.; Yushenkoff, V.; Schiffer, S.; Chang, D.; McCall, E. and Bagby, G. (1988):**  
Vascular endothelial cells granulopoiesis IL-1 stimulate release of G-CSF and GM-CSF.  
Blood, 71: 103.
- Zwierzina, H.; Schollenberger, S.; Herold, M. and Besemer, J. (1992):**  
Endogenous serum levels and surface receptor expression of granulocyte-macrophage colony stimulating factor and interleukin-3 in patients with myelodysplastic syndrome.  
Leuk. Res., 16: 1181.