

## **REFERENCES**

---

## **REFERENCES**

**Agency for Healthcare Research and Quality.** Anaesthesia management during cataract surgery. Evidence Report/Technology Assessment: Number 16. Available at: <http://www.ahcpr.gov/clinic/epcsums/anestsum>.

**Allman K & Wilson I** - Eds, Oxford Handbook of Anaesthesia, Second Edition, Oxford University Press, 2006

**Assia EI, Pras E, Yehezkel M, Rotenstreich Y, Jager-Roshu S.** Topical anesthesia using lidocaine gel for cataract surgery. J Cataract Refract Surg 1999;25:635-9.

**Atkinson WS.** Retrobulbar injection of anesthetic within the muscular cone (cone injection). Arch Ophthalmol 1936;16:495-503.

**Bardsley H, Gristwood R, Baker H, et al.** A comparison of the cardiovascular effects of levobupivacaine and rac-bupivacaine following intravenous administration to healthy volunteers. Br J Clin Pharmacol 1998;46:245– 9.

**Behndig A.** Sub-Tenon's anesthesia with a retained catheter in ocular surgery of longer duration. J Cataract Refract Surg 1998; 24: 1307–1309.

**Bergman L, Berglin L, Algvere PV, Laurell CG, Stenkula S.** Limbal sub-Tenon's administration of retrobulbar anesthesia using a blunt irrigating cannula. Ophthalmic Surg Lasers 1996; 27: 106–112.

**Bloomberg LB.** Administration of periocular anesthesia. J Cataract Refract Surg 1986;12:677-679.

## **REFERENCES**

---

**Brahma AK, Pemberton CJ, Ayeko M, Morgan LH.** Single medial injection peribulbar anaesthesia using prilocaine. *Anaesthesia* 1994;49:1003-5.

**Bron AJ, Tripathi R, Tripathi B.** Wolf's Anatomy of the Eye and Orbit, 8th edn. London: Chapman and Hall, 1997.

**Brown S, Brooks S, Mazow M, et al.** Cluster of diplopia cases after periocular anesthesia without hyaluronidase. *J Cataract Refract Surg* 1999;25:1245–9.

**Budd J, Hardwick M, Barber K, Prosser J.** A single-centre study of 1000 consecutive peribulbar blocks. *Eye* 2001;15:464-8.

**Centers for Medicare and Medicaid Services.** Available at: <http://www.cms.hhs.gov>. Accessed October 25, 2005

**Chang D, Ladd L, Copeland S, et al.** Direct cardiac effects of intracoronary bupivacaine, levobupivacaine and ropivacaine in sheep. *Br J Pharmacol* 2001;132:649– 58.

**Courtney K.** Mechanism of frequency-dependent inhibition of sodium currents in frog myelinated nerve by the lidocaine derivative GEA 968. *J Pharmacol Exp Ther* 1975;195:225–36.

**Davis DB II, Mandel MR.** Posterior peribulbar anesthesia. an alternative to retrobulbar anesthesia. *J Cataract Refract Surg* 1986;12:182-4.

**Eagle K, Berger P, Calkins H, et al.** ACC/AHA Guideline update for perioperative cardiovascular evaluation for noncardiac surgery, executive summary. A report of the American College of Cardiology/ American Heart Association Task Force on Practice Guidelines (Committee to update the 1996 Guidelines on Perioperative Cardiovascular Evaluation for Non-cardiac Surgery). *J*

## **REFERENCES**

---

Am Coll Cardiol 2002;39:542– 53 Available at: <http://www.acc.org>. Accessed October 11, 2005

**Eghbal MH, Tabei H, Taregh SA, Razeghinejad MR**, The effect of addition of low dose atracurium to local anesthetic in retrobulbar block for cataract surgery. Middle East J Anesthesiol;2010

**Eke T, Thompson JR**. Serious complications of local anaesthesia for cataract surgery: a 1 year national survey in the United Kingdom. Br J Ophthalmol 2007; 91: 470-5.

**Elfituri.K, Arthurs G, Gemmell.L.**, MCQs for the Primary FRCA, Cambridge University Press, 2008

**Elhakim M, Sadek RA**. **Addition** of atracurium to lidocaine for intravenous regional anaesthesia. Acta .Anaesthesiol Scand 1994;38:542-544.

**Feldman H, Arthur G, Pitkanen M, et al**. Treatment of acute systemic toxicity after rapid intravenous injection of ropivacaine and bupivacaine in the conscious dog. Anesth Analg 1991;73:373–84.

**Finucane B**. Allergies to local anesthetics - the real truth. Canadian Journal of Anesthetics 2003;50:869–74.

**Fischer H & Pinnock C**, Fundamentals of Regional Anaesthesia, Cambridge University Press, 2004

**Friedman DS, Reeves SW, Bass EB, Lubomski LH, Fleisher LA, Schein OD**. Patient preferences for anaesthesia management during cataract surgery. Br J Ophthalmol 2004;88:333-5.

## **REFERENCES**

---

**Frieman BJ, Friedberg MA.** Globe perforation associated with subtenon's anesthesia. Am J Ophthalmol 2001;131:520-1.

**Gerson MC, Hurst JM, Hertzberg VS, et al.** Prediction of cardiac and pulmonary complications related to elective abdominal and noncardiac thoracic surgery in geriatric patients. Am J Med 1990;88:101– 7.

**Ghosh YK, Goodall KL.** Postoperative pain relief in vitreoretinal surgery with subtenon bupivacaine 0.75%. Acta Ophthalmol Scand 2005; 83: 119–120.

**Glantz L, Drenger B, Gozal Y.** Perioperative myocardial ischemia in cataract surgery patients: general versus local anesthesia. Anesth Analg 2000;91:1415– 9.

**Gold BS, Young ML, Kinman JL, et al.** The utility of preoperative electrocardiograms in the ambulatory surgical patient. Arch Intern Med 1992;152:301–5.

**Gouws P, Galloway P, Jacob J, English W, Allman KG.** Comparison of articaine and bupivacaine/lidocaine for sub-Tenon's anaesthesia in cataract extraction. Br J Anaesth 2004; 92: 228–230.

**Guise P, Laurent S.** Sub-tenon's block: the effect of hyaluronidase on speed of onset and block quality. Anaesth Intens Care 1999; 27: 179–181.

**Guise PA.** Sub-Tenon anesthesia: a prospective study of 6,000 blocks. Anesthesiology 2003; 98: 964–968.

**Hamilton RC.** Complications of ophthalmic regional anesthesia. Ophthalmol Clin North Am 1998;11:99–114.

## **REFERENCES**

---

**Hamilton RC.** Complications of ophthalmic regional anesthesia. In: Finucaine BT, editor. Complications of regional anesthesia. Philadelphia7 Churchill Livingstone; 2000. p. 39–55.).

**Hammond E, McIndoe A, Base Q;et al;** Anaesthesia 2, Greenwich Medical Media, 2005.

**Heavner J.** Cardiac dysrhythmias induced by infusion of local anesthetic into the lateral ventricles of cats. *Anesth Analg* 1986;65:133–8.

**Hollman M, Wieczorek K, Berger A.** Local anesthetic inhibition of G protein-coupled receptor signaling by interference with Galpha(q) protein function. *Mol Pharmacol* 2001;59:294–301

**Howell SJ, Sear JW, Foex P.** Hypertension, hypertensive heart disease and perioperative cardiac risk. *Br J Anaesth* 2004;92:570– 83 htm. Accessed November 18, 2004.

**Huang Y, Pryor M, Mather L, et al.** Cardiovascular and central nervous system effects of intravenous levobupivacaine and bupivacaine in sheep. *Anesth Analg* 1998;86:797–804

**Kanai Y, Katsuki H, Takasaki M.** Lidocaine disrupts axonal membrane of rat sciatic nerve in vitro. *Anesth Analg* 2000;91:944–8.

**Kim SK, Andreoli CM, Rizzo III JF, Golden MA,Bradbury MJ.** Optic neuropathy secondary to sub-Tenon anesthetic injection in cataract surgery. *Arch Ophthalmol* 2003; 121: 907–909.

**Kim Chishti, Andrey Varvinskiy, Consultant** -Torbay Hospital,UK; Anaesthesia For Ophthalmic Surgery- Regional Techniques ,Anaesthesia Tutorial Of The Week 135 ;25th May 2009

## **REFERENCES**

---

**Kluger MT, Tham EJ, Coleman NA, et al.** Inadequate pre-operative evaluation and preparation: a review of 197 reports from the Australian incident monitoring study. *Anaesthesia* 2005;55:1173– 8.

**Kumar CM, Williamson S, Chabria R.** A simple method of sub-Tenon anaesthesia delivery. *Anaesthesia* 2000; 55:612–613.

**Kumar CM, Dodds C.** A disposable plastic sub-Tenon cannula. *Anaesthesia* 2001; 56: 399–400.

**Kumar CM, Dodds C.** Evaluation of Greenbaum sub-Tenon's block. *Br J Anaesth* 2001; 87: 631–633.

**Kumar CM, Dodds C, Faanning GL,** editors. *Ophthalmic Anaesthesia*. The Netherlands: Swets and Zeitlinger, 2002.

**Kumar CM, McNeela BJ.** Ultrasonic localization of anaesthetic fluid using sub-Tenon's cannulae of three different lengths. *Eye* 2003; 17: 1003–1007.

**Kumar CM, Dowd TC, Dodds C, Boyce R.** Orbital swelling following peribulbar and sub-Tenon's anaesthesia. *Eye* 2004; 18: 418–420.

**Kumar CM, Williamson S, Manickam B.** A review of sub-Tenon's block: Current practice and recent development. *Eur J Anaesth*. 2005; 22:567-77.

**Kumar CM, Dodds C.** Sub-Tenon's Anaesthesia. *Ophthalmol Clin North Am* 2006; 19: 209-19.

**Kumar CM, Dowd TC, Hawthorne M.** Retrobulbar alcohol injection for orbital pain relief under difficult circumstances: a case report. *Ann Acad Med Singapore* 2006; 35: 260-5.

## **REFERENCES**

---

- Kumar CM, Dowd TC.** Ophthalmic regional anaesthesia. *Current Opin Anaesthesiol* 2008;oct 21(5):632-7
- Kurt N, Kurt I, Aygünes B, Oral H, Tulunay M.** Effects of adding alfentanil or atracurium to lidocaine solution for intravenous regional anaesthesia. *Eur J Anaesthesiol* 2002;19:522-525.
- Lambert L, Lambert D, Strichartz G.** Irreversible conduction block in isolated nerve by high concentration of local anesthetic. *Anesthesiology* 1994;260:121–8
- Leaming DV.** Practice styles and preferences of ASCRS members – 2003 survey. *J Cataract Refract Surg* 2004; 30:892–900
- Lee C:** Structure, conformation, and action of neuromuscular blocking drugs. *Br J Anaesth* 2001;87:755. [[PMID: 11878528](#)]
- Mathew Alan ;** Sub-Tenon’s Anaesthesia for Ophthalmic Procedures, anaesthesia UK; 2010
- McGlone R, Heyes F, Harris P.** The use of muscle relaxant to supplement local anaesthetics for Bier’s blocks. *Archives of Emergency Medicine* ;1988;5(2):79–85
- McLure HA, Rubin AP.** Review of local anaesthetic agents. *Minerva Anesthesiol* 2005;71:59-74.
- Mein CE, Woodcock MG.** Local anesthesia for vitreoretinal surgery. *Retina* 1990;10:47-9.
- Monk W.** Skin reactions to apothetin and quinine (sic) in susceptible persons. *Arch Dermatol* 1920;1:651– 5

## REFERENCES

---

**Morrison S, Dominguez J, Frascarolo P, et al.** A comparison of the electrocardiographic effects of racemic bupivacaine, levobupivacaine, and ropivacaine in anesthetized swine. *Anesth Analg* 2000;90:1308–14.

**Moshfeghi DM, Lowder CY, Roth DB, et al.** Retinal and choroidal vascular occlusion after posterior sub-tenon triamcinolone injection. *Am J Ophthalmol* 2002; 134:132–4.

**Nancarrow C, Rutten A, Runciman W, et al.** Myocardial and cerebral drug concentrations and the mechanism of death after fatal intravenous doses of lidocaine, bupivacaine, and ropivacaine in sheep. *Anesth Analg* 1989;69:276–83.

**Niemi-Murola L, Krootila K, Kivisaari R, Kangasmaki A, Kivisaari L, Maunuksela EL.** Localization of local anesthetic solution by magnetic resonance imaging. *Ophthalmology* 2004; 111: 342–347.

**Nouvellon E, L'Hermite J, Chaumeron A, Mahamat A, Mainemer M, Charavel P, Mahiou P, Dupeyron G, Bassoul B, Dareau S, Eledjam JJ, Ripart J.** Ophthalmic regional anesthesia: Medial canthus episcleral (sub-Tenon's) single injection block. *Anesthesiology* 2004;100:370-374.

**Parkar Tansneem, Gogate Parikshit, Deshpande Madan, Adenwala Arif, Maske Amar, Verappak.** Comparison of Subtenon anaesthesia with peribulbar anaesthesia for manual small incision cataract surgery. *Indian journal of ophthalmology* 2005, 5 255-259.

**Pasternak LR, Arens JF, Caplan RA, et al.** *Practice advisory for preanesthesia evaluation: a report by the American Society of Anesthesiologists Task Force on preanesthesia evaluation.* *Anesthesiology* 2002;96:

## REFERENCES

---

**Quantock CL, Goswami T.** Death potentially secondary to sub-Tenon's block. *Anaesthesia* 2007; 62: 175-7.

**Rahman I, Ataullah S.** Retrobulbar hemorrhage after sub-Tenon's anesthesia. *J Cataract Refract Surg* 2004; 30:2636–2637.

**RCOA&RCOO** (Royal collage of anaesthesia and royal college of ophthalmologists;2001)

**RCOA&RCOO** (Royal collage of anaesthesia and royal college of ophthalmologists,2002).

**Reah G, Bodenham AR, Braitwait P, Esmond J, Ménage MJ.** Peribulbar anaesthesia using a mixture of local anaesthetic and vecuronium. *Anaesthesia* 1998; 53:551-554.

**Ripart J, Metge L, Prat-Pradal D, Lopez FM, Eledjam JJ.** Medial canthus single-injection episcleral (sub-tenon anesthesia): computed tomography imaging. *Anesth Analg* 1998; 87: 42–45.

**Ripart J, Prat-Pradal D, Vivien B, Charavel P, Eledjam JJ.** Medial canthus episcleral (sub-Tenon) anesthesia imaging. *Clin Anat* 1998;11:390-5.

**Ripart J, Lefrant JY, Vivien B, Charavel P, Fabbro-Peray P, Jaussaud A, et al.** Ophthalmic regional anesthesia: medial canthus episcleral (sub-tenon) anesthesia is more efficient than peribulbar anesthesia: a double-blind randomized study. *Anesthesiology* 2000;92:1278-85.

**Ripart J, Lefrant JY, de La Coussaye JE, Prat-Pradal D, Vivien B, Eledjam JJ.** Peribulbar versus retrobulbar anesthesia for ophthalmic surgery. An anatomical comparison of extraconal and intraconal injections. *Anesthesiology* 2001;94:56-62.

## REFERENCES

---

**Roman SJ, Chong Sit DA, Boureau CM, et al.** Sub-Tenon's anaesthesia: an efficient and safe technique. Br J Ophthalmol 1997;81:673– 6.

**Rosenthal KJ.** Rosenthal deep topical, fornix, applied, pressurized,“nerve block” anesthesia. Ophthalmol Clin North Am 1998;11:137-143.

**Rous SM.** Simplified sub-Tenon's anesthesia: miniblock with maxiblock effect. J Cataract Refract Surg 1999; 25:10–15.

**Rubin A.** Eye blocks. In: Wildsmith JAW, Armitage EN, McLure JH, editors. Principles and Practice of Regional Anaesthesia. London:Churchill Livingstone, 2003

**Ruschen H, Bremner FD, Carr C.** Complications after sub-Tenon's eye block. Anesth Anal 2003;96:273-7.

**Rutten A, Nancarrow C, Mather L, et al.** Hemodynamic and central nervous system effects of intravenous bolus doses of lidocaine, bupivacaine, and ropivacaine. Anesth Analg 1989;69:291–9.

**Olschewski A, Hemplemann G, Vogel W.** Blockade of Na<sup>+</sup> currents by local anesthetics in the dorsal horn neurons of the spinal cord. Anesthesiology 1998;88:172–9.

**Sandler G.** The importance of the history in the medical clinic and the cost of unnecessary tests. Am Heart J 1980;100(6 Pt 1):928– 31.

**Schein OD.** Adverse intraoperative medical events and their association with anesthesia management strategies in cataract surgery. Ophthalmology 2001;108:1721-1726.

## REFERENCES

---

**Scott D, Lee A, Fagan D, et al.** Acute toxicity of ropivacaine compared with that of bupivacaine. *Anesth Analg* 1989;78:1125–30.

**Scott IU, Gayer S, Voo I, et al.** Regional anesthesia with monitored anesthesia care for surgical repair of selected open globe injuries. *Ophthalmic Surg Lasers Imaging* 2005;36:122–8.

**Shammas HJ, Milkie M, Yeo R.** Topical and subconjunctival anesthesia for phacoemulsification: prospective study. *J Cataract Refract Surg* 1997;23:1577–80.

**Soufia Farrukh, D.O.M.S, FCPS;** Sub-Tenon anaesthesia; an efficient and safe technique for cataract surgery. *Professional Med J* Sept. 2005; 12(3): 300–303.

**Standring S, Ellis H, Healy J, Johnson D, Williams A.** *Gray's Anatomy: The Anatomical Basis of Medicine and Surgery*, 39th edn. Harlow: Churchill Livingstone, 2004.

**Sztark F, Thiocoip M, Favarel-Garrigues JF, Lassie P, Petitjean ME, Dabadie P.** The use of 0.25% lidocaine with fentanyl and pancuronium for intravenous regional anaesthesia. *Anaesth Analg* 1997; 84:777–779.

**Stevens JD.** A new local anesthesia technique for cataract extraction by one quadrant sub-Tenon's infiltration. *Br J Ophthalmol* 1992;76:670–4.

**Stewart J, Kellet N, Castro D.** The central nervous system and cardiovascular effects of levobupivacaine and ropivacaine in healthy volunteers. *Anesth Analg* 2003;97:412–

**Sudoh Y, Cahoon E, Gerner P, et al.** Tricyclic antidepressants as long-acting local anesthetics. *Pain* 2003;103:49–55.

## REFERENCES

---

**Thind GS, Rubin AP.** Local anaesthesia for eye surgery – no room for complacency. Br J Anaesth 2001;86:473-6.

**Torrance JM, Nikki P, Paloheimo M.** Orbicular muscle akinesia in regional ophthalmic anaesthesia with pH- adjusted bupivacaine: Effects of hyaluronidase and epinephrine. Can J Anaesthesia 1993; 40:1028-1034.

**Tucker G.** Local anesthetic drugs: mode of action and pharmacokinetics. In: Nimmo W, Rowbotham D Smith G, editors. Anesthesia, vol. 2. Oxford7 Blackwell Scientific Publications; 1994. p. 1371.

**Turnbull CS.** The hydrochlorate of cocaine, a judicious opinion of its merits. Med Surg Rep 1884;29:628-9.

**Vanhoutte F, Vereecke J, Verbeke N, et al.** Stereoselective effects of the enantiomers of bupivacaine on the electrophysiological properties of the guinea-pig papillary muscle. Br J Pharmacol 1991;103:1275– 81

**Varvinskiy A, Eltringham E,** Anaesthesia for Ophthalmic Surgery, Update in Anaesthesia, Issue 6 (1996), Article 3

**Waterman H, Mayer S, Lavin MJ, Spencer AF, Waterman C.** An evaluation of the administration of sub-Tenon local anaesthesia by a nurse practitioner. Br J Ophthalmol 2002;

**Winder S, Walker SB, Atta HR.** Ultrasonic localization of anesthetic fluid in sub-Tenon's, peribulbar, and retrobulbar techniques. J Cataract Refract Surg 1999;25:56-9.

**Xiong Z, Strichartz G.** Inhibition by local anesthetics of Ca<sup>2+</sup>channels in rat anterior pituitary cells. Eur J Pharmacol 1998;363:81–90

## **REFERENCES**

---

**Zuhal Kucukyavuz** ;Effects of atracurium added to local anesthetics on akinesia in peribulbar block; 2002)by American Society of Regional Anesthesia and Pain Medicine.