

REFERENCES

REFERENCES

- Ahima, R.S.; Prabakaran, D.; Mantzoros, C. et al. (1996):**
Role of leptin in the neuroendocrine response.
Nature, 382: 250-252.
- Altmann, P. (1989):**
Disturbance of cerebral function by aluminum in hemodialysis patients without overt aluminum toxicity.
Lancet, 11: 7-12.
- Arcol (1989):**
ISB 15: 121-124.
- Audi, L.; Mantzoros, C.S.; Vidal-Plug, A. et al. (1998):**
Leptin in relation to the resumption of menses in women with anorexia nervosa.
Mol. Psychiatry, 3: 544-547.
- Auwerx, J.; Staels, B. (1998):**
Leptin.
Lancet, 351: 737-42.
- Ayus, J.C., Frommer, J.P. and Young J.B. (1981):**
Cardiac and circulatory abnormalities in chronic renal failure.
Semin. Nephrol. 1:112-123.
- Baily, G.L. (1977) :**
The sick kidney and sex.
N. Eng. J. 296:1288-1291.
- Banks, W.A.; Kastin, A.Y.; Huang, W.(1996):**
Leptin enters the brain by a saturable system independent of insulin.
Peptides, 17:305-311.

Bazzic, Y. (1991):

Uraemic polyneuropathy.
Clin. Nephrol. 35:176-180.

Becker, J. (1992) :

Aetiology of chronic renal failure in the kidney (Prischilla Kincaid-smith ed.) Blackwell scientific publication P. 393.

Bennett, B.D.; Solar, G.P.; Yuan, J.Q. et al. (1996):

A role for leptin and its cognate receptor in hematopoiesis.
Curr. Biol., 6: 1170-1180.

Bergstrom, J. (1995):

Why are dialysis patients malnourished?
Am. J. Kidney. Dis., 26 (1):229-241.

Bilbery, G.L. (1972):

Potassium deficiency in chronic renal failure.
Kid. Intern., 4, 423-430.

Bjorbaek, C.; Elmquist, J.K.; Frantz, J.D. et al. (1998):

Identification of SOCS-3 as a potential mediator of central leptin resistance.
Mol. Cell., 1: 619-625.

Blacke, P.; Schmidt, P.; Zazgornik, J. et al. (1983)

Reduction of elevated plasma oxalic acid levels by pyridoxine therapy in patients on regular dialysis therapy.
Kidney Int., 24 (suppl. 16):5346. Abstract.

Blagg, C.R. (1991) :

Importance of nutrition in dialysis patients.
Am. J. Kidney. dis., 17:458-462.

Blumenkrantz, M.J. (1994):

Special problems in the dialysis patient : nutrition. In :
Handbook of dialysis. Little Brown and Company, Boston, pp
374-400.

- Borah, M.F.; Schoenfeld, P.Y.; Gotch, F.A. et al. (1978):**
Nitrogen balance during intermittent dialysis therapy of uraemia.
Kidney. Int, 14:491-500.
- Boyle, J.M. and Johnston, B. (1983):**
Acute upper gastrointestinal haemorrhage in patients with chronic renal disease.
Am. J. Med., 75: 409.
- Brzechffa, P.R.; Jakimiuk, A.J.; Agarwal, S.K. et al. (1996):**
Serum immunoreactive leptin concentrations in women with polycystic ovary syndrome.
J. Clin. Endocrinol. Metab., 81:4166-4169.
- Campfield, L.; Smith, F.; Guisez, Y. et al. (1995):**
Recombinant mouse OB protein: Evidence for a peripheral signal linking adiposity and central neural networks.
Science, 269: 546-549.
- Cano, N.; Fernandez, J.P.; LaCombe, P. et al. (1987):**
Statistical selection of nutritional parameters in haemodialyzed patients.
Kidney Int 37 (suppl 22): S 178-S180.
- Caro, J.F.; Sinha, M.K.; Kolaczynski, J.W. et al. (1996a):**
Leptin : the tale of an obesity gene.
Diabetes, 45:1455-1462.
- Caro, J.F.; Kolaczynski, J.W.; Nyce, M.R. et al. (1996b):**
Decreased cerebrospinal-fluid/serum leptin ratio in obesity: a possible mechanism for leptin resistance.
Lancet, 348: 159-161.
- Carro, E.; Senaris, R.; Considine, R.V. et al. (1997):**
Regulation of in vivo growth hormone secretion by leptin.
Endocrinology, 138: 2203-2206.

- Casabiell, X.; Pineiro, V.; Tome, M.A. et al. (1997):**
Presence of leptin in colostrum and/or breast milk from lactating mothers: a potential role in the regulation of neonatal food intake.
J. Clin. Endocrinol. Metab., 82: 4270-4273.
- Chapman, L.M.; Wittert, G.A.; Nouman, R.J. (1997):**
Circulating leptin in polycystic ovary syndrome: relation to anthropometric and metabolic parameters.
Clin. Endocrinol. Metab., 82: 1692-1696.
- Chehab, F.F.; Lim, M.E.; Lu, R. (1996):**
Correction of the sterility defect in homozygous obese female mice by treatment with the human recombinant leptin.
Nat. Gene., 12(3):318-320.
- Chen, H.; Chariot, O.; Tartaglia, L.A. et al. (1996):**
Evidence that the diabetes gene encodes the leptin receptor identification of a mutation in the leptin receptor gene in db/db mouse.
Cell, 84:491-495.
- Christs, S. and Mantzoros, M.D. (1999):**
The role of leptin in human obesity and disease: A review of current evidence.
Ann. Intern. Med., 130: 671-80.
- Chua, S.C.; Chung, W.K.; Wu-Peng, X.S. et al. (1996):**
Phenotypes of mouse diabetes and rat fatty due to mutation in the ob (leptin) receptor.
Science:271:994-996.
- Clark, A.S.; Mitch, W.E. (1983):**
Muscle protein turnover and glucose uptake in acutely uraemic rat effects of insulin and the duration of renal insufficiency.
J. Clin. Invest., 72:836-845.
- Clement, K.; Garner, C.; Hager, J. et al. (1996):**
Indication for linkage of the human OB gene region with extreme obesity.
Diabetes, 45: 687-690.

- Considine, R.V.; Considine, E.L.; Williams, C.J. et al. (1995a):**
Evidence against either a premature stop codon or the absence of obese gene mRNA in human obesity.
J. Clin. Invest., 95:2986-2988.
- Considine, R.V.; Sinha, M.K.; Heiman, M.L. et al. (1995b):**
Serum irrununoreactive-leptin concentrations in normal-weight and obese humans.
N. Eng. J. Med., 334:292-295.
- Considine, R.V.; Considine, E.L.; Williams, C.J. et al. (1996):**
Mutation screening and identification of a sequence variation in the human Ob gene coding.
Biochem. Biophysics Res. Common, 220:735-739.
- Colernan, D.L.; Hummel, K.P. (1969):**
Effects of parabiosis of normal with genetically diabetic mice.
Am. J. Physiol, 217:1298-1304.
- Collins, S.; Kuhn, C.M.; Petro, A.E. et al. (1996):**
Role of leptin in fat regulation.
Nature, 380: 677-684.
- Conway, G.S.; Jacobs, H.S. (1997):**
Leptin: a hormone of reproduction.
Hum. Reprod., 12: 633-635.
- Couttenye, M.M. (1992):**
Productive value of low serum bone alkaline phosphatase (B AP) in the diagnosis of dynamic bone disease.
J.Am. Soc. Nephrol., 3: 672-680.
- Cumin, F.; Baum, H.P.; Levens, N. (1996):**
Leptin is cleared from the circulation primarily by the kidney.
Int. J. Obesity, 20:1120-1126.
- Cumin, F.; Baum, H.P.; De Gasparo, M. et al. (1997):**
Removal of endogenous leptin from the circulation by the kidney.
Int. J. Obesity, 21: 6: 495-504.

Cusin, I.; Sainsbury, A.; Doyle, P.; Rohner-Jeanrenaud, F. et al. (1995):

The Ob gene and insulin: a relationship leading to clues to the understanding obesity.

Diabetes 44:1467-1470.

Cusumano, A.; Lambardo, M.; Milano, C. et al. (1996):

Nutritional status of patient on chronic haemodialysis.

Medicana, 56 (6): 643-649.

Degoul, P.; Legrain, M.; Reachaine, F. and Deries, P. (1982):

Mortality risk factors in patient treated by chronic haemodialysis.

Nephron 31:103-110.

DeKlerk, G.; Wilnik, J.M.; Rosengarten, P.C.J. et al. (1982):

Serum erythropoietin (ESF) titers in anemia of chronic renal failure.

J.Clin. Lab. Med. 100 (5): 729-734.

Descombes, E. et al. (1993):

Water soluble vitamins in chronic haemodialysis patients and need for supplementation.

Kidney int., 43:13-19.

Deuschle, M.; Blum, W.F.; Englaro, P. et al. (1996):

Plasma leptin in depressed patients and healthy controls.

Horm. Metab. Res., 28: 714-717.

DeVos, P.; Saladin, R.; Auwerx, J.; Stoels, B. (1995):

Induction of ob gene expression by corticosteroids is accompanied by body weight loss and reduced food intake.

J. Biol. Chem., 270:15958-15961.

Dinosa, V.P.; Murthy, S.N.S. and Saris, A.L. (1982):

Gastric and pancreatic function in patients with end-stage renal disease.

J.Clin. Gastroenterol., 4: 321-328.

- Dimkovic, N. (1992):**
Uremic pruritus and skin mast cells.
Nephron 62: 5-11.
- Dobblestein, H.W.; Korner, W.F.; Mempel, W. et al. (1974):**
Vitamin B6 deficiency in uraemia and its implications for the depression of immuneresponses.
Kidney Int., 5:233-239.
- Dreiling, D.A. (1975):**
Pancreatic secretory testing in 1974 symposium in diagnosis of pancreatic disease.
Gut, 16: 647-652.
- Dumas, B.T.; Watson, W.A.; Biggs, H.G. (1971):**
Clin. Chem. Acta., 31: 87-89.
- Dumler, F. (1997):**
Use of bioelectric impedance analysis and dual-energy X-ray absorptiometry for monitoring the nutritional status of dialysis patients.
ASA. J., 43 (3): 256-260.
- Dunbar, J.C.; Hu, Y.; Lu, H. (1997):**
Intracerebroventricular leptin increase lumbar and renal sympathetic nerve activity and blood pressure in normal, rats.
Diabetes, 46: 2040-2043.
- Erickson, J.C.; Hollopeter, G.; Palmiter, R.D. (1996):**
Attenuation of the obesity syndrome of ob/ob mice by the loss of neuropeptide Y.
Science, 274(5293):1704-1707.
- Farron, F.; Considine, R.V.; Pelno, R. et al. (1997):**
Serum leptin concentrations in patients with anorexia nervosa, bulimia nervosa and non-specific eating disorders correlate with the body mass index but are independent of the respective disease.
Clin. Endocrinol. (Oxf), 46: 289-293.

- FEI, H.; Okano, H.; Li, C. et al. (1997):**
Anatomic localization of alternatively spliced leptin receptors (Ob-R) in mouse brain and other tissues.
Proc. Natl. Acad. Sci. USA, 94: 7001-7005.
- Flamenbaum, W. and Hamburger (1982):**
An approach to patients with renal disease.
J. B. Lippincott company, 1st edition, PP 544-550.
- Foley, R.N.; Parfrey, P.S.; Harnett, J.D et al. (1996):**
Hypoalbuminaemia, cardiac morbidity, and mortality in end stage renal disease.
J. Am. Soc. Nephrol., 7(5): 728-736.
- Fraser, C. L. and Arieff, A.I. (1988):**
Nervous system complications in uraemia.
Ann. Intern. Med., 109: 143-153.
- Frisch, R.E.; McArthur, J.W. (1974):**
Menstrual cycles, fatness, as a determinant of minimum weight for height necessary for their maintenance or onset.
Science, 185: 949-951.
- Gabriel, M. (1985):**
Biochemical changes and internal haemostasis in chronic renal failure.
Am. J. Physiol. 271: F1362-368.
- Gallieni, M.; Koren, J.W. and Saris, A.D. (1992):**
Low dose intravenous calcitriol treatment of secondary hyperparathyroidism in haemodialysis patients.
Kidney Int., 42:1191-1199.
- Garcia-Mayor, R.V.; Andrade, M.A.; Rios, M. et al. (1997):**
Serum leptin levels in normal children: relationship to age, gender, body mass index, pituitary-gonadal hormones, and pubertal stage.
J. Clin. Endocrinol. Metab., 82: 2849-2855.

- Gilmour, E.R.; Hartley, G.M.; Goodship, T.M.J. (1993):**
Trace elements and vitamins in renal disease. In Mitch, W.E.; Kloth, S. (eds): Nutrition and the kidney, Vol.2, Little Brown, Boston, pp 114-131.
- Golden, A. and Malier, J.F. (1977):**
In the kidney (structure and function) Monograph series (Abner Golden, series ed.) The Williams and Wilkins Company second edition. p. 32.
- Golper, T.A.; Ralbianni, J.M. and Lew, M. (1990):**
Multicenter trial of L-carnitine in maintenance haemodialysis patients. I. Carnitine concentrations and lipid effects.
Kidney Int, 38:904-909.
- Goodship, T.M.J.; Lloyd, S.; Clague, M.B. et al. (1987):**
Whole body leucine turnover and nutritional status in continuous ambulatory peritoneal dialysis.
Clin. Sci., 73:463-469.
- Grinspoon, S.; Gullick, T.; Askari, H. et al. (1996):**
Serum leptin levels in women with anorexia nervosa.
J. Clin. Endocrinol., Metab., 81: 3861-3863.
- Grunfeld, C.; Zhao, C.; Fuller, J.; Pollack, A.; Moser, A. et al. (1996):**
Endotoxin and cytokines induce expression of leptin, the ob gene product, in hamsters.
J. Clin. Invest., 97 (9):2152-2157.
- Gutierrez, A. et al. (1992) :**
Protein catabolism in shamhaemodialysis: the effect of different membranes.
Clin. Nephrol., 38 : 20-25.
- Halaas, J.; Gajiwala, K.; Maffei, M. et al. (1995):**
Weight reducing effects of the plasma protein encoded by the obese gene.
Science, 269: 543-546.

- Hamilton B. S.; Paglia, D.; Kwan, A.Y.M.; Deitel, M. (1995):**
Increased obese mRNA expression in omental fat cells from
massively obese humans.
Nature Med., 1:953-956.
- Hamilton, B.S. (1996):**
A new role for a fat actor.
Nature Med., 2: 272-273.
- Hassink, S.G.; Sheslow, D.V.; de Lancey, E. et al. (1996):**
Serum leptin in children with obesity: relationship to gender
and development.
Pediatrics., 98 (2 pt 1): 201-319.
- Hauseberger, F.X. (1959) :**
Parabiosis and transplantation experiments in hereditarily
obese mice.
Anal Rec., 130:313.
- Haynes, W.; Morgan, D.; Walsh, S. et al. (1997):**
Receptor-mediated regional sympathetic nerve activation by
leptin. *J. Clin. Inv.*, 100: 2: 270-278.
- He, Y.; Chen, H.; Quen, M.Y.; Reitman, M. (1995):**
The mouse obese gene.
J. Biol. Chem., 270:2887-2889.
- Hebebrand, J.; Bileum, W.F.; Barth, N. et al. (1997):**
Leptin levels in patients with anorexia nervosa are reduced in
the acute stage and elevated upon short-term weight
restoration.
Mol. Psychiatry, 2: 330-334.
- Heimbürger, O.; Lonnquist, F.; Danielsson, A. et al. (1997):**
Serum immunoreactive leptin concentration and its relation to
the body fat content in chronic renal failure.
J. Am. Soc. Nephrol., 8(9):1423-1430.

Hervey, G.r. (1958) :

The effects of lesions in the hypothalamus in parabiotic rats.
J. Physiol., 145:336-352.

Hoggard, N.; Mercer, J.; Rayner, D. et al. (1997):

Localization of leptin receptor mRNA splice variants in murine peripheral tissues by RT-PCR and in situ hybridization. *Biochem. Biophys. Res. Comm.*, 232: 387-390.

Holliday, M.A.; Chantler, C.A.; MacDonnell, R. et al. (1977):

Effect of uraemia on nutritionally induced variations in protein metabolism.
Kidney Int., 11:236-245.

Holworth, S.; Atkins, R.C. and Dekrester, D.M. (1977):

The pituitary testicular axis in end chronic renal failure.
N. Eng. J. Med., 299, 1245-1250.

Houseknecht, K.L.; Chris, S.M.; Kuliawat, R. j.S. et al. (1996):

Evidence for leptin binding to proteins in serum of rodents and humans: modulation with obesity.
Diabetes, 45:1638-1643.

Howard, J.K.; Lord, G.M.; Cluterbuck, E.J. et al. (1997):

Plasma immunoreactive leptin concentrations in end-stage renal disease.
Clin. Science, 93:119-126.

Iida, M.; Murakani, T.; Yamada, M. et al. (1996):

Hyperleptinaemia in chronic renal failure.
Horm. Metab. Res., 28 (12): 724-727.

Isse, N.; Ogawa, Y.; Tamura, N. et al. (1995):

Structural organization and chromosomal assignment of the human obese gene.
J. Biol. Chem., 270:27728-27733.

- Jackson, R.S.; Creemers, J.W.; Ohagi, S. et al. (1997):**
Obesity and impaired prohormone processing associated with mutations in the human prohormone convertase. J. gene. Nat. Genet., 16: 303-306.
- Jackson, E.; Li, P. (1997):**
Human leptin has natriuretic activity in the rat. Am. J. Physiol., 272: 41: F333-F338.
- Jenkins, A.B.; markovic, T.P.; Fleury, A.; Campbell, L.V. (1997):**
Carbohydrate intake and short-term regulation of leptin in humans. Diabetologia 40(3):348-351.
- Josselson, J.; Kyser, B.A.; Weir, M.R. and Sadler, R.LI. (1987):**
Hepatitis B surface antigenemia in a chronic hemodialysis program: lack of influence on morbidity and mortality. Am. J. Kidney. Dis., 9: 456-461.
- Kaufmann, P.; Swalle, K.H.; Morine, J.H. et al. (1994):**
Impact of long term haemodialysis on nutritinal status in patients with end-stage renal failure. J. Clin. invest., 72 (10): 754-761.
- Karlsson, C.; Lindell, K.; Svensson, L. et al. (1997):**
Expression of functional leptin receptors in the human ovary. J. Clin. Endocrinol. Metab., 82: 4144-4148.
- Kaysen, G.A. (1998):**
Albumin turnover in renal disease. Miner Electrolyte Metab., 24 (1): 55-63.
- Kennedy, G.C. (1953) :**
The role of depot fat in the hypothalamic control of food intake in the rat. Proc. R. Soc., 140:578-592.

- Kieffer, T.J.; Meller, R.S.; Habener, J.F. (1996):**
Leptin receptors expressed on pancreatic beta-cells.
Biochem. Biophys. Res. Commun., 224(2):522-527
- Kiess, W.; Englaro, P.; Hanitsch, S. et al. (1996):**
High leptin concentrations in serum of very obese children are further stimulated by dexamethasone.
Horm. Metab. Res., 28: 708-10.
- Kishimoto, T.; Taga, T.; Akira, S. (1994):**
Cytokine signal transduction.
Cell, 76:253-262.
- Klein, S.; Coppack, S.W.; Mohamed-Ali, V. and Landt, M. (1996):**
Adipose tissue leptin production and plasma leptin kinetics in humans.
Diabetes, 45: 984-987.
- Knochel, J.P. and Seldin, D.W. (1991):**
The pathophysiology of uraemia. In the kidney, ed. by Brenner B.M. and Rector, F.C. 4th ed. by Saunders Company vol 1, chap. 37 P. 1587.
- Kolaczynski, J.W.; Considine, R.V.; Ohannesian, J. (1996):**
Responses of leptin to short-term fasting and refeeding in humans: a link with ketogenesis but not ketones themselves.
Diabetes, 45:1511-1515.
- Kopple, J.D.; Mercuno, K.; Blumerkrantz, M.J. et al. (1981) :**
Daily requirement for pyridoxine supplement in chronic renal failure.
Kidney Int., 19:694-704.
- Kopple, J.D. (1984):**
Causes of catabolism and wasting in acute or chronic renal failure. In Robinson, R.R. (ed.): *Nephrology*, Springer-Verlag, New York, pp.498-15114.

- Kopple, J.D.; Ber, R.; Heuren, J. et al. (1989):*
Nutritional status of patients with different levels of chronic renal failure.
Kidney Int., 36 (suppl. 27): S184-S194.
- Kopple, J.D. (1994):*
Effect of nutrition on morbidity and mortality in maintenance dialysis patients.
Am. J. Kidney Dis., 24: 1002-1009.
- Krude, H.; Biebermann, H.; Ludeke, J. et al. (1998):*
Severe early-onset obesity, adrenal insufficiency and red hair pigmentation caused by POMC mutations in humans.
Nat. Genet., 19: 145-152.
- Kumar, P. and Clark, M. (1994):*
Complications of chronic renal failure. In *Clinical Medicine Textbook*, 3rd. ed., 1994.
- Kurtin, P.S. et al. (1990):*
Volume status and lung composition of chronic dialysis patients, utility of bioelectric impedance plethysmography.
Am. J. Nephrol., 10: 362-369.
- Lee, G.M.; Proenca, R.; Montez, A.M. et al. (1996):*
Abnormal splicing of the leptin receptor in diabetic mice.
Nature, 379: 632-635.
- Legrand, G.; Emerson, C.H.; Ahima, P.S. et al. (1997):*
Leptin prevents fasting-induced suppression of prothyrotropin-releasing hormone messenger ribonucleic acid in neurons of the hypothalamic paraventricular nucleus.
Endocrinology, 138: 2560-2576.
- Liach, F. and Massry, S.C. (1985):*
On the mechanism of hyperparathyroidism in moderate renal insufficiency.
J. Clin. Endocrinol. Metab., 61: 601-610.

Li, X.; Nickios, RB. (1995):

Mitotic forces control a cell cycle checkpoint.
Nature, 373:630-632.

Licinlo, J.; Mantzoros, C.; Negrao, A.B. et al. (1997):

Human leptin levels are pulsatile and inversely related to pituitary-adrenal function.
Nat. Med., 3: 575-579.

Licinlo, J.; Negrao, A.B.; Mantzoros, C. et al. (1998):

Synchronicity of frequently sampled, 24-h concentrations of circulating leptin, luteinizing hormone, and estradiol in healthy women.
Proc. Natl. Acad. Sci., USA, 95: 2541-2546.

Lindsay, R.; Micheal, I. and Nyce, R. (1991):

The influence of dialysis membrane upon protein catabolic rate.
ASAJO Trans, 37:M134-M140.

Liscovitch, M.; Cantley, L.C. (1994) :

Lipid second messengers.
Cell. 77:329-334.

Lonnquist, F.; Amer, P.; Nordford, L.; Schalling, M. (1995):

Overexpression of the obese (ob) gene in adipose tissue of human obese subjects.
Nature Med., 1:950-953.

Lowrie, E.G. and Lew, N.L. (1990) :

Death risk in haemodialysis patients: The predictive value of commonly measured variables and evaluation of death rate differences between facilities.
Am. H. Kidney. Dis., 15:458-482.

Lucas, P.A.; Meadows, J.h.; Roberts, D.E. et al. (1986):

The risks and benefits of a low protein-essential amino acid-keto acid diet.
Kidney Int., 29:995-1003.

Lynn, R.B.; Cao, G.Y.; Considine, R.V. (1996):

Auto-radiographic localization of lep tin binding in the choroid plexus of Ob/Ob and dbldb mice.
Biochem. Biophys. Res. Commun., 219 (3):884-889.

Macdougall, O.A.; Hwang, C.S.; Fan, H.P Lane, M.D. (1996):

Regulated expression of the obese gene product (leptin) in white adipose tissue and 3T3-Li adipocytes.
Proc. Natl. Acad. Sci. USA. 93:873-877.

Maffei, M.; Malaas, J.; i~ivussin, E.; Partley, R.E. (1995):

Leptin levels in human and rodent: measurement of plasma leptin and ob RNA in obese and weight-reduced subjects.
Nature Med., 1.1155-1161.

Maffei, M.; Stoffel, M.; Barone, M. et al. (1996):

Absence of mutations in the human OB gene in obese/diabetic subjects.
Diabetes, 45: 679-682.

Mahajan, S.K.; Prasad, A.8.; Rahbani, P. et al. (1982):

Zinc deficiency : A reversible complication of uraemia.
Am. J. Clin. Nutr., 36:1177-1183.

Makoff, R. (1991):

Water soluble vitamin status in patients with renal disease treated with haemodialysis or peritoneal dialysis.
J. Renal. Nutrion., 1:56-62.

Mantzoros, C.S.; Flier, J.S. (1995):

Insulin resistance: the clinical spectrum.
Adv. Endocrinol. Metab., 6: 193-232.

Mantzoros, C.S.; Rosen, H.N.; Greenspan, S.L. et al. (1997a):

Short-term hyperthyroidism has no effect on leptin levels in man.
J. Clin. Endocrinol. Metab., 82: 497-499.

Mantzoros, C.S.; Filler, J.S.; Rogol, A.D. (1997b):

A longitudinal assessment of hormonal and physical alterations during normal puberty in boys. V. Rising leptin levels may signal the onset of puberty.
J. Clin. Endocrinol. Metab., 82: 1066-1070.

Mantzoros, C.S.; Varvarigou, A.; Kaklamanl, V.G. et al. (1997c):

Effect of birth weight and maternal smoking on cord blood leptin concentrations in full-term and preterm newborns.
J. Clin. Endocrinol. Metab., 82: 2856-2861.

Mantzoros, C.S.; Moschos, S.; Avramopoulos, I. et al. (1997d):

Leptin concentrations in relation to body mass index and the tumor necrosis factor-alpha system in humans.
J. Clin. Endocrinol. Metab., 82: 3408-3413.

Maroni, B.J. (1993):

Requirements for protein, calories, and fat in the predialysis patients. In Mitch, W.E.; Klahi, S. (eds): Nutrition and the kidney, Vol.2, Little Brown, Boston, pp.985-212.

Massry, S.G. and Campbell, R.A. (1989):

Pathogenesis of uremic toxicity. In Textbook of Nephrology, vol. 2 (2nd ed.) S.G. Massry and R.J.G; Assock editors. Williams and Wilkins, Baltimore, 1126-1153.

McGregor, G.P.; Desaga, J.F.; Ehlenz, K. et al. (1996):

Radioimmunological measurement of leptin in plasma of obese and diabetic human subjects.
Endocrinology, 137: 1501-1504.

McFarlane, H.; Ogbcide, M.J. ; Peddy, S. et al. (1969):

biochemical assessment of protein-calorie malnutrition.
Lancet, 1:392-394.

Mehls, O.; Tonshoff, B.; Tonshoff, C. et al. (1992):

Therapeutic value of recombinant human growth hormone in children with chronic renal failure.
Min. Electrol. Metab., 18: 320-324.

- Merabet, E.; Dagogo-Jack, S.; Coyne, D.W. et al. (1997):**
Increased plasma leptin concentrations in end-stage renal disease.
J. Clin. endocrinol. Metab., 82 (3):847-850.
- Merrill, R.H. (1979):**
Iron metabolism in end-stage renal disease.
Dial Transplant., 8:890-904.
- Michael, J. (1991):**
Chronic renal failure: end stage management.
Med. International, PP3569-3577.
- Mikhail, A.A.; Beck, E.X.; Shafer, A. et al. (1997):**
Leptin stimulates fetal and adult erythroid and myeloid development.
Blood., 89: 1507-1517.
- Mitch, W.E. et al. (1989):**
Protein and amino acid metabolism in uraemia: influence of metabolic acidosis.
Kidney Int., 27 (suppl. 18):S205-S210.
- Mitch, W.E.; Walser, M. (1996):**
Nutritional therapy for the uraemic patients. In Brenner and Rector's The kidney, Vol.2. W.B. Saunders, Philadelphia, pp.2382-2423.
- Montague, C.T.; Farooqi, I.S.; Whitehead, J.P. et al. (1997):**
Congenital leptin deficiency is associated with severe early-onset obesity in humans.
Nature, 387: 903-908.
- Muolo, D.M.; Dohm, G.L.; Fledoruk, T.F. et al. (1997):**
Leptin directly alters lipid partitioning in skeletal muscle.
Diabetes, 46: 1360-1363.
- Murakami, T.; Iida, M.; Shima, K. (1995):**
Dexamethasone regulates obese gene expression in isolated rat adipocytes.
Biochem. Biophys Res. Commun., 214:1260-1267.

Nolan, J.J.; Olefsky, J.M.; Nyce, M.R. et al. (1996):

Effect of troglitazone on leptin production : studies in vitro and in human subjects.

Diabetes, 45:1276-1278.

Oettinger, C.W.; William, S.; Levens, W.J. (1992):

The effects of calcium carbonate as the sole phosphate binder in combination with low calcium dialysate and calcitriol therapy in chronic haemodialysis patients.

J. Am. Soc. Nephrol, 3:995-1001.

Ooi, B.S.; Parocy, A.F.; Pollak, V.E. (1972) :

Serum transferrin levels in chronic renal failure.

Nephron, 9: 200-208

Parfrey, P.S. (1991):

The natural history of myocardial disease in dialysis patients.

J. Am. Soc. Nephrol. 2:2-9.

Patton, A. and Crouch, W. (1977):

Anal. Chom., 49: 464-469.

Petersdorf, R.G. (1995):

Chronic renal failure: Pathophysiology and clinical considerations. In Harrison's principles of internal medicine By McGraw – Hill book company (ed.) sec. 4 P. 1595.

Rafoth, R.S.; Onstad, G.R. (1975):

Urea synthesis after oral protein loading in man.

J. Clin. Invest., 56:1170-1174.

Raja, R.; Ijelu, G.; Goldstein, M. (1992):

Influence of Kt/V and protein catabolic rate on hemodialysis morbidity. A long term study.

ASAIO, 38: M179-M180.

Reed, D.R.; Ding, Y.; Xu, W. et al. (1996):

Extreme obesity may be linked to markers flanking the human OB gene.

Diabetes, 45: 691-694.

Roitt, I. (1991):

Essential immunology. Blackwell Scientific Publication, Chapter 8, P. 169-170.

Rouru, J.; Anttila, L.; Koskineen, P. et al. (1997):

Serum leptin concentrations. In women with polycystic ovary syndrome.

J. Clin. Endocrinol. Metab., 82: 1697-700.

Rutsky, E.A., Robards, M., VanDyke, J.A. and Rostand, S.G. (1986):

Acute pancreatitis in patients with end-stage renal disease without transplantation.

Arch. Intern. Med., 146:1791-1800.

Rylance, P.B.; Brown, I.R.F.; Morrells, D.W. et al. (1984):

Relationship between vitamin A and bone disease in chronic renal failure.

Nephron, 36:131-135.

Sanaka, T.; Shinobe, M.; Ando, M. et al. (1994):

IGF-I as an early indicator of malnutrition in patients with end-stage renal disease.

Nephron, 67(1): 72-77.

Sargent, J.; Gotch, F.; Borob, M. et al. (1978):

Urea kinetics: A guide to nutritional management of renal failure.

Am. J. Clin. Nutr., 31:1696-1702.

Sarraf, P.; Frederick, R.C.; Turner, E.M. et al. (1997):

Multiple cytokines and acute inflammation raise mouse leptin levels: potential role in inflammatory anorexia.

J. Exp. Med., 185 (1):171-175.

Schieff, R T.; Zuckernian, G. and Harter, H. (1980):

Diverticular disease in patients with chronic renal failure due to polycystic kidney disease.

Ann. Intern. Med., 92 (Part 1): 202-209.

- Schirmenster, (1964):*
Ptsch. Med. Wschr., 89: 1018-1021.
- Schmicker, R. (1995):*
Nutritional treatment of hemodialysis and peritoneal dialysis patients.
Artif. Organs, 19: 8: 837-841.
- Schoenfeld, P.Y.; Henery, R.R.; Laird, N.M. and Roxe, D.M. (1983):*
Assessment of nutritional status of the national cooperative dialysis study population
Kidney. Int., 23 (suppl. 13) P. 80.
- Schubring, C.; Kiess, W.; Englaro, P. et al. (1997):*
Levels of leptin in maternal serum, amniotic fluid, and arterial and venous cord blood: relation to neonatal and placental weight.
J. Clin. Endocrinol. Metab., 82: 1480-1483.
- Schwartz, M.W.; Baskin, D.G.; Buckowski, T.R. (1996):*
Specificity of leptin action on elevated blood glucose levels and hypothalamic neuropeptide Y gene expression in ob/ob mice.
Diabetes, 45: 531-535
- Seaworth, W.W. (1984):*
Non A, Non B hepatitis and chronic dialysis. Another dilemma.
Am. J. Nephrol., 4: 235-241.
- Segal, K.R.; Landt, M.; Klein, S. (1996):*
Relationship between insulin sensitivity and plasma leptin concentration in lean and obese men.
Diabetes, 45: 987-991.
- Serradeil, L.E.; Gal, C.; Raufaste, D. et al. (1997):*
Characterization and localization of leptin receptors in the rat kidney.
FEBS Lett., 404: 185-191.

Sharma, K.; Considine, R.V.; Mielech, B. et al. (1997):

Plasma leptin is partly cleared by the kidney and is elevated in haemodialysis patients.

Kidney Int., 51 (6): 1980-1985.

Sharma, K. and Considine, R.V. (1998):

The ob protein (leptin) and the kidney.

Kid. Int., 53: 1483-1487.

Shoji, T.; Nishizawa, Y.; Emoto, M. et al. (199):

Renal function and insulin resistance as determinants of plasma leptin level in patient with NIDDM.

Diabetologia, 40: 676-679.

Sinhu, M.K.; Ohannesian, J.P.; Heiman, M.L.; Kriauciunas, A. et al. (1996):

Nocturnal rise of leptin in lean, obese, and non-insulin dependent diabetes mellitus subjects.

J. Clin. Invest., 97: 5: 1344-1347.

Sivitz, W.I.; Walsh, S.A.; Morgan, D.A. et al. (1997):

Effects of leptin on insulin sensitivity in normal rats.

Endocrinology, 138: 3395-401.

Slieker, L.J.; Sloop, K.W.; Surface, P.L. (1996):

Regulation of expression of ob mRNA and protein by glucocorticoids and cAMP.

J. Biol. Chem., 271: 5301-5304.

Stenvinkel, P. (1998):

Leptin : a new hormone of definite interest for the nephrologist.

Nephrol Dial. Transplant., 13:1099-1101.

Snitker, S.; Pratley, R.E.; Nicolson, M. et al. (1997):

Relationship between muscle sympathetic nerve activity and plasma leptin concentration.

Obese, Res., 5: 338-340.

- Smith, F.R.; Goodman, D.S. (1971):**
The effects of diseases of the liver, thyroid and kidneys on transport of vitamin A in human plasma.
J. Clin Invest, 50:2426-2436.
- Spicer, U.; Franciso, C.C. (1997):**
The adipose obese gene product, leptin: evidence of a direct inhibitory role in ovarian function.
Endocrinology, 138: 3374-3379.
- Stephens, T.W.; Basinski, M.; Bristow, P.K. et al. (1995):**
The role of neuropeptide Y in the antiobesity action of the obese gene product.
Nature 377:530-532.
- Strobel, A.; Issad, T.; Camoin, L. et al. (1990):**
A leptin missense mutation associated with hypogonadism and morbid obesity.
Nat. Gene., 18: 213-215.
- Stone, W.J.; Warnock, L.G.; Wagner, C. (1975):**
Vitamin B6 deficiency in uraemia.
Am. J. Clin. Nutr., 2&950-957.
- Susulic, V.S.; Lowell, B.B. (1996):**
Brown adipose tissue and the regulation of body fat stores.
Curr. Opin Endo Diabetes, 3:44-50 (Abstract).
- Tartaglia, L.A.; Dembeski, M.; Werg, X. (1995):**
Identification and expression cloning of a leptin receptor, OB-R.
Cell, 83:1263-1271.
- Teitz, N.W. (1996):**
Nitrogen metabolites and renal function. Fundamentals of clinical chemistry. W.B. Saunders Company, Philadelphia-London- Mentril- sydeny-Tokyo., P. 576-577.

Trayhurn, P.; Duncan, J.S.; Rayner, D.V. (1996):

Acute cold-induced suppression of ob (obese) gene expression in white adipose tissue of mice: mediation by the sympathetic system.
Biochem.J.; 311: 729-733.

Trayhurn, P.; Thomas, M.E.A.; Duncan, J.S.; Rayner, D.V. (1995):

Effects of fasting and refeeding on ob gene expression in white adipose tissue of lean and obese mice.
FEBS Lett., 368: 488-490.

Triger, D.R. and Jockes, A.M. (1972):

Severe muscle cramps due to acute hypo-magnesemia in haemodialysis.
Br. Med. J., 2: 804-810.

United States Renal Data System (1993):

A combined report on dialysis and transplantation.
Nephrol. Dial. Transplant., 4: 509-513.

Ursea, N.; Stancu, S.; Mircescu et al. (1998):

Nutritional status of patients undergoing chronic haemodialysis.
Nephrol. Dial., Transplant., 13 (6): A712-173.

Vahlquist, A.; Berne, B.; Berne, C. (1982):

Skin content and plasma transport of vitamin A and carotene in chronic renal failure.
Eur. J. Clin. invest., 12: 63-67.

Valcavi, R.; Zini, M.; Peino, R. et al. (1997):

Influence of thyroid status on serum immunoreactive leptin levels.
J. Clin. Endocrinol. Metab., 82: 1632-1634.

Van Wyck, D.B.; Stivelman, J.C.; Reidy et al. (1989):

Iron status in patients receiving erythropoietin for dialysis associated anaemia.
Kidney Int., 35: 712-716.

Vogel, G. (1996):

Leptin: a trigger for puberty?
Science, 274: 1466-1468.

Walser, M. (1993):

Does prolonged protein restriction preceding dialysis lead to protein malnutrition at the onset of dialysis?
Kidney Int., 44:1139-1144.

Waterlow, J.C. (1969):

The assessment of protein nutrition and metabolism in the whole animal with special reference to man. In Munro, H.N. (ed) *Mammalian protein metabolism*. Academic Press, New York, PP.326-390.

Wolfson, M. (1992):

Use of nutritional supplements in dialysis patients.
Semin. Dialysis., 5:285-291.

WHO Physical Status (1995):

The use and interpretation of anthropometry. Geneva Ch.
 WHO Technical Report, 854. World Health Organization.

York, P.; Daniel, C. and Celia G. (1992):

Predicting chronic renal insufficiency in idiopathic membranous glomerulonephritis.
Kidney Int., vol. 42 pp. 960-966.

Young, G.A.; Woodrow, G.; Kendall, S. et al. (1997):

Increased plasma leptin/ fat ratio in patients with chronic renal failure: a cause of malnutrition?
Nephrol. Dial. Transplant, 12 (11): 2318-2323.

Yu, V.L. (1986):

Staphylococcus aureus nasal carriage and infection in patients on hemodialysis.
N. Eng. J. Med., 315:91

Yu, W.H.; Kimura, M.; Walczewska, A. et al. (1997):

Role of leptin in hypothalamic-pituitary function.
Proc. Natl. Acad. Sci. USA, 94: 1023-1028.

Zachow, R.J.; Magoffin, D.A. (1997):

Direct intraovarian effects of leptin: impairment of the synergistic action of insulin-like growth factor-1 on follicle-stimulating hormone-dependent estradiol-17 beta production by rat ovarian granulosa cells.
Endocrinology, 138: 847-850.

Zakrzewska, K.E.; Cusin, I.; Sainsbury, A. et al. (1997):

Glucocorticoids as counterregulatory hormones of leptin: toward an understanding of leptin resistance.
Diabetes, 46: 717-722.

Zarate, A.R.; Hood, A.F. and Hardagen, G.L. (1978):

Hyperkeratosis penetrans (HKP): A rare dermatologic disease with high incidence in patients on haemodialysis.
Dial. and Transplant. 17: 99-104.

Zhang, Y.; Proenca, R.; Maffei, M. et al. (1994):

Positional cloning of the mouse obese gene and its human homologue.
Nature, 372:425-432.