

References:

Abdalah MG, (1988): Study of some forms of extrapulmonary tuberculosis. M. Sc. Thesis (Chest diseases), Cairo University.

Abdel-Hameed AA, (1981): Review of the results of Radiology and sputum examination in Sohag dispensary in the years 1976 - 1980. M.Sc. Thesis (Chest diseases), Ain Shams University.

Al-Azem A, Kaushal SM, Turenne C, Hoban D, Hershfield E, MacMorran J, Kabani A, (1998): Interscience Conference on Antimicrobial Agents and Chemotherapy. Abstr Intersci Conf Antimicrob Agents Chemother Intersci Conf Antimicrob Agents Chemother. 1998 Sep 24-27; 38: 555 (abstract no. L-27).

Agarwal R, Malhotra P, Awasthi A, Kakkar N, Gupta D, (2005): "Tuberculous dilated cardiomyopathy: an under-recognized entity?". BMC Infect Dis 5 (1): 29.

Agy, M.B.; Warris, C.K.; Plorde, J.J. (1989):

Evaluation of four mycobacterial blood culture media.
BACTEC 13A, Isolator/ BACTEC 12B,
Isolator/Middlebrook Agar and a biphasic medium.

Diagn. Microbial. Infect. Dis., 12:303-8.

Aiken AM, Hill PC, Fox A, et al, (2006): Reversion of the ELISPOT test after treatment in Gambian tuberculosis cases. BMC Infect Dis; 30: 66.

Alam U, Biomndo K, Davis J, Odonde W, Shanks GD, Elbeik T, Bell C, Wasunna KM, Rothel J, Mason C, (2000): Comparison of tuberculin skin testing with the quantiferon-TB assay in a tuberculosis and HIV endemic population in Kericho, Kenya. Int Conf AIDS. 2000 Jul 9-14; 13: abstract no. WePeC4441.

Al-Helaly A, Youssef S, Arafa R, El-Mallah M and Abou Seif N, (1993): Tuberculosis of the tonsils among Qualubia Children, Pathology and typing. Egypt, J. Chest Dis. And Tubere., 42 (2): 12.

American Society for Microbiology, (1998):

Bacteriophage. From the web site created by Division M, which is responsible for its contents.

Created 1.25.98, revised 6.8.01.

American Thoracic Society, (1981): The tuberculin skin test. Am. Rev. Respir. Dis., 1981, 124: 356.

American Thoracic Society, (1997): "Diagnosis and treatment of disease caused by nontuberculous mycobacteria. This official statement of the American Thoracic Society was approved by the Board of Directors, March 1997. Medical Section of the American Lung Association". Am J Respir Crit Care Med 156 (2 Pt 2): S1–25.

American Thoracic Society, (2000): Diagnostic standards and classification of tuberculosis in adults and children. Joint statement of the American Thoracic Society and the Centers for Disease Control and Prevention, endorsed by the

Council of the Infectious Diseases Society of America in September 1999. *Am. J. Respir. Crit. Care Med.* 161:1376-1395.

American Thoracic Society, (2000): Targeted tuberculin testing and treatment of latent tuberculosis infection. Joint statement of the American Thoracic Society and the Centers for Disease Control and Prevention, endorsed by the Council of the Infectious Diseases Society of America in September 1999. *Am. J. Respir. Crit. Care Med.* 161:S221-S247.

American Thoracic Society, (2003): CDC, and Infectious Diseases Society of America. Treatment of tuberculosis, *MMWR*; 52(No. RR-11):1--74.

Andersen P, Munk ME, Pollock JM, and Doherty TM, (2000): Specific immune-based diagnosis of tuberculosis. *Lancet* 356:1099-1104.

Arend SM, Thijsen SFT, Leyten EMS, Bouwman JJM, Franken WPJ, Koster BFP, Cobelens FGJ, van Houte AJ and Bossink AWJ, (2007): Comparison of Two Interferon- γ Assays and Tuberculin Skin Test for Tracing Tuberculosis Contacts. *American Journal of Respiratory and Critical Care Medicine* Vol 175. pp. 618-627.

Arend SM, van Meijgaarden KE, de Boer K, et al, (2002): Tuberculin skin testing and in vitro T cell responses to ESAT-6 and culture filtrate protein 10 after infection with *Mycobacterium marinum* or *M. kansasii*. *J Infect Dis*; 186: 1797–807.

Ariga H and Harada N, (2008): Evolution of IGRA researches, *Kekkaku*;83(9):641-52.

Ayad SS, (1993): The type of tubercle Bacilli in patient with pulmonary tuberculosis. M.Sc. Thesis (Chest diseases), Ain Shams University.

Barclay L and Vega C, (2007): Interferon-Gamma Tests May Be More Effective Than Tuberculin Skin Test. American Journal of Respiratory & Critical Care Medicine. Medscape, Medical News.

Parlet, R.C.; Youman, G.P.; Rehl, G.P. and Laster, V. (1958)

The detection of antibodies in the serum of tuberculous patients by agar double diffusion precipitation technique.

Am. Rev. Tuberc.,77:462.

Barnes PF, (2001): Diagnosing latent tuberculosis infection: the 100-year upgrade. Am J Respir Crit Care Med;163:807–808.

Barnes PF, (2004): Diagnosing latent tuberculosis infection: turning glitter to gold. Am J Respir Crit Care Med; 170:5–6.

Baron, E.J.; Peterson, L.R. and Finegold,S.M. (1994):

Methods for identification of etiological agents of infectious disease. In: Diagnostic microbiology, 9th ed. Baily and Scott's. Chapter 4, P. 611.

Batra, H.V.; Arya, A. and Jain, N.K. (1990):

Monoclonal antibody fluorescent assay for detection of antigens in pulmonary tuberculosis.

Am. Rev. Respir. Dis., 141(4): 805.

Bellete B, Coberly J, Barnes GL, et al, (2002): Evaluation of a Whole-Blood Interferon-Gamma Release Assay for the Detection of Mycobacterium tuberculosis Infection in 2 Study Populations. Clin Infect Dis;34:1449-1456.

Berthet FX, Rasmussen PB, Rosenkrands I, Andersen P, and Gicquel B, (1998): A Mycobacterium tuberculosis operon encoding ESAT-6 and a novel low-molecular-mass culture filtrate protein (CFP-10). Microbiology 144(Pt. 11):3195-3203.

Biron CA and Gazzanelli RT, (1995): Effects of IL-12 on immune responses to microbial infection: A key mediator in regulating disease outcome. Curr. Opin. Immunol.,7:485-496.

Brett,J.L. and Humble, M.W. (1991):

The Bactec TB. System in the laboratory diagnosis of mycobacterial infection.

N-Z-Med. J. Oct. 9; 104(921): 422-4.

Britton WJ, Gilbert GL, Wheatley J, Leslie D, Rothel JS, Jones SL, Bradley P, (2005): Sensitivity of human gamma interferon assay and tuberculin skin testing for detecting infection with Mycobacterium tuberculosis in patients with culture positive tuberculosis. Tuberculosis (Edinb). 85(3):137-45.

Brock I, Munk ME, Kok-Jensen A, Andersen P, (2001): Performance of whole blood IFN-gamma test for tuberculosis diagnosis based on PPD or the specific antigens ESAT-6 and CFP-10. Int J Tuberc Lung Dis; 5: 462-7.

- Brock I, Ruhwald M, Lundgren B, Westh H, Mathiesen LR, Ravn P, (2006):** Latent tuberculosis in HIV positive, diagnosed by the M. tuberculosis specific interferon-gamma test. *Respir Res*; 7: 56.
- Brock I, Weldingh K, Leyten EM, Arend SM, Ravn P, Andersen P, (2004):** Specific T-cell epitopes for immunoassay-based diagnosis of Mycobacterium tuberculosis infection. *J Clin Microbiol*; 42: 2379–87.
- Brock I, Weldingh K, Lillebaek T, Follmann F and Andersen P, (2004):** Comparison of Tuberculin Skin Test and New Specific Blood Test in Tuberculosis Contacts. *American Journal of Respiratory and Critical Care Medicine* Vol 170. pp. 65-69.
- Bruins J, Gribnau JH and Bwire R, (1995):** Investigation into Typical and Atypical Tuberculin Sensitivity in the Royal Netherlands Army, Resulting in a More Rational Indication for Isoniazid Prophylaxis. *Tuber Lung Dis*;76:540-544
- CDC, (2000):** Targeted tuberculin testing and treatment of latent tuberculosis infection. *MMWR*;49(No. RR-6):1--54.
- CDC, (2005):** Guidelines for preventing the transmission of Mycobacterium tuberculosis in health-care settings. *MMWR*;54(No. RR-17):1--47.
- CDC, (2005):** Guidelines for the investigation of contacts of persons with infectious tuberculosis: recommendations from the National Tuberculosis Controllers Association and CDC. *MMWR*;54(No. RR-17):1--47.

Cellestis Limited, (2007): QuantiFERON-TB Gold (in-tube method), package insert.

Cellistis limited, (2009): manufacturer of QuantiFERON diagnostics and research products: Mycobacterium Tuberculosis Novel Approaches to TB Diagnosis, Interferon Gamma Release Assays; and Image.

Centers for Disease Control and Prevention (CDC), (2009): TB: Mantoux Tuberculosis Skin Test Facilitator Guide.

Centers for Disease Control and Prevention (CDC), (2003): Division of Tuberculosis Elimination. Core Curriculum on Tuberculosis: What the Clinician Should Know. 4th edition.

Centers for Disease Control and Prevention (CDC), (2006): Surveillance Slide Set.

Centers for Disease Control, (2005): Fact Sheet: Tuberculosis in the United States.

Chaisson RE and Martinson NA, (2008): "Tuberculosis in Africa—combating an HIV-driven crisis". N Engl J Med 358 (11): 1089–1092.

Chaparas SD, Vandiviere HM, Melvin I, Koch G, Becker C, (1985): Tuberculin test. Variability with the Mantoux procedure. Am Rev Respir Dis; 132: 175–7.

Chapman AL, Munkanta M, Wilkinson KA, et al, (2002): Rapid detection of active and latent tuberculosis infection in HIV-positive individuals by enumeration of Mycobacterium tuberculosis-specific T cells. AIDS; 16: 2285–93.

Coker R, (2004): Compulsory screening of immigrants for tuberculosis and HIV. BMJ 328:298-300.

- Coker RJ, Bell A, Pitman R, Hayward A, and Watson J, (2004):** Screening programmes for tuberculosis in new entrants across Europe. *Int. J. Tuberc. Lung Dis.* 8:1022-1026.
- Cole E and Cook C, (1998):** "Characterization of infectious aerosols in health care facilities: an aid to effective engineering controls and preventive strategies". *Am J Infect Control* 26 (4): 453–64.
- Connell TG, Rangaka MX, Curtis N, Wilkinson RJ, (2006):** QuantiFERON-TB Gold: state of the art for the diagnosis of tuberculosis infection? *Expert Review of Molecular Diagnostics*, Volume 6, Number 5, pp. 663-677(15).
- Corbett EL, Watt CJ, Walker N, et al, (2003):** The growing burden of tuberculosis: global trends and interactions with the HIV epidemic. *Arch Intern Med*; 163: 1009–21.
- Cox R, (2004):** "Quantitative relationships for specific growth rates and macromolecular compositions of *Mycobacterium tuberculosis*, *Streptomyces coelicolor* A3(2) and *Escherichia coli* B/r: an integrative theoretical approach". *Microbiology* 150 (Pt 5): 1413–26.
- Davies PD, (2003):** "The world-wide increase in tuberculosis: how demographic changes, HIV infection and increasing numbers in poverty are increasing tuberculosis". *Annals of medicine* 35 (4): 235–43.
- Davies PDO, Yew WW, Ganguly D, et al, (2006):** "Smoking and tuberculosis: the epidemiological association and pathogenesis". *Trans R Soc Trop Med Hyg* 100: 291–8.

Davis L, (1995): "Vegetarian diet and tuberculosis in immigrant Asians". *Thorax*; 50 (8): 915–6.

Democratis Ja, Bailey S, Barer M, Pereira N, Stephenson I, Nicholson KG, Wiselka MJ, Democratis J, (2006): The value of the QuantiFERON-Gold TB test in the management of patients with suspected active tuberculosis. 16th European Congress of Clinical Microbiology and Infectious Diseases, Nice, France, April 1-4 2006, Abstract number: p1112.

Dewan PK, Grinsdale J, Liska S, Wong E, Fallstad R, Kawamura LM, (2006): Feasibility, acceptability, and cost of tuberculosis testing by whole-blood interferon-gamma assay. *BMC Infect Dis*; 6: 47.

Dhir V. K. and Stupple M. (1997):

New approach to laboratory diagnosis.

Tuberculosis: Nov.: 23-24

Diel R, Loddenkemper R, Meywald-Walter K, Niemann S, Nienhaus A, (2008): Predictive Value of a Whole Blood IFN- γ Assay for the Development of Active Tuberculosis Disease after Recent Infection with *Mycobacterium tuberculosis*. *Am J Respir Crit Care Med*; 177:1164–1170

Diel R, Nienhaus A, Lange C, Meywald-Walter K, Forssbohm M, Schaberg T, (2006): Tuberculosis contact investigation with a new, specific blood test in a low-incidence population containing a high proportion of BCG-vaccinated persons. *Respir Res*; 7: 77.

Dinnes J, Deeks J, Kunst H, Gibson A, Cummins E, Waugh N, Drobniewski F, Lalvani A, (2007): "A systematic review of rapid diagnostic tests for the detection of tuberculosis infection". *Health Technol Assess* 11 (3): 1–314.

Dogra S, et al, (2006): Comparison of a whole blood interferon- γ assay with tuberculin skin testing for the detection of tuberculosis infection in hospitalized children in rural India. *J Infect.*

Donia TO, (1976): Planning of National Tuberculosis Programme in Developing Countries *Egy. J. Chest Dis. Tuberc.*, 18 (1)

Dye C, (2006): Global epidemiology of tuberculosis. *Lancet*; 367: 938–40.

Dye C, Scheele S, Dolin P, Pathania V, and Raviglione MC, (1999): Consensus statement: global burden of tuberculosis: estimated incidence, prevalence, and mortality by country. WHO Global Surveillance and Monitoring Project. *JAMA* 282:677-686.

El-Batawi YA, (1981): Manual of microbial infection of man. 4th ed., El-naser Modern Bookshop, Cairo, Part III P. 85.

El-Daly MO, Hafez M, Abuzaid MA, El-Sabah, BS and Wagdan AA, (1998): Tuberculosis in Children in Sharkia. *Egypt, J. Med. Sci.*, 9 (1): 22.

El-shair, O. S.; El-Gazzar, A. G.; Khattab, N. A. and El-Hendy, M. A. (1995):

Clinical evaluation of rapid serodiagnosis of pulmonary tuberculosis by enzyme linked immunosorbent assay(ELISA)

Egy. J. Chest Dis. And Tuberculosis, 44(2):63-72.

Elzinga G, Raviglione M, Maher D, (2004): Scale up: meeting targets in global tuberculosis control. The Lancet, Volume 363, Issue 9411, Pages 814-819.

Ernst JD, Trevejo-Nuñez G, Banaiee N, (2007): "Genomics and the evolution, pathogenesis, and diagnosis of tuberculosis". J. Clin. Invest. 117 (7): 1738–45. Zink A, Sola C, Reischl U, Grabner W, Rastogi N, Wolf H, Nerlich A (2003). "Characterization of Mycobacterium tuberculosis complex DNAs from Egyptian mummies by spoligotyping". J Clin Microbiol 41 (1): 359–67..

Ewer K, Deeks J, Alvarez L, Bryant G, Waller S, Andersen P, Monk P, Lalvani A, (2003): "Comparison of T-cell-based assay with tuberculin skin test for diagnosis of Mycobacterium tuberculosis infection in a school tuberculosis outbreak". Lancet 361 (9364): 1168–73.

Ewer K, Deeks J, Alvarez L, et al, (2006): Comparison of T-cell-based assay with tuberculin skin test for diagnosis of Mycobacterium tuberculosis infection in a school tuberculosis outbreak. Lancet; 361: 1168–73.

Ferrara G, Losi M, D'Amico R, Roversi P, Piro R, Meacci M, Meccugni B, Marchetti Dori I, Andreani A, Bergamini B, et al, (2006): Use in routine clinical practice of two commercial blood tests for diagnosis of infection with Mycobacterium tuberculosis: a prospective study. Lancet; 367:1328–1334.

- Ferrara G, Losi M, Meacci M, et al, (2005):** Routine hospital use of a new commercial whole blood interferon-gamma assay for the diagnosis of tuberculosis infection. *Am J Respir Crit Care Med*; 172: 631–65.
- Fine P, Floyd S, Stanford J, Nkhosa P, Kasunga A, Chaguluka S, Warndorff D, Jenkins P, Yates M, Ponnighaus J, (2001):** "Environmental mycobacteria in northern Malawi: implications for the epidemiology of tuberculosis and leprosy". *Epidemiol Infect* 126 (3): 379–87.
- Fishman AP, (1998):** Microbiology of tuberculosis, pulmonary disease and disorders, 2nd edition, McGraw-HillBook Company, Vol.3, P.1811.
- Fishman AP, Elias JA, Fishman JA, Grippi MA, Kaiser LR and Senior RM, (1998):** Microbiology of tuberculosis. In: Fishman`s pulmonary disease and disorders, 3rd edition, McGraw-Hill, Health Professions Division, New York, San Francisco, London and Madrid. Chapter 159: 2441-2445.
- Franken WPJ, Timmermans JF, Prins C, et al, (2007):** Comparison of Mantoux and QuantiFERON TB Gold Tests for Diagnosis of Latent Tuberculosis Infection in Army Personnel. *Clin Vaccine Immunol*;14:477-480
- French, G.L.; Teoh,R. and Chan, (1987):**
Diagnosis of tuberculous meningitis by detection of tuberculostearic acid in cerebrospinal fluid. *Lancet* II: 117-119.
- Frieden TR, Sterling TR, Munsiff SS, Watt CJ, Dye C, (2003):** Tuberculosis. *Lancet*; 362: 887–99.

- Furie MB and Randolph GJ, (1995):** Chemokines and tissue injury. *Am.J. Pathol.*, 146:1287-1301.
- Gandhi NR, Moll A, Sturm AW, et al, (2006):** Extensively drug-resistant tuberculosis as a cause of death in patients co-infected with tuberculosis and HIV in a rural area of South Africa. *Lancet*; 368: 1575–80.
- Geluk A, van Meijgaarden KE, Franken KL, et al, (2002):** Identification and characterization of the ESAT-6 homologue of *Mycobacterium leprae* and T-cell cross-reactivity with *Mycobacterium tuberculosis*. *Infect Immun*; 70: 2544–8.
- Golden MP and Vikram HR, (2005):** Extrapulmonary Tuberculosis: An Overview. *Am Fam Physician*. 1;72(9):1761-8.
- Goletti D, Vincenti D, Carrara S, et al, (2005):** Selected RD1 peptides for active tuberculosis diagnosis: comparison of a gamma interferon whole-blood enzyme-linked immunosorbent assay and an enzyme-linked immunospot assay. *Clin Diagn Lab Immunol*; 12: 1311–6.
- Graham AW, (2004):** Tuberculosis edited by Madkour MM and Warrell DA, chp.9: The Immunology and Pathogenesis of Tuberculosis, p. 133-152.
- Grange J, (1984):** The mycobacteria. In Topley and Wilson`s principles of bacteriology, virology and immunology, 7th ed., Vol. 2, Edited by M.T.Parker, 1984, Ch.24, p.61.
- Grosset J, (2003):** "Mycobacterium tuberculosis in the extracellular compartment: an underestimated adversary". *Antimicrob Agents Chemother* 47 (3): 833–6.

Grosset J, Mouton Y.(1995).

IS PCR a useful tool for the diagnosis of tuberculosis in Tubercle Lung Dis.; 75:183.

Guide for the national tuberculosis control program. (1990):

Ministry of health, 1st version, Arab Republic of Egypt, p.5.

Gupta R, Espinal MA, and Raviglione MC, (2004):
Tuberculosis as a Major Global Health Problem in the 21st:
Main Issues for 21st Century Tuberculosis Control. World
Health Organization, Geneva, Switzerland. Semin Respir
Crit Care Med. ;25(3). Medscape Today.

Hanna BA, (1996): Diagnosis of tuberculosis by microbiologic
techniques in Tuberculosis 1st ed. (Rom, W.N. and Garay,
S.M.). Little Brown and Company, Boston/ New York/
Toronto/ London, P. 147.

**Hanna, B.A.; Walters, S.B.; Kodosi, S.E.; Stitt, D.T; Tierno,
P.M. and Tick, L.J. (1994):**

Detection of Mycobacterium tuberculosis directly from
patients specimens with Mycobacteria Growth Indicator
Tube; a new rapid method, abstr. C-112, p.510. In Program
and abstracts of 94th General Meeting of the American
Society for Microbiology 1994.

Herrmann J and Lagrange P, (2005): "Dendritic cells and
Mycobacterium tuberculosis: which is the Trojan horse?".
Pathol Biol (Paris) 53 (1): 35–40.

Hill PC, Brookes RH, Fox A, et al, (2004): Large-scale
evaluation of enzyme-linked immunospot assay and skin

test for diagnosis of *Mycobacterium tuberculosis* infection against a gradient of exposure in The Gambia. *Clin Infect Dis*; 38: 966–73.

Hotta K, Ogura T, Nishii K, Kodani T, Onishi M, et al, (2007): Whole Blood Interferon-Gamma Assay for Baseline Tuberculosis Screening among Japanese Healthcare Students. *PLoS ONE* 2(8): e803.

Houben E, Nguyen L, Pieters J, (2006): "Interaction of pathogenic mycobacteria with the host immune system". *Curr Opin Microbiol* 9 (1): 76–85.

Iademarco MF and Castro KG, (2003): "Epidemiology of tuberculosis". *Seminars in respiratory infections* 18 (4): 225–40.

Janeway CA and Travers P, (1996): The immune system in health and disease, 2nd ed. New York, Current biology, Ltd. London/ Garland Publishing.

Jha P, Jacob B, Gajalakshmi V, et al, (2008): "A nationally representative case–control study of smoking and death in India". *N Engl J Med* 358 (11): 1137–1147.

Kang YA, Lee HW, Yoon HI, et al, (2007): Discrepancy between the tuberculin skin test and the whole-blood interferon g assay for the diagnosis of latent tuberculosis infection in an intermediate tuberculosis-burden country. *JAMA*; 293:2756--61.

Kaufmann S, (2002): "Protection against tuberculosis: cytokines, T cells, and macrophages". *Ann Rheum Dis* 61 Suppl 2: ii54–8.

- Kim J, Park Y, Kim Y, Kang S, Shin J, Park I, Choi B, (2003):** "Miliary tuberculosis and acute respiratory distress syndrome". *Int J Tuberc Lung Dis* 7 (4): 359–64.
- Kobashi Y, Mouri K, Yagi S, Obase Y, Miyashita N, Okimoto N, Matsushima T, Kageoka T, Oka M, (2008):** Clinical utility of the QuantiFERON TB-2G test for elderly patients with active tuberculosis. *Chest*. 133(5):1196-202.
- Koch R, (1882):** "Die Aetiologie und die bekämpfung der tuberkulose". *Medical Classic* 2: 853-80. Quoted from *clinical tuberculosis*, edited by ADO davies, 1994, 1: 1-15.
- Kochi A, Vareldzis B and Styblo K, (1994):** Multidrug resistant tuberculosis and its control. *Res. Microbiol.*, 144: 104-110.
- Kumar V, Abbas AK, Fausto N, & Mitchell RN, (2007):** *Robbins Basic Pathology* (8th ed.). Saunders Elsevier. pp. 516-522.
- Lalvani A, (2003):** "Spotting latent infection: the path to better tuberculosis control". *Thorax* 58 (11): 916–8.
- Lalvani A, (2004):** Counting antigen-specific T cells: a new approach for monitoring response to tuberculosis treatment? *Clin Infect Dis*; 38: 757–9.
- Lalvani A, Nagvenkar P, Udwadia Z, Pathan AA, Wilkinson KA, Shastri JS, Ewer K, Hill AV, Mehta A, Rodrigues C, (2001):** Enumeration of T cells specific for RD1-encoded antigens suggests a high prevalence of latent *Mycobacterium tuberculosis* infection in healthy urban Indians. *J Infect Dis*;183:469–477.

Lalvani A, Pathan AA, Durkan H, Wilkinson KA, Whelan A, Deeks JJ, Reece WH, Latif M, Pasvol G, Hill AV, (2001): Enhanced contact tracing and spatial tracking of Mycobacterium tuberculosis infection by enumeration of antigen-specific T cells. *Lancet*;357:2017–2021.

Lalvani A, Pathan AA, McShane H, Wilkinson RJ, Latif M, Conlon CP, Pasvol G, Hill AV, (2001): Rapid detection of Mycobacterium tuberculosis infection by enumeration of antigen-specific T cells. *Am J Respir Crit Care Med*;163:824–828.

Lee E, Holzman RS, (2002): Evolution and current use of the tuberculin test. *Clin Infect Dis*; 34: 365–70.

Lee JY, Choi HJ, Park IN, Hong SB, Oh YM, Lim CM, Lee SD, Koh Y, Kim WS, Kim DS, Kim WD, Shim TS, (2006): Comparison of two commercial interferon gamma assays for diagnosing Mycobacterium tuberculosis infection. *Eur Respir J*;28:24–30.

Ieven M, Goossens H. (1997):
Relevance of nucleic acid amplification techniques for diagnosis of respiratory tract infections in the clinical laboratory. *Clin Micro Rev*; 10: 242

Liebeschuetz S, Bamber S, Ewer K, Deeks J, Pathan AA, Lalvani A, (2004): Diagnosis of tuberculosis in South African children with a T-cell-based assay: a prospective cohort study. *Lancet*; 364: 2196–203.

Lighter J, Rigaud M, Eduardo R, Peng CH, Pollack H, (2009):
Latent tuberculosis diagnosis in children by using the

QuantiFERON-TB Gold In-Tube test. *Pediatrics*; 123(1): 30-7.

Lönnroth K and Raviglione M, (2008): "Global epidemiology of tuberculosis: prospects for control". *Seminars in respiratory and critical care medicine* 29 (5): 481–91.

Madariaga MJ, Jalali Z, and Swindells S, (2007): Evidence-Based Clinical Medicine : Clinical Utility of Interferon Gamma Assay in the Diagnosis of Tuberculosis. *The Journal of the American Board of Family Medicine* 20 (6): 540-547.

Madison B, (2001): "Application of stains in clinical microbiology". *Biotech Histochem* 76 (3): 119–25.

Mahairas GG, Sabo PJ, Hickey MJ, Singh DC, and Stover CK, (1996): Molecular analysis of genetic differences between *Mycobacterium bovis* BCG and virulent *M. bovis*. *J. Bacteriol.* 178:1274-1282.

Martin C, (2006): "Tuberculosis vaccines: past, present and future". *Curr Opin Pulm Med* 12 (3): 186–91.

Mazurek GH, Jereb J, LoBue P, Iademarco MF, Metchock B, Vernon A, (2005): CDC Guidelines for Using the QuantiFERON®-TB Gold Test for Detecting *Mycobacterium tuberculosis* Infection, United States Guidelines for the investigation of contacts of persons with infectious tuberculosis: recommendations from the National Tuberculosis Controllers Association and CDC. *MMWR*; 54(No. RR-15):1-47; 49-55. Division of Tuberculosis Elimination, National Center for HIV, STD, and TB Prevention.

Mazurek GH, LoBue PA, Daley CL, et al, (2001): Comparison of a Whole-Blood Interferon Gamma Assay with Tuberculin Skin Testing for Detecting Latent Mycobacterium tuberculosis Infection. JAMA; 286:1740-1747

Meier T, Eulenbruch HP, Wrighton-Smith P, Enders G, Regnath T, (2005): Sensitivity of a new commercial enzyme-linked immunospot assay (T SPOT-TB) for diagnosis of tuberculosis in clinical practice. Eur J Clin Microbiol Infect Dis; 24: 529–36.

Middlebrook, G. and Dubos, R.J. (1948):

Specific serum agglutination of erythrocytes sensitized with extracts of tubercle bacilli.

J.Exp. Med., 88:521.

Mohsen T, Abou Zeid A, Haj-Yahia S, (2007): Lobectomy or pneumonectomy for multidrug-resistant pulmonary tuberculosis can be performed with acceptable morbidity and mortality: A seven-year review of a single institution's experience. The Journal of Thoracic and Cardiovascular Surgery, Volume 134, Issue 1, Pages 194-198.

Molicotti P, Bua A, Mela G, Olmeo P, Delogu R, Ortu S, Sechi LA, Zanetti S, (2008): Performance of QuantiFERON-TB testing in a tuberculosis outbreak at a primary school. J Pediatr.;152(4):585-6.

Morbidity and mortality weekly report (MMWR), (2003):
"Update: adverse event data and revised American Thoracic

Society/CDC recommendations against the use of rifampin and pyrazinamide for treatment of latent tuberculosis infection- United States". 52 (31): 735–9.

Morbidity and mortality weekly report (MMWR), (2006):
"Emergence of Mycobacterium tuberculosis with extensive resistance to second-line drugs-worldwide, 2000–2004". 55 (11): 301–5.

Morell V, (1994): Mummy settle tuberculosis antiquity depot.

Mori T, Sakatani M, Yamagishi F, et al, (2004): Specific detection of tuberculosis infection: an interferon-gamma-based assay using new antigens. Am J Respir Crit Care Med; 170: 59–64.

Nahid P, Pai M, Hopewell P, (2006): "Advances in the diagnosis and treatment of tuberculosis". Proc Am Thorac Soc 3 (1): 103–10.

National Institute of Allergy and Infectious Diseases (NIAID), (2005): "According to the World Health Organization (WHO), nearly 2 billion people, one-third of the world's population, have TB."

National Tuberculosis Control Programme of Egypt (NTP). (1998): Epidemiology of Tuberculosis. P. 8-12.

Negm, M.F.; El-Gazzar, A.G.; Abdel-Rahman, A.; Abdel-Rahman, S. and El-Desoky, S. (2002):

Role of mycobacterial growth indicator tube (MGIT) in the diagnosis and assessment of drug sensitivity of pulmonary tuberculosis.

M.D. Thesis (Chest diseases) Benha faculty of medicine,
Zagazig University.

Nicas M, Nazaroff WW, Hubbard A, (2005): "Toward understanding the risk of secondary airborne infection: emission of respirable pathogens". J Occup Environ Hyg 2 (3): 143–54.

Niemann S, Richter E, Dalügge-Tamm H, Schlesinger H, Graupner D, Königstein B, Gurath G, Greinert U, Rüsç-Gerdes S, (2000): "Two cases of Mycobacterium microti derived tuberculosis in HIV-negative immunocompetent patients". Emerg Infect Dis 6 (5): 539–42.

Niemann S, Rüsç-Gerdes S, Joloba ML, et al, (2002): "Mycobacterium africanum subtype II is associated with two distinct genotypes and is a major cause of human tuberculosis in Kampala, Uganda". J. Clin. Microbiol. 40 (9): 3398–405.

Niobe-Eyangoh SN, Kuaban C, Sorlin P, et al, (2003): "Genetic biodiversity of Mycobacterium tuberculosis complex strains from patients with pulmonary tuberculosis in Cameroon". J. Clin. Microbiol. 41 (6): 2547–53.

Nnoaham KE and Clarke A, (2008): "Low serum vitamin D levels and tuberculosis: a systematic review and meta-analysis". International journal of epidemiology 37 (1): 113–9.

O'Brien R, (1994): "Drug-resistant tuberculosis: etiology, management and prevention". *Semin Respir Infect* 9 (2): 104–12.

Office for National Statistics, (2006): Notification rates of tuberculosis: by NHS Regional Office area, 1990-2001: *Regional Trends* 37. Rural outbreaks of mycobacterium tuberculosis in a Canadian province.

Onyebujoh P, Onyebujoh R, Graham AW, (2006): World Health Organization Disease Watch: Focus: Tuberculosis.

Orme IM, Roberts AD, Griffin JP, Abrams JS, (1993): Cytokine secretion by CD4 T lymphocytes acquired in response to Mycobacterium tuberculosis infection. *J Immunol*; 151: 518–25.

Pai M and Lewinsohn DM, (2005): Interferon-gamma assays for tuberculosis: is anergy the Achilles' heel? *Am J Respir Crit Care Med*; 172: 519–21.

Pai M, Gokhale K, Joshi R, et al, (2005): Mycobacterium tuberculosis infection in health care workers in rural India: comparison of a whole-blood interferon gamma assay with tuberculin skin testing. *JAMA*; 293: 2746–55.

Pai M, Joshi R, Dogra S, et al, (2006): Persistently elevated T cell interferon-gamma responses after treatment for latent tuberculosis infection among health care workers in India: a preliminary report. *J Occup Med Toxicol*; 1: 7.

Pai M, Joshi R, Dogra S, et al, (2006): Serial testing of health care workers for tuberculosis using interferon-gamma assay. *Am J Respir Crit Care Med*; 174: 349–55.

Pai M, Riley LW, and Colford JM, (2004): Interferon-gamma assays in the immunodiagnosis of tuberculosis: a systematic review. *Lancet Infect. Dis.* 4:761-776.

Pai M, Zwerling A, Menzies D, (2008): Systematic review: T-cell-based assays for the diagnosis of latent tuberculosis infection:an update.*Ann Intern Med.*;149(3):p:177-84;p:1-9.

Palaci, M.; Ueki, S.Y.M.; Sato, D.N. et al (1996):

Evaluation of Mycobacterial growth indicator tube for recovery and drug susceptibility testing of mycobacterium tuberculosis isolates from respiratory specimens.

J.Clin. Microbiol., 34: 762-4.

Parish T and Stoker N, (1999): "Mycobacteria: bugs and bugbears (two steps forward and one step back)". *Mol Biotechnol* 13 (3): 191–200.

Pathan AA, Wilkinson KA, Klenerman P, McShane H, Davidson RN, Pasvol G, Hill AV, Lalvani A, (2001): Direct ex vivo analysis of antigen-specific IFN-gamma-secreting CD4 T cells in Mycobacterium tuberculosis-infected individuals: associations with clinical disease state and effect of treatment. *J Immunol*;167:5217–5225.

Pathan AA, Wilkinson KA, Wilkinson RJ, Latif M, McShane H, Pasvol G, Hill AV, Lalvani A, (2000): High frequencies of circulating IFN-gamma-secreting CD8 cytotoxic T cells specific for a novel MHC class I-restricted Mycobacterium tuberculosis epitope in M. tuberculosis-infected subjects without disease. *Eur J Immunol*;30:2713–2721.

Pearce-Duvet J, (2006)P: "The origin of human pathogens: evaluating the role of agriculture and domestic animals in the evolution of human disease". *Biol Rev Camb Philos Soc* 81 (3): 369–82.

Pottumarthy S, Morris AJ, Harrison AC, Wells VC, (1999): Evaluation of the Tuberculin Gamma Interferon Assay: Potential to Replace the Mantoux Skin Test. *J Clin Microbiol*;37:3229-3232

QuantiFERON[®]-TB Gold (In Tube Method) package insert, (2007). Cellestis limited, www.cellestis.com

Randall, K.H.; and Michael,G.J. (1991):

Bacteriophages, In *Medical Microbiology*, third edition, chapter 5; pp:96-98.

Churchill livingstone. Inc., 1560 broadway, New York, NY 10036

Raok N, Viswanthan R and Sen PK, (1982): Textbook of tuberculosis, the tuberculosis association of India 2nd ed., Viskas publishing House, P.207.

Rashwan AA, (1991): Statistical data analysis for patients follow up in the chest clinic of the new paediatric hospital. M.Sc. Thesis (Paediatrics) Cairo University.

Ravn P, Munk ME, Andersen AB, Lundgren B, Lundgren JD, Nielsen LN, Kok-Jensen A, Andersen P and Weldingh K, (2005): Prospective evaluation of a whole-blood test using Mycobacterium tuberculosis-specific antigens ESAT-6 and CFP-10 for diagnosis of active tuberculosis. *Clin. Diagn. Lab. Immunol.* 12:491-496.

- Reddy JR, Kwang J, Lechtenberg KF, Khan NC, Prasad RB, Chengappa MM, (2002):** "An immunochromatographic serological assay for the diagnosis of Mycobacterium tuberculosis". *Comp. Immunol. Microbiol. Infect. Dis.* 25 (1): 21–7.
- Restrepo BI, (2007):** "Convergence of the tuberculosis and diabetes epidemics: renewal of old acquaintances". *Clin. Infect. Dis.* 45 (4): 436–8.
- Richeldi L, (2006):** An update on the diagnosis of tuberculosis infection. *Am J Respir Crit Care Med*; 174: 736–42.
- Richeldi L, Ewer K, Losi M, Bergamini BM, Roversi P, Deeks J, Fabbri LM, and Lalvani A, (2004):** T cell-based tracking of multidrug resistant tuberculosis infection after brief exposure. *Am. J. Respir. Crit. Care Med.* 170:288-295.
- Richeldi L, Ewer K, Losi M, Hansell DM, Roversi P, Fabbri LM, Lalvani A, (2004):** Early diagnosis of subclinical multidrug-resistant tuberculosis. *Ann Intern Med*; 140:709–713.
- Rossman DM and Mayock LR, (1995):** Pulmonary tuberculosis
In: Rossman, D.M. and MacGregor R.R. (eds). *Tuberculosis Clinical Management and New Challenges*, 1st edition, New York, MacGraw Hill Press, ch.9. P.145-154.
- Rothel J and Andersen P, (2005):** "Diagnosis of latent Mycobacterium tuberculosis infection: is the demise of the Mantoux test imminent?". *Expert Rev Anti Infect Ther* 3 (6): 981–93.

Salem, E.E.; Sobhy, K.E.; Ali, G.; El-Ansary, M. and Fanous, N.V. (1990):

HLA-phenotyping in adult tuberculosis patients.

Am. Rev. Resp. Dis., 14(4): 792.

Salem ES, Al-Helaly AH, Abdel-Rahman H, Sharaf Eldin M, Negm M and Okab A, (1992): Annual tuberculosis infection rate in Qualiubia Governorate. Medical J. Cairo University, 60 (1): 55.

Schaible UE and Kaufmann SH, (2007): "Malnutrition and infection: complex mechanisms and global impacts". PLoS medicine 4 (5): e115.

Science, 263:1686-1687.

Shams H, Weis SE, Klucar P, Lalvani A, Moonan PK, Pogoda JM, Ewer K, Barnes PF, (2005): Enzyme-linked immunospot and tuberculin skin testing to detect latent tuberculosis infection. Am J Respir Crit Care Med;172:1161–1168.

Smith T, (1898): A comparative study of mycobacterium bovis T.B. from sputum. T-exper. Med. 3: 451. Quated from Clinical Tuberculosis, edited by P.D.O. Davis, 1994, 1: 1-15.

Sobero R and Peabody J, (2006): "Tuberculosis control in Bolivia, Chile, Colombia and Peru: why does incidence vary so much between neighbors?". Int J Tuberc Lung Dis 10 (11): 1292–5.

Somoskovi, A.; Magyar, P. (1999):

Comparison of the Mycobacteria Growth Indicator Tube with MB redox, Lowenstein-Jensen, and Midllbrook 7H11 media for recovery mycobacteria in clinical specimens.

J. Clin. Microbiol., 37(5): 1366-9.

Sorensen AL, Nagai S, Houen G, Andersen P and Andersen AB, (1995): Purification and characterization of a low-molecular-mass T-cell antigen secreted by Mycobacterium tuberculosis. Infect. Immun. 63:1710-1717.

Soysal A, Millington KA, Bakir M, Dosanjh D, Aslan Y, Deeks JJ, Efe S, Staveley I, Ewer K, Lalvani A, (2005): Effect of BCG vaccination on risk of Mycobacterium tuberculosis infection in children with household tuberculosis contact: a prospective community-based study. Lancet;366:1443–1451.

Spence DP, Hotchkiss J, Williams CS, Davies PD, (1993): "Tuberculosis and poverty". BMJ (Clinical research ed.) 307 (6907): 759–61.

Spitznagel J,(1989):Mycobacterium tuberculosis and leprosy. Chap. 22, p.303-19 cited from schaechter, M.; Medoff, G. and Schlessinger, D., 1989, Mechanisms of microbial disease. Puble. William and wilkin: 428 Eastt preston st., Baltimore MD 21202, U.S.A.

Stitt, D.T.; Parks, T.L.;Mellarkey,J.C. and Tanner,J.E. (1994):

A rapid non radiometric method for detection of the growth of mycobacteria. P 159. In program and abstracts of the6th

European Congress of Clinical Microbiology and infectious Diseases. European Society of Clinical Microbiology and infectious Diseases. Seville, Spain.

Strachan DP, Powell KJ, Thaker A, Millard FJ, Maxwell JD, (1995-02): "Vegetarian diet as a risk factor for tuberculosis in immigrant south London Asians". *Thorax*; 50 (2): 175–80.

Streeton JA, Desem N, Jones SL, (1998): Sensitivity and specificity of a gamma interferon blood test for tuberculosis infection. *Int J Tuberc Lung Dis*; 2: 443–50.

Styblo K, (1976): Epidemiological Basis of Tuberculosis Control. In: Donia, T.O., Recent trends in tuberculosis control, 1976, 222.

Sutherland R, Yang H, Scriba TJ, et al, (2006): Impaired IFN-gamma-secreting capacity in mycobacterial antigen-specific CD4 T cells during chronic HIV-1 infection despite long-term HAART. *AIDS*; 20: 821–9.

Swierzewski SJ, (2007): pulmonologychannel.com, Tuberculosis, Types.

Taylor JP, Vitek I, Enriquez V, Smedley JW, (1999): A continuing focus of Hansen's disease in Texas. *Am J Trop Med Hyg*; 60: 449–52.

Thoen C, Lobue P, de Kantor I, (2006): "The importance of *Mycobacterium bovis* as a zoonosis". *Vet. Microbiol.* 112 (2-4): 339–45.

Topley WW and Wilson GS, (1990): Principles of bacteriology, virology and immunity, 8th. Arnold London. Vol. (1): 76,

Vol.(2): 74, Vol.(3): 105.

Trinchieri G, (1997): Cytokines acting on or secreted by macrophages during intracellular infection (IL-10, IL-12, IFN γ). *Curropin. Immuno.*, 9: 17-23.

Tully G, Kortsik C, Hohn H, Zehbe I, Hitzler WE, Neukirch C, Freitag K, Kayser K, Maeurer MJ, (2005): Highly focused T cell responses in latent human pulmonary *Mycobacterium tuberculosis* infection. *J Immunol*;174:2174–2184.

Ustianowski A, Shaffer R, Collin S, Wilkinson RJ, Davidson RN, (2005): "Prevalence and associations of vitamin D deficiency in foreign-born persons with tuberculosis in London". *The Journal of infection* 50 (5): 432–7.

Van Pittius NCG, Gamiieldien J, Hide W, Brown GD, Siezen RJ, Beyers AD, (2001): The ESAT-6 gene cluster of *Mycobacterium tuberculosis* and other high G+C Gram-positive bacteria. *Genome Biol*; 2: 44.

Van-Pinxteren LA, Ravn P, Agger EM, Pollock J and Andersen P, (2000): Diagnosis of tuberculosis based on the two specific antigens ESAT-6 and CFP10. *Clin. Diagn. Lab. Immunol.* 7:155-160.

Vooren, J.P.; Farber, B.M. and Serruys, E. (1990):

Immunological detection of mycobacterial antigens in infected fluid, cells and tissues by latex agglutination.

Am. Rev. Respir. Dis., 14 (4):805.

Welch, D.A.; Sides, G.S.; Shaw, C. and Gilchrist, M. (1993):

Timely culture of mycobacteria which utilizes
a microcolony method.

J. Clin. Microbiol., 31: 2178-84.

Wikipedia, the free encyclopedia, (2008): "How T-SPOT.TB
Works". Tuberculosis diagnosis.

**Wilkinson KA, Wilkinson RJ, Pathan A, Ewer K, Prakash M,
Klenerman P, Maskell N, Davies R, Pasvol G and
Lalvani A, (2005):** Ex vivo characterization of early
secretory antigenic target 6-specific T cells at sites of active
disease in pleural tuberculosis. Clin. Infect. Dis. 40:184-187.

World Health Organization(WHO) report, (2006): Stop TB
Partnership.

World Health Organization(WHO) report, (2006):
Tuberculosis Fact sheet N°104 - Global and regional
incidence.

World Health Organization(WHO) Report, (2006): Global
Tuberculosis Control Report, - Annex 1 Profiles of high-
burden countries.

World Health Organization(WHO) Report, (2006): Global
tuberculosis control - surveillance, planning, financing.

World Health Organization (WHO) report, (2007): Global
tuberculosis control: surveillance, planning, financing.

World Health Organization; Geneva,
(WHO/HTM/TB/2007.376).

Yamamoto Y, Saito H, Tomioka H, Sato K, Yamane T, Yamashita K, Hosoe K and Hidaka T, (1992): In vitro and in vivo activities of KRM 1648, a newly synthesized benzoxazinorifamycin, against mycobacterium marinum. Zentralbl, Bakteriologie, 554:625-631.

Yi, S.; Yan, S. and Yan, Y. (1990):

Detection of serum IgG antibody to PPD in tuberculous children.

Am. Rev. Respir. Dis., 141 (4): 788.