

REFERENCES

A

Abdulla, S.M.; Hassan, F.B.; Mohamed, E.K. and Mahmoud H. (2009): Gram stain-acridine orange leukocyte cytospin test for rapid diagnosis of central venous catheter-related blood stream infection. Thesis submitted for practical fulfillment of master degree in clinical and chemical pathology. Faculty of medicine Ain Shams University.

Andreumont, A.; Paulet, R.; Nitenberg, G. and Hill, C. (1988): Value of semi quantitative cultures of blood drawn through catheter hubs for estimating the risk of catheter tip colonization in cancer patients. Service de microbiologie medicale, unite de Reanimation and Department de statistique medicale. France. Antimicrob. Agents Chemother. 36: 1208-14.

Anwar, H.; Strap, J.L. and Chen, K. (1992): Dynamic interactions of biofilms of mucoid *Pseudomonas aeruginosa* with tobramycin and piperacillin.

Appelgren, P.; Hellstroem, I.; Weitzberg, E.; Soderlund, V. And Bindsvlev, L. (2001): Risk factor for nosocomial intensive care infection: a longterm prospective analysis. Acta. Anaesthesiol. Scand. 45:710-719.

Assadian, O. (2004): Skin antiseptic in reducing the risk of central venous catheter-related infections. Crit. Care Med. 32:887-888.

**B**

Baron, E. J.; Peterson, L.R.; and Finegold, S.M. (1994): Microorganisms encountered in the blood In: Bailey & Scott's Diagnostic Microbiology, ed 9. Mobsy, Missouri. P: 193-208.

Bassetti, S.; Hu, J.; D'Agostino, R.B. Jr and Sherertz, R.J. (2001): Prolonged antimicrobial activity of a catheter containing chlorhexidine- silver sulfadiazine extends protection against catheter infections in vivo. Antimicrob Agents Chemother; 45:1535-1538.

Baumgartner, T.G.; Schmidt, G.L. and Thakker, K.M. (1986): Bacterial endotoxin retention by inline intravenous filters. Am J Hosp Pharm; 43:681-684.

Beathard, G.A. (1999): Management of bacteremia associated with tunneledcuffed hemodialysis catheters. J Am Soc Nephrol; 10: 1045-1`049.

Bloemberg, G.V.; O'Toole, G.A.; Lugtenberg, B.J. and Kolter, R. (1997): Green fluorescent protein as a marker for Pseudomonas spp. Appl. Environ. Microbiol. 63: 4543-4551.

Blot, F.; Nitenbert, G. and Brun- Buisson, C. (2000): New tools in diagnosing catheter- related infections. Support care cancer; 8: 287-292.

Blot, F.; Schmidt, E.; Nitenberg, G. And et al., (1998): Earlier positivity of central venous versus peripheral cultures is highly predictive of catheter- related sepsis. J. Clin. Microbiol.; 36:105–109.



- Bonawitz, S.C.; Hammell, E.J. and Kirkpatrick, J.R. (1991):** Prevention of central venous catheter sepsis: a prospective randomized trial. *Am Surg*; 57: 618-623.
- Bong, J.; Kite, P.; Ammori, B. And et al. (2003):** The use of a rapid in situ test in the detection of central venous catheter-related bloodstream infection: A prospective study. *J. Parenter. Enteral Nutr.*; 27(2):146-150.
- Boraks, P.; Seale, J. and Price, J. (1998):** Prevention of central venous catheter associated thrombosis using minidose warfarin in patients with haematological malignancies. *Br J Haematol*; 101: 483-486.
- Bouza, E.; Burillo, A. and Munoz, P. (2002):** Catheter- related infections: diagnosis and intravascular treatment. *Clin. Microbial. Infect.* 8:265-274.
- Brian, A.; Lauer, L.; Barth, R. and Stanley, M. (1981):** Comparison of acridine orange and gram stains for detection of micro organisms in cerebrospinal fluid and other clinical specimens.
- Brismar, B.; Jordahl, L.; Nystrom, B.; and Petterson, N. (1982):** Bacterial contamination of intravenous line side ports of different designs. *Clin. Nutr.*; 6:31-40.
- Brun- Buisson, C.; Abrar, F.; Legrand, P.; Huet, Y.; Larabi, S. and Rapin, M. (1987):** Diagnosis of central venous catheter related sepsis. Critical level of quantitative tip cultures. *Arch. Intern. Med.* 147: 873-877.



Brun-Buisson, C.; Doyon, F. and Carlet, J. (1996): Bacteremia and severe sepsis in adults: a multicenter prospective survey in ICUs and wards of 24 hospitals. *Am J Respir Crit Care Med*; 154:617-624.

Burke, (1922): Notes on Gram stain with description of a new method. *J. of Bacteriology* 7:159.

Butler, D.L.; Munson, J.M. and DeLuca, P.P. (1980): Effect of inline filtration on the potency of low- dose drugs. *Am J Hosp Pharm*; 37: 935-941.

C

Cappelli, G.; Tetta, C. and Banaud, B. (2005): Is biofilm a cause of silent chronic inflammation in haemodialysis patients? A fascinating working hypothesis. *Nephrol Dial Transplant*; 19: 1-4.

Casewell, M.W. (1998): The nose: an underestimated source of staphylococcus aureus causing wound infection. *J Hosp Infect*; 40 (suppl): S 3- S11.

Catton, J.A.; Dobbins, B.M.; Kite, P.; and Wood, J.M.; et al., (2005): In situ diagnosis of intravascular catheter related blood stream infection: A comparison of quantitative culture, differential time to positivity, and endoluminal brushing. *Crit Care Med*. 33; 787-791.



Centers for Disease Control and Prevention (1995): Recommendations for preventing the spread of vancomycin resistance. Recommendations of the Hospital Infection Control Practices Advisory Committee (HICPAC). MMWR Recomm Rep; 44, RR 12) 1-13.

Centers for Disease Control and Prevention (1996): CDC definitions of nosocomial infections. APIC Infection Control and Applied Epidemiology: Principles and Practice; A-1-A-20.

Centers for Disease control and Prevention (2002): Guide lines for the prevention of intravascular catheter related infections. MMWR; 51(No.RR-10):1-30.

Chan, D.C.W.; Chan, H.B. and Ip L.S.(2005):Occasional survey diagnosis and treatment of catheter-related blood stream infection in neonates and childrens. HK J. Paediatr.;10:26-33.

Chan, D.C.W.; Chan, H.B. and Ip L.S. Chatzinikolaou, L. and Raad, I. (2006): Catheter- related infections Iranian Journal of Clinical Infectious Disease 1:5-10.

Chatzinikolaou, L. and Raad, I. (2006): catheter related infection Iranian Journal of clinical Infectious Disease 1:5-10.

Cheesbrough, M.(2000): Culturing blood. In: District laboratory practice in Tropical countries. Part 2. Cambridge university press p: 124-130.

Cicalini, S.; Palmieri, F. and Noto, P. (2002): Diagnosis of intra vascular catheter- related infection. J Vasc Access; 3:114-119.



- Cicalini, S.; Palmieri, F. and Petrosillo, N. (2004):** New technologies for prevention of intravascular catheter- related infections. Crit Care; 8 (3): 157-162.
- Clemence, M.A.; Walker, D. and Farr, B.M. (1995):** Central venous catheter practices: results of a survey. Am J Infect Control; 23:5-12.
- Cleri, D.J.; Corrado, M.L. AND Seligmen, S.J. (1980):** Quantitive culture of intravenous catheter and other insert. J Infect 141; 781-786.
- Cobb, D.K.; High, K.P. and Sawyer, R.G. (1992):** A controlled trial of scheduled replacement of central venous and pulmonary- artery catheters. N Engl J Med; 327:1062-1068.
- Coello, R.; Charlett, A.; Ward, V. and et al., (2003):** Device-related sources of bacteraemia in English hospitals-opportunities for the prevention of hospital-acquierd bacteraemia, The Journal of Hospital Infection; 53 : 46–57.
- Cook, D.; Randolph, A. and Kernerman, P. (1997):** Central venous catheter replacement strategies: a systematic review of the literature. Crit Care Med; 25: 1417-1424.
- Costerton, B. and Lang, E. (1998):** Center for Biofilm Engineering (CBE), Montana State University, www.erc.montana.edu.
- Costerton, J.W. (1999):** Introduction to biofilm. Int J Antimicrob agents; 11: 217-221.



Costerton, W.; Veeh, R.; Shirtliff, M.; Pasmore, M.; Post, C. and Ehrlich, G. (2003): The application of biofilm science to the study and control of chronic bacterial infections. J Clin Invest.; 112: 1466-1477.

Culpepper, C. and Allison, D. (2004): Intravascular Catheter Infections.
Available at:
https://www.utmem.edu/obgyn/student_pres/Peripheral_Venous_Accessel.ppt

D

Darouiche, R.O.; Raad, I.I. and Heard, S.O. (1999): A comparison of two antimicrobial- impregnated central venous catheters. Catheter study Group. N Engl J Med; 340:1-8.

Digiovine, B.; Chenoweth, C. and Watts, C. (1999): The attributable mortality and costs of primary nosocomial bloodstream infections in the intensive care unit. Am J Respir Crit Care Med; 160:976-981.

Dimick, J.B.; Swoboda, S.; Talamini, M.A. And et al. (2003): Colonization of Central Venous Catheters: Catheters for Total Parenteral Nutrition Vs Other Catheters. American Journal of Critical Care; 12: 328-335.



Dimick, J.B.; Swoboda, S.; Talamini, M.A.; Pelz, R.K.; Hendrix, C.W. and Lipsett, P.A. (2003): Risk of colonization of central venous catheters: catheters for total parenteral nutrition vs other catheters. *American Journal of Critical Care*, 12: 328-335.

Donalan, R.M. (2002): Biofilms: microbial life on surfaces. *Emerg. Infect. Dis.*; 8:881-890.

Donalan, R.M. and Costerton, J.W. (2002): Biofilms: survival mechanisms of clinically relevant microorganisms. *Clin. Microbiol. Rev.*; 15:167-193.

Donlan, R.M. (2001) a: Biofilms and device- associated infections. *Emer. Infect. Dis.* 7:277-281.

Donlan, R.M. (2001) b: Biofilm formation: a clinically relevant microbiological process. *Clin. Infect. Dis.*; 33:1387-1392.

Donlan, R.M. and Costerton, J.W. (2002): Biofilms: survival mechanisms of clinically relevant microorganisms. *Clin Microbiol Rev*; 15: 167-193.

E

Eggimann, P. and Pittet, D. (2001): Infection Control in the ICU. *Chest*; 120: 2059-2093.

Eggimann, P. and Pittet, D. (2002): Overview of catheter- related infections with special emphasis on prevention based on educational programs *CMI* 8; 295-309.



Eggimann, P.; Harbarth, S.; Constantin, M.N.; Touveneau, S.; Chevrolet, J.C. and Pittet, D. (2000): Impact of a prevention strategy targeted at vascular access care on incidence of infections acquired in intensive care. *Lancet*; 335: 1864-1868.

El- Mahallawy, H.; Sidhom, I.; Ali El-Din, N.H.; Zam Zam, M. and El-Lamie, M.M. (2005): Clinical and microbiologic determinants of serious blood stream infection in Egyptian Pediatric Cancer patients: a one year study. *International Journal of infectious disease* 9; 43-51.

Elliott, T.S.J.; O'Grady, N.P. and Pittet, D. (2004): The management and treatment of intravascular catheter-related infections in the critically ill. Kluwer Academic publishers Group, The Netherlands ; 5:113-126.

Eyer, S.; Brummitt, C.; Crossley, K.; Siegel, R. and Cerra, F. (1990) : Catheter- related sepsis: prospective, randomized study of three methods of long- term catheter maintenance. *Crit Care Med*; 18: 1073-1079.

F

Farr, B.M. (2004): Catheters, microbes, time, and gold standards. *Ann Intern Med*; 140:62-64.

Fatkenheuer, G.; Cornely, O. and Seifert, H. (2002): Clinical management of catheter- related infections. *Clin. Microbiol. Infect.* 8: 545-50.



- Flemming, H.C.; Wingender, J. and Mayer, C. (2000):** Physico-chemical properties of biofilms. In: Evans, L.V. (ed.) Biofilms: recent advances in their study and control. Amsterdam: Harwood Academic Publishers; p: 19-34.
- Fowler, V.G.; Jr, Li, J. and Corey, C.R. (1997):** Role of echocardiography in evaluation of patients with staphylococcus aureus bacteremia: experience in 103 patients. J Am Coll Cardiol; 30:1072-1078.
- Fridkin, S.K.; Pear, S.M.; Williamson, T.H.; Galgiani, J.N. and Jarvis, W.R. (1996):** The role of understaffing in central venous catheter-associated bloodstream infections. Infect control Hosp Epidemiol; 17:150-158.
- Friedman, N.B.; Kaye, K.S. and Stout, J.E. (2002):** Health care-associated bloodstream infections in adults. Ann. Intern. Med. 137: 791-797.
- Furfaro, S.; Gauthier, M.; Lacroix, J. And et al. (1991):** Arterial catheter-related infections in children. Am. J. Dis. Child; 145:1037–1042.
- Fux, C.A.; Wilson, S. and Stoodley, P. (2004):** Detachment: characteristics and oxacillin resistance of Staphylococcus aureus biofilm emboli in an in vitro catheter infection model. J Bacteriol; 186: 4486-4491.



G

- Garland, J.S.; Buck, R.K. and Maloney, P. (1995):** Comparison of 10% povidone- iodine and 0.5% chlorhexidine gluconate for the prevention of peripheral intravenous catheter colonization in neonates: a prospective trial. *Pediatr infect Dis J*; 14:510-516.
- Garner, J.S.; Emori, T.G. and Hughes, J.M. (2005):** Definition of central line. From CDC National Nosocomial Infections Surveillance (NNIS) System. Available at: <http://www.jenympro.org/showthread.php?t=350>.
- Garner, J.S.; Jarvis, W.R.; Emori, T.G.; Horan, T.C. and Huges, T.M. (1988):** CDC definition for nosocomial infections. *Am. J. Infect. Control*. 16: 128-140.
- Garner, J.S.; Jarvis, W.R.; Emori, T.G.; Horan, T.C. and Huges, T.M. (1996):** CDC definition for nosocomial infections. In: Olmsted, R.N. (ed.) *Infection Control and Applied Epidemiology: Principle and Practice*. St Louis, MO: Mosby, P: 1-20.
- Gast, P.; Hagley, M.; Martin, B. And et al. (1992):** Infectious and mechanical complications of central venous catheters placed by percutaneous venipuncture and over guide wires. *Crit. Care Med.*; 20:1426–1430.
- Gastmeier, P.; Weist, K. and Ruden, H. (1999):** Catheter- related primary blood stream infection: Epidemiology and methods of prevention, *Infection* 27; 51-56.



- Gaynes, R. (2002):** Health care- associated bloodstream infections: a change in thinking. *Ann. Intern. Med.* 137: 850-851.
- Gilbert, P.; Maira-Litran, T.; McBain, A.J.; Rickard, A.H. and Whyte, F.W. (2002):** The physiology and collective recalcitrance of microbial biofilm communities. *Adv Microbial Physiol*; 46: 203-256.
- Goetz, A.M.; Wagener, M.M.; Miller, J.M. and Muder, R.R. (1998):** Risk of infection due to central venous catheters. Effect of site of placement and catheter type. *Infect Control Hosp Epidemiol*; 19: 843-845.
- Goldmanni, D.A. and Pier, G.B. (1993):** Pathogenesis of infections related to intravascular catheterization. *Clin. Microbiol. Rev.* 6: 176-192.
- Goldstein, A.M.; Weber, J.M. and Sheridan, R.L. (1997):** Femoral venous access is safe in burned children: an analysis of 224 catheters. *J Pediatr*; 130-442-446.
- Gowardman, J.; Montgomery, C.; Thirlwell, S. And et al. (1998):** Central venous catheter-related bloodstream infections: an analysis of incidence and risk factors in a cohort of 400 patients. *Intensive care med.*; 24:1034-1039.
- Grohskopf, L.A.; Roth, V.R. and Feikin, D.R. (2001):** *Serratia liquefaciens* bloodstream infections from contamination of epoetin alfa at a hemodialysis center. *N Engl J Med*; 344: 1491-1497.



Gupta, S.; Batra, Y.K.; Puri, G.D; And et al. (1995): Infection rates in single- and double-lumen central venous catheters in critically ill patients. Natl. Med. J. India; 8:114–7.

H

Hall, K. and Farr, B. (2004): Diaganosis and management of longterm central venous catheter infections. J Vasc Interv Radiol; 15: 327-334.

Hanna, H.A. and Raad, I.I. (2001): Blood products: a significant risk factor for longterm catheter- related bloodstream infections in cancer patients. Infect Control Hosp Epidemiol; 22: 165-166.

Heard, S.; Wagle, M.; Vijayakumar, E. And et al., (1998): Influence of triple lumen central venous catheters coated with chlorhexidine and silver sulfadiazine on the incidence of catheter related bacteremia. Arch. Intern. Med.; 158:81-87.

Heard, S.O.; Wagle, M. and Vijayakumar, E. (1998): Influence of triple-lumen central venous catheters coated with chlorhexidine and silver sulfadiazine on the incidence of catheter- related bacteremia. Arch Intern Med; 158:81-87.

Henrickson, K.J.; Axtell, R.A. and Hoover, S.M. (2000): Prevention of central venous catheter- related infections and thrombotic events in immunocompromised children by the use of vancomycin/ciprofloxacin/ heparin flush solution: a randomized, multicenter, double- blind trial. J Clin Oncol; 18: 1269-1278.



Henry, B.; Plante- Jenkias, C. and Ostrowska, K. (2001): An outbreak of *serratia marcescens* associated with the anesthetic agent propofol. *Am J Infect Control*; 29: 312-315.

Hodge, D.; and Puntis, J.(2002): Diagnosis, prevention, and management of catheter related bloodstream infection during long term parenteral nutrition. *Arch. Dis. Child Fetal Neonatal Ed.*; 87:21-24.

Hoiby, N.; Johansen, H.K. and Moser, C. (2001): *Pseudomonas aeruginosa* and the in vitro and in vivo biofilm mode of growth . *Microbes Infect*; 3:23-35.

Holmes, A.; Dore, C.J.; Saraswatula, A.; Bamford K.B.; et al., (2008): Risk factors and recommendations for not stratification for surveillance of neonatal health care- associated blood stream infection. *Journal of hospital infection*; 68: 66-72.

Humar, A.; Ostromecki, A. and Drenfeld, J. (2000): Prospective randomized trial of 10% povidone- iodine versus 0.5% tincture of chlorhexidine as cutaneous antisepsis for prevention of central venous catheter infection. *Clin infect Dis*; 31: 1001-1007.

I

Inoue, Y.; Nezu, R. and Mastuda, H. (1992): Prevention of catheter-related sepsis during parenteral nutrition: effect of a new connection device. *J Parenter Enteral Nutr*; 16: 581-585.



J

Jeremy, M.Y.; Douglas, J.B.; Esther, M.V. and Greenberg, E.P. (2004): Quorum sensing in *staphylococcus aureus* biofilms. J. bacterial. 186:1838-1850.

Josephson, A.; Gombert, M.E.; Sierra, M.F.; Karanfil, L.V. and Tansino, G.F. (1985): The relationship between intravenous fluid contamination and the frequency of tubing replacement. Infect control; 6:367-370.

Joynt, G.M.; Kew, J.; Gomersail, C.D.; Leung, V.Y. and Liu, E.K. (2000): Deep venous thrombosis caused by femoral venous catheters in critically ill adult patients. Chest; 117-178-183.

K

Karchmer, A.W. (2000): Nosocomial bloodstream infections: organisms, risk factors, and implications. Clin. Infect. Dis. 31: 139-143.

Keren, I.; Shah, D.; Spoering, A.; Kaldalu, N. And Lewis, K. (2004): Specialized persister cells and the mechanism of multidrug tolerance in *Escherichia coli*. J Bacteriol; 186:8172-8180.

Kite, P.; Dobbins, B.; Wilcox, M. et al. (1997): Evaluation of a novel endoluminal brush method for in situ diagnosis of catheter related sepsis. J. clin. Pathol.; 50:278-82.

Kite, P.; Dobbins, B.M.; Wilcox, M.K. and McMahon, M.J. (1999): Rapid diagnosis of central venous catheter-related bloodstream infection without catheter removal. Lancet; 354: 1504-1507.



Kite, P.; Eastwood, K.; Sugden, S. and Percival, S.L. (2004): Use of in vivo- generated biofilms from hemodialysis catheters to test the efficacy of a novel antimicrobial catheter lock for biofilm eradication in vitro. J. Clin. Microbiol. 42: 3073-3076.

L

Lai, K.K. (1998): Safety of prolonging peripheral cannula and i.v. tubing use from 72 hours to 96 hours. Am J infect Control; 26:66-70.

Larson, E.L.; Rackoff, W.R. and Weiman, M. (1995): APIC guideline for handwashing and hand antisepsis in health care settings. Am J Infect control; 23: 251-269.

Lauer, B.A.; Reller, L. B. and Mirrett, S. (1981): Comparison of acridine orange and Gram stains for detection of microorganisms in cerebrospinal fluid and other clinical specimens. J. Clin. Microbiol. 14:201-205.

Leid, J.G.; Shirtliff, M.E.; Costerton, J.W. and Stoodley, A.P. (2002): Human leukocytes adhere to, penetrate, and respond to *Staphylococcus aureus* biofilms. Infect Immun; 70: 6339-6345.

Levin, A.; Mason, A.J.; Jindal, K.K.; Fong, I.W. and Goldstein, M.B. (1991): Prevention of hemodialysis subclavian vein catheter infections by topical povidone- iodine. Kidney Int; 40: 934-938.

Lewis, K. (2001): Riddle of biofilm resistance Antimicrob Agents. Chemother; 45:999-1007.



- Linares, J.; Sitges-Serra, A.; Garau, J.; Perez, J.L. and Martin, R. (1985):** Pathogenesis of catheter sepsis: a prospective study with quantitative and semiquantitative cultures of catheter hub and segments. *J. Clin Microbiol*; 21:357-360.
- Ljungman, P.; Hagglund, H.; Bjorkstrand, B.; Lonnqvist, B. and Ringden, O. (1997):** Perioperative teicoplanin for prevention of gram positive infections in neutropenic patients with indwelling central venous catheters: a randomized, controlled study. *Support Care Cancer*; 5:485-488.
- Longfiled, R.N.; Smith L.P.; Longfiled, J.N.; Coberly, J and Cruess, D.(1985):** Multiple-dose vials: persistence of bacterial contaminants and infection control implications. *Infect Control*;6:194-199.
- Lorente, L.; Henry, C.; Martín, M.M. And et al., (2005):** Central venous catheter-related infection in a prospective and observational of 2,595 catheters. *Crit. Care*; 9(6):631-635.
- Lorente, L.; Villegas, J.; Martín, M. And et al. (2004):** Catheter- related infection in critically ill patients. *Journal of Intensive Care Medicine*; 30(8):1681-1684.
- Lowy, F.D. (1998):** Staphylococcus aureus infection. *N Engl J Med*; 339:520-532.
- Lugauer, S.; Regenfus, A.; Boswald, M. and Martus, P. et al., (1999):** A new scoring system for the clinical diagnosis of catheter- related infection. *Infection* 27 S 49- S 53.



M

- Maddox, R.R.; John, J.F.; Jr, Brown, L.L. and Smith, C.E. (1983):** Effect of inline filtration on postinfusion phlebitis. Clin Pharm; 2: 58-61.
- Maki, D.G. and Band, J.D. (1981):** A comparative study of polyantibiotic and iodophor ointments in prevention of vascular catheter-related infection. Am J Med; 70 :739-744.
- Maki, D.G. and Crinch, J. (2003):** Line sepsis in the IC4: Prevention, diagnosis and management, semin Respir crit care Med 24; 23-36.
- Maki, D.G. and Mermel, L.A. (1991):** Infections due to infusion therapy. In: Bennett, J.V. and Brachman, P.S. (eds). Hospital Infections. 4th ed. Philadelphia: Lippencott- Raven, P: 689-724.
- Maki, D.G. and Ringer, M. (1987):** Evaluation of dressing regimens for prevention of infection with peripheral intravenous catheters. J. Am. Med. Assoc. 258: 2396-2403.
- Maki, D.G. and Ringer, M. (1991):** Risk factors for infusion-related phlebitis with small peripheral venous catheters: a randomized controlled trial. Ann intern Med; 114:845-854.
- Maki, D.G.; Botticelli J.T.; Leroy, M.L. and Thielke, T.S. (1987):** Prospective study of replacing administration sets for intravenous therapy at 48-vs 72-hour intervals: 72 hours is safe and cost-effective. JAMA; 258:1777-1781.



- Maki, D.G.; Cobb, L.; Garman, J.K.; Shapiro, J.M.; Ringer, M. and Helgerson, R.B. (1988):** An attachable silver- impregnated cuff for prevention of infection with central venous catheters: a prospective randomized multicenter trial. *Am J Med*; 85: 307-314.
- Maki, D.G.; Goldman, D.A. and Rhame, F.S. (1973):** Infection control in intravenous therapy, *Annals of Internal Medicine*; **79**: 867–887.
- Maki, D.G.; Mermel, L.A. and Klugar, D. (2000):** The efficacy of a chlorhexidine impregnated sponge (Biopatch) for the prevention of intravascular catheter- related infection-a prospective randomized controlled multicenter study [Abstract]. Presented at the Interscience Conference on Antimicrobial Agents and Chemotherapy. Toronto, Ontario, Canada: American Society for Microbiology.
- Maki, D.G.; Ringer, M. and Alvarado, C.J. (1991):** Prospective randomized trial of povidone- iodine, alcohol, and chlorhexidine for prevention of infection associated with central venous and arterial catheters. *Lancet* 338: 339-43.
- Maki, D.G.; Stolz, S.M.; Wheeler, S. and Mermel, L.A. (1997):** Prevention of central of central venous catheter-related bloodstream infection by use of an antiseptic-impregnated catheter. A randomized, controlled trial. *Am. Intern Med*; 127:257-266.
- Maki, D.G.; Weise, C. and Sarafin, H. (1977):** A semiquantitative culture method for identifying intravenous- catheter- related infection. *N. Engl. J. Med.* 296: 1305-1309.



- Martinez, J.A.; DesJardin, J.A.; Aronoff, M. and et al.(2002):** Clinical utility of blood drawn from central venous or arterial catheters in critically ill surgical patients. Crit Care Med ; 30 : 7-13.
- McCarthy, M.C.; Shives, J.K.; Robison, R.J.; And et al. (1987):** Prospective evaluation of single and triple lumen catheters in total parenteral nutrition. J. Parenter. Enteral Nutr.; 11:259–62.
- McGee, D.C. and Gould, M.K. (2003):** Preventing Complications of Central venous Catheterization. N Engl J Med; 348: 1123-1133.
- McKibben, L.; Horan, T.; Tokars, J.I.; Fowler, G.; Cardo, D.M.; Pearson, M.L.; Brennan, P.J. and the Healthcare Infection Control Practices Advisory Committee (2005):** Guidance on Public Reporting of Healthcare-Associated Infections: Recommendations of the Healthcare Infection Control Practices Advisory Committee. Am J Infect Control; 33:217-226.
- Mendelson, M.H.; Short, L.J. and Schechter, C.B. (1998):** Study of a needleless intermittent intravenous- access system for peripheral infusions: analysis of staff, patient, and institutional outcomes. Infect Control Hosp Epidemiol; 19: 401-406.
- Mermel, L.A. (2000):** Prevention of intravascular catheter- related infections. Ann Intern Med; 132: 391-402.
- Mermel, L.A.; Farr, B.M.; Sheretz, R.J.; Read, I.I. et al., (2001):** Guidelines for the management of intra vascular catheter – related infection/ clin infect Dis 32; 1249-1272.



Mermel, L.A.; Parenteau, S. and Tow, S.M. (1995): The risk of midline catheterization in hospitalized patients. A prospective study. Ann Intern Med; 123: 841-484.

Mermel, L.A.; Stolz, S.M. and Maki, D.G. (1993): Surface antimicrobial activity of heparin- bonded and antiseptic- impregnated vascular catheters. J Infect Dis; 167: 920-924.

Merrer, J.; De Jonghe, B. and Golliot, F. (2001): Complications of femoral and subclavian venous catheterization in critically ill patients: a randomized controlled trial. JAMA; 286: 700-707.

Miller, M.A.; Dascal, A.; Portnoy, J. and Mendelson, J. (1996): Development of mupirocin resistance among methicillin-resistant staphylococcus aureus after widespread use of nasal mupirocin ointment. Infect Control Hosp Epidemiol; 17: 811-813.

N

Nandakumar, S. (2006): Noscomial infections. Available at: www.sunmed.org.

Nitenberg, G. and Blot, F. (2004): Diagnosis of catheter- related infection. In: N.p. O'Grady and Pittet, D, Editors, Catheter related infections in the critically ill, kluwer Academic Publishers Group the Nether lands 5-76.



o

Oda, T.; Hamasaki, J.; Kanda, N. and Mikami, K. (1997): Anaphylactic shock induced by an antiseptic- coated central venous catheter. *Anesthesiology*; 87: 1242-1244.

Odetola, F.O.; Moler, F.W.; Dechert, R.E.; VanDerElzen, K. and Chenoweth, C. (2003): Nosocomial catheter- related bloodstream infections in a pediatric intensive care unit: Risk and rates associated with various intravascular technologies. *Pediatr. Crit. Care Med.* 4:432-436.

O'Grady, N.P.; Alexander, M. and Dellinger, E.P. (2002): Guidelines for the prevention of intravascular catheter- related infections. *Infect Control Hosp Epidemiol*; 23:759-769.

Oncu, S. and Sakarya, S. (2003): Central Venous Catheter Related Infections: An Overview with Special Emphasis on Diagnosis, Prevention and Management. *The Internet Journal of Anesthesiology*; 7(1). Available at:
<http://www.ispub.com/ostia/index.php?xmlFilePath=journals/ija/vol7nl/cvc.ml>.

Oncu, S.; Ozsut, H.; Yildirim, A.; Ay, P.; Cakar, N.; Eraksoy, H. and Calangu, S. (2003): Central venous catheter related infections: Risk factors and the effect of glycopeptide antibiotics. *Ann. Clin. Microbiol. Antimicrob.* 2:3-9.



Oncu, S.; Ozsut, H.; Yildirim, A.; Ay, P.; Cakar, N.; Eraksoy, H. and Calangu, S. (2003): Central venous catheter related infections: Risk factors and the effect of glycopeptide antibiotics. *Annals of Clinical Microbiology and Antimicrobials*, 2:3 available at: <http://www.annclinmicrob.Com/content/2/1/3>.

O'Toole, G. (2003): To build a biofilm. *J Bacteriol*; 185:2687-2689.

P

Parsek, M.R. and Singh, P.K. (2003): Bacterial biofilms: an emergig link to disease pathogenesis. *Annu Rev Microbiol*; 57:677-701.

Pascual, A. (2002): Pathogenesis of catheter- related infections: lessons for new designs. *Clin Micrbiol Infect*; 8:256-264.

Pierce, C.M.; Wade, A. and Mok, Q. (2000): Heparin- bonded central venous lines reduce thrombotic and infective complications in critically ill children. *Intensive Care Med*; 26: 967-972.

Pittet, D.; Hugonnet, S. and Harbath, S. (2000): Effectiveness of a hospital- wide program to improve compliance with hand hygiene. *Lancet*; 356: 1307-1309.

Pittet, D.; Li, N.; Woolson, R.F. and Wenzel, R.P. (1997): Microbiological factors influencing the outcome of nosocomial bloodstream infectors: a 6- year validated, population- based model. *Clin. Infect. Dis.* 24: 1068-78.



Pittet, D.; Tarara, D. and Wenzel, R.P. (1994): Nosocomial blood stream infection in critically ill patients. Excess length stay, extra cost, and attributable mortality. JAMA; 271:1598-1601.

Q

Quilici, N.; Audibert, G. and Conroy, M.C.; et al., (1997): Differential quantitative blood cultures in the diagnosis of catheter-related sepsis in intensive care units Clin Infect Dis; 25 (1066-1070).

Quirynen, M.; Breck, M. and Steenberghe, D. (2000): Biofilms in the oral cavity: impact of surface characteristics. In: Evans, L.V. (ed.) Biofilms: recent advances in their study and control. Amsterdam: Harwood Academic Publishers. P: 167-187.

R

Raad, I. (1998): Intravascular-catheter-related infections. Lancet; 351: 893-898.

Raad, I.; Umphrey, J.; Khan, A. And et al. (1993): The duration of placement as a predictor of peripheral and pulmonary arterial catheter infections. J. Hosp. Infect.; 23:17-26.

Raad, I.I ; Hanna, H. and Maki, D. (2007): **Intravascular catheter-related infections: advances in diagnosis, prevention, and management.** The Lancet Infectious Diseases; 7: 645-657.

Raad, I.I. and Bodey, G.P. (1992): Infectious complications of indwelling vascular catheters. Clin Infect Dis; 15:197-208.



Raad, I.I. and Darouiche, R.O. (1996): Catheter – related septicemia: Risk reduction. Infect. Med. 13: 807-823.

Raad, I.I.; Darouiche, R. and Dupuis, J. (1997): Central venous catheters coated with minocycline and rifampin for the prevention of catheter- related colonization and bloodstream infections: a randomized, double- blind trial. The texas Medical Center Catheter Study Group. Ann Intern Med; 127: 267-274.

Raad, I.I.; Darouiche, R.; Hachem, R.; Mansouri, M. and Bodey, G.P. (1996): The broad spectrum activity and efficacy of catheters coated with minocycline and rifampin. J Infect Dis; 173: 418-424.

Raad, I.I.; Hanna, H.A. and Awad, A. (2001): Optimal frequency of changing intravenous administration sets: is safe to prolong use beyond 72 hours? Infect control Hosp Epidemiol; 22: 136-139.

Raad, I.I.; Hohn, D.C. and Gilbreath, B.J. (1994): Prevention of central venous catheter related infections by using maximal sterile barrier precautions during insertion. Infect Control Hosp Epidemiol; 15: 231-238.

Raad, I. Hanna, H.A. Alakech, B. Chatzinikolaou, L. Johnson, M.M and Tarrand, j. (2004): Differential time to positivity: A useful method for diagnosing catheter –related blood stream infection without catheter removal. The Lancet 354; 1487-1488.



- Rackoff, W.R.; Welman, M.; and Jakobowski, D. (1995):** A randomized, controlled trial of the efficacy of a heparin and vancomycin solution in preventing central venous catheter infections in children. *J pediatr*; 127: 147-151.
- Randolph, A.G.; Cook, D.J.; Gonzales, C.A. and Andrew, M. (1998):** Benefit of heparin in central venous and pulmonary artery catheters: a meta- analysis of randomized controlled trials. *Chest*; 113: 165-171.
- Randolph, A.G.; Cook, D.J.; Gonzales, C.A. and Pribble, C.G. (1996):** Ultrasound guidance for placement of central venous catheters: a meta- analysis of the literature. *Crit Care Med*; 24: 2053-2058.
- Rangel-Frausto, M.S. Wiblin, T. and Blumberg, H.M. (1999):** National epidemiology of mycoses survey (NEMIS): Variations in rates of bloodstream infections due to *Candida* species in seven surgical intensive care units and six neonatal intensive care units. *Clin Infect Dis*; 29:253-258.
- Rani, S.A.; Pitts, B. and Stewart, P.S. (2005):** Rapid diffusin of fluorescent tracers into *Staphylococcus epidermidis* biofilms visualized by time lapse microscopy. *Antimicrob Agenst Chemotehr*; 49: 728-732
- Rao, S.D.; Preetha, M.J.; Lavi, R. and Macaden, R.(2005) :** Infections related to vascular catheters in a pediatric intensive care unit. *Indian Pediatrics*. 42 :667-672.



- Rello, J.; Coll-Pand Prats, G. (1991):** Laboratory diagnosis of Catheter-related bacteraemia infect. Dis, 23: 583-588.
- Riu, S.; Ruiz, C.G.; Martinez, Vea, A.; Peralra, C. and Oliver, J.A. (1998):** Spontaneous rupture of polyurethane peritoneal catheter: a possible deleterious effect of mupirocin ointment. Nephrol Dial Transplant; 13: 1870-1871.
- Roberts, M.E. and Stweart, P.S. (2005):** Modeling protection from antimicrobial agents in biofilms through the formation of persister cells. Microbiology; 151: 75-80.
- Rowin, M.E.; Patel, V.V. and Christenson, J.C. (2003):** Pediatric intensive care unit nosocomial infections. Crit. Care Clin. 19: 473-487.
- Rushforth, J.A.; Hot, C.M. Kite, P. and Puntis, J.W.L. (1993):** Rapid diagnosis of central venous catheter sepsis. Lancet; 342: 402-403.
- Rushforth, J.A.; Hoy, C.M.; Kite, P. and Puntis, J.W. (1993):** Rapid diagnosis of central venous catheter sepsis. The Lancet; 342:402-403.
- Ryder, M. (2001):** The role of biofilm in vascular catheter- related infections. N Dev Vasc Dis; 2:15-25.
- Ryder, M.A. (2005):** Catheter- Related Infections: It's All About Biofilm. Topics in Advanced Practice Nursing eJournal; 5(3). Available at: http://www.medscape.com/viewarticle/508109_2.



S

Sachdev, A.; Gupta, D. and Soni, A. (2002): Central venous catheter colonization and related bacteremia in pediatric intensive care unit. *Indian Pediatrics* 39: 752-760.

Safdar N., Fine J. and Maki D. (2005): Meta-analysis: methods for diagnosing intravascular device-related bloodstream infection. *Ann. Intern. Med.*; 142:451–466.

Safdar, N. and Maki, D.G. (2002): Inflammation at the insertion site is not predictive of catheter-related bloodstream infection with short-term, noncuffed central venous catheters, *Critical Care Medicine*; 30: 2632–2635.

Safdar, N. and Maki, D.G. (2004): The pathogenesis of catheter- related bloodstream infection with noncuffed short-term central venous catheters. *Intensive Care Med.*; 30:62-67.

Safdar, N.; Fine.; J.P. and Maki, D.G. (2005): Meta- analysis: methods for diagnosing intravascular device-related bloodstream infection. *Ann Intern Med.*; 142:451-466.

Saiman, L.; Ludington, E. and Dawson, J.D. (2001): Risk factors for *Candida* species colonization of neonatal intensive care unit patients. *Pediatr. Infect. Dis.* 20:1119-1124.



- Seifert, H.; Cornely, O.; Seggewiss, K.; Decker, M.; et al., (2003):** Blood stream infection in neutropenic cancer patients related to short-term non tunneled catheters, determined by quantitative blood cultures, differential time to positive and molecular epidemiological typing with J Clin microbial 41; 118-123.
- Seigman-Igra, Y.; Anglim, A.M.; Shapiro, D.E. et al. (1997):** Diagnosis of Vascular Catheter-Related Bloodstream Infection: a Meta-Analysis. J. Clin. Microbiol.; 35(4):928-936.
- Sesso, R.; Barbosa, D. and Leme, I.L. (1998):** Staphylococcus aureus prophylaxis in hemodialysis patients using central venous catheter: effect of mupirocin ointment: J Am Soc Nephrol; 9: 1085-1092.
- Seymour, V.M.; Dhallu, T.S.; Moss, H.A.; Tebbs, S.E. and Elliot, T.S. (2000):** A prospective clinical study to investigate the microbial contamination of a needleless connector. J Hosp Infect; 45: 165-168.
- Sherertz, R.J.; Ely, E.W. and Westbrook, D.M. (2000):** Education of physicians in- training can decrease the risk for vascular catheter infection. Ann Intern Med; 132: 641-648.
- Sheth, N.K.; Franson, T.R.; Rose, H.D.; Buckmire, F.L. Cooper, J.A. and Sohnle, P.G. (1983):** Colonization of bacteria on polyvinyl chloride and Teflon intravascular catheters in hospitalized patients. J Clin Microbiol; 18: 1061-1063.



- Singh- Naz, N.; Sprague, B.M.; Patel, K.M. and Pollack, M.M.(1996):** Risk factors for nosocomial infection in critically ill children: a prospective cohort study. Crit. Care Med. 24:875-878.
- Slaughter, S.E. (2004):** Intravascular catheter-related infections: strategies for combating this common foe. Postgrad Med; 116(5):59-6.
- Slonim, A.D.; Kurtines, H.C. and Sprague, B.M.(2001):** The costs associated with nosocomial bloodstream infections in the pediatric intensive care unit. Pediatr. Crit. Care Med.; 2:170-174.
- Snydman, D.R.; Donnelly- Reidy, M.; Perry, L.K. and Martin, W.J. (1987):** Intravenous tubing containing burettes can be safely changed at 72 hour intervals. Infect Control; 8: 113-116.
- Soifer, N.E.; Borzak, S.; Edlin, B.R. and Weinstein, R.A. (1998):** Prevention of peripheral venous catheter complications with an intravenous therapy team: a randomized controlled trial. Arch Intern Med; 158: 473-477.
- Spafford, P.S.; Sinkin, R.A.; Cox, C.; Reubens, L. and Powell, K.R. (1994):** Prevention of central venous catheter- related coagulase-negative staphylococcal sepsis in neonates. J Pediatr; 125: 259-263.
- Stewart, P.S. (2002):** Mechanisms of antibiotic resistance in bacterial biofilms. Int. J Med Microbiol; 292:107-113.
- Stewart, P.S. and Costerton, J.W. (2001):** Antibiotic resistance of bacteria in biofilms. Lancet; 358:135-138.



Stickler, D.J. (1996): Bacterial biofilms and the encrustation of urethral catheters. *Biofouling* 94: 293-305.

Stickler, D.J.; Morris, N. and Moreno, M.C. (1998): Studies on the formation of crystalline bacterial biofilms on urethral catheters *Eur. J. Clin. Microbiol. Infect. Dis.* 17: 649-652.

Stoodley, P. and Dirckx, P. (2003): Center for Biofilm Engineering (CBE), Montana State University. www.erc.Montana.Edu.

Stewart, P.I. and Dirckx, P. (2001): Center for Biofilm Engineering (CBE), Montana State University. [www. Erc. Montana. Edu](http://www.Erc.Montana.Edu).

T

Tanguy, M.; Seguin, P.; Laviolle, B.; Desbordes, L. and Malledant, Y. (2005): Hub qualitative blood culture is useful for diagnosis of catheter- related infections in critically ill patients. *Intensive care med.*

Templeton, A.; Rettenmund, G.; Schobi, B.; Diethelm, K.; Schlegel, M.; et al., (2007): Prospective evaluation of risk factors for catheter-related blood stream infections in two Swiss hospitals. *International journal of antimicrobial agents*; 29: 5252.

Templeton, A.; Schlegel, M.; Fleisch, F. And et al. (2008): Multilumen Central Venous Catheters Increase Risk for Catheter-Related Bloodstream Infection: Prospective Surveillance Study. *Journal of infection*; 36:322-327.



- Tenover, P.C. and Schaberg, D.R. (1998):** Molecular biology of resistance. In: Bennett JV, Brachman, PS, eds. Hospital Infections, 4th ed. Philadelphia, Pa: Lippincott- Raven: 237-247.
- Timsit, J.F.; Farkas, J.C. and Boyer, J.M. (1998):** Central vein catheter-related thrombosis in intensive care patients: incidence, risk factors, and relationship with catheter-related sepsis. Chest; 114: 207-213.
- Tony worthington and Tom S.J. Elliott. (2005):** Diagnosis of central venous catheter related infection in adult patients. Journal of infection; 51:267-280.
- Touyama, M. and Saito, A. (2002):** Bloodstream infection. Nippon Rinsho; 60(11): 2187-2193.
- Tully, J.L.; Friedland, G.H.; Baldini, L.M. and Goldman, D.A. (1981):** Complications of intravenous therapy with steel needles and Teflon ® catheters: a comparative study. Am J Med; 70: 702-706.
- Tunney, M.M.; Jones, D.S. and Gorman, S.P. (1999):** Biofilm and biofilm-related encrustations of urinary tract devices. In: Doyle RJ, ed. Methods in enzymology, vol. 310. Biofilms. San Diego: Academic Press, 558-66.
- Turco, S.J. and Davis, N.M. (1973):** Particulate matter in intravenous infusion fluids- phase 3. Am J Hosp Pharm; 30: 611-613. Quoted from: CDC (2002).



U

Uldall, P.R.; Merchant, N.; Woods, F.; Yarworski, U. and Vas, S. (1981): Changing subclavian haemodialysis cannulas to reduce infection. *Lancet*; 1: 1373.

V

Valles, J.; Anoro, D.; Fontanals, M. and et al. (2008): Bloodstream infection in adults :Importance of health care associated infections *Journal of Infection*; 56:27-34.

Vandecasteele, S.J.; Peetermans, W.E.; Merckx, R. And Van Eldere, J. (2003): Expression of biofilm- associated genes in *Staphylococcus epidermidis* during in vitro and in vivo foreign body infections. *J Infect Dis*; 188:730-737.

Veenstra, D.L.; Saint, S.; Saha, S.; Lumley, T. and Sullivan, S.D. (1999): Efficacy of antiseptic- impregnated central venous catheters in preventing catheter- related bloodstream infection: a meta- analysis. *JAMA*; 281: 261-267.

Vicca, A.F. (1999): Nursing staff workload as a determinant of methicillin- resistant *Staphylococcus aureus* spread in an adult intensive therapy unit. *J Hosp Infect*; 43:109-113.

Von Eiff, C.; Becker, K.; Machka, K.; Stammer, H. and Peters, G. (2001): Nasal carriage as a source of *Staphylococcus aureus* bacteremia. *N Engl J Med*; 344: 11-16.

**W**

Warren, D.K.; Yokoe, D.S.; Climo, M.W.; Herwaldt, L.A.; Noskin, G.A.; Zuccotti, G.; Tokars, J.I.; Perl, T.M. and Fraser, V.J. (2006): Preventing Catheter- Associated Bloodstream Infections: A Survey of policies for insertion and Care of Central Venous Catheters from Hospitals in the Prevention Epicenter Program. *Infection Control Hosp. Epidemiol*; 27: 8-13.

Warren, D.K.; Zack, J.E. and Elward, A.M. (2001): Nosocomial primary bloodstream infections in intensive care unit patients in a nonteaching community medical center: a 21-months prospective study. *Clin Infect Dis*; 33: 1329-1335.

Wilcox, H.M.; Kite, P.; Mills, K.;and Sugden, S.(2001): In-situ measurement of linezolid and vancomycin concentrations in intravascular catheter-associated biofilm.*Journal of Antimicrobial Chemotherapy*.; 47:171-175.

Wilson, J.N.; Grow, J.B.; Demong, C.V.; Prevedel, A.E. and Owens, J.C. (1962): Central venous pressure in optimal blood volume maintenance. *Arch. Surg.* 85:563-578.

Worthington, T.; Traube, A.; Lambert, A.; and Elliott, T.S.J. (2002): a Rapid ELISA for the diagnosis of intravascular catheter- related sepsis caused by coagulate negative staphylococci. *J Clin path* 55; 41-43.



Worthington, T.; Tom, S.J. and Elliott, T.S.J. (2005): Diagnosis of central venous catheter-related infection in adult patients. *Journal of Infection*. ;51:267-280.

Y

Yamamoto, A.J.; Solomon, J.A. and Soulen, M.C. (2001): Sutureless securement device reduces complications of peripherally inserted central venous catheters. *J. Vasc Interv Radiol*; 12.

Yogaraj, J.S.; Elward, A.M. and Fraser, V.J. (2002): Rate, risk factors, and outcomes of nosocomial primary bloodstream infection in pediatric intensive care unit patients. *Pediatrics*. 110:481-485.

Z

Zing, W. ;Vanessa, C.F.;and Walder, B. (2008): Central venous catheter related infection. *Best practice and research clinical anaesthesiology*.22; 407-421.