

REFERENCES

Akhtar G, Gilbert CJ, and Wolf FG. (1979): Retrograde conduction in the His- Purkinje system. *Circulation*, 59: 1252-1265.

Arruda M, Mcclelland, JH, Wang H, et al. (1998): Development and validation of an ECG algorithm for identifying accessory pathway ablation site in WPW syndrome. *Journal of Cardiovascular electrophysiology*, 9: 2 – 11.

Arruda M, Mauricio S, James H, et al. (1998): Development and validation of an ECG algorithm for identifying accessory pathway ablation site in WPW syndrome. *J Cardiovasc Electrophysiol*, 9: 2-12.

Bardy GH, Packer DL, German LD, et al. (1984): Preexcited reciprocating tachycardia in patients with WPW syndrome, incidence and mechanisms. *Circulation*, 70: 377: 391.

Bardy GH, Poole JE, Coltorti F, et al. (1984): Catheter ablation of a concealed accessory pathway. *Am. J. Cardiol*, 54: 366.

Bashir Y, Heald SC, O'Nunain S, et al. (1993): Radiofrequency current delivery by way of a bipolar tricuspid annulus mitral annulus electrode configuration for ablation of posteroseptal accessory pathways. *J Am Coll Cardiol*, 22: 550-556.

Belhassen B, Rogowski O, Glick A, et al. (2007): *Radiofrequency ablation of accessory pathways: a 14 year experience at the Tel Aviv Medical Center in 508 patients. Isr Med Assoc J. Apr,9(4):265-70*

Benditt DG, Durrer D, and Bremne S. (1979): *Ventriculoatrial intervals: diagnostic use in paroxysmal supraventricular tachycardia. Ann intern Med, 91: 161-166.*

Blomstrom LC, Scheinman MM, Aliot EM, et al.(2003):
ACC/AHA/ESC guidelines for the management of patients with supraventricular arrhythmias executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines. (Writing Committee to Develop Guidelines for the Management of Patients with Supraventricular Arrhythmias). Circulation,108:1871–909.

Bockeria LA, Kupadatze NI, Saprigin DB, et al. (1991): *Surgical treatment of the Wolff – Parkinson – White syndrome by epicardial electrical ablation. Ann. Thorac Surg. 51: 563, .*

Bogun F, Kalusche D, Li YG, et al (1999). *Septal Q waves in surface electrocardiographic lead V6 exclude minimal ventricular preexcitation. Am J Cardiol;84:101–4, A9.*

Brechenmacher C, Laham J, Iris L, et al. (1974): *Histological study of abnormal conduction pathways in the Wolff-Parkinson-White syndrome and Lown-Ganong-Levine syndrome. Arch Mal Coeur Vaiss 1974, 67:507–519*

Broggreffe M, Budde T, Podcizek A, et al. (1987): *High frequency alternating current ablation of an accessory pathway in humans. J. Am. Coll. Cardiol, 10: 576.*

Borggreffe M, Budde T, and Podcizek A. (1987): *High frequency alternating current ablation of an accessory pathway in humans. J Am Coll Cardiol, 10: 576-582.*

Calkins H, Kim YN, and Schmaltz S. (1992): *Electrogram criteria for identification of appropriate target sites for radiofrequency catheter ablation of accessory atrioventricular connections. Circulation, 85: 565-573.*

Calkins H, Langberg J, Sousa J, et al.(1992): *Radiofrequency catheter ablation of accessory atrioventricular connections in 250 patients: abbreviated therapeutic approach to Wolff-Parkinson-White. Circulation, 85:1337–46.*

Calkins H. (1998): *Catheter ablation of anteroseptal and midseptal accessory pathways. In: singer I, Barold SS, Camm AJ: Nonpharmacological therapy of arrhythmias for the 21st century, the state of the art, future publishing company, New York. Chap 6 117-137.*

Calkins H, Yong P, Miller JM, et al.(1999):

Catheter ablation of accessory pathways, atrioventricular nodal reentrant tachycardia, and the atrioventricular junction: final results of a prospective, multicenter clinical trial. The Atakr Multicenter Investigators Group. Circulation, 99:262–70.

Calkins H, Berger RD.(2001): *Catheter ablation of anteroseptal and*

midseptal accessory pathways. In: Singer I, ed. Interventional Electrophysiology, 2nd ed. Philadelphia: Lippincott, Williams & Wilkins, 237–54.

Chen SA, and Tai CT.(2001): *Ablation of atrioventricular accessory pathways. Current technique-state of the art..Pacing Clin Electrophysiol 24:1795-1809.*

Chiang, CE, Chen, SA, and Tai, CT. (1994): *Major Coronary sinus abnormalities: identification of occurrence and significance in radiofrequency ablation of supraventricular tachycardia. Am Heart J, 127: 1279-1289.*

Chiang CE, Chen SA, Teo WS, et al. (1995): *An accurate stepwise electrocardiographic algorithm for localization of accessory pathways in patients with Wolf-Parkinson-White syndrome from a comprehensive analysis of delta waves and R/S ratio during sinus rhythm. Am J Cardiol, 76: 40–46*

Cobb FR, Blumenschein SD, Sealy WC, et al. (1968): Successful surgical interruption of the bundle of Kent in a patient with Wolff – Parkinson – White syndrome. *Circulation*, 38 : 1018.

Compell MJ, and Machin D.(1993): Medical statistics: a common source approach 2nd edition. John Wiley and sonf chiechester, New York, pris bone, Toronto and thing apore.

Deal, B.J.; Keana, J.F.; Gillett, P.C. and Garson, A.J. (1985): WPW syndrome and supraventricular tachycardia during infancy: Management and follow- up J Am Coll cardiol, 5: 130.

Dean JW, Ho SY, and Rowland, E. (1994): Clinical anatomy of the atrioventricular junctions. *J Am Coll Cardiol*, 24: 1725-1731.

Della Bella P, Brugada P, Talajic M, et al. (1991): Atrial fibrillation in patients with an accessory pathway: importance of the conduction properties of the accessory pathway. *J Am Coll Cardiol*, 17:1352–1356.

De la Fuente D, Sasyniuk B, and Moe GK.(1971): Conduction through a narrow isthmus in isolated canine atrial tissue. A model of the W-P-W syndrome. *Circulation*, 44:803–809.

Dennis, WX. (1997): Radiofrequency catheter ablative therapy for anteroseptal and midseptal accessory atrioventricular pathways. In: *Singer 1: interventional Electrophysiology*, Williams & Wilkins, Balimore, Maryland. Chapter 9, P: 253-273.

Dhala AA, Deshpande SS, Bremner S, et al.(1994): Transcatheter ablation of posteroseptal accessory pathways using a venous approach and radiofrequency energy. *Circulation*, 90: 1799–810

Engelstein, ED, Wilber D, and Wadas M. (1993): Limitations of adenosine in assessing efficacy of radiofrequency catheter ablation of accessory pathways. *Circulation*, 8: 1-295.

Fananapazir L, German LD, Gallagher JJ, et al. (1990): Importance of pre – excited QRS morphology during induced atrial fibrillation to the diagnosis and localization of multiple accessory pathways. *Circulation*, 81 : 578.

Fisch C. (1990): Clinical electrophysiological studies and the Wolff – Parkinson – White pattern. *Circulation*, 82: 1872.

Fitzpatrick AP, Gonzales RP, Lesh MD, et al .(1994): New algorithm for the localization of accessory atrioventricular connections using a baseline electrocardiogram. *J Am Coll Cardiol*, 23:107–116.

Fitzpatrick AP. (1995): The ECG in Wolf – Parkinson – White syndrome. *PACE*, 18: 1469.

Gallagher LA, Sealy WC, and Anderson W. (1977): Cryo-surgical ablation of accessory atrioventricular connections a method for correction of the preexcitation syndrome: *Circulation*, 55: 471-478.

Gallagher JJ, Pritchett FM., Sealy WC, et al. (1978): The preexcitation syndromes. *Prog Cardiovasc Dis*, 20: 285.

Gallagher JJ, Sealy WC, Cox JL, et al. (1984): Results of surgery for pre – excitation caused by accessory aotrioventricular pathways in 267 consecutive cases. In: Josephson ME, Wellens HJJ (eds). *Tachycardias: Mechanisms, Diagnosis, Treatment*. Philadelphia, PA, Lea & Febiger, pp. 259.

Gallagher JJ, Selle, JG, and Sealy WC. (1986): Intermediate septal accessory pathways a subset of preexcitation at risk for complete heart block/ failure during WPW surgery. *Circulation*, 74: 11-387.

Gallagher JJ. (1987): Accessory pathway tachycardia technique of electrophysiologic study and mechanisms. *Circulation*, 75 (suppl III): III – 31 – 6.

Gallagher JJ, Sealy WC, and Kasell J. (1979): Intraoperative mapping studies in the Wolff – Parkinson – White syndrome. *PACE*, 2 : 523.

Chang SL, Lee SH, Tai CT, et al. (2005): . Electrocardiographic and electrophysiologic characteristics of midseptal accessory pathways. *J Cardiovasc Electrophysiol*. Mar,16(3):237-43.

Goudevenos JA, Katsoura CS, Graeka G, et al. (2000): Ventricular pre-excitation in the general population: a study on the mode of presentation and clinical course *Heart* ,83:29–34

Guiraudon GM, Klein GJ, and Sharma A.D. (1986): Surgical ablation of posteroseptal accessory pathways in the WPW syndrome y a closed heart technique. *J Thorac cardiovasc Surg*, 92: 406 – 413.

Guiraudon GM, Guiraudon CM, and Klein GJ. (1988): *The coronary sinus diverticulum : a pathologic entity associated with the WPW syndrome. Am J Cardiol, 62: 733- 735.*

Guiraudon GM, Klein GJ, and Sharma AD. (1988): *Atypical posteroseptal accessory pathways in WPW syndrome. J Am Coll Cardiol, 2: 1605- 1608.*

Guize L, Soria R, Chaouat JC, et al. (1985): *Prevalence and course of Wolff-Parkinson-White syndrome in population of 138,048 subjects. Ann Med Interne (Paris), 136:474– 89.*

Haghjoo Majid, Ali Kharazi, Amir Farjam Fazelifar, et al. (2007): *Electrocardiographic and electrophysiologic characteristics of anteroseptal, midseptal, and posteroseptal accessory pathways Heart Rhythm,4:1411–1419*

Haissaguerre M, Dartigues JF, Warin JF. (1991): *Electrogram patterns predictive of successful catheter ablation of accessory pathways, value of unipolar recording mode. Circulation, 84: 188- 202.*

Haissaguerre M, Fischer B, Warin J, et al. (1992): *Electrogram patterns predictive of successful radiofrequency catheter ablation of accessory pathways. PACE, 15 : 1000.*

Haissaguerre M, Macus F, Pouet F, et al. (1994): *Electrocardiographic characteristics and catheter ablation of para – Hisian accessory pathways. Circulation, 90 : 1124*

- Haissaguerre M, Cauchemez B, Marcus F, et al. (1995):** *Characteristics of the ventricular insertion sites of accessory pathways with antegrade decremental conduction properties. Circulation 1995, 91:1077–1085*
- Huang SK, Graham AR, Hyot RH, et al. (1987):** *Transcatheter dissection of the canine left ventricle using radiofrequency energy: pilot study. Am Heart J. 14-42-49.*
- Huang SK. (1989):** *Radio – frequency catheter ablation of cardiac arrhythmias: Appraisal of an evolving therapeutic modality. Am. Heart. J, 118: 1317.*
- Iturralde P, Araya-Gomez V, Colin L, et al. (1996):** *A new ECG algorithm for the localization of accessory pathways using only the polarity of the QRS complex. J Electrocardiol, 29:289 –299.*
- Jackman, WM, Froidoay KJ, Scherlag BJ, et al. (1983):** *Direct endocardial recording from an accessory atrioventricular pathway : Localization of the site of block, effect of antiarrhythmic drugs, and attempt of nonsurgical ablation. Circulation, 68 : 906.*
- Jackman WM, Friday KJ, Yeung L , Wah JA, et al. (1988):** *New catheter technique for recording left free wall accessory atrioventricular pathway activation – identification of pathway orientation. Circulation, 78 : 598.*

Jackman WM, Bowman AJ, and Lazzara R. (1989): Localization of left free wall and posteroseptal accessory pathways by direct recording of accessory pathways activation. *PACE*, 12: 204-214.

Jazayeri MR, Caceres J, Tchou P, et al. (1989): Electrophysiologic characteristics of sudden QRS axis deviation during orthodromic tachycardia: role of functional fascicular block in localization of accessory pathway. *J. Clin Invest*, 83: 952.

Jazayeri JM, Deshpande S, and Dhala A. (1994): Transcatheter mapping and radiofrequency ablation of cardiac arrhythmias. *Curr Probl Cardiol*, 292-395.

Jackman WM, Wang X, Friday KJ. et al. (1991): Catheter ablation of accessory atrioventricular pathways (Wolff – Parkinson – White syndrome) by radiofrequency current. *N. Engl. J. Med*, 324 : 1605.

John Miller, Mithilesh K Das, Anil V, et al. (2006): Catheter ablation of superparaseptal and Midseptal accessory pathways. In catheter ablation of cardiac arrhythmias. Philadelphia : Saunders, 413-425.

Josephson, M.E. (1993): Pre – excitation syndrome. In: Josephson ME (2nd). *Clinical cardiac electrophysiology: Techniques and interpretations, Second Edition*. Philadelphia. Lea & Febiger, P. 311.

Josephson ME. (2008): *Clinical Cardiac electrophysiology. In: Preexcitation syndromes 4th edition Philadelphia: Lippincott Williams and Wilkins, 340-398).*

Kay GN, Pressley JG, Packer DL, et al. (1987): *Value of the 12 – lead electrocardiogram in discriminating atrioventricular nodal reciprocating tachycardia from circus movement atrioventricular tachycardia utilizing a retrograde accessory pathway. Am. J. Cardiol, 59 : 296.*

Kay GN, Epstein AE, and Dailey SM. (1993): *Role of radiofrequency ablation in the management of supraventricular arrhythmias: experience in 760 consecutive patients. J Cardiovasc Electrophysiol, 4: 371-389.*

Keim S, Curtis AB, Belardinelli L, et al. (1992): *Adenosine – induced atrioventricular block : A rapid and reliable method to assess surgical and radiofrequency catheter ablation of accessory atrioventricular pathways. J. Am. Coll. Cardiol, 19 : 1005.*

Kerr CR, Gallagher JJ, and German LD. (1982): *Changes in ventriculoatrial intervals with bundle branch block aberration during reciprocating tachycardia in patients with accessory atrioventricular pathways. Circulation, 66: 196.*

Klein GJ, Prystowsky EN, Yee R, et al. (1989): *Asymptomatic Wolff-Parkinson-White: should we intervene? Circulation, 80:1902–5.*

Klein GJ, Yee R, and Sharma AD. (1989): A longitudinal electrophysiologic assessment of asymptomatic patients with the Wolff – Parkinson – White electrocardiographic pattern. *N. Engl. J. Med*, 320 : 1229

Klein GJ, and Prystowsky EN (1997): Wide QRS Tachycardia. In: *Clinical electrophysiology review*, McGraw-Hill company, New York. Chap 4, P, 148.

Klein GJ, Guiraudon GM, Kerr CR, et al. (1988): Nodoverricular” accessory pathway: evidence for a distinct accessory atrioventricular pathway with atrioventricular node-like properties. *J Am Coll Cardiol* ,11: 1035–1040.

Konstantinos G, Theodoros A, Xenofon C, et al .(2004): Paraseptal accessory connections in the proximity of the atrioventricular node and the His bundle. Additional observations in relation to the ablation technique in a high risk area. *Europace*. 2004 Jan;6(1):1-9.

.Krahn AD, Manfreda J, Tate RB, et al. (1992): The natural history of electrocardiographic pre- excitation in men : The Manitoba follow-up study. *Ann. Intern. Med*, 116 : 456.

Krahn AD, Klein GJ, Guiraudon GM, et al. (1997): Ablation of posteroseptal accessory pathways. In : *Singer I : interventional Electrophysiology*, Williams & Wilkins, Baltimore, Maryland. Chapter 8, P: 231 – 252.

- Kuck KH, Schluter M, and Guroy S. (1992):** *Preservation of atrioventricular nodal conduction during radiofrequency current catheter ablation of mid-septal accessory pathways. Circulation, 86: 1743-1752.*
- Kuck K, Schluter M, Cappato R, et al. (1994):** *Ablation of septal accessory pathways . in: Zipes DP: Catheter ablation of arrhythmias, Future Publishing company, Armonk, New York. Chapter 9, P: 231- 257.*
- Kuck KH, Ouyang F, Goya M, et al. (2002):** *Ablation of anteroseptal and midseptal accessory pathways. In: Zipes DP, Haïssaguerre M, eds. Catheter Ablation of Arrhythmias, 2nd ed. Armonk, NY: Futura, 2002: 305–20.*
- Langberg JJ, Calkins H, and El-Atassi R. (1992):** *Temperature monitoring during radiofrequency catheter ablation of accessory pathways. Circulation, 86: 1469- 474.*
- Langberg JJ, Calkins H, and Klm YN. (1992):** *Recurrence of conduction in accessory atrioventricular connection after initially successful RF ablation J AM coll Cariol, 19: 1588- 1592.*
- Langberg JJ, Chnig MK, and Vorperian VR. (1993):** *Recognition and catheter ablation with very large distal electrodes. Circulation, 88: 245-249.*

- Lesh MD, VanHare GF, and Schamp DJ. (1992):** Curative percutaneous catheter ablation using radiofrequency energy for accessory pathways in all location. *J AM Coll Cardiol*, 19: 1303-1309.
- Leitch JW, Klein GJ, Yee R, et al. (1990):** Prognostic value of electrophysiology testing in asymptomatic patients with Wolff-Parkinson-White pattern. *Circulation*, 82:1718–23.
- Leitch JW, Klein GJ, Yee R, et al. (1992):** Syncope associated with supraventricular tachycardia. An expression of tachycardia rate or vasomotor response? *Circulation*, 85: 1064.
- Lorga Filho A, Sosa E, Scanavacca M, et al. (1996):** Electrocardiographic identification of midseptal accessory pathways in close proximity to the atrioventricular conduction system. *Pacing Clin Electrophysiol*, 19:1984 –1987.
- Miles, W.M.; Yee, R.M.; Klein, G.J. et al. (1986):** The pre – excitation index: An aid in determining the mechanism of supraventricular tachycardia and localizing accessory pathways. *Circulation*, 74 : 493.
- Miller JM, Rosenthal ME, Gottlieb CD, et al. (1987):** A new criterion reliably distinguishes atrioventricula reentrant tachycardia from septal bypass tract tachycardia. *J Am Coll Cardiol* , (9): 12A.
- Milstein S, Sharma D, and Guiraudon G. (1987):** An algorithm for the electrocardiographic localization of accessory pathways in the WPW syndrome. *PACE*, 10: 555-653.

- Morady F, Scheinman MM, Kou WH, et al . (1989):** Long-term results of catheter ablation of a posteroseptal accessory atrioventricular connection in 48 patients. *Circulation* 79: 1160- 1170.
- Morady F ,Aman Chugh, and Frank Bogun.(2008):** Catheter ablation of accessory pathways.In *Catheter Ablation of Cardiac Arrhythmias Basic Concepts and Clinical Applications* ,3rd edition.Blackwell publishing.New York. 149-172.
- Morgan-Hughes, N.J.; Griffith, M.J. and McComb, J.M. (1993):** Intravenous adenosine can reveal accessory pathways not revealed by routine electrophysiologic testing. *PACE*, 16: 2059-2063.
- Milstein S, Sharma AD, and Klein GJ. (1986):** Electrophysiologic profile of asymptomatic Wolff – Parkinson – White patient. *Am. J. Cardiol*, 57: 1097.
- Montoya PT, Brugada P, Smeets J, et al. (1991):** Ventricular fibrillation in the Wolff – Parkinson – White syndrome. *Eur. Heart, J.* 12 : 144.
- Moradi F, and Scheinman MM. (1984):** Transvenous catheter ablation of a posteroseptal accessory pathway in a patient with the Wolff – Parkinson – White syndrome. *N. Engl. J. Med*, 310 : 705.
- Munger TM, Packer DL, Hammil SC, et al. (1993):** A population study of the natural history of Wolff-Parkinson-White syndrome in Olmsted County, Minnesota, 1953–1989. *Circulation* ,87:866–73.

Nakagawa H, and Warren M Jackman .(2007): *Catheter Ablation of Paroxysmal Supraventricular Tachycardia. Circulation, 116, 2465-2478.*

Packer DL, Gallagher JJ, and Prystowsky EN. (1992): *Physiological substrate for antidromic reciprocating tachycardia : prerequisite characteristics of the accessory pathway and atrioventricular conduction system. Circulation, 85: 574.*

Pappone C, Santinelli V, Rosanio S, et al.(2003): *Usefulness of invasive electrophysiologic testing to stratify the risk of arrhythmic events in asymptomatic patients with Wolff-Parkinson-White pattern: results from a large prospective long-term follow-up study. J Am Coll Cardiol, 41:239–44.*

Pappone C ,Santinelli V, Radinovic A, et al .(2009) : *The natural history of asymptomatic ventricular pre-excitation a long-term prospective follow-up study of 184 asymptomatic children. J Am Coll Cardiol. 2009 Jan 20;53(3):275-80.*

Paul T, Guccione P, and Garson A. (1990): *Relation of syncope in young patients with Wolff – Parkinson – White syndrome to rapid ventricular response duringatrial fibrillation. Am. J. Cardiol, 65 : 318.*

Petersen AH, Andersen ED, and Sandoe E. (1992): *Atrial fibrillation in the Wolff-Parkinson-White syndrome. Am J Cardiol,*

70:38–43A.

Priori SG, Aliot E, Blomstrom-Lundqvist C, et al. Task force on sudden cardiac death of the European Society of Cardiology. *Eur Heart J* 2001;22:1374–450.

Pritchett EL, Tonkin AM, Dugan FA ,et al . (1976): Ventriculo-atrial conducton time during reciprocating tachycardia with intermittent bundle branch block in WPW syndrome. *Br heart J*, 38: 1058.

Prystowsky EN, Pritchett EL, and Smith WM (1979): Electrophysiologic assessment of the atrioventricular conduction system after surgical correction of ventricular preexcitation. *Circulation*, 59: 789- 796.

Prystowsky EN , Browne KF, and Zipes DP. (1988): Intracardiac recording by catheter electrode of accessory pathway depolarization. *J. Am. Coll. Cardiol*, 1 : 468 – 470.

Prystowsky EN, and Klein GJ. (1994): Pre-excitation syndromes: In *Cardiac Arrhythmias : An Integrated Approach for the Clinician*. New York, McGraw – Hill, Inc, P. 131.

Rinne, C.; Klein, G.J.; Sharma, A.D. et al. (1987): Relation between clinical presentation and induced arrhythmias in the Wolff – Parkinson – White syndrome. *Am. J. Cardiol.*, 60 : 576.

Robinson K, Rowland E, and Krikler DM.(1988): Wolff-Parkinson-White syndrome: atrial fibrillation as the presenting Arrhythmia. *Br Heart J*, 59:578–80.

Rodriguez LM, Smeets JL, and De Chillouctal (1993): *The 12 lead electrocardiogram in midseptal, anteroseptal, posteroseptal and right free wall accessory pathways. Am. J. Cardiol, 72: 1274.*

Satake S, Okishige K, and Aegami K. (1996): *Radiofrequency ablation for WPW syndrome with monitoring the local electrogram at the ablation site. JPN Heart J, 37: 741-750.*

Schluter M, Geiger M, Siebels J, et al.(1991): *Catheter ablation using radiofrequency current to cure symptomatic patients with tachyarrhythmias related to an accessory atrioventricular pathway. Circulation,324:1606–11.*

Schluter M, and Kuck KH.(1992): *Catheter ablation from right atrium of anteroseptal accessory pathways using radiofrequency current. J Am Coll Cardiol, 19: 663- 670.*

Schumacher B, Eick O, Wittkamp F, et al. (1989): *Temperature response following non-traumatic low power radiofrequency application. Pacing Clin Electrophysiol, 22: 339-343.*

Schumacher B, Tebbenjohanns J, and Schimpf R. (1995): *Retrograde venography of the coronary sinus in 125 patients with accessory pathways: impact on ablation procedures (Abstract). Circulation, 92 (8): 2918.*

- Sheinman MM, Wang YS, Vanharc GF, et al. (1992):** *Electrocardiographic and electrophysiologic characteristics of anterior, midseptal and right anterior free wall accessory pathways. JACC, 20 : 1220.*
- Sealy WC (1983):** *Kent bundles in the anterior septal space. Ann thorac surg, 36: 180-186.*
- Sharma, A.D.; Klein, G.J.; Guiraudon, G.M. et al. (1985):** *Atrial fibrillation patients with Wolff – Parkinson – White syndrome : Incidence after surgical ablation of the accessory pathway. Circulation, 72 : 161.*
- Sun Y, Arruda M, Otomo K, et al.(2002):***Coronary sinus-ventricular accessory connections producing posteroseptal and left posterior accessory pathways.Circulation 106:1362-1367.*
- Swartz JF, Tracy CM, and Fletcher RD. (1993):** *Radiofrequency endocardial catheter ablation of accessory atrioventricular pathway atrial insertion sites. Circulation,87:487–99.*
- Tai CT, Chen SA, Chiang CE, et al .(1996):** *Radiofrequency catheterablation midseptal, and para-Hisian accessory pathways. Implication for radiofrequency catheter ablation. Chest,109:730 – 740.*
- Thakur, R.K.; Bishof, J.K. and Downing, D.F. (1996):** *The risk of AV nodal block during ablation of posteroseptal accessory pathways (Abstract) Submitted to American College of cardiology 45th annual scientific session, March 24-27.*

- Timmermans C, Smeets J, and Rodriguez LM, et al.(1995):** Aborted sudden death in the Wolff-Parkinson-White syndrome. *Am J Cardiol*,76:492–4.
- Tonkin AM, Wanger GS, Gallagher JJ, et al. (1975):** Initial forces of ventricular depolarization in the Wolff – Parkinson – white syndrome. *Circulation*, 52 : 1030 – 1036.
- Topaz O, Perin E, Cox M, et al.(1989):** Young adult survivors of sudden cardiac arrest: analysis of invasive evaluation of 22 subjects. *Am Heart J*,118:281–7.
- Twidale N, Wang X, and Beckman KJ. (1991):** Factors associated with recurrence of accessory pathway conduction after radiofrequency catheter ablation *PACE*, 14: 2042-2048.
- Yamane T, Jais P, Shah DC, et al. (2000):** Efficacy and safety, of an irrigated- tip catheter for the ablation of accessory pathways resistant to conventional radiofrequency ablation. *Circulation*. 102: 2565- 2568.
- Yang Y, Cheng J, Glatter K, et al. (2000):** Quantitative effects of functional bundle branch block in patients with atrioventricular reentrant tachycardia. *Am J Cardiol*, 85: 826- 831.
- Wathen M, Natale A, Wolfe K, et al. (1993):** Initiation of atrial fibrillation in the Wolff – Parkinson – White syndrome: The importance of the accessory pathway. *Am. Heart J*, 125: 753.

Weiss C, Cappato R, Schluter M, et al. (1995): *Anomalies of the coronary venous system in patients with and without accessory pathways [abstract]. J Am Coll Cardiol 25:18 A.*

Wellens H, and Durrer D. (1974): *Patterns of ventriculoatrial conduction in the Wolff – Parkinson – White syndrome. Circulation, 49 : 22.*

Winters SL, and Gomes, JA. (1986): *Intracardiac electrode catheter recordings of atrioventricular bypass tracts in Wolff – Parkinson – White syndrome: Techniques, electrophysiologic characteristics and demonstration of concealed and decremental propagation. J. Am. Coll. Gardiol, 7 : 1392.*

Xie B, Heald SC, Bashir Y, et al .(1994): *Localization of accessory pathways from the 12-lead electrocardiogram using a new algorithm. Am J Cardiol, 74:161–165.*

Zardini M, Yee R, Thakur R.K., et al. (1994): *Risk of sudden arrhythmic death in the Wolff – Parkinson – White syndrome : Current Perspectives. PACE, 17 : 966.*

