

REFERENCES

- Abdel Rhim, J. Youssef and Chareeb J.A. (1995):** ADA activity and sialic acid in exudative pleural effusions. *Egyptian Journal of Chest diseases and tuberculosis*. 44 (1):93-100.
- Adelman M., Albelda S.M., Gottlieb J. and Haponik C. (1984):** Diagnosis utility of pleural eosinophilia. *Am J Med*; 77:914-20.
- Agostoni E(1986):** Mechanics of the pleural space. In:Fishman A. P. (edi): *Handbook of physiology: The Respiratory System*; Bethesda MD. American physiology society; 531-59.
- Ahn, H.J; Maruo, S.; Tomura, M.; Mu, J.; Hamaoka, T.; Nakanishi, K.; Clark, S.; Kurimoto, M.; Okamura, H and Fujiwara, H. (1997):** A mechanism underlying synergy between IL-12 and INF- γ inducing factor in enhanced production of INF- γ . *J.Immunol*.159:2125.
- Alagiakrishnan K. and Patel P.J. (1999):** Left-sided hepatic hydrothorax with ascites. *Int J Clin Praet*; 53:225-6.
- Al-Harby A., Al-Furayh O., Al-Dayel F. and Al-Mobeireek A. (2006):** Pleural effusion in a patient with end-stage renal disease. *Ann Saudi Med*; 26(2):145-146.
- Allan M., Stephen R.B., Lawrence G.M. and Andrew J.P. (1998):** Investigation of pleural effusion: an evaluation of the new Olympus semi-flexible bronchofiberscope and comparison with Abram's needle biopsy. *Chest*; 114(1):150-154.
- Andrews B.S. (1981):** The role of immune complexes in the pathogenesis of pleural effusions. *Am Rev Respir Dis*; 115- 124.

- Anne C., Tarn B. and Ruth L.W. (2001):** Pleural effusions. Evaluation and management. *Ann Clin Biochem*; 38:311-322.
- Antman K.H. (1993):** Natural history and epidemiology of malignant mesothelioma. *Chest*; 103:3735-65.
- Antoniou, K. M.; Ferdoulsis, E. and Bouros D. (2003):** Interferons and their applications in diseases of the lung. *Chest*; 123:209-16.
- Antony V.B. (1998):** Drug induced pleural diseases. *Clin Chest Med*; 19(2):331-340.
- Antony V.B., Loddenkemper R., Astoul P., Boutin C., Goldstraw P., Hott J., Rodriguez F. and Sahn S.A. (2001):** Management of malignant pleural effusions. *Eur Respir J*; 18:402-419.
- Aoe. K, HiraKi A, Muakami and Eda R (2003):** Diagnositic significance of gamma interferon in tuberculous pleural effusions. *Chest*. 123(3): 740-744.
- Bahr H. (1991):** Carcinoembryoniv antigen as a diagnostic aid in pleural effusions. M.Sc. Faculty of Medicine , Tanta University. Egypt.
- Barnes P.F. and Wizel B. (2000):** Type 1 cytokines and the pathogenesis of tuberculosis. *Am J Respir Crit Care Med*; 161 (6): 1773-1774.
- Beach C. and Manthey E. D. (1998):** Absorption of fluid from pleural cavity. *J Emerg Med*; 16 (1):33-36.
- Beauchamp H.D., Kundra N.K., Aranson R., Chong F. and Mac Donnell K.F. (1992):** The role of closed pleural needle biopsy in the diagnosis of malignant mesothelioma of the pleura. *Chest*; 102:1110-1112.

- Beck, J.L.; Colgrave, M.L.; Ralph, S.F. and Sheil, M.M. (2001):** Mass Spectrometry Review; 20(2):61-87.
- Becklake M.R. (2000):** Pneumoconioses. In: Murray J.F. and Nadel J.A. (eds): Textbook of Respiratory Medicine. Philadelphia. W.B. Saunders; Pp.1955-2001.
- Benoit P., Maguric D., Plavec I., Koocher H., Tovey M. and Meyer F. (1993):** A monoclonal antibody to recombinant human IFN- α receptor inhibits biologic activity of several species of human IFN- α & IFN- β . J Immunol; 150:5401-5407.
- Bernard F., Doniel H.S., Robin J.S., Robin J.S., Larry R.K. and Steven M.S. (1998):** Metabolic imaging of malignant pleural mesothelioma. Chest; 114(3):713-722.
- Boutin C., Viallat J.R., Cargnino P. and Fraisse P.(1991):** Thoracoscopy in malignant pleural effusion. Am Rev Respir Dis; 124:588-92.
- Bradley J., Howard J., Wallace E. and Elborn S. (2000):** Fungal empyema thoracis: an emerging clinical entity. Chest; 117(6):113-21.
- Broadbuss V.C. and Light R.W. (1992):** What is the origin of the pleural transudates and exudates. Chest; 102:658-671.
- Broadbuss V.C. and Light R.W. (2000):** Disorders of the pleura: general principles and diagnostic approach. Chest; 102:658-671.
- Brod S.A., Benjamin D. and Haffler D.A. (1991):** Restricted T cell expression of IL-2/IFN- γ mRNA in human inflammatory disease. J Immunol; 147:810-815.
- Byrd R.P. and Roy T.M. (1998):** Pleural fluid PH determination. Chest; 113(5):1426-1427.

- Camero A.O., Pharm D., Vargas F.S. and Onishi R. (1999):** Cholesterol profile in pleural diseases. *Chest*; 116(4):283-286.
- Cardenas A., Gines P., Arroyo V. and Rodes J. (2004a):** Management of cirrhosis and ascites. *N Engl J Med*; 350:1646-54.
- Cardenas A., Kellerher T. and Chopra S. (2004b):** Hepatic hydrothorax. *Aliment Phamaeol Ther*; 20:271-279.
- Chaignaud B.E.,Bonsack T.A. and Kazakewich H.P. (1998):** Pleural effusion in lymphoblastic lymphoma. *J Pediatr Surg*; 23(9):1355-1367.
- Chang S, Hsu Y Chen Y, Lin Y(1994):** Usefulness of SIL-2R in differentiating tuberculos and carcinomatous effusions. *Arch Intern. Med*; 145.
- Chen, Y. M.; Yang, W.K.; Whang-Peng, J.; Tsai, C.H. and Perng, R. P. (2001):** An analysis of cytokine status in the serum and effusions of patients with tuberculous and lung cancer. *Lung Cancer*; 31: 25-30.
- Collazas J, Espana J, Martiniz F, Lzquerdo F (1998):** Sequential evaluation of serum ADA in patients treated for tuberculosis. *Chest*; 114 (2):432-435.
- Collins T.R. and Sahn S.A. (1987):** Thoracocentesis, complications, patients experience and diagnostic values. *Chest*; 91:817-22.
- Colt H.G.(1999):** Tumor of the pleura. In: Colt H. G., Mathur P.N. (eds); pleural procedure. Philadelphia, Lippincott&Williams, Pp.3-15.
- Connros F.A. and Ahose D.M.(1994):** Pleural inflammation and pleural effusion. In: Baum G.L., Wolinisky E. (eds); Textbook of pulmonary Diseases. NewYork, Marcel Dekker, Pp. 1853-1866.

- Daganou M., Dintopoulou I., Michalopoulos N. and Papa Dopoulas K. (1998):** Respiratory complications after coronary artery bypass. Chest; 113(5):1285-1289.
- Dalton D.K., Pitts-Meek S. and Keshav S. (1993):** Multiple defects of immune cell function in mice with disrupted interferon- γ genes. Science; 259:1739-1742.
- Daniil, Z. D.; Zintzaras, E.; Kiropoulos, T.; Papaioannou, A. I.; Koutsokera, A.; Kastanis, A. and Gouroulanis, K. I. (2007):** Discrimination of exudative pleural effusions based on multiple biological parameters. Eur. Respir. J; 30(5):957-64.
- Darenskaia, S. D.; Makarava, N. V.; Vladimrski, M. A.; Borisov, S. E.; Emelianov, M. N.; Koltsov, K. I.; Efremov, E. E. and Ignashenkova, G.V. (2008):** Significance of determination of gamma- interferon in the diagnosis of tuberculous exudative pleurisy. Probl Tuberk Bolezn Legk.; (2):29-32.
- David B.D., Dulbecco R., Fisen H.N. and Cinsherg H.S. (1990):** Interferons: effects, production after viral infection, relation to cells and chemical induction. J Microbiology; 49:872-877.
- Davies R.D. and Glesson F.M. (1998):** Diagnosis and management of pleural empyema. Curr Opin Infect Dis; 11(2):163-168.
- De Groot R.E., Willem L.N. and Dijkman J.H. (1996):** Interstitial lung disease with pleural effusion caused by simvastin. J Intern Med; 239:361-363.
- Djeu T.Y., Heinbaugli J.A., Hoolden H.T. and Herberman R.B. (1996):** Augmentation of mouse natural killer cell activity by interferon and interferon inducers. J Immunol; 122:175-181.

- Edward F. and Patz J.R.(1998):** Malignant pleural effusions: recent advances. Chest; 133(1):745-775.
- El azab M.E (2003):** Diagnositic value of interferon gamma in tuberculous pleural effusion. MD, Thesis, Faculty of Medicine, Tanta University.
- EL-Arag A.S, Shaheen A M, Saleh M A (1999):** Diagnostic utility of adenosine deaminase activity with lymphocyte/neutrophil ratio in tuberculous pleural effusion.Egy J Chest Dis T uber ;48 (1):65-72.
- El-Bouhy M S and Abdel Wahab I A (1999):** Pleural abrasions through vido-assisted thoracoscopy by a newly developed pleural abrader.Egy J Chest and Tuber;48(2):165-176.
- EL-Helaly S , Ashmawy S and Nagaty M (1999):** pleural fluid to serum bilirubin concentration ratio for differentiation of transudates from exudates.Egy.J of Chest Diseases and tuberculosis:22:2.
- El-Shahat H.M., Abdel-Aziz L.R. and Goher O.A. (2000):** Value of lactate dehydrogenase isoenzyme pattern in the diagnosis of pleural effusion. Egypt J Chest and Tub; 49(1):146-152.
- Esmat A and Fatama A (1999):** Astudy of interferon gamma in pleural effusion.A marker for tuberculosi. Egy J Cest Dis Tuber; 48(1):61-64.
- Fields B.N., Knipe D.M., Chqunock R.M., Molnick F.C., Roizman B.and Shope R.E. (1985):** Virology,Revan press. New York.
- Fisher, R.S. (1964):** Statistical methods for research workers. Edin burgh. Dliver and Boyer.

- Frank, W. (2002):** Tuberculous pleural effusions. *Eur Resp Mon.*; 22:219-33.
- Freeman, W. M.; Walker, S. J. and Vrana, K. E. (1999):** Quantitative RT-PCR: pitfalls and potential. *Biotechniques*, 26, 112–122.
- Gallardo X., Castaner E. and Mata J.M. (2000):** Benign pleural diseases. *Eur J Radiol*; 34:87-97.
- Galvani E.W., Netherselll .B. and Cawley J.C. (1989):** The beneficial effects of alpha interferon in chronic myeloid leukemia. *Br J Haemato*; 111(2):233-237.
- Garcia-Pachon E., Sanchez F.J., Jimaraz B. and Custrody J. (1996):** Pleural fluid to serum cholinesterase ratio for the separation of transudate and exudates. *Chest*; 110:96-101.
- Geo, F.; Janet, B. and Stephen, A. (2001):** Pathogenesis and control of viral diseases. In *Medical Microbiology*. 2nd .ed; 349-351.
- George, R. B. (1990):** Chest Medicine. Essentials of Pulmonary and Critical Care Medicine. Second Edition. Williams & Wilkins. Baltimore, Hong Kong, London; Pp.127-137.
- Goldman J.A., Wang K.P. and Keith F.M. (2000):** Diagnostic procedures. In: Murray J.F. and Nadel J.A. (eds): *Textbook of Respiratory Medicine*. Philadelphia. W.B. Saunders; Pp.711-782.
- Good J.T., Taryle T.A. and Sahn S.A. (1985):** The pathogenesis of low glucose, low PH in malignant effusions. *Am Rev Respir Dis*; 131:737-41.

- Gottelhrer A., Taryle T.A., Reed C.E. and Sahn S.A. (1991):** Pleural fluid analysis in malignant mesothelioma. Prognostic implications. Chest; 100:1003-6.
- Gottelhrer A., Toa J., Stanford G., Chernow B. and Sahn S.A. (1990):** Hypothyroidism and pleural effusions. Chest; 98:1130-1132.
- Greenspan R.H. and Sostman H.D. (2000):** Radiographic techniques. In: Murray J.F. and Nadel J.A. (eds): Textbook of Respiratory Medicine. Philadelphia. W.B. Saunders, Pp. 610-681.
- Gresser I (1985):** Antitumor effects of interferon. Acta Oncologica; 28:347-353.
- Grunewald J., Berlin M. and Olerup O. (2000):** Clinical features of pulmonary sarcoidosis. Am J Cdt Care Med; 161(3):814-898.
- Guterman J.U. (1994):** Cytokine therapeutics: lessons from interferon γ . Proc Natl Acad Sci USA; 91: 1198-1205.
- Hamm H., Pfalzer B. and Fabel H. (1991):** Lipoprotein analysis in chyloform pleural effusion. Implications for pathogenesis and diagnosis. Resp; 58:294-300.
- Hassan A.R., Abdel salam H.M. and Hassan M.R. (2001):** Interferon gamma and tumour necrosis factor alpha as amarker for diagnosis of tuberculous pleural effusion. Egyptian Journal of Chest 50(1): 98-102.
- Heffner J.E. (1998):** Evaluating diagnostic tests in pleural space: differentiating transudates from exudates. Clin Chemist Med; 19(2):261-271.

- Heffner J.E., Nietert P.J. and Barbieri C. (2000):** Pleural fluid PH as a predictor of survival for patients with malignant pleural effusion. *Chest*; 117(1):87-95.
- Hiraki, A.; Aoe, K. and Eda, R. (2004):** Comparison of six biological markers for the diagnosis of tuberculous pleuritis. *Chest*; 125: 987-989
- Hoheisel G, Izbicki G, Roth M, Goodman H (1998):** compartmentalization of pro-inflammatory cytokines in tuberculous pleurisy. *Respir Med*; 92 (1) 14-17.
- Hsu C. (1987):** Cytologic detection of malignancy in pleural effusion; a review of 5,255 samples from 3,811 patients. *Diag Cytopathol*; 3:8-12.
- Hubbard A.T.(1997):** The etiology of mesothelioma: are risk factors other than asbestos exposure. *Thorax*; 52(6):496-498.
- Hurlbut D., Myers M.L. and Lefcoe M. (1990):** pleuropulmonary morbidity. Internal Thoracic artery versus saphenous vein graft. *Ann Thorac Surg*; 50(6):959-64.
- Hyates D.H., Corrin B. and Stidolph P.N.(1997):** Malignant mesothelioma. *Thorax*; 52(6):507-512.
- Imokawa S., Colbym T.V., Leslie K.O. and Helmers R.A. (2000):** Methotrexate pneumonitis: review of the literature and histopathological findings in nine patients. *Eur Respir J*; 15:373-1 .
- Iqbal M., Rossoff L.J., Marzouk K.A., Steinberg H.N. (2000):** Yellow nail syndrome: resolution of yellow nails after successful treatment of breast cancer. *Chest*; 117(5):1516-1517.

- Issacs A. and Lindenman (1957):** Virus interference I. The interferons. Proc R. Soc; 147:258-267.
- Ivan G., Valdimir S., Marija B. and Davor J. (2002):** Pleural pathophysiology. Acta Clin Croat; 41:57-63.
- Jaume S., Ferrer M. and Rannon M. (1996):** Evolution of idiopathic pleural effusion; a prospective long term follow up study. Chest; 109:509-13.
- Jenkinson B. and Banschbach R. (1982):** Prostaglandin E as a parameter for differentiating between transudative and exudative pleural effusion. J Appl Physiol; 105-458.
- Jenkinson S.G. and Sam J. (1993):** Pleural effusion complicating pleural adhesions. Ann Intern Med; 59:177-256.
- Jing, J.; Huan-Zhong S.; Qiu-Li L.; Shou-Ming Q. and Xue-Jun Q. (2007):** Diagnostic Value of Interferon- γ in Tuberculous Pleurisy. Chest. 131:1133-1141.
- Jo E.K, Kim H.J., Min J.H, Mm D., Song Y., Song C.H., Paik T.H. and Suhr J.W. (2000):** Dysregulated production of IFN- γ , interleukins in early tuberculus patients. Scand J Immunol; 51(2): 209-217.
- Johnson C.C. and Finegold (2000):** Pyogenic bacterial pneumonia, lung abscess, and empyema. IN: Murray J.F. and Nadel J.A. (eds): Textbook of Respiratory Medicine. Philadelphia. W.B. Saunders, Pp.1036-1093.
- Johnson J. (2000):** Pleural effusions in cardiovascular disease. Postgrad Med; 107:95–101.

- Jones D., Lieb T., Narita M., Hoollnder E.S. and Pitchenika F. (2000):** Mesothelial cells in tuberculous pleural effusion of HIV-infected patients. *Chest*; 117(1):289-291.
- Jose C., Raed A. and Dwei K. (2005):** Pleural effusions: evaluation and management. *Cleveland Clinic J*; 72;854-872.
- Kazielski J. (1999):** The value of transthoracic biopsy of visceral pleura in diagnosis of neoplastic pleural effusion. *Chest*; 116(4):356s-357s.
- Keane M.P. and Lynch J.P. (2000):** Pleuropulmonary manifestations of systemic lupus erythematosus. *Thorax*, 55(2):159-166.
- Keshk A.A. (2002):** Gamma interferon level in pleural fluid patients with tuberculous pleural effusion. MD. Thesis, Faculty of Medicine, Cairo University.
- Kim YC, Park KO, Bom HS, Lim SC, Park HK, Na HJ, Park JH(1998):** Combining ADA, Protein and IFN- γ best allows discrimination between tuberculosis and malignant pleural effusion. *Korean J Inter Med*; 12(2): 225-231.
- Kimchi A. (1992):** Cytokine triggered molecular pathways that control cell cycle arrest. *J Cell Biochem*; 50:1-9.
- Kinasewitz G.T. (1997):** Pleural fluid dynamics and effusions. In: Fishman A.R., Elias J.A., Fishman J.A., Grippi M.A., Kaiser L.R. and senior M.R. (eds): *Fishman's Pulmonary Diseases and Disorders*. New York, McGraw Hill, Pp. 1381-1409.
- Klein J. (1990):** Lymphokines and their receptors. In: *Immunology*, ed 1, Edinburgh, Melbourne.

- Kohno, K.; Kataoka, J.; Ohtsuki, T.; Suemoto, Y.; Okamoto, I.; Usui, M.; Ikeda, M. and Kurimoto, M. (1997):** INF- γ inducing factor (IGIF) is a costimulatory factor on the activation of Th1 but not Th2 cells. *J. Immunol*; 148:1143.
- Kramer, M.R.; Saldana, M.J. and Cepero, R.J., (1989):** High amylase levels in neoplasm-related pleural effusion. *Ann Intern Med*: 110,567-569.
- Krenke, A.; Safianowska, M.; Nasilowski, B. and Dmowska-Sobstyl, E. (2008):** pleural fluid adenosine deaminase and interferon gamma diagnostic tools in tuberculous pleurisy. *Journal of physiology and pharmacology*.59, Suppl 6, 349–360
- Landrneau R.J., Hazerlrigg S.R, Mack M.J. Keenan R.J. and Ferson P.F. (2000):** Video-assisted thoracic surgery for pulmonary and pleural disease. In: Shields T.W. (ed). *General Thoracic Surgery*. Baltimore, Williams& Wilkins; Pp.508-528.
- Last's Anatomy (1999):** Pleura, In: Chwmy Sinatenly (edi), *Last's Anatomy*. London. Churchill Livingstone, Pp.205-217.
- Lazaridis K.N., Frank J.W., Krowka M.J. and Kamatb P.S. (1999):** Hepatic hydrothorax: pathogenesis, diagnosis and management. *Am J Fed*; 107:262-7.
- Lee LN, Wang JY, Hsu HL and Yang PC (2009):** Enzyme Linked immunospot assay for interferon-gamma in the diagnosis of tuberculous pleurisy. *Clinical Microbiology and infectious diseases*.15(2): 173-179.
- Lee, J.H. and Chang, J.H. (2005):** Diagnostic utility of serum and pleural fluid carcinoembryonic antigen, neuron-specific enolase, and

cytokeratin 19 fragments in patients with effusions from primary lung cancer. *Chest*.:128(4):2298-303.

Light R.W. (1995): In: Light R.W.(edi), *Pleural Diseases*. Baltimore, Williams, Willkins; Pp. 23-47.

Light R.W. (1997): Diagnostic principles in pleural disease. *Eur Respir J*;10(2):476-481.

Light R.W. (1998a): Management of parapneumonic effusions. *Clin Chest Med*; 19 (2):373-382.

Light R.W. (1998b): Establishing the diagnosis of tuberculous pleuritis. *Arch Intern Med*; 158(18):1735-1744.

Light R.W. (2000c): Clinical pulmonary function testing. In: George R.B., Light R.W., Matthy M.A. Matthy R.A. (eds): *Chest Medicine*, Williams & Willkins, Pp.132-162.

Light R.W.(2000a): Pleural effusion. In: Murray J.F. and Nadel J.A.(eds): *Textbook of Respiratory Medicine*. Philadelphia. W.B. Saunder, Pp. 2164-2192.

Light R.W., Stansbury D.W. and Brown S. E. (1986): The relationship between pleural pressures and changes in pulmonary function after therapeutic thoracentesis. *Am Rev Respir Dis*; 133,658-661.

Light, R.W; Mc Gregor M.I. and Luchsinger, P.C. (1972): Pleural effusions: The diagnostic separation of transudates and exudates. *Ann Intern Med*. 77:507-13.

Light, R. W. (2000): Management of pleural effusions. *J. Formos. Med. Assoc.*; 99(7):523-31.

- Lim T.K. and Yim A.P. (1999):** Management of pleural empyema Chest; 166(3):845-846.
- Magnani Z.I., Confetti C., Besozzi G., Codecasa L.R., Bardignon P., Lang R., Rossi G.A., Pardi R. and Burastero S.E. (2000):** Circulating mycobacterium tuberculosis-specific lymphocytes from PPD skin test-negative patients with tuberculosis don't secrete IFN- γ and lack the cutaneous lymphocyte antigen skin-selective homing receptors. Clin Exp Immunol; 199(1):99-106.
- Marina Costa, Sapunar M. and Montero J. (1989):** Rendimiento de los criterios de light on la diferencia de transudates, exudato pleurales. Intern Cir Thorac; 51:24-7. Quoted from Chest (1997); 97:70-80.
- Marten, N. W.; Burke, E. J.; Hayden, J. M. and Straus, D. S. (1994):** Effect of amino acid limitation on the expression of 19 genes in rat hepatoma cells. *FASEB J.*, 8, 538–544.
- Mathur, P.C.; Tiwari, K.K.; Tripathi, S. and Tiwari, D. (2006):** Diagnostic value of adenosine deaminase (ADA) activity in tubercular serositis. *Indian J Tuberc*; 53:92-95.
- Mc Loud T.C. (1998):** CT and MRI in pleural diseases. Clin Chest Med; 19(2):261-271.
- Medford A. and Maskell L.A.N. (2005):** Pleural effusion. Postgraduate Medical Journal 81:702-710.
- Meisel S., Shamiss A., Thaler M., Nussinovitch N. and Rasenthal T. (1990):** Pleural fluid to serum bilirubin concentration ratio for the separation of transudates from exudates. Chest; 98:141-144.

- Miyamoto S., Zellca T., Murata S. and Mabuchi H.(1998):** Detection of diaphragmatic defect as the cause of sever hepatic hydrothorax with magnetic resonance imaging. *Am J Gastroenterol*; 93:2288-9.
- Mobasher A.A, Nafee R.M, Abdel Moneim A.M., Salah A. and Al-Attar H.I.(1997):** Gamma interferon in pleural effusion cases. *Egyptian Journal of Chest Disease and Tuberculosis*. 46 (2):1-9.
- Mobasher AM, EL-Naggar IZ, Ghonicm AH, Yousef AR, Saleh A and Hassan KH (1995):** Neopterin a novel approach to cell mediated immunity assessment in pulmonary tuberculosis and malignant patients. *Egyptian J of Chest disease and tuberculosis*, 44(2): 57-61.
- Monti G.,Jaurand M.C., Monnet I., Chetein P., Saint L., Portiey A., Pevillieux Galanaud P. and Bingnon J. (1994):** Interpleural production of interleukin-6 during mesothelioma and its modulation by gamma interferon. *Cancer Res*; 54(16):4419-4423.
- Moodley Y.P., Dorasamy T., Venketasany S., Naicker V. and Lalla U (2000):** A paired comparison of CD4:CD8 ratio and TNF- α levels in sarcoidosis. *Thorax*; 55(8):676-700.
- Morrison, T.; Weis, J. J. and Wittwer, C. T. (1998):** Quantification of low-copy transcripts by continuous SYBR Green I monitoring during amplification. *Biotechniques*, 24, 954–962.
- Murray J.F. (2000):** History and physical examination. In:Murray J.F. and Nadel J.A. (eds): *Textbook of Respiratory Medicine*. Philadelphia. W.B. Saunders; Pp.563-584.

- Narseen N., Mohamed K., Dowling P., Word M., Galffy G. and Antony V. (2000):** Talc induces apoptosis in human malignant mesothelioma. *Am J Respir Crit Care Med*; 161(2): 595-600.
- Nazan, D.; Ya, S.; Ebubekir, B. and_Irfan, K. (2000):** Zinc Determination in Pleural Fluid. *Turk J Chem*: 24, 377-380.
- Neysman M., Elist K.V., Noppen M., Saeddeleer B.D. and Magnolet M (1999):** Lactate dehydrogenase isoenzymes analysis is not useful in differentiation between malignant and non-malignant. *Chest*; 116(4):343s-345s.
- Nicklson A. and Carrin B. (1998):** Non-Hodgkin Lymphoma. *Thorax*;53 (12):1094-1099.
- Numez M., Peacosk J. and Chin R.C. (1998):** Pulmonary cryptococcosis in immunocompetent host. *Chest*; 118(2):527-534.
- Nunn F.J. (2002):** Physiology of pleural space. In: Lumb A.B., Nunn J.F. (eds), *Nunn's Applied Respiratory Physiology*. Oxford, Butterworth Heinennan, Pp.370-429.
- Ocana I, Martinez-Vasques J M,and Segura M (1983):** ADA in pleural fluid: test for diagnosis of tuberculous effusion. *Chest*; 84 (1):51.
- Oda S. and Sami M. (1990):** Platelet activating factor in exudative pleural effusion. *Chest Department Alex. University Hospital. Egyptian J of Chest Diseases and Tuberculosis*;16.
- O'Donovan P.B. and Eng P. (1994):** Pleural changes in malignant pleural effusions: appearance on computed tomography. *Cleve Clin J Med*; 61:127-131.
- Ogawa K, Koga H, Hirakata Y Tomono K, Tashiro Kohnos (1998):** Differential diagnosis of tuberculous pleurisy by measurement

of cytokine concentrations in pleural effusion. *Tuber Lung Dis*; 78 (1): 29-34.

Okamoto, M.; Kawabe, T.; Wasaki, Y.; Hara, T.; Hashimoto, N.; Imaizumi, K.; Hasegawa, Y and Shimokata, K (2005): Evaluation of interferon- γ , interferon- γ -inducing cytokines, and interferon- γ -inducible chemokines in tuberculous pleural effusions. *J Lab Clin Med*; 145:88–93.

Ostensen M.E., Thiele E.L. and Lipsky P.F. (1989): Enhancement human natural killer cells function by combined effects of tumor necrosis factor alpha. *J Biol Response*; (8):53-61.

Oudkark M.(1999): Pulmonary embolism; Epidemiology. In: Oudkark M.,Van Beck E., Tencate J.W. (eds), *Pulmonary Embolism: Epidemiology, Diagnosis and Treatment*. Berlin, Blackwell Science, Pp.9-30.

Owens M.W. and Milligan S.A.(1995): Pleuritis and pleural effusions. *Curr Opin Pulm Med*; 1:318.

Padimini U., Moffett M., Bryan K., Moffett M., and Damian A.(2009): Diagnosing and managing suspected malignant pleural effusions. *The Journal of supportive oncology* 7 (4): 178-205.

Peek G. J., Morcos S. and Cooper G. (2000): The pleural cavity. *BMJ*; 320(5): 1318-1321.

Peterman T.A. and Brothers S.K. (1983): Pleural effusions in congestive heart failure and in pericardial disease. *N Eng J Med*; 309(5):313.

- Petterson C. and Mill J. (2000):** Parasitic infections. In Murray J.F. and Nadel J.A. (eds). Textbook of Respiratory Medicine. Philadelphia. W.B. Saunders; Pp.1201-1244.
- Petterson T., Klockers M. and Hellstorm P.E. (1982):** Chemical and immunological features of pleural effusion. Comparison between rheumatoid arthritis and other diseases. Thorax; 37:354-61.
- Pfaffl, M. W. (2001):** A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Research. (9) 29 :45.
- Porcel J.M. and Vives M. (1999):** Classic, abbreviated and modified Light's criteria: the end of the story. Chest; 116(6):1833-1835.
- Poyraz, B.; Kaya, A.; Ciledag, A.; Oktem, A. and Gonullu, u. (2004):** Diagnostic significance of gamma – interferon in tuberculous pleurisy. Tuberculous in toraks dergisi 52(3):211-217.
- Pritikin J.D., Jensen W.A., Yenokida G.G., Kirsch C.M. and Fainstat M. (1990):** Respiratory failure due to a massive rheumatoid pleural effusion. J. Rheumatoid; 17:673-4.
- Riantawan R., Chaouwalit P., Wongasaangien M. and Rojanaraweewong P. (1999):** Diagnostic value of pleural fluid ADA in tuberculous pleuritis with reference to HIV co-infection and a Bayesian analysis Chest; 116(1):97-103.
- Ribera I.O., Vazquez J.M., RAuiz I., Bezarano E. and Pigrau C. (1998):** ADA activity in rheumatoid pleural effusion. Ann Rheum Dis; 47: 394-397.
- Rinderknecht E., O'Connor B.H. and Lopez M. (1984):** Natural human interferon gamma: complete amino acid sequence and

determination of sites of glycosylation. J Biol Chem; 259: 6790-6797.

Rodriguez P., Borderas F. and Lopez M. (1998): Pleural metastatic tumors and effusions. Frequency and pathogenic mechanisms in a post-mortems series. Eur Respir J; 2:366-369.

Romero S., Fernandez C., Arriero M., Espasa A., Candela C. and Martin J. (1996): CEA, CA 15-3 and CYFRA 21-1 in serum and pleural fluid of patients with pleural effusions. Eur Respir J; 9, 17-23.

Rubens M.B. (1998): The pleura: collapse and consolidation. In: Murifilt J., Gregson R.H. and Hyde I. (eds). Textbook of Radiology and Imaging. New York, Livingstone; Pp.365-390.

Sadikat R.T., Fredi F.L. and Light R.W.(2000): A43year old man with a large recurrent right side pleural effusion. Chest; 117(4):1191-1194.

Sahn S. A. (1985): Pleural fluid PH in the normal state and diseases affecting the pleural space. In: J Chretien J., Bijron J. and Hirsch A. (eds): Health and Disease. New York; Marcel Decker; Pp.253-66.

Sahn S.A. (1987): Malignant pleural effusion. Seminar Respir Med;9:43-53.

Sahn S.A. (1988): The pleura. Am Rev Respir Dis; 138:184-234.

Sahn S.A. (1997): Malignant pleural effusion. In: Fishman A.P., Elias J.A., Fishman J.A., Grippi M.A., Kaisr L.R. and Sensor R.M. (eds), Fishman's Pulmonary Diseases and Disorders. New York, Mc Graw Hill, Pp.1428-1438.

Sahn S.A. (1998): Malignancy metastastize to the pleura. Clin Chest Med; 19(2):351-361.

- Sahn S.A. and Antony V.B.(1984):** Asbestosis pneumoconiosis; occupational lung disease. *Ann Intern Med*; 102:1002-5.
- Sahn S.A. and Cood J.T. (1988):** Pleural fluid PH in malignant effusions; diagnostic, prognostic and therapeutic implications. *Am Rev Respir Dis*; 126: 121-2.
- Salomaa E.R., Viender M. and Saresranta T. (1998):** Complement component and their activation products in pleural fluid. *Chest*; 114(3):723-730.
- San Jose E., Valdes L., Sarandes A. and Alvarez D. (1997):** Diagnostic value of adenosine deaminase and lysozyme in tuberculous pleurisy. *Clinical Chem Acta*; 209:73-81.
- Sasse S., Nguyen T., Teixeira L. and Light R. (1999):** The utility of daily therapeutic thoracocentesis for the treatment of early empyema. *Chest*; 116(6): 1703-1709.
- Sasson C. S. and Light R.W. (1985):** Chylothorax and pseudochylothorax. *Clin Chest Med*; 6:163-71.
- Seaton A. (2000):** Diseases of the pleura. In: Seaton A., Seaton D. and Leitch A. (eds): *Crofton and Douglas's Respiratory Diseases*. Oxford, Blackwell Scientific, Pp.1152-1181.
- Sharma, S. K. and Banga, A. (2004):** Diagnostic utility of pleural fluid IFN-gamma in tuberculosis pleural effusion. *J. Interferon Cytokine Res.*; 24(4):213-7.
- Shimokata K. (1997):** Cytokines and local cellular immunity. *Nagoya J Med Sci*; 60(3):101-108.
- Simpson G., Roones D. and Davies R. (1997):** Treatment of pleural empyema. *Thorax*; 52(12):1009-1018.

- Singh V., Kharh S. and Ghalaut B. (1998):** Serum ADA in pleural effusion. Thorax; 53(9):813-814.
- Sinnatamby C.S. (2006):** Last's Anatomy: Regional and Applied. 11th edition. London: Churchill Livingstone; Pp. 221-225; 504-506.
- Soderblom T., Nyberg P., Teppo A.N., Klockers M., Riska H. and Pettersson T. (1996):** Pleural fluid interferon gamma and tumor necrosis factor alpha in tuberculous and rheumatoid pleurisy. Eur Resp J; 9:1652-1655.
- Steele P.C. and Rudd R.M. (2000):** Malignant mesothelioma: predictors of diagnosis and clinical trials. Thorax; 55(6):725-729.
- Stefanovski I., Kaeva B. and Kovkarova F. (1999):** The role of blind pleural biopsy in the diagnosis of unconfined pleural effusion. Chest; 116(4):357s.
- Strauss R.M. and Boyer T.D. (1997):** Hepatic hydrothorax. Semin Liver Dis; 17:227-32.
- Tag EL-Din M, Ashmawi S, Tash M, Abdel Fatah N (2000 a):** Estimation of interleukin-2 levels in tuberculous pleural effusion. Egy J Chest and Tuber; 49(1) 158-164.
- Tag EL-Din M, Gohars, A shamawis, Amin M (2000b):** Pulmonary complications in hemodialysis patients. Egy. J Chest and Tuber; 49 (1) 230-237.
- Teklu B. (1999):** Cytology on transudative pleural effusion. Chest; 166(3):846-847.
- Thomas J. (1993):** Management of malignant pleural effusion. Chest; 103:3855-3871.

- Ubright TM and Katazenstein AA (1980):** Solitary necrotizing granuloma of the lung. Differentiating features and aetiology Am. J. Pathol.,4:13.
- Valdes L., Alvarez E., San Jose E., Penel P., Valle J., Garc J., Su J. and Pase A. (1998):** Tuberculous pleurisy, a study of 254 patients. Arch Intern Med; 158:2017-2021.
- Valdes L., Jose E.S., Alvarez D., Sarandeses A., Pose A., Chomon B., Dobano J.M., Salgueiro M. and Sudrez J.R. (1991):** Cholesterol; a useful parameter for distinguishing between pleural exudates and transudates. Chest; 99:1097-1101.
- Valdes, L.; Pose, A. and Jose, E.S. (2003):** Tuberculous pleural effusions. Eur. J. Int. Med.14:77-88.
- Vaz M.A., Antonangelo L., Carbal M. and Onishi R. (1999):** Pleural effusion due to lymphoma. Chest; 116(4):358s-362s.
- Vilcek, J. and Sen, G.C. (1996):** Interferon and other cytokines. In: Fields Virology, 3rd.ed. Field BN. Lippincott-Raven; 2-301.
- Villegas, M. V.; Labrada, L. A. and Saravia, N.G. (2000):** Evaluation of polymerase chain reaction, adenosine deaminase, and interferon-gamma in pleural fluid for the differential diagnosis of pleural tuberculosis. Chest.:118(5):1355-64.
- Villegas, M.V.; Labrada, L.A. and Saravia, N.G. (2000):** Evaluation of polymerase chain reaction, adenosine deaminase and interferon- γ in pleural fluid for the differential diagnosis of pleural tuberculosis. Chest; 118:1355-64.
- Villena V., En Cuetra A.L. and Rodriguez F.P. (1998):** False positive results of CEA in pleural effusion. Chest; 113(4):1143-1144.

- Villena, V.; Pérez, V.; Pozo, F.; López-Encuentra, A.; Echave-Sustaeta, J.; Arenas, J. and Escribano, P. (2002):** Amylase Levels in Pleural Effusions A Consecutive Unselected Series of 841 Patients. *Chest*: 121 (2) 470-474.
- Wachter H, Fuchs D, Hausen A, Reibnegger G and Werner E (1989):** Neopterin as amarker for activation of cellular immunity, Immunologic basis and clinical application. *Advan. In. Clin. Chem.*, 27: 81-140.
- Wallach E., Fellous M. and Revel M. (1982):** Preferential effect of gamma interferon on syntheses of HLA antigens and their mRNA in human cells. *Nature*; 49:299-333.
- Warwich R. and Williams P.L. (2005):** *Gray's Anatomy*. Pp. 362-367; 1059-1062.
- Weight D.A., Langford M.P. and Fleildmann W.R. (1983):** Potentiation of lymphocyte natural killer by mixtures of alpha or beta interferon with recombinant gamma interferon. *Infect immune*; 40:35-39.
- Weiss J.M. and Spodick D.H. (1984):** Laterality of pleural effusion in CHF. *Am J Cardiol*; 53(7):951.
- Weynard B., Collard P. and Galant C. (1998):** The utility of open biopsy in patients with hematologic malignancies. *Am J Respir Crit Care Med*; 161(3):723-790.
- White D.A, Wong P.W, Douney R.(2000):** The utility of open biopsy in patients with hematologic malignancies. *AM. J. Respir Cirt Care Med*; 161 (3): 723-790.
- Wiener-Kronish J.P., Goldstein R., Mathy R.A., Biondi J.W., Broadus V.C., Chatterjee K. and Mattay M.A. (1993):** Lack of

association between pleural effusion with chronic pulmonary arterial and right atrial hypertension. *Chest*; 92: 967-70.

Winterbauer R.H. (1997): Non-malignant pleural effusions. In: Fishman A.P., Elias J.A., Fishman J.A., Grippi M.A., Kaiser L.R. and Senior M.R. (eds): *Fishman's Pulmonary Diseases and Disorders*. New York, Mc Graw Hill; Pp. 1411-1427.

Wipa R., Thipawan B. and Eakkapol U. (2010): Chronic Pancreatitis Presenting with Right Pleural Effusion: A Case Report *J Med Assoc Thai*; 93 (3): 378-82.

Wonga, C.; Yewa, W.; Leunga, S.; Chanb, C.; Huib, M.; Yeangb, C. and Augustine Fun-bun (2003): Assay of pleural fluid interleukin-6, tumour necrosis factor-alpha and interferon-gamma in the diagnosis and outcome correlation of tuberculous effusion. *Respiratory Medicine* 97, 1289–1295.

Wongtin S., Silachmroon U., Ruxrungthan K. (1999): Interferon gamma for diagnosing tuberculous pleural effusions. *Thorax*; 54 (10): 921-924.

Xu , F.; Chen, J.; Shen, H.; Wang, X. and Shan, J. (2004): The diagnostic significance of the detection of cytokeratin 19 mRNA by quantitative RT-PCR in benign and malignant pleural effusions. *J Zhejiang Univ Sci.*; 5(10): 1286–1289.

Yigla M., Simsolo C. and Goralnik L. (2002): the problem of empyematous pleural effusion in rheumatoid arthritis: report of two cases and review of literature. *Clin Rheumatol*; 3(2):125-129.

Younge H.A. and Hardy K.J. (1990): Producer cells, activation, stimuli and molecular genetics regulation. *Pharm Ther*; 45:137-143.

- Yousef A.A., El-shaer O.S, Eissa S.A., Yousef S.A., Enan A. and Morghan M.A. (1997):** Diagnostic value of neopterin in pleural effusion. Egy. J. Chest Dis and Tuber; 46 (1): 53-62.
- Zocchi, L. (2002):** Physiological and Pathophysiology of pleural fluid, Eur. Respir. J; 20:1545.