

References

A

- Aasen, J. and Oeding, P. (1971):** Acta Pathologica et Microbiologica Scandinavica B79:287. Cited from Easmon, C.S.F. and Goodfellow, M. Staphylococcus and Micrococcus in Topley and Wilson s vol 2.ch 2.8. Edward Arnold.168.
- Acland, K.M. Darvay, A. and Griffin, C. (1999):** Staphylococcal scalded skin syndrome in an adult associated with methicillin-resistant, Staphylococcus aureus. Br J Dermatol 140, 518–520.
- Aguita and Maria. (2009):** "Combatting Bacterial Infection". LabNews.co.uk.http://www.labnews.co.uk/prINTER_FRIENDLY.php/2912/co_bating-bacterial-infection. Recieved on 2009-05-05.
- Ameisen, J.C. and Capron, A. (2007):** Aspirin sensitive asthma. Clin. Exp. Allergy.; 20: 127-129.
- Amyes, S.G.B. Paton, R.H. and Miles, R.S. (1992):** Ofloxacin as the standard 4- quinolone in sensitivity tests. Journal of Antimicrobial Chemotherapy. 30:101-102.
- Andersen, K. White, I. and Goossens, A. (2006):** Allergens from the standard series. In: Frosch PJ, Menné T, Lepoittevin J-P, eds. Contact Dermatitis, 4th edn. Berlin: Springer, 466–468.
- Aoki, T. Kojima, M. and Horiko, T. (2006):** Acute urticaria; history and natural course of 50 patients J. Dermatol.; 21: 73-77. Quoted from Zuberbier T.; Ifflander J. Semmler C. and HenzB.M. (2007) : Acute urticaria: Clinical aspects and therapeutic responsiveness. Acta. Dermatol. Venereol.; 76: 2957.
- Arias, E. Cesar, A. and Barbara, E.(2009) :** Antibiotic-Resistant Bugs in the 21st Century- A Clinical super-challenge " NEJM 360(5): 439-443.

- Armenaka, M. and Rosenstreich, D.L. (2008) :** The pathophysiology of chronic urticaria .Clinical Reviews. Allergy; 10:257-272.
- Asero, R. (2008):** Leukotriene receptor antagonists may prevent NSAID-induced exacerbations in patients with chronic urticaria. Ann. Allergy Asthma immunol.; 85(2): 150-157.
- August, P.J. and O'Driscoll, J. (2006):** Urticaria successfully treated by desensitization with grass pollen extract. Br. J. Dermatol.; 120: 409-410.

B

- Baird, D. (1996) :** Staphylococcus: cluster-forming gram-positive cocci. in Collee J.G. Fraser, A. G. Marmion, B. P.(eds) Mackie& McCartney Practical Medical Microbiology.Churchill Livingstone. 245-274.
- Barber, M. and Rozwadowska-Dowzenko, M. (1948):** Infection by penicillin resistant staphylococci. Lancet. 252: 641-644.
- Barlow, R.J. Warburton, F. and Watson, K. (2008):** The incidence of delayed pressure urticaria in patient with chronic idiopathic urticaria. J. Am. Acad. Dermatol.; 29: 954-958.
- Bayston, R. Ashraf, W. and Smith, T. (2007):** "Triclosan resistance in methicillin-resistant Staphylococcus aureus expressed as small colony variants: a novel mode of evasion of susceptibility to antiseptics".J.Antimicrob.Chemother.59(5): 848-853.
- Beal, G.N. (2007):** Urticaria : a review of laboratory and clinical observation. Medicine.; 43: 131-151.
- Beck, M.H. and Wilkinson, S.M.(2010) :** **Contact Dermatitis** in (eds) Burns T, Breathnach S , Cox N: Rook's Textbook of Dermatology , 8th edition, Blackwell Publishing Ltd ,ch 1:26-84.
- Beltrani, V.S. (1996):** Urticaria and angio-oedema. Dermatol. Clin.; 14:171-198.

- Bergdoll, M .S. (1983):** In Easmon C.S.F, Adlam C (eds) Staphylococci and staphylococcal infections , vol.2. Academic press, London, 559.
- Bhakdi, S. and Trantum-Jensen, j. (1991):** Alpha-toxin of Staphylococcus aureus. Microbiol Rev 55:733.
- Black, A.K. (2007):** The pathogenesis of urticaria. Keio. Med. J.; 46: 37-39.
- Bohach, G. A. Fast, D. J. and Nelson, R. D. (1990):** Staphylococcal and streptococcal pyrogenic toxins involved in toxic shock syndrome and related illnesses. Crit. Rev. Microbiol. 17:251-272.
- Bohach, G. A. Dinges, M. M. and Mitchell, D. T. (1997):** Exotoxins, In Crossley, K. B. Archer, G. L. (ed.), The staphylococci in human disease. Churchill Livingstone, Inc., New York, 83:111.
- Bohach, G. A. Jablonski, L. and Roggiani, M. (1998):** Biological activity of pyrogenic toxins delivered at the mucosal surface. In Arbuthnott, J. ; Furman, B. (ed.), European Conference on Toxic Shock Syndrome. International Congress and Symposium Series 229. Royal Society of Medicine Press Ltd, New York, N.Y. 170-172.
- Bohmeyer, J. Heller, A. and Harting, E. (1996):** Association of chronic urticaria with staphylococcal infection.: Ann. Rev.Med .43:135-145.
- Bressler, R.B. Sowell, K. and Huston, D.P. (1998) :** Therapy of chronic idiopathic urticaria with nifedipine: demonstration of beneficial effect in double-blinded placebo controlled cross over trial. J. Allergy Clin. Immunol.; 83: 756-763.
- Brestel, E.P. and Thrush, L.B. (2002):** The treatment of glucocorticosteroid dependent chronic urticaria with stanazolol. J. Allergy Clin. Immunol.; 82: 265-269.
- Brooks,G.F, Carroll,K.C, and Butel,J.S. (2007):** Antimicrobial chemotherapy.ch 10 in Jawetz, Melink and Adelbergs(eds), medical microbiology, (24) edition Mc Graw Hill: 161-196.

- Buhner, D. and Grant, J.A. (2005):** Serum sickness. *Dermatol. Clin.*; 3: 107-117.
- Burrall, B.A. Halpan, G.M. and Huntly, A.C. (2008):** Chronic urticaria. *West. J.Med.*; 152:268-276.

C

- Campli, D.C. Gasbarrin, A. and Nucera, E.(1998):** Beneficial effects of staphylococcus eradication in chronic idiopathic urticaria .;43(6):1226-1229.
- Cesar, A . Murray, C. and Barbara, E. (2009):** "Antibiotic-Resistant Bugs in the 21st Century - A Clinical Super-Challenge". *New England Journal of Medicine* 360 (5): 439–443.
- Champion, R.H. Greaves, M.W. and Kobza Black, A. (2006):** The urticarias. *J. Clin. Invest.*; 96: 2606-2612.
- Champion, R.H. Roberts, S.O.B. and Carpenter, R.G. (2008):** Urticaria and angioedema a review of 554 patients. *Br. J. Dermatol.*; 81: 588-597.
- Champion, R.H. (2009):** Urticaria. In: *Textbook of Dermatology*. R.H. Champion, J.L. Burton and F.L.G. Ebling. (eds) Fifth edition. Black well Scientific publication. 2133 - 2139.
- Clauditz, A . Resch, A. and Wieland, K . P . (2006):** "Staphyloxanthin plays a role in the fitness of Staphylococcus aureus and its ability to cope with oxidative stress". *Infection and immunity* 74 (8): 4950–4953.
- Coleman, D. C. Arbuthnott, J. P. and Pomeroy, H. M. (1986):** Cloning and expression in *Escherichia coli* and *Staphylococcus aureus* of the beta-lysin determinant from *Staphylococcus aureus*: evidence that bacteriophage conversion of beta-lysin activity is caused by insertional inactivation of the beta-lysin determinant. *Microb. Pathog.* 1:549-564.

- Cosgrove, S.E. Vigliani, G.A. and Campion, M. (2009):** "Initial low dose gentamicin for Staphylococcus aureus bacteremia and endocarditis is nephrotoxic". Clin Infect Dis. 48 (6): 713–721.
- Coulson, C. J. (1994):** "Bacterial RNA-Polymerase - Rifampin as Antimycobacterial." Molecular Mechanisms of Drug Action. 2nd (ed). Bristol, PA: Taylor Francis, 40-41.
- Cox, N.H. and Coulson, I.H. (2010):** Diagnosis of Skin Disease in (eds) Burns T, Breathnach S, Cox N: Rook's Textbook of Dermatology , 8th edition, Blackwell Publishing Ltd ,ch:5.22 .

D

- Davis, S.L. McKinnon, P.S. and Hall, L.M. (2007):** Pharmacotherapy 27 (12): 1611–1618.
- Dinges, M. M. Orwin, P. M. and Schlievert, P. M.(2000):** Exotoxins of Staphylococcus aureus. Clin Microbiol Rev. 13:16–34.
- Dover, J.S. Black, A.K. and Greaves, M.W. (2008):** Delayed pressure urticaria. J. Am. Acad. Dermatol.; 18: 1289-1298.

E

- Earhart, C. A. Mitchell, D. T. and Murray, D. L. (1998):** Structures of five mutants of toxic shock syndrome toxin-1 with reduced biological activity. Biochemistry 37:7194-7202.
- Easmon, C.S.F. and Adlam, C.(1983) :** Staphylococci and staphylococcal infections. In Easmon and Adlam vol,1 and 2 Academic Press, London. 20-39.
- Egan, C.A. and Rallis, T.M. (2001) :** Treatment of chronic urticaria with Ketotifen. Arch. Dermatol.; 133: 147-149.

- Ellings, A.R. and Pedersen, K.T. (1999):** Treatment of chronic idiopathic urticaria with topical steroids. *Acta. Dermatol. Venereol.*; 76: 43-44.
- Erel, F. Sener, O. and Erdil, A.(2000):** Impact of staphylococcus aureus and Giardia lamblia infection on chronic urticaria .*J. Investig.Allergol. Clin. Immunol.*;10(2):94-97.
- Eriksen, N.H.F. Espersen, V.T. R. and Jensen, K. (1995):** Carriage of staphylococcus aureus among 80 healthy persons during a 19 month period.*Epidmiol.Infect.* 115:51-60.
- Ertam, I. Biyikli, S.E.and Yazkan, F.A.(2007):** The frequency of nasal and throat carriage in chronic urticaria patients.*JEAD*,21, 777-780.

F

- Fischer , W. Rösel , P. and Koch , H. U. (1981):** Effect of alanine ester substitution and other structural features of lipoteichoic acids on their inhibitory activity against autolysins of *Staphylococcus aureus*. *J. Bacteriol.* 146:467-475.
- Forsgren, A. (1970) :** Significance of protein A production by staphylococci . *J.Infect. Immun.*2:672-673.
- Francois, P. and Schrenzel, J. (2008):** "Rapid Diagnosis and Typing of *Staphylococcus aureus*". *Staphylococcus: Molecular Genetics*. Caister Academic Press. 978-981.
- Frain-Bell, W. and Magnus, I . A. (2005):** A study of the photosensitivity factor in cutaneous lymphocytoma. *Br. J. Dermatol.* 84: 25-31.
- Free man, B. A. (2005):** (Eds) Text book of Microbiology, Twenty-first edition. W.B. Saunders Company Philadelphia - London - Toronto. 429 – 440.

G

- Gala Ortiz, G., Cuevas, A.M. and Martinez, E.P. (2001):** Chronic urticaria and staphylococcus aureus . Ann. Allergy Asthma Immunol.;86(6): 696-698.
- Gedebou, M. (1982):** Staphyloococcus aureus strains from a teaching hospital, clinical Bourses and anti biogram, E. Afr. Med. J., Vol(1): 810 - 815.
- Geier, J. Gefeller, O. and Wiechmann, K. (1999):** Patch test reactions at D4, D5 and D6. Contact Dermatitis; 40: 119–126.
- Gemmell, C. G. King, S. L. and Sturrock, R. D. (1987):** Teichoic acid antibody measurement as an aid to the diagnosis of septic arthritis due to Staphylococcus aureus. Serodiagnosis and Immuntherapy. 1:201-207.
- Genigeorgis, C. Savoukidis, M. and Martin, S. (1971):** Initiation of staphylococcus growth in processed meat environments. Appl. Microbiol. 21: 940.
- Gentry, W.C. and Walverton, S.E. (2005):** Drug reaction. Presentation drug reaction workshop. Academy of Dermatology Anni Meeting. Quoted from Beltrani V.S. (1996) : Urticaria a angio-oedema. Dermatol. Clin.; 14: 171-198.
- George, J.J. and Adam, H.(1995) :** Staphylococcus aureus. In : Clinicians Manual on staphylococcus aureus :93-98.
- Ghuysen, J.M. and Hakenbeck, R.(1994):** Structure of the cell wall of staphylococcus aureus , Copenhagen.II .Separation and structure of disaccharides. Biochemistry 2,1119-1125.
- Gilani, S. J. Gonzalez, M. and Hussain, I. (2005):** Staphylococcus aureus re-colonization in atopic dermatitis: beyond the skin. Clin.Exp. Dermatol. 30: 10-13.
- Gilligan, P. H. Gage, P. A. and Welch, D. F. (1987):** Prevalence of thymidine-dependent Staphylococcus aureus in patients with cystic fibrosis. Journal of Clinical Microbiology. 25:1258-1261.

- Graham, S. and Mae, F. (1993):** *Helicobacter pylori*, Current status. Gastroenterol.; 105: 179-182.
- Grattan, C.E.H. O'Donnell, B.F. and Francis, D.M. (2000):** Randomized double blind study of cyclosporine in chronic idiopathic urticaria. Br. J. Dermatol.; 143: 365-372.
- Grattan, C.E.H. and Warin, R.P. (2006):** A serological mediator in chronic idiopathic urticaria, a clinical immunological and histological evaluation. Br. J. Dermatol.; 114:583-590.
- Grattan, C.E.H. Francis, D.M. and Greaves, M.W. (2007):** Plasmapheresis for severe, unremitting chronic urticaria the Lancet; 339(2): 1075-1080.
- Greaves, M.W. (1981):** New Approaches to investigation and management. Austral. J. Dermatol.; 22: 44-52. Quoted from Burrall B.A.; Halpan G.M. and Huntley A.C. (2007) : Chronic urticaria. West. J.Med.; 152:268-276.
- Greaves, M.W. (1991):** Urticaria and angioedema. Diagnosis and evaluation. J. Am. Acad. Dermatol.; 25: 166-176.
- Greaves, M.W. (1995):** Chronic urticaria. N. Eng. J. Med.; 332: 1767-1772.
- Greaves, M.W. and Sabroe R.A. (1998):** Allergy and skin (urticaria). B.M.J.;316: 1147-1150.
- Greaves, M.W. (2000):** Chronic urticaria. J, Allergy Clin. Immunol.; 105: 664-672.
- Greaves, M.W. (2002):** Pathophysiology of chronic urticaria. Int. Arch. Allergy Immunol.; 127(1): 3-9.
- Greaves, M.W, Kaplan, A.P , Grattan, C. (2009):** Diagnostic Techniques for Urticaria and Angioedema in(eds) Kaplan AP and Greaves MW: Urticaria and Angioedema ,2nd edition , Informa Healthcare USA, Inc. ch:8, 141-152
- Green, G.R. Koelsche, G.A. and Kierland, R.R. (2004):** Aetiology and pathogenesis of chronic urticaria. Ann. Allergy.; 23: 30-36.

Griffith, R.S. (1981): "Introduction to vancomycin". Rev Infect Dis 3(suppl.).S200- S2004.

Gruber, B.L. Beza, M.L. and Marchese, M.J. (2006): Prevalence and functional role of anti IgE autoantibodies in urticaria syndromes. I.Invest. Dermatol.; 90: 213 - 217.

Grundmann, H . Aires-de-Sousa, M . and Boyce, J . (2006): Emergence and resurgence of methicillin-resistant Staphylococcus aureus as a public-health threat. Lancet. 368:874-885.

H

Handley, D . A. Magnetti, A. and Higgins, A. (1998) : Therapeutic advantages of third generation antihistamines. Exp. Opin. Invest. Drugs. : 7 : 1045-1054.

Harving, H. Korsgard, J. and Dah, R. (2000): Successful management of chronic urticaria. Allergy. 10: 371-390.

Hawk, J. L. (2006): Cutaneous photobiology; Solar urticaria In Textbook of Dermatology.R.H.Champion, J.L. Burton and F.J.G. Ebling(eds) Fifth edition. Black-well scientific publications .861-863.

Hide, M. Francis, D. M. and Grattan, C. E. (2007): The pathogenesis of chronic urticaria, new evidence suggests an autoimmune basis and implication of treatment, Clin. Exp. Allergy. 25: 624-627.

Hidvegi, B. Cabello, G.R. and Temesvari, E. (2001): The effect of heat inactivated S. aureus on the blastogenic response of peripheral blood mononuclear cells of patients with chronic urticaria Int.Arch.Allergy Immunol.; 126(2): 167-172.

Hiramatsu, K . Hanaki, H . and Ino,T. (1997): Methicillin- resistant Staphylococcus aureus clinical strains with reduced vancomycin susceptibility. J, Antimicrob. Chemother. 40:135-136.

- Hirschmann, J.V. Lawer, F. and Grattan, C. E. (2007):** Chronic urticaria. A clinical and histologic study. Arch. Dermatol.; 123:462-465.
- Hochkeppel, H. K. Braun, W. F. and Vischer, W. (1987):** Serotyping and electron microscopy studies of Staphylococcus aureus clinical isolates with monoclonal antibodies to capsular polysaccharide types 5 and 8. J. Clin. Microbiol. 25:526-530.
- Howden, B.P. and Grayson, M.L. (2006):** "Dumb and dumber—the potential waste of a useful antistaphylococcal agent: emerging fusidic acid resistance in Staphylococcus aureus". Clin Infect Dis 42 (3): 394–400.
- Huang, Y. C. SU, L.H. and Lin, T.Y. (2004).** Nasal carriage of methicillin-resistant Staphylococcus aureus in contacts of an adolescent with community-acquired disseminated disease . Pediatric Infect Dis J;23:919-922.

J

- James, J.and Warin, R.P. (1971) :** An assessment of the role of candida albicans and food yeasts in chronic urticaria.Br. J. Dermatol;84:227-235.
- Jawetz , E. Melnick, J. L. and Adelberg, E.A.(2004):** The staphylococci in Medical Microbiology, international edition (23ed). New York Mc Graw Hill (14): 223-228. .
- Jeffries, C. D. (1961):** Comparison of six physiological characteristics of staphylococci from laboratory specimens. American Journal of Clinical Pathology. 36:114-118.
- Joklik, W.K. Willett, H.D. and Almos, D.B.(1984):** Staphylococci in :Zinsser Microbiology, 18th edition , Applton-century-Croft-Norwalk-Connecticut. 444.
- Jones, R.R. Bhogal, B.and Schfferli, J.(2003):**Urticaria and vasculitis: histological and immunopathological changes Br. J. Dermatol. ;108:194-201.

- Jonker, M.J. and Bruynzeel, D.P. (2000):** The outcome of an additional patch-test reading on days 6 or 7. *Contact Dermatitis*; **42**: 330-335.
- Jorgenson, J. H. and Kloos, W. E. (1987):** Staphylococcal Infections. In B. B. Wentworth (eds), *Diagnostic procedures for bacterial infections*. (7th) ed. American Public Health Association. Washington, D. C.377-382.
- Jorrizo, J.L. and Smith, E.B.(2002):** The physical urticarias.*Arch Dermatol*;118:194-201.
- Julianelle, L.A. and Meghord, C.W. (2005):** Immunological specificity of staphylococci, occurrence of serological types. *J. exp. med* 62, 11-21.
- Juhlin, L. Michaelsson, G. and Zetterstrom, O. (2006):** Urticaria and asthma induced by food and drug additives in patients with Aspirin sensitivity, *J. Allergy Clin. Immunol.*;50:92-98.
- Jungk, J . Como-Sabetti, K. and Stinchfield , P. (2007):** Epidemiology of Methicillin-Resistant *Staphylococcus aureus* at a Pediatric Healthcare System, 1991-2003. *Pediatr Infect Dis J*. 26:339-344.

K

- Kamio, Y. Rahman, A. and Izaki, K. (1993):** The two staphylococcal bi-component toxins, leukocidin and gamma- hemolysin, share one component in common. *FEBS Lett*. 321:15-18.
- Karakawa, W. W. and Vann, W. F. (1982):** Capsular polysaccharides of *Staphylococcus aureus*. *Semin. Infect. Dis*. 4:285-293.
- Kasbekar, N. (2006):** "Tigecycline: a new glycyclcycline antimicrobial agent.". *Am J Health Syst Pharm* 63 (13): 1235–1243.
- Kasten, B. and Reski, R. (1997) :** β -lactam antibiotic inhibit chloroplast division in a moss(*Physocmitrella patens*) but not in tomato(*Lycopersicon esculentum*). *Journal of Plant Physiology* 150(1): 137-140.

- Kauppinen, K. Juntunen, K. and Lank, H. (2007):** Urticaria in children. Retrospective evaluation and follow up. *Allergy*; 39: 469-472.
- Kenneth, T. (2005):** Edited from www.textbookofbacteriology.net.
- Kennard, C. D. and Ellis, C. N. (1998) :** Pharmacological therapy for urticaria. *J. Am. Acad. Dermatol.*; 25: 176-187.
- Kirby, W .M. (1944):** Extraction of a high potent penicillin inactivator from penicillin resistant staphylococci. *Science*. 99:452-453.
- Kirchner, J.T. (2000):** Drug therapy for chronic urticaria post graduate. *Med. J*; 100. (4) :48-58.
- Klefeanoff, S.J. (2000):** Myeloperoxidase: Contribution of the microbicidal activity of intact leucocytes. *Science*, 1069-1075.
- Klebanoff, S.J. (2003):** Oxygen metabolism and the toxic properties of phagocytes. *Ann. int. Med.*, Kennard CD. and Ellis C.N. (1991): Pharmacological therapy for urticaria. *J. Am. Acad. Dermatol.*; 25: 176-187.
- Kloos, W.E. and Schleifer, K.H. (1975):** Simplified scheme for routine identification of human staphylococcus species. *J. Clin. Microbiol.* 1:82-88.
- Kloos, W. E. and Schleifer, K .H. (1981):** In Starr, MP and Stolp, H (eds) *The prokaryotes*, vol.2, Springer, Berlin. 1548.
- Kloos, W.E. and Jorgensen, J.H.(1985):** Staphylococci . In: *Manuals of Clinical Microbiology*. Balowas, W.J.; Hausler H.J.(eds), 4th edition, American Society for Microbiology, Washington. 143-153.
- Khuytmans, J. van Belkum, A. and Verbrugh, H. (1997):** "Nasal carriage of Staphylococcus aureus: epidemiology, underlying mechanisms, and associated risks". *Clin. Microbiol. Rev.* 10(3): 505–520.
- Knieh, E. Becker, A. and Forster, D.H. (2005):** Bed, bath and beyond :pitfalls in prompt eradication of methicillin-resistant staphylococcus aureus carrier status in healthcare workers. *J Hosp Infect*; 59: 180- 187.

Kobza-black, A. (2007): Cold urticaria. *Seminars in Dermatology*; 6(4):292-300.

Kohli, R.M. Walsh, C.T. and Burkart, M.D. (2002): "Biomimetic synthesis and optimization of cyclic peptide antibiotics". *Nature* 418 (6898): 658–661.

Kontinen, V.P. Saris, P. and Sarvas, M.(1991): Ultrastructure of gram-positive cell walls, In Fishetti, V. A. Novick, R.P. and Feretti, J. J. (eds), *Gram-positive pathogens*. ASM Press, Washington, D.C.3-10.

L

Labenz, J. Gyres, E. and Rahl, G. (1992) : *Helicobacter pylori* infection and clinical course of peptic ulcer disease. *Eur. J. Gastroenterol. Hepatol.*;4:893-896.

Ladhani, S. Joannou, C. L. and Lochrie, D. P. (1999): Clinical, microbial, and biochemical aspects of the exfoliative toxins causing staphylococcal scalded-skin syndrome. *Clin. Microbiol. Rev.* 12:224-242.

Lahti, A. VonKrogh, G. and Maibach, H.L. (2006): Contact urticaria syndrome; an expanding phenomenon. In: *Dermatologic immunology and allergy*. Stone J. St. Louis (ed) Mosby. 379.

Lahti, A. and Maibach, H.L. (2008): Immediate contact urticaria: Contact Urticaria syndrome. *Seminars in Dermatology*; 6(4): 313.

Laurence, A.B. and Robert, L.B. (2001): Defects in neutrophil leucocyte function. In: *Clinical Immunology update. Reviews for physicians*. (Eds.) Edward C. Franklin, M.D.Editor in chief. Churchill livingstone, Edinburgh, London and Melbourne., 327.

Lawrence, C.M. Jorrizo J.L. and Kobza-Black, A. (2004) : Cholinergic urticaria with associated angio-oedema. *Br. J. Dermatol.*; 105: 543-548.

- Lee, K.H. Fitton, J.E. and Wuthrich, K. (1987):** Nuclear magnetic resonance investigation of the conformation of D-haemolysin bound to dodecylphosphocholine micelles. *Biochim. Biophys. Acta* 911:144-153.
- Lee, C. Y.(1995):** Association of staphylococcal type-1 capsule-encoding genes with a discrete genetic element. *Gene* 167:115-119.
- Lee, A. and Rouke, J. (2003):** Nasal and throat bacteria other than staphylococcus aureus . *Microbiol. Clin .North AM.;*22:21-42.
- Leenutaphong, K. Holzle, E. and Plewing, G. (2009):** Pathogenesis and classification of solar urticaria: A new concept. *J. Am. Acad. Dermatol.;* 210: 237-240.
- Leigh, I.M. and Ramsay, C.A. (2005):** Localized heat urticaria treated by inducing tolerance to heat. *Br. J. Dermatol.;* 92: 191-194.
- Leznoff, A. Josse, R.G. and Dolovich, J. (2003):** Association of chronic urticaria and angioedema with thyroid autoimmunity. *Arch. Dermatol.;* 119: 636.
- Leznoff, A. and Sussman, G.L. (2005):** Syndrome of idiopathic chronic urticaria and angioedema with thyroid autoimmunity. A study of 90 patients. *J. Allergy Clin. Immunol.;* 84: 66-71.
- Linda, M. W. Don, B. C. and Steven, R. G. (2003):** Genetic Analysis of a High-Level Vancomycin-Resistant Isolate of *Staphylococcus aureus*. *Science* 302 (5650): 1569 – 1571.
- Lindsay, J.A. Kurepina, N. and Novick, R.P. (1998) :** Clinical isolates of *Staphylococcus aureus* encode TSST-1 on genetic elements related to *S. aureus* pathogenicity island-1 (SaPI1). *European Conference on Toxic Shock Syndrome*. Arbutnott, J., and Furman, B. (eds). Royal Society of Medicine, London .1650-1654.
- Liutu, M. Kalimo, H. and Kalimo, H. (2007):** Etiological aspects of chronic urticaria. *Int. J. Drematol.;* 37: 515-519.
- Lowy, F. D. (1998):** *Staphylococcus aureus* infections. *N. Engl. J. Med.* 339:520-532.

Lu, P.L. Chin, L.C. and Peng, C.F.(2005): Risk factors and molecular analysis of community methicillin-resistant *Staphylococcus aureus* carriage. *J.Clin Microbiol*;43: 132-139.

M

Manzella, J.P. (2001): "Quinupristin-dalfopristin: a new antibiotic for severe gram-positive infections.", *Am Fam Physician*. 1;64(11):1863-1866.

Matsuzaki, S. Yasuda, M. and Nishikawa, H. (2003): "Experimental protection of mice against lethal *Staphylococcus aureus* infection by novel bacteriophage phi MR11". *J. Infect. Dis*. 187 (4): 613–624.

Matthews, K. R. Roberson, J. and Gillespie, B.E. (1997): "Identification and Differentiation of Coagulase-Negative *Staphylococcus aureus* by Polymerase Chain Reaction". *Journal of Food Protection* 60 (6): 686–688.

Matthews, K.P. (2005): Exploiting the cold urticaria model. *N.Eng. J. Med.*; 305:1090-1091.

May, L.M. (2005): Evaluation of urticaria In Primary Care Medicine Office evaluation and management of adult patient. A.H. Gorall, L.A. May, AG. Mullery (eds). 3rd edition. Philadelphia lippincott. 895-899.

Mehregan, D.R. Hall, M.J. and Gibson, L.E. (2008): Urticarial vasculitis: A histopathologic and clinical review of 72 cases. *J. Am. Acad. Dermatol.*; 26: 441-448.

Melish, M. Fukunaga, C. and Murata, S. (1998): TSST-1 dissemination in a vaginal model for TSS. In J. Arbuthnott, and B. Furman (ed.), *European Conference on Toxic Shock Syndrome. International Congress and Symposium Series 229*. Royal Society of Medicine Press Ltd, New York, N.Y.169.

Mellor, I. R. Thomas, D. H. and Sansom, M. S. (1988): Properties of ion channels formed by *Staphylococcus aureus* -toxin. *Biochim. Biophys. Acta*. 942:280-294.

- Mittman, R.J. Bernstein, D.L. and Steinberg, D.R. (2008):** Progesterone responsive urticaria and eosinophilia. *J. Allergy Clin. Immunol.*; 84: 304-309.
- Mollby, R. (1983):** in Easmon, C.S.F. Adlam, C. (eds) *Staphylococci and staphylococcal infections*, vol. 2. Academic Press, London, 619.
- Monecke, S. Slickers, P. and Ellington, M.J. (2007):** High diversity of Panton-Valentine leukocidin-positive, methicillin-susceptible isolates of *Staphylococcus aureus* and implications for the evolution of community-associated methicillin-resistant *S.aureus*. *Clin Microbiol Infect.* 13(12):1157-1164.
- Mongkolrattanothai, K. Boyle, S. and Kahana, M.D. (2003):** Severe *Staphylococcus aureus* infections caused by clonally related community-associated methicillin-susceptible and methicillin-resistant isolates. *Clin. Infect. Dis.* 37 (8): 1050–1058.
- Monroe, E.W. Cohen, S.H. and Schulz, C.I.(1999):** Combined H1 and H2 antihistamine therapy in chronic urticaria. *Arch. Dermatol.*; 117:404-407.
- Monroe, E.W. (2005):** Treatment of urticaria. *Dermatol. Clin.*; 4: 51-55.
- Monroe, E.W. (2007):** Urticarial vasculitis : *Seminars of dermatology.*; 6 (4): 388.

N

- Nelson, J.M. Chiller, T.M. and Powers, J.H. (2007):** "Fluoroquinolone-resistant *Campylobacter* species and the withdrawal of fluoroquinolones from use in poultry: a public health success story". *Clin. Infect. Dis.* 44 (7): 977–980.
- Nikanne, H. Varionen, E. and Kouunen, T.U. (2000):** Is *staphylococcus aureus* infection associated with chronic urticaria. *Acta Dermatol.Venereol.*;80(6):425-426.
- Noble, W .C . Virani, Z . and Cree, R .G. (1992):** Co-transfer of vancomycin and other resistance genes from *Enterococcus faecalis* NCTC 12201 to *Staphylococcus aureus*. *FEMS Microbiol Lett.* 72:195-198.

Nouwen, J.L. Fieren, M.W. and Snijders, S. (2005): Persistent(not intermittent) nasal carriage of *Staphylococcus aureus* is the determinant of CPD- related infections. *Kidney Int* ;67:1084-1092.

Novick, R. Lindsay, J. and Ruzin, A. (1998): Molecular genetics of the test element, a mobile pathogenicity island in *Staphylococcus aureus*., In J. Arbuthnott, and B. Furman (ed.), *European Conference on Toxic Shock Syndrome. International Congress and Symposium Series 229*. Royal Society of Medicine Press Ltd, New York.109-112.

O

Odom, R.B. James, W.D. and Berger, T.G. (2006): Erythma & Urticaria In *Andrew's Diseases of the Skin Clinical Dermatology*. Ninth Edition. Saunders Company. Volume 1. 161 - 171.

O'Donnel, B.F. Barlow, R.J. and Greaves, M.W. (2001): Response of severe chronic urticaria to intravenous immunoglobulin (IVIG).*Br.j.Dermatol.*;131:23-24.

Orwin, P.M. Fitzgerald, J. R. and Leung, D.Y (2003): Characterization of *Staphylococcus aureus* enterotoxin L. *Infect Immun*. 71:2916–2919.

Oteifa, N. M. Moustafa, M. A. and Elgozamy, B. M. (1998): Toxocariasis as a possible cause of allergic diseases in children. *J. Egypt Soc. Parasitol.*; 28: 365-372.

P

Parfitt, T. (2005): "Georgia: an unlikely stronghold for bacteriophage therapy". *Lancet* 365 (9478): 2166–2167.

Parker, M.T. Tomlinson, A.J.H. and Williams, R.E.O. (2005): *J. Hyg. Camb.* 53,458.

- Parrish, J.A. Jaenicke, K.F. and Shea , C. (2006):** Solar urticaria treatment with PUVA and mediator inhibitors. *Br. J. Dermatol.*; 106: 575-580.
- Parsonnet, J. (1998):** Case definition of staphylococcal TSS: A proposed revision incorporating laboratory findings. In Arbuthnott, J. and Furman, B. (ed.), *European Conference on Toxic Shock Syndrome. International Congress and Symposium Series 229.* Royal Society of Medicine Press Ltd, New York, N.Y. 15.
- Pawlowicz, R. Panaszek, B.(2004):**Pathogenic and clinical significance of *Helicobacter pylori* infection in chronic urticaria and angioedema. *Pol Merkuriusz lek*,17:399-402.
- Peters, M.S. and Winkelman, R.K. (2005) :** Neutrophilic urticaria. *Br. J. Dermatol.*; 113:25-30.
- Peter, N. and Torvald, R.(2006):** *Staphylococcus aureus* throat colonization is more frequent than colonization in the anterior nares; *J Clin Microbiol.* 44(9): 334-339.
- Pirisi, A. (2000):** "Phage therapy—advantages over antibiotics?". *Lancet* 356 (9239): 1418.
- Prasad, G. S. Radhakrishnan, R. and Mitchell, D. T.:** Refined structures of three crystal forms of toxic shock syndrome toxin-1 and of a tetramutant with reduced activity. *Protein Sci.* 6:1220-1227.
- Projan, S. J. Kornblum, J. and Kreiswirth, B. (1989):** Nucleotide sequence: the β –hemolysin gene of *Staphylococcus aureus*. *Nucleic Acids Res.* 17:3305.
- Projan, S. J. and Novick, R. P. (1997):** The molecular basis of pathogenicity, In K. B. Crossley and G. L. Archer (ed.), *The staphylococci in human disease.* Churchill Livingstone, New York. 55-81.

Q

Qehiling, A. Neffen, H. and Siibim, M. (2006): Humoral and cellular immunity in chronic urticaria, *Allergol Immunopathol*; 7:351-356.

R

Rogers, H. J. Perkins, H. R. and Ward, J. B. (1980): Microbial cell walls and membranes. Chapman and Hall, ltd., London, United Kingdom. 437-460.

Root, R.R. and Metcalf, J.A. (2008): H₂ O₂ release from neutrophils during phagocytosis in relation to superoxide ion,formation and cellular metabolism of H₂O₂. *J. Clin. Invest.* 60: 1266.

Rosenbach, A.E. (2004): Cited by Cohen,J.O.:The staphylococci,Wiley Interscience New York,1-20.

S

Sabroe, R.A. and Greaves, M.W. (1997): The pathogenesis of chronic idiopathic urticaria. *Arch. Dermatol.*; 133: 1003-1008.

Sabroe, R.A. Seed, P.T. and Francis, D.M. (2006): Chronic idiopathic urticaria, *J, Invest. Dermatol.*; 106: 1001-1006.

Sabroe, R.A. Seed, P.T. and State, C. (2008): Chronic idiopathic urticaria: Comparison of the clinical features of patient with and without anti-Fc. Epsilon R.I. or anti IgE autoantibodies. *J. Am. ^x Acad. Dermatol.*; 40; 443-450.

Sakurane, M. Shotani, A. and Fiirukawa, F. (2002): Therapeutic effect of antibacterial treatment for intractable skin diseases in helicobacter pylori positive Japanese patients. *Dermatol.*; 29(1): 23-27.

Sanad, M.M. and Sanad, E.M. (1997): Toxocara canis as a cause of chronic urticaria. *The New Egyp. J. Med.*; 16 (2): 101-107.

- Sanchez, J.L. and Benmann, O. (2006):** Clinicopathological correlation in chronic urticaria. *Am. J. Dermatopathol.*; 143: 220-223.
- Saxena, A.K, Panhotra, B.R, and Chopra, R.(2004):** Advancing age and the risk of nasal carriage of staphylococcus aureus among patients on long-based term hospital –based hemodialysis. *Ann Saudi Med*;24:337-342.
- Schito, G.C. (2006):** "The importance of the development of antibiotic resistance in Staphylococcus aureus". *Clin Microbiol Infect* 12 Suppl 1: 3–8.
- Schleifer, K.H. and Kloos, W.E. (1975):** A Simple test for separation of staphylococci from micrococci. *J. Clin. Microbiol.* 1:337-338.
- Schlievert, P. M. (1982) :** Staphylococcal Pyrogenic Exotoxin C: Further Characterization. *Ann. Intern. Med Infection and Immunity* 36: 123.
- Schlievert, P. M. (1983) :** Identification and characterization of an exotoxin from Staphylococcus aureus associated with toxic shock syndrome. *I Infect Dis.*;143:509–516.
- Schwalbe, R.S. Stapleton, J.T. and Gilligan, P.H.(1987):** Emergence of vancomycin resistance in coagulase-negative staphylococci. *N Engl. J. Med.* 316: 927-931.
- Schuenken, R.P. Dadalti, P.E. and Silva, M.G. (2008):** Oxacillin and mupirocin –resistant staphylococcus aureus :in vitro activity of silver sulphadiazine and cerium nitrate in hospital strain. *J Chemother*; 16:453-458.
- Seifert, H. Schnabel, P. and von Eiff, C.(2003):** Small colony variants of Staphylococcus aureus and pacemaker-related infection. *Emerg Infect Dis.* 9:1316–1318.
- Sensi, P. Margalith, P. and Timbal, M.T. (1959):** "Rifomycin, a new antibiotic-preliminary report". *Farmaco Ed Sci* 14: 146–147.
- Sewell, C.M. Clarridage, J.E. and Guthrie, R.K.(1982):** Clinical significance of coagulase-negative staphylococci. *J. Clin. Microbiol.* 236-239.

- Sheldon, J.M.L. Matthews, K.P. and Lovell, R.G. (2008):** The vexing urticaria problem present concept of etiology and mangement. J Allergy; 35: 525-560.
- Shelley, W.B. and Rawnsley, H.M. (1964):** Aquagenic urticaria; contact sensitivity to water J. Am. Acad Dermatol; Quoted from Textbook of Dermatology (2001); 189: 895-908.
- Shelley, W.B. Shelley, E.D. and Talanin, N.Y. (2005):** Estrogen dermatitis. J. Am. Acad. Dermatol.; 32: 25-31.
- Shelley, W.B. and Shelley, E. (2007):** Acyclovir therapy for angioedema and chronic urticaria. Cutis.; 59: 185-188.
- Shertzer, E.N. and Lookingbill, D.P. (2006):** Effects of relaxation therapy and hypnotizability in chronic urticaria. Sheldon J.M.165.
- Shiomori, T. Yoshida, S . and Miyamoto, H.(2005):** Relationship of nasal and throat carriage of staphylococcus aureus to pathogenesis of perennial allergic rhinitis .J Allergy Clin Immunol ;105 :499-454.
- Sibbald, R.G. Chemma, A.S. and Lozinski, A. (2007):** Chronic urticaria: evaluation of the role of physical, immunologic and other contributory factors. Int. J. Dermatol.; 30:381-386.
- Sicherer, S.H. and Leung, D.Y. (2009):** Advances in allergic skin disease, anaphylaxis, and hypersensitivity reactions to foods, drugs, and insects. J Allergy Clin Immunol. Jan 2010; 125(1):85-97.
- Simons, F. E. and Simons, K. J. (2003) :** The pharmacology and use of H1 receptor antagonist drugs. N . Eng. J. Med.; 330: 381-386.
- Small, P. Barrett, D. and Champlin, E. (2008):** Chronic urticaria and angioedema. Clin. Allergy; 12:131-136.
- Soriano, A. Martinez, J.A. and Mensa, J. (2000):** Pathogenic significance of methicillin resistance for patients with Staphylococcus aureus bacteremia. Clin Infect Dis. 30:368-373.
- Soter, N.A. (2007):** Physical urticaria and angio-oedema. Seminars in Dermatology; 6(4): 302.

- Soter, N.A. (2009):** Urticaria : current therapy. J. Allergy Clin. Immunol.; 80: 1009-1014.
- Soter, N.A. and Wasserman, S.J. (2007):** Cutaneous changes in disorders of altered reactivity; IgE-Dependent urticaria and angio-oedema. In Dermatology in General Medicine. Thomas B.; Fitzpatrick; Arthur Z. Eisen, Klaus Wolff, Twin M. Freedbergs K. Frank. Austen (eds), 3rd edition. Me. Graw Hill. Information. Service Company. 1282-1293.
- Stephens, C.J.M. and Black, M.M. (2008):** Premenstrual eruption autoimmune progesterone dermatitis. Semin Dermatol.; 8: 26-29.
- Stokes, E.J. and Ridgway, G.L. (2003):** Clinical Bacteriology, Fifth Edition. Staphylococci, Edward Arnold. 110-114.
- Sundin, D.P. Sandoval, R. and Molitoris, B.A. (2001):** Gentamicin Inhibits Renal Protein and Phospholipid Metabolism in Rats: Implications Involving Intracellular Trafficking. J. Am. Soc. Nephrol 12:114-123.
- Sveum, R.J. (2006):** Urticaria. The diagnostic challenge of the hives. Post graduate Medicine; 100 (7): 81-82.
- Sykes, R.B. (2008):** The classification and terminology of enzymes that hydrolysis. B-Lactam antibiotic . J . inf. Dis. Vol, 145. No. S. May. 762.

T

- Tebbe, B. Christoph, C . and Geilen, J.(1996):** Staphylococcus aureus infection and chronic urticaria .J.Am.Acad. Dermatol.;39.(4) : 685-686.
- Tharp, M.D. (1996):** Cetirizine: A new therapeutic alternative for chronic urticaria. Cutis.; 58:94-98.
- Thomas, D.H. Hice, D.W. and Fitton, J.E. (2007):** J. Mol. Biol. Dec, 5, 192 (3): 625 -626.

Tong, L.J. Balakrishnan, G. and Kochan, J.P. (1997): Assessment of autoimmunity in patients with chronic urticaria. J. Allergy Clin. Immunol.;99:461-465.

Touissaints, S. and Kamino, H. (2006): Non infectious erythematous papular and squamous diseases. In: Lever's histopathology of the skin. Eight edition. David Elder (ed). Lippincott Company. Philadelphia. 151-179.

V

Valeva, A. Palmer, M. and Bhakdi, S. (1997): Staphylococcal exotoxin:formation of the heptameric pore is cooperative and proceeds through multiple intermediate stages. Biochemistry 36:13298-13304.

Valsecchi, R. and Pigatto, P.(2008): Chronic urticaria and staphylococcus aureus .Acta.Dermatol.Venereol.;78:440-442.

Varga, L. Losceiz, D.Z..and Mercetey, K.(2005) :Staphylococcus aureus infection and chronic urticaria. Orv.Hetil.; 45:812-815 .

Venzor, J. Lee, W.L. and Huston, D.P. (2002): Urticarial vasculitis. Clin Rev, Allergy Immunol. ; 23(2):201-216.

Verwey, (2000): Cited by Free man, B.A. (1999). Textbook of microbiology. Staphylococci, twenty-first(ed.) W.B. Chap 15;431.

Viada, G.A. Goldman, M.A. and Block, K.J. (2003): Testing for hepatitis B virus in patients with chronic urticaria and angioedema. J. Allergy Clin. Immunol.; 72: 193-198.

Vigan M, Sauvage C, Adessi, B.(1994): Pourquoi et comment réaliser une batterie standard chez les enfants? Nouv Dermatol; **13**: 12–5.

Von, C. Becker, K. and Machka, K. (2001) :Nasal carriage as a source of staphylococcus aureus bacteremia. N Engl J Med; 344: 11-16.

W

- Wadlvogel, F. A. (2000):** Staphylococcus aureus (including staphylococcal toxic shock). In: Mandell GL, Bennett JE, Dolin R, (eds). Principles and practice of infectious diseases. 5th ed. Philadelphia: Churchill Livingstone. 2072-2073 .
- Wahlgren, C.F. Tengval, L. and Under, H. (2000):** Itch and inflammation induced by intradermally injected interleukin-2 in atopic dermatitis patients and healthy subject. Arch. Dermatol.; Res.; 287: 272-280.
- Warin, R.P. and Smith, R.J. (2006):** Challenge test battery in chronic urticaria. Br. J. Dermatol.; 44: 4016-4017.
- Wedi, B. and Kappa, A.(2002) :** Staphylococcus infection and skin disease .J.Physiol.Pharmacol.; 50(5) : 753-776.
- Wedi, B. and Kappa, A.(2004) :** Staphylococcus infection in skin disease: a critical appraisal .Am.J.Clin.Dermatol.; 3(4) :273-282.
- Wilkinson, P.C. (1983):** Non specific cells: Granulocytes, Mono-unclear Phagocytes and Mast cells. In Immunology in Medicine: A comprehensive guide to clinical immunology. Edited by: E.J. Holborow and W.G. Reeves. Academic press, London, Grune and Stratton. Hew York, 231.
- Witlock, R.P. (1981):** Factitious urticaria and dermatographism. Br. J. Dermatol.; 104:285-288.
- Wolfgang, K. Joklik ,W. and Hilda, P. (2006):** (Ed) Zinsser Micribiology. & 16th edition. Appletion - Century Crofts. 412-430.
- Wolform, E. Chene, G. and Boissmu, H. (2005) :** Chronic urticaria and toxocara canis. Lancet.; 345: 196-200.
- Wood, S.M. Mann, R.D. and RawLins, M.M.(2007):** Angio-oedema and urticaria associated with angiotensin-converting enzyme inhibitors .Br.Med.j.294: 91-93.
- Woodin, A. M.(1960) :** Purification of the two component of leukocidin from staphylococcus aureus .Biochem. J. 75:158-165.

Woodin, A. M.(1970): In Montie, T. C.Kadis, S. Ajl, S. J.(eds) Microbial toxins, vol. 3.Academic Press, New York.27.

Wransa, K. Wahlberg, J.E. and Axelson, I.G.K. (1998): IgE mediated allergy to natural rubber in 30 patient with contact urticaria.Dermatol.; 19:264-271.

Wright, H. T. and Reynolds, K.A. (2007): Antibacterial targets in fatty acid biosynthesis. Current Opinion in Microbiology .10,447-453.

Z

Zierdt, C .H. (1982) : Blood-lysing solution nontoxic to pathogenic bacteria. Journal of Clinical Microbiology 16: 517.

Zuberbier, T. Ifflander, J. and Henz, B.M. (2006): Acute urticaria. Clinical aspects and therapeutic responsiveness. Acta. Dermatol. Venereol.; 76:295-297.

Zuberbier,T. and Maurer, M (2007): Urticaria: current opinions about etiology, diagnosis and therapy. Acta. Dermatol. Venereol. 87(3):196-205.