

References

(A)

- 1) **Adebamowo, C.A.; Spiegelman, D.; Berkey, C.S.; Danby, F.W.; Rockett, H.H. and Colditz, G.A. (2008):**

Milk consumption and acne in teenaged boys.

J. Am. Acad. Dermatol.; 58: 787-93.

- 2) **Adebamawo , C.A.; Spiegelman, D.; Danby, F.W.; Frazier, A.L. ;Willett, W.C. and Holmes, M.D. (2005) :**

High school dietary dairy intake and teenage acne.

J. Am. Acad. Dermatol.; 52: 207-14.

- 3) **Ahima, R.S. and Oseis, Y. (2008):**

Adipokines in obesity.

Front. Horm. Res.;36: 182–97.

- 4) **Appelhans, B.M. (2010):**

Circulating leptin moderates the effect of stress on snack intake independent of body mass.

EAT BEH (Eating Behaviors)-00443; No of Pages 4.

- 5) **Arbesman, H. (2005):**

Dairy and acne—the iodine connection.

J. Am. Acad. Dermatol.; 53: 1102-05.

- 6) **Ashby, R.D., Zerkowski, J.A., Solaiman, D.K. and Liu, L.S. (2011):**

Biopolymer scaffolds for use in delivering antimicrobial sophorolipids to the acne-causing bacterium *Propionibacterium* acnes. New Biotechnology Volume 28, Number 1.

J. Am. Acad. Dermatol. 54, 420–426.

(B)

7) *Badman, M.K. and Flier, J.S. (2007):*

The adipocyte as an active participant in energy balance and metabolism.

Gastro-enterology; 132:2103–15.

8) *Batya, B.; Davidovici, M.D. and Ronni Wolf (2010):*

The role of diet in acne: facts and controversies.

Clinics in Dermatology; 28, 12–16.

9) *Bernier, V.; Weill, F.X .and Hirigoyen, V. (2002):*

Skin colonization by *Malassezia* species in neonates.

Arch. Dermatol.; 138: 215–18.

10) *Bernotiene, E.; Palmer, G. and Gabay, C. (2006):*

The role of leptin in innate and adaptive immune responses.

Arthritis Res. Ther.; 8: 217-18.

11) *Berra, B. and Rizzo, A.M. (2009):*

Glycemic index, glycemic load, wellness and beauty:

the state of the art

Prog. Nutr.; 27, 230–235.

12) *Berra, B.; Montorfano, G.; Berselli, P. and Rizzo, A.M. (2007):*

Diet, exercise, long chain polyunsaturated omega-3 fatty acids and the metabolic syndrome.

Prog. Nutr.; 9: 124-33.

13) *Beylot, C. (1993):*

Seborrhoea and its complications.

Rev. Prat; 43: 2320–7

14) *Bigby, M. (2001):*

Challenges to the hierarchy of evidence: does the emperor have no clothes?

Arch. Dermatol.; 137: 345-6.

- 15) Bjorbaek, C.; Elmquist, J.K.; Frantz, J.D.; Shoelson, S.E. and Flier, J.S. (1998):**

Identification of potential mediator of central leptin resistance.
Mol. Cell.; 1: 619–25.

- 16) Brabant, G.; Muller, G.; Horn, R.; Anderwald, C.; Roden, M. and Nave, H. (2005):**

Hepatic leptin signaling in obesity.
J. Faseb.; 19: 1048–50.

- 17) Brabant, G.; Nave, H.; Horn, R.; Anderwald, C.; Muller, G. and Roden, M. (2004):**

In vivo and in vitro evidence for a hepatic modulation of the leptin.
J. Eur. Clin. Invest. ; 34:831–7.

- 18) Brabant, G.; Nave, H.; Mayr, B.; Behrend, M.; Van Harmelen, V. and Arner, P. (2002):**

Secretion of free and protein-bound leptin from subcutaneous adipose tissue of lean and obese women.
J. Clin. Endocrinol Metab.; 87: 3966–70.

- 19) Berra, B and Rizzo, A.M. (2009):**

Glycemic index, glycemic load, wellness and beauty: the state of the art.
Clinics in Dermatology 27, 230-235.

- 20) Brynes, A.; Edwards, C. and Jadhav, A. (2000):**

Diet-induced change in fatty acid composition of plasma triacylglycerols is not associated with change in glucagon-like peptide 1 or insulin sensitivity in people with type 2 diabetes.
Am. J. Clin. Nutr.; 72: 1111-8.

21) Burkhart, C.G. and Burkhart, C.N. (2007):

Expanding the microcomedone theory and acne therapeutics: Propionibacterium acnes biofilm produces biological glue that holds corneocytes together to form plug.

J. Am. Acad. Dermatol.; 57: 722-4.

(C)

22) Capitanio, B.; Sinagra, J.L.; Ottaviani, M.; Bordinon, V.; Amantea, A. and Picardo, M. (2007):

‘Smoker’s acne’: a new clinical entity?

Br. J. Dermatol.; 157: 1070-1.

23) Cerman, A.A.; Bozkurt, S.; Sav, A.; Tulunay, A.; Elbas, M.O. and Ergun, T. (2008):

Serum leptin levels, skin leptin and leptin receptor expression in psoriasis.

B. J. Dermatology 11 May, 1365-2133.

24) Chehab, F.F. (2002):

Leptin as a regulator of adipose mass and reproduction.

Trends Pharmacol.Sci. ; 21: 309-314.

25) Chen, W.; Thiboutot, D. and Zouboulis, C.C.(2002):

Cutaneous androgen metabolism: basic research and clinical perspectives.

J. Invest. Dermatol.; 119: 992-1007.

26) Chiu, A.C.; Chon, S.Y. and Kimball, A.B.(2003):

The response of skin disease to stress: changes in the severity of acne vulgaris as affected by examination stress.

Arch. Dermatol.; 139: 897-900.

27) Coates, P.; Vyakrnam, S. and Eady, E.A. (2002):

Prevalence of antibiotic resistant proprionibacteria on the skin of acne patients: 10-year surveillance data and snapshot distribution study.

Br. J .Dermatol.; 146: 840-48.

28) Cordain, L.; Lindeberg, S.; Hurtado, M.; Hill, K.; Eaton, S.B. and Brand- Miller, J. (2002):

Acne vulgaris: a disease of Western civilization.

Arch .Dermatol;. 138: 1584-90.

29) Cunliffe, W.J.;Baron, S.E. and Coulson, I.H. (2009):

A clinical and therapeutic study of 29 patients with infantile acne.

Br. J. Dermatol.; 145: 463–66.

30) Cunliffe, W.J.; Caputo, R. and Dreno, B. (2010):

Efficacy and safety comparison of adapalene (CD271) gel and tretinoin gel in the topical treatment of acne vulgaris. A European multicentre trial.

J. Dermatolog Treat.; 8: 173–8.

31) Cunliffe, W.J. ;Holland, D.B. and Clark, S.M. (2000):

Comedogenesis: some new aetiological, clinical and therapeutic strategies.

Br. J. Dermatol.; 142: 1084-91.

(D)

32) Dahl, L.; Opsahl, J.A.; Meltzer, H.M. and Julshamn, K.(2003):

Iodine concentration in Norwegian milk and dairy products.

Br. J .Nutr.; 90: 679-85.

33) Danby, F.W.(2008):

Diet and acne.

Clin. Dermatol.; 26:93-6.

34) Deplewski, D. and Rosenfield, R.L.(2000):

Role of hormones in pilosebaceous unit development.

Endocr Rev.; 21: 363-92.

35) Dessinioti, C. and Katsambas, D.A. (2010):

The role of *Propionibacterium acnes* in acne pathogenesis: facts and controversies.

Clinics in Dermatology; 28,2–7

36) Dixit, V.D. (2008):

Hormonal, neural, and metabolic regulation of immunity.

J. Leukoc. Biol.; 84:881.

37) Dobbins, R.; Szczepaniak, L. and Myhill, J. (2002):

The composition of dietary fat directly influences glucose-stimulated insulin secretion in rats.

Diabetes; 51: 1825-33.

38) Doyon, C. ; Drouin, G.; Trudeau, V.L. and Moon, T.W. (2001):

Molecular evolution of leptin, general and comparative endocrinology; 124, 188–198.

(E)

39) Eady, E.A.; Farmery, M.R. and Ross, J.I.(2007):

Effects of benzoyl peroxide and erythromycin alone and in combination against antibiotic-sensitive and resistant skin bacteria from acne patients.

Br .J .Dermatol.; 131: 331-6.

- 40) Elinav, E.; Pappo, O.; Sklair-Levy, M.; Margalit, M.; Shibolet, O.; Gomori, M.; Alper, R.; Thalenfeld, B.; Engelhardt, D.; Rabbani, E. and Ilan, Y.(2006):**

Adoptive transfer of regulatory NK-T lymphocytes ameliorates non-alcoholic steato-hepatitis and glucose intolerance in ob/ob mice and is associated with hepatic CD8 trapping.

J. Pathol.; 209: 121–8.

(F)

- 41) Ferrante Jr, A.W. (2007):**

Obesity-induced inflammation: a metabolic dialogue in the language of inflammation.

J. Intern. Med.; 262:408–14.

- 42) Ford, E.S. and Liu, S.(2001):**

Glycemic index and serum high-density lipoprotein cholesterol concentration among adults.

Arch. Intern. Med.; 161: 572-6.

- 43) Foster-Powell, K.; Holt, S. and Brand-Miller, J.(2002):**

International table of glycemic index and glycemic load values.

Am. J. Clin. Nutr.; 76: 5-56.

(G)

- 44) Garner, S.E.; Eady, E.A. and Popescu, C. (2003):**

Minocycline for acne vulgaris: efficacy and safety.

Cochrane Database Syst Rev.; 1: 20-86.

- 45) Gollnick, H. (2003):**

Current concepts of the pathogenesis of acne: implication for drug treatment.

Drugs; 63: 1579-96.

46) Gollnick, H.;Finlay, A.Y. and Shear, N. (2008):

Global alliance to improve outcomes in acne. Can we describe acne as a chronic disease? If so, how and when?

Am. J. Clin .Dermatol.; 9: 279-84.

47) Gollnick, H. and Schramm, M. (2007):

Topical drug treatment in acne.

Dermatology; 196: 119–25.

48) Graham G.M., Farrar M.D., Cruse-Sawyer J.E., Holland K.T. and Ingham E.(2004):

Proinflammatory cytokine production by human keratinocytes stimulated with *Propionibacterium acnes* and *P. acnes*.

Br. J. Dermatol.; 150:421-8.

(H)

49) Hamminga E.A., Van der Lely A.J., Neumann H.A.and Thio H.B.(2006):

Chronic inflammation in psoriasis and obesity: implication for therapy.

Med. Hypotheses.; 67:768–73.

50) Hunter, N. and Shaker, O. (2009):

Diet and body mass index, do they have a role In acne vulgaris? a study using serum leptin and serum Insulin like growth factor-1 Egypt.

J. Derm. & Androl. Vol. 29, No (2)

51) Harper, J.C. and Thiboutot, D.M. (2003):

Pathogenesis of acne: research advances.

Adv. Dermatol.; 19: 1-10.

52) Herrom, M.D. ;Hinckley, M. and Hoffman, M.S. (2005):

Impact of obesity and smoking on psoriasis presentation and management.

Arch Dermatol.; 141:1527–34.

53) Heymann, W.R. (2006):

Toll-like receptors in acne vulgaris.

J.Am. Acad .Dermatol.; 55: 691-2.

54) Hongcharu W., Taylor C.R., Chang Y., Aghassi D., Suthamjariya K. and Anderson R.R.(2000):

Topical ALA-photodynamic therapy for the treatment of acne vulgaris.

J. Invest. Dermatol .2000; 115: 183-92.

55) Hotamisligil, G.S. (2006):

Inflammation and metabolic disorders.

Nature; 444: 860–7.

(J)

56) Jansen T, Romiti R and Woitalla S. (2000):

Eruptive acne vulgaris with infectious mononucleosis.

Br. J. Dermatol.; 142: 837–8.

57) Jarrousse, V.; Castex-Rizzi, N. and Khammari, A. (2007):

Modulation of integrins and filaggrin expression by Priopionibacterium acnes extracts on keratinocytes.

Arch. Dermatol. Res.; 299: 441-7.

58) Jenkins, D.J. (1981):

Glycemic index of foods: a physiological basis for carbohydrate exchange.

Am. J. Clin .Nutr.; 34: 362-6.

- 59) **Jeremy, A.H.; Holland, D.B. and Roberts, S.G. (2003):**
Inflammatory events are involved in acne lesion initiation.
J. Invest. Dermatol.; 121: 20-7.
- 60) **Jiménez , M.A.; Sánchez, E.M.; Haro-Gabaldón, V.; Solís, E. and Moreno, A. (2004):**
Acné grave pruriginoso en anciano con insuficiencia renal crónica.
Act. Dermosifiliogr.; 95: 652-57.
- 61) **Jugeau, S.; Tenaud, I. and Knol, A.C. (2005):**
Induction of Toll-like receptors by *Propionibacterium acnes*.
Br. J. Dermatol.; 153: 1105-13.
- 62) **Juzeniene, A.; Nielsen, K.P. and Zhao, L.(2008):**
Changes in human skin after topical PDT with hexyl aminolevulinate.
Photodiagn. Photodyn. Ther.; 5: 176-81.
- (K)**
- 63) **Kavazarakis, K. ; Maria M. ;Dimitrios G. ; Antonios D. ; Apostolos B. ; Petros M.; Androniki X. and Themistocles K.(2001):**
Relation of serum leptin levels to lipid profile in healthy children.
Br. J .Dermatol.; pp 1091-1094.
- 64) **Kaymak, Y.; Adisen, E.; Ilter, N.; Bideci, A.; Gurler, D. and Celik, B. (2007):**
A Dietary Glycemic Index and glucose, insulin-like growth factor-1, insulin-like growth factor binding protein 3 and leptin levels in patients with acne.
J. Am. Acad. Dermatol.; 57: 819-23.

- 65) **Kennedy A, Gettys TW, Watson P, Wallace P, Ganaway E, Pan Q & Garvey WT. (1997):**

The metabolic significance of leptin in humans: gender-based differences in relationship to adiposity, insulin sensitivity, and energy expenditure.

Journal of Clinical Endocrinology and Metabolism ;82 :1293–300.

- 66) **Kılıc, M.; Taskın, E.; Ustundag, B. and Aygun, A.D. (2004):**

The evaluation of serum leptin level and other hormonal parameters in children with severe malnutrition.

Clin. Biochem.; 37:382-7.

- 67) **Kim, J.; Ochoa, M.T; Krutzik, S.R;Takeuchi, O.; Uematsu, S. and Legaspi, A. (2002):**

Activation of toll-like receptor 2 in acne triggers inflammatory cytokine responses.

J. Immunol.; 169: 1535- 41.

(L)

- 68) **La Cava, A. and Matarese, G. (2004):**

The weight of leptin in immunity.

Nat. Rev. Immunol.; 4:371–9.

- 69) **Lanier, L.L. (2005):**

NK cell recognition.

Annu. Rev. Immunol.; 23: 225–74.

- 70) **Lee, D.H., ; Choi, Y.S.; Seong Uk Min, ;Mi Young Yoon, and Dae Hun Suh.(2009):**

Comparison of a 585-nm pulsed dye laser and a 1064-nm Nd:YAG laser for the treatment of acne scars: A randomized split-face clinical study.

J. Am. Acad. Dermatol.; 60: 801-11.

71) Li, J.; Li, F. and Zhao, A.(2006):

Inflammation and leptin.

Drug Discov. Today; 3:387–93.

72) Lindsay S. (2007):

The treatment of acne.

Annu. Rev. Immunol.; 17:10-15.

73) Liu, P.T.; Krutzik, S.R.; Kim, J. and Modlin, R.L.(2005):

Cutting edge: all trans-retinoic acid down-regulates TLR2 expression and function.

J. Immunol.; 174: 2467-70.

74) Liu, S. and Willett, W.C.(2002):

Dietary glycemic load and atherothrombotic risk. Curr Atheroscler Rep.; 4: 454-61.

75) Liu,Y.; Zhonga, Y.; Pei, J.; Zhu, Y.; Hu, Y.; Chi, S. and Alan, D.(2009):

Inhibitory effect of leptin on growth hormone secretion of GH3 cells: Involvement of cell proliferation, apoptosis and intracellular free Ca²⁺.

Cytokine; 46: 245–50.

76) Lo, J.C.; Wang, Y.; Tumanov, A.V.; Bamji, M.; Yao, Z.; Reardon, C.A.; Getz, G.S. and Fu, Y.X.(2007):

Lymphotoxin beta receptor- dependent control of lipid homeostasis.

Science; 316: 285–8.

77) Lovejoy, J. (2002):

The influence of dietary fat on insulin resistance. Curr Diabetes Rep.; 2: 435-40.

78) Lovejoy, J.; Champagne, C. and Smith, S. (2001):

Relationship dietary fat and serum cholesterol ester and phospholipid fatty acids to markers of insulin resistance in men and women with a range of glucose tolerance.

Metabolism; 50: 86-92.

79) Lovejoy, J.; Smith, S. and Champagne, C. (2002):

Effects of diets enriched in saturated (palmitic), monounsaturated (oleic), or trans (elaidic) fatty acids on insulin sensitivity and substrate oxidation in healthy adults.

Diabetes Care; 25: 1283-8.

80) Lu, H. and Li, C. (2000):

Leptin a multifactorial hormone .

Cell Res.; 10:81-87.

81) Lucky, A.W. (1998):

Hormonal correlates of acne and hirsutism.

Am. J. Med.; 98: 89S-94S.

82) Ludwig, D.S. (2003):

Dietary glycemic index and the regulation of body weight.

Lipids; 38: 117-21.

(M)

83) McLane, J. (2009):

Analysis of common side effects of isotretinoin.

J. Am. Acad. Dermatol.; 45: S188–94.

84) Matarese, G.; Moschos, S. and Mantzoros, C. (2005):

Leptin in immunology.

J. Immunol.; 173:3137–42.

85) Meyer, K.; Kushi, L. and Jacobe, D. (2000):

Carbohydrates, dietary fiber, and incidence of type 2 diabetes in older women.

Am. J. Clin. Nutr.; 71: 921-30.

86) Mills, O.H. and Kligman, A.M. (1978):

Ultraviolet phototherapy and photochemotherapy of acne vulgaris.

Arch Dermatol.; 114: 221-3.

87) Ming Zhao; Jia-Qing Shao and Hong Du. (2009):

Effects of three antidiabetics on insulin and leptin in Chinese type 2 diabetes Diabetes & Metabolic Syndrome

Clinical Research & Reviews; 80: 4-40.

88) Mitra, A. and Stables, G.I.(2006):

Topical photodynamic therapy for non-cancerous skin conditions.

Photodiagn. Photodyn. Ther.; 3: 116-27.

89) Mizuta, E.; Kokubo, Y.; Yamanaka, I.; Miyamoto, Y.; Okayama, A. and Yoshimasa, Y. (2008):

Leptin gene and leptin receptor gene polymorphisms are associated with sweet preference and obesity.

Hypertension Research, 31: 1069-77.

90) Mori, A.; Sakurai, H.; Choo, M.K.; Obi, R.; Koizumi, K.; Yoshida, C.; Shimada, Y. and Saiki, I.(2006):

Severe pulmonary metastasis in obese and diabetic mice.

Int. J. Cancer; 119:2760-7.

91) Morton, C.A.; McKenna, K.E. and Rhodes, L.E. (2008):

Guidelines for topical photodynamic therapy.

Br. J. Dermatol.; 159:1245-66.

92) Muurling, M.; Jong, M. and Mensink, R.(2002):

A low-fat diet has a higher potential than energy restriction to improve high-fat diet-induced insulin resistance in mice. *Metabolism*; 51: 695-701.

(N)

93) Nagy, I.; Pivarcsi, A.; Koreck, A.; Sz  ll, M.; Urb  n, E. and Kem  ny, L. (2005):

Distinct strains of *Propionibacterium acnes* induce selective human beta-defensin-2 and interleukin-8 expression in human keratinocytes through toll-like receptors.

J. Invest. Dermatol.; 124: 931-8.

94) Nakatsuji, T.; Liu, Y.T. and Huang, C.P.(2008):

Vaccination targeting a surface sialidase of *P. acnes*: implications for new treatment of acne vulgaris.

J. Invest. Dermatol.; 3: 1551.

95) Nave, H.; Mueller, G.; Siegmund, B.; Jacobs, R.; Stroh, T.; Schueler, U.; Hopfe, M.; Behrendt, P.; Buchenauer, T.; Pabst, R. and Brabant, G.(2008):

Resistance of Janus kinase-2 dependent leptin signaling in natural killer (NK) cells: a novel mechanism of NK cell dysfunction in diet-induced obesity.

Endocrinology; 149:3370–8.

96) Nouri, K. and Ballard, C. (2006):

Laser therapy for acne.

Clin. Dermatol.; 24: 26-32.

(P)

97) Pappas, A.; Anthonavage, M. and Gordon, J.(2002):

Metabolic fate and selective utilization of major fatty acids in human sebaceous gland.

J. Inves. Dermatol.;118:164-71.

98) Perry, A.L. and Lambert, P.A.(2006):

Propioni-bacterium acnes.

J Invest. Dermatol.; 42: 185–8.

99) Pi-Sunyer, F. (2002):

Glycemic index and disease.

Am. J. Clin. Nutr.; 76 (suppl): 290S-8S.

100) Pivarcsi, A.; Bodai, L. and Rethi, B. (2003):

Expression and function of Toll-like receptors 2 and 4 in human keratinocytes.

Int. Immunology; 15: 721-30.

101) Pollock, B.; Turner, D. and Stringer, M.R. (2004):

Topical amino levulinic acid-photodynamic therapy for the treatment of acne vulgaris: a study of clinical efficacy and mechanism of action.

Br. J. Dermatol.; 151: 616-22.

(R)

102) Ramsaya, T.G.; Bushb, .JA.; McMurtrya, J.P.; Thiviergeb, M.C. and Davisb, T.A. (2004):

Peripheral leptin administration alters hormone and metabolite levels in the young pig.

Comp. Biochem. Physiol.;138:17–25.

- 103) Rapini P .R., Callen J.P., Horn T.D., Mancini A.G., Salasche S.J., Schwarz J.V., Stingl G. and Stone M.S. (2008):**

Acne vulgaris. In: Bologna J. L. and Jorizzo L. J. editors.

Bologna Dermatology (2nd edn), Spain.

Dermatology, New York: Mosby p; 531-543.

- 104) Rivera, R. and Guerra, R. (2009):**

Management of acne in women over 25 years of age.

Am. J. Clin. Nutr.; 100: 33-7.

- 105) Rigopoulos, D.; Gregoriou, S. and Ifandi, A. (2007):**

Coping with acne: beliefs and perceptions in a sample of secondary school Greek pupils.

J. Eur. Acad. Dermatol. Venereol.; 21: 806-10.

- 106) Rosen, T. (2007):**

The Propionibacterium acne genome: from the laboratory to the clinic.

J. Drugs Dermatol.; 6:582–86.

- 107) Robyn, N. ;Smith, M.;Anna Braue; George, A.; Varigos and Neil, J.(2008):**

The effect of a low glycemic load diet on acne vulgaris and the fatty acid composition of skin surface triglycerides.

J. Dermatological Science; 50: 41-52.

- 108) Rosen, E.D. and Spiegelman, B.M.(2006):**

Adipocytes as regulators of energy balance and glucose homeostasis.

Nature; 444: 847–53.

- 109) Rosenfield, R. (1999):**

Ovarian and adrenal function in polycystic ovary syndrome.

Endocrinol Metab Clin North Am; 28: 265-93.

110) Rosenfield, R. (2001):

Polycystic ovary syndrome and insulin resistant hyperinsulinemia.
J. Am. Acad. Dermatol.; 45: S95-S104.

111) Roth, J.; Mobarhan, S. and Clohisy, M.(2002):

The metabolic syndrome: where are we and where do we go? Nutr
Rev.; 60: 335-7.

(S)

**112) Saleri , R.;Giustina , A.; Tamanini, C.; Valle , D.;Burattin, A. and
Weluenberg, W.B.(2004):**

Leptin stimulates growth hormone secretion via a direct pituitary
effect combined with a decreased somatostatin tone in a median
eminence-pituitary physiology study.

Neuroendocrinology; 79(4): 221-8.

**113) Sano, S.; Chan, K.S.; Carbajal , S.; Clifford, J.; Peavey, M. and
Kiguchi, Kl.(2005):**

Stats links activated keratinocytes and immunocytes required for
development of psoriasis in a novel transgenic mouse model.

Nat. Med.; 11: 43–9.

**114) Savastano, S.; Di Somma, C.; Belfiore, A.; Guida, B.; Orio, F.J. and
Rota, F.(2006):**

Growth hormone status in morbidly obese subjects and correlation
with body composition.

J. Endocrinol .Invest.; 29(6): 536–43.

115) Saygılı, F.; Oge, A. and Yılmaz , C.(2005):

Hyperinsulinemia and insulin insensitivity in women with
nonclassical congenital adrenal hyperplasia due to 21-hydroxylase
deficiency: the relationship between serum leptin levels and
chronic hyperinsulinemia.

Horm. Res.; 63: 270-4.

116) Schafer, T., Nienhaus, A. and Vieluf, D. (2001):

Epidemiology of acne in the general population: the risk of smoking.

Br. J. Dermatol.; 45: 100–4.

117) Shalita, A. (2009):

The integral role of topical and oral retinoids in the early treatment of acne.

J. Eur. Acad .Dermatol. Venereol.; 15: 43–9.

118) Shaw, J.C. and White, L.E. (2001):

Persistent acne in adult women.

Arch. Dermato.; 137: 1252-3.

119) Sinha, M.K.; Opentanova, I.; Ohannesian, J.P.; Kolaczynski, J.W.;Heiman, M.L.;Hale , J.; Becker, G.W.; Bowsher, R.R.;Stephens, T.W. and Caro, J.F.(1996):

Evidence of free and bound leptin in human circulation. Studies in lean and obese subjects and during short-term fasting.

J. Clin. Invest.;98:1277–82.

120) Smith, K.R. and Thiboutot, D.M. (2008):

Sebaceous gland lipids: friend or foe?

J. Lipid Res.; 49: 271-81.

121) Smith, R.N.; Mann, N.J.; Braue, A.; Mächeläinen, H. and Varigos, J.A. (2007):

A low glycemic-load diet improves symptoms in acne vulgaris patients: a randomized controlled trial.

Am. J. Clin. Nutr.; 86: 107-15.

- 122) Smith, R.N., Mann, N.J., Braue, A., Makelainen, H. and Varigos, G.A.(2007):**

The effect of a high-protein, low glycemic-load diet versus a conventional, high glycemic-load diet on biochemical parameters associated with acne vulgaris: a randomized, investigator-masked, controlled trial.

J. Am. Acad. Dermatol.; 57: 247-56.

- 123) Smithard, A.; Glazerbrook, C. and Williams, H. (2001):**

Acne prevalence, knowledge about acne and psychological morbidity in mid-adolescence: a community-based study.

Br. J. Dermatol.; 145: 274-9.

- 124) So"derquist, B.; Holmberg, H. and Unemo, M. (2009):**

Propionibacterium acnes as an etiological agent of arthroplastic and osteosynthetic infections – Two cases with specific clinical presentation including formation of draining fistulae

Br. J. Dermatol.; 85: 107-13.

- 125) Stephen, J.; Schueler, M.D.; John, H. ;Beckett, M.D. and Scott Gettings, D . (2007):**

Acne therapy and predisposing factors.

Br. J. Dermatol.; 14: 143-52

- 126) Steven, J. ;Greco; Sraboni Sarkar; Jane, M.; Johnston; Xiongwei Zhu; Bo Su; Gemma Casadesus; J. Wesson Ashford and Mark, A. (2008):**

Leptin reduces Alzheimer's disease-related to phosphorylation in neuronal cells.

Biochemical and Biophysical Research Communications; 376 :536-541.

127) Sutton Jr, R.L. (1956):

Diseases of the Skin. 11th ed.
St. Louis (Mo): Mosby Co;p. 683-90.

128) Szeimies, R.M.; Matheson, R.T. and Davis, S.A.(2009):

Topical methyl aminolevulinate photodynamic therapy using red light-emitting diode light for multiple actinic keratoses: a randomized study.
Dermatol. Surg.; 35: 586-92.

(T)

129) Takeda, K.; Kaisho, T. and Akira, S. (2003):

Toll-like receptors.
Annu. Rev. Immunol.;21:335-76.

130) Tan, J.; Vasey, K. and Fung, K. (2001):

Beliefs and perceptions of patients with acne.
J. Am. Acad. Dermatol.; 44: 439-45.

131) Tartaglia, L.A.(1997):

The leptin receptor.
J. Biol. Chem.;272: 6093–6.

132) Taylor, M.N. and Gonzalez, M.L. (2009):

The practicality of photodynamic therapy in acne vulgaris.
Br .J. Dermatol.; 160: 1140-8.

133) Thiboutot , D.M. and Strauss, J.S.(2002):

Diet and acne revisited [comment].
Arch. Dermatol.; 138: 1591-2.

134) Thielitz, A.;Sidou, F. and Gollnick, H.(2007):

Control of microcomedone formation throughout a maintenance treatment with adapalene gel, 0.1%.
J. Eur. Acad. Dermatol. Venereo.; 21: 747-53.

135) Tian, Z.;Sun, R.;Wei, H. and Gao, B. (2002):

Impaired natural killer (NK) cell activity in leptin receptor deficient mice: leptin as a critical regulator in NK cell development and activation.

Biochem. Biophys. Res. Commun.; 298: 297–302.

(V)

136) Van de Kerkhof, P.C.; Kleinpenning, M.M.; de Jong, E.M.; Gerritsen, M.J.;van Dooren-Greebe, R.J. and Alkemade, H.A. (2006):

Current and future treatment options for acne.

J. Dermatolog. Treat.; 17: 198-204.

137) Vans, D.M.; Kirk, K.M. and Nyholt, D.R. (2005):

Teenage acne is influenced by genetic factors.

Br. J. Dermatol.; 152: 565–95.

138) Vivier, E.;Tomasello, E.; Baratin, M.;Walzer, T. and Ugolini, S. (2008):

Functions of natural killer cells.

Nat. Immunol.; 9: 503–10.

139) Vora, S.;Ovhal, A. and Jerajani, H.(2008):

Correlation of facial sebum to insulin like growth factor-1 in patients with acne.

Br. J. Dermatol.; 159: 990-1.

(W)

140) Webster GF. (2005):

The pathophysiology of acne.

Cutis; 76 (Suppl):4–7.

141) Webster, G.F. (2008):

Commentary: diet and acne.

J. Am. Acad. Dermatol.; 58: 794-5.

142) Webster, G.F.; Leyden, J.J .and Nilsson, U.R.(1979):

Complement activation in acne vulgaris: consumption of complement by comedones.

Infect. Immun.; 26: 183-6.

143) Willett, W.; Manson, J. and Liu, S. (2002):

Glycemic index, glycemic load, and risk of type 2 diabetes.

Am. J. Clin. Nutr.;76:274S-80S.

144) Williams, C. and Layton, A.M. (2006):

Persistent acne in women. Implications for the patient and for therapy.

Am. J. Clin. Dermatol.; 7: 281-90

145) Williams, M., Cunliffe, W.J. and Gould, D.(1974):

Pilo-sebaceous duct physiology. Effect of hydration on pilosebaceous duct orifice.

Br. J. Dermatol.; 90: 631–5.

146) Wolf, R.; Matz, H. and Orion, E. (2004):

Acne and diet.

Clin. Dermatol.; 22: 387-93.

147) Wolters, M. (2005):

Diet and psoriasis: experimental data and clinical evidence.

Br. J. Dermatol.; 153:706–14.

148) Wynniss, L.; Tom and Victoria, R. (2008):

New insights into adolescent acne.

Current Opinion in Pediatrics ; 20:436–40.

(X)

- 149) Xiu-Li Wang; Hong-Wei Wang; Ling-Lin Zhang; Ming-Xia Guo and Zheng Huang. (2010):**

Topical ALA PDT for the treatment of severe acne vulgaris, Photodiagnosis and Photodynamic Therapy; 7, 33-38.

(Y)

- 150) Yee, K.C. and Cunliffe, W.J. (2010):**

Itching in acne an unusual complication of therapy. In: Burns, T., Breathnach, S., Griffiths, C. and Cox, N. editors. In Rook's Textbook of Dermatology. Dermatology; 189: 117–9.

- 151) Youn, S.W. (2010):**

The role of facial sebum secretion in acne pathogenesis facts and controversies. Clinics in Dermatology; 28: 8–11.

(Z)

- 152) Zaenglein, A.; Graber, E.M. and Thiboutot, D.M.(2008):**

Acne vulgaris and acneiform eruptions. In: Wolff K, Goldsmith LA, Katz SI, et al, editors. Fitzpatrick's dermatology in general medicine. New York: McGraw Hill; p. 690-703.

- 153) Zammit, V.A.; Waterman, I.J.; Topping, D. and McKay, G.(2001):**

Insulin stimulation of hepatic triacylglycerol secretion and the etiology of insulin resistance. J. Nutr.; 131: 2074-7.

- 154) Zhang, Y.; Proenca, R.; Maffei, M.; Barone, M.; Leopold, L. and Friedman, J.M.(1994):**

Positional cloning of the mouse obese gene and its human analogue.

Nature.; 372: 425-32.

- 155) Zolins, M.; Eady, E.A. and Avery, A.J. (2004):**

Comparison of five antimicrobial regimens for treatment of mild to moderate inflammatory facial acne vulgaris in the community: randomised controlled trial.

Lancet; 364: 2188–95.

- 156) Zorrilla, E.P.; Inoue, K.; Valdez, G.R.; Tabarin, A. and Koob, G.F. (2005):**

Leptin and post-prandial satiety: Acute central leptin more potently reduces meal frequency than meal size in the rat. Psychopharmacology, 177: 324–35.

- 157) Zouboulis, C.C. and Bohm, M. (2004):**

Neuroendocrine regulation of sebocytes pathogenetic link between stress and acne.

Exp. Dermatol.; 13(Suppl): 31-5.

- 158) Zouboulis, C.C.; Baron, J.M.; Bohm, M.; Kippenberger, S.; Kurzen, H. and Reichrath, J. (2008):**

Frontiers in sebaceous gland biology and pathology.

Exp. Dermatol.; 17: 542-51.

- 159) Zouboulis, C.C.; Chen, W.C.; Thornton, M.J.; Qin, K. and Rosenfield, R.(2007):**

Sexual hormones in human skin.

Horm. Metab. Res.; 39: 85-95.

- 160) Zouboulis, C.C.; Eady, A. and Philpott, M. (2005):**

What is the pathogenesis of acne?

Exp. Dermatol.; 14: 143-52.