

References

(1) Adamson, J.W. (2005):

Iron deficiency and other hypoproliferative anemia.

In: Kasper, D.L.; Braunwald, E.; Fauci, A.S.; Hauser, S.L.; Longo, D.L. and Jameson, J.L., (eds.): Harrison's principles of internal medicine. 16th edition.

New York: McGraw-Hill.; PP. 586-592.

(2) Adamson, J.W. and Longo, D.L. (2005):

Anemia and polycythemia.

In: Kasper, D.L.; Braunwald, E.; Fauci, A.S.; Hauser, S.L.; Longo, D.L. and Jameson, J.L., (eds.): Harrison's principles of internal medicine. 16th edition.

New York: McGraw-Hill.; PP. 329-336.

(3) Ahluwalia, N.; Lonnerdal, B.; Lorenz, S.G. and Allen L.H. (1998):

Spot ferritin assay for serum samples dried on filter paper.

Am. J. Clin. Nutr.; 67:88-92.

(4) Apache, P.G. (1986):

Eczematous dermatitis of the scalp.

In: Zviak, C., (ed.) : The science of hair care.

New York: Marcel Dekker.; 513-521.

(5) Arosio, P. and Levi, S. (2002):

Ferritin, iron homeostasis, and oxidative damage.

Free. Radic. Biol. Med.; 33: 457- 463.

(6) Artantas, S.; Gul, U.; Kilic, A. and Guler, S. (2009):

Skin finding in thyroid diseases.

E. J. I. M.; 20 : 158-161.

(7) Bardelli, A. and Rebora, A. (1989):

Telogen effluvium and minoxidil.

J. Am. Acad. Dermatol.; 21: 572-573.

(8) Barraud-Klenovsek, M.M. and Trueb, R.M. (2000):

Congenital hypotrichosis due to short anagen.

Br. J. Dermatol.; 143: 612- 617.

(9) Barron, B. A.; Hoyer, J.D. and Tefferi, A. A. (2001):

Bone marrow report of absent stainable iron is not diagnostic of iron deficiency.

Ann. Hematol.; 80:166-169.

(10) Bergfeld, W.F. (2008):

Telogen effluvium.

In: Michael, J. and Hordin, M. K., (eds.): Hair and Scalp Diseases: Medical, Surgical, and Cosmetic Treatments.

London, UK: Informa Health Care.; 9: 119-136.

(11) Bergfeld, W.F. and Mulinari-Brenner, F. (2001):

Shedding: How to manage a common cause of hair loss.

Cleve. Clin. J. Med.; 68: 256-261.

(12) Blume-Peytavi, U. and Orfanos, C.E. (2006):

Microscopy of the hair: the trichogram.

In: Serup, J.; Jemec, G.B.E. and Grove, G.L., (eds.): Handbook of non-invasive methods and the skin. Second edition .

Boca Raton, Fla: CRC Press.; PP. 875-881.

(13) Blume-Peytavi, U.; Hillmann, K. and Marcella, G. (2008):

Hair growth assessment techniques.

In: Blume-Peytavi, U.; Whiting, D.A.; Tosti, A. and Trub, R.M.,
(eds.): Hair growth and disorders.

Berlin, Heidelberg : Springer.; 8:125-157.

(14) Bregy, A. and Treub, R.M. (2008):

No association between serum ferritin levels >10µg/l and hair loss
activity in women.

Dermatology.; 217:1- 6.

(15) Brittenham, G.M. (2000):

Disorders of iron metabolism: iron deficiency and overload.

In: Hoffman, R. (ed.): Hematology: basic principles and practice.
3rd edition.

New York : Churchill-Livingston.; PP. 397- 428.

(16) Burman, K.D. and McKinley-Grant, L. (2006):

Dermatological aspects of the thyroid.

Clin. Dermatol.; 24:247-255.

(17) Burton, J.W.; Harlow, C. and Theil, E.C. (1998):

Evidence for reutilization of nodule iron in soybean seed
development.

J. Plant. Nutr.; 21: 913-927.

(18) Campbell, M.J. and Machin, D. (1993):

Medical statistics, a common sense approach.

In Wiley, J.S. (ed).

New York, PP. 266.

(19) Canfield, D. (1996):

Photographic documentation of hair growth in androgenetic alopecia. *Dermatol. Clin.*; 14(4):713-721.

(20) Chua, E.; Clague, J.E.; Sharma, A.K.; Horan, M.A. and Lombard, M. (1999):

Serum transferrin receptor assay in iron deficiency and anaemia of chronic disease in the elderly.

Q. J. Med.; 92:587-594.

(21) Chuong, C.M.; Hou, L.H.; Chen, P.J.; Wu, P.; Patel, N. and Chen, Y.P. (2001):

Dinosaur's feather and chicken's tooth? Tissue engineering of the integument .

Eur. J. Dermatol.; 11: 286-292.

(22) Dawber, R.P.R.; Simpson, N.B. and Barth, J.H. (1997):

Diffuse alopecia : endocrine, metabolic and chemical influences on the follicular cycle.

In: Dawber, R. P. R., (ed.): *Diseases of the follicular cycle.*

Oxford : Blackwell Science.; PP. 123-150.

(23) Decassia, W.R.; Kiyohara, P.K. and Silveira, M. (2007):

Electron microscopic observations of human hair medulla.

J. Microsc.; 226.(1): 54-63.

(24) Deloche, C.; Bastien, P.; Chadoutaud, S.; Galan, P. ; Bertrais, S.; Hercberg, S. and Delacharriere, O. (2007):

Low iron stores: a risk factor for excessive hair loss in non-menopausal women.

Eur. J. Dermatol.; 17 (6): 507-512.

- (25) Deloche, C.; Delacharriere, O. and Misciali, C.; Piraccini, B.M.; Vincenzi, C.; Bastien, P.; Tardy, I.; Bernard, B.A. and Tosti, A. (2004):**

Histological features of peripilar signs associated with androgenetic alopecia.

Arch. Dermatol. Res.; 295(10):422-428.

- (26) Elledge, S.J.; Zhou, Z. and Allen, J.B. (1992):**

Ribonucleotide reductase: Regulation.

Trends. Biochem. Sci.; 17: 119-123.

- (27) Elston, D.M.; McCollough, M.L. and Angeloni, V.L. (1995):**

Vertical and transverse sections of alopecia biopsy specimens:

Combining the two to maximize diagnostic yield.

Jam. Acad. Dermatol.; 32: 454- 457.

- (28) Feidler, V.C. and Gray, A.C. (2003):**

Diffuse alopecia: telogen hair loss.

In: Olsen, E.A., (ed.): Disorders of hair growth: diagnosis and treatment. 2nd edition.

New York: McGraw-Hill.; PP. 303-320.

- (29) Feidler, V.C. and Hafeez, A. (1994):**

Diffuse alopecia. telogen hair loss.

In: Olsen, E.A., (eds.) : Disorders of hair growth, diagnosis and treatment.

New York: McGraw-Hill.; PP. 241-255.

- (30) Ferreira, C.; Bucchini, D.; Martin, M.E.; Levi, S.; Arosio, P.; Grandchamp, B. and Beaumont, C. (2000):**
Early embryonic lethality of H ferritin gene deletion in mice.
J. Biol. Chem.; 275: 3021-3024.
- (31) Garcia-Hernandez, M.J. and Rodrioguez-Pichardo, A. (1999):**
Multivariate analysis in alopecia areata : Risk factors and validity of clinical forms.
Arch. Dermatol.; 135 : 998-999.
- (32) Goddard, A.F.; McIntyre, A.S. and Scott, B.B. (2000):**
Guidelines for the management of iron deficiency anemia.
B. S. G.; 46:1-5.
- (33) Goette, D.K. and Odum, R.B. (1976):**
Alopecia in crash dieters.
J. A. M. A.; 235: 2622-2623.
- (34) Guarrera, M.; Semino, M.T. and Rebora, A. (1997):**
Quantitating hair loss in women: a critical approach.
Dermatology.; 194(1):12-16.
- (35) Gupta, A.K.; Goldfarb, M.T.; Ellis, C.N. and Vorhees, J.J. (1989):**
Side-effect profile of acitretin therapy in psoriasis.
J. Am. Acad. Dermatol.; 20: 1088-1093.
- (36) Guyatt, G.H.; Oxman, A.D.; Ali, M.; Willan, A.; McIlroy, W. and Patterson, C. (1992):**
Laboratory diagnosis of iron-deficiency anemia: an overview.
J. Gen. Intern. Med.; 7:145-153.

- (37) Guyatt, G.H.; Patterson, C.; Ali, M.; Singer, J.; Levine, M.; Turpie, I. (1990):**
Diagnosis of iron-deficiency anemia in the elderly.
Am. J. Med.; 88:205-209.
- (38) Hadjur, C.; Daty, G. and Madry, G. (2002):**
Cosmetic assessment of the human hair by confocal microscopy.
Scanning.; 24:59-64.
- (39) Hanakawa, Y.; Li, H.; Lin, C.; Stanley, J.R. and Cotsarelis, G. (2004):**
Desmogleins 1 and 3 in the companion layer anchor mouse anagen hair to the follicle.
J. Invest. Dermatol.; 123: 817-822.
- (40) Hard, S. (1963):**
Non-anemic iron deficiency as an etiologic factor in diffuse loss of hair of the scalp in women.
Acta. Dermatol. Venereol.; 43:562-569.
- (41) Harrison, P.M.; Treffry, A. and Lilley, T.H. (1986):**
Ferritin as an iron-storage protein: mechanisms of iron uptake.
J. Inorg. Biochem.; 27: 287-293.
- (42) Harrison, S. and Bergfeld, W. (2009):**
Diffuse hair loss: its triggers and management.
Cleve. Clin. J. Med.; 76 (6): 361-367.
- (43) Harrison, S. and Sinclair, R. (2002):**
Telogen effluvium.
Clin. Exp. Dermatol.; 27: 389-395.

(44) Hassan, E.O.; El-Hussinie, M. and El-Nahal, N. (1999):

The prevalence of anemia among clients of family planning clinics in Egypt.

Contraception.; 60 : 93-99.

(45) Headington, J.T. (1993):

Telogen effluvium. New concepts and review.

Arch. Dermatol.; 129: 356–363.

(46) Heath, A.L.M. and Worwood, M. (2001):

If it□'s not broken, don't fix it.

Br. Med. J.; 323: 806.

(47) Herbert, V. (1992):

Everyone should be tested for iron disorders.

J. Am. Diet. Assoc.; 92: 1502-1509.

(48) Hillmann, K. and Blume-peytavi, U. (2009):

Diagnosis of hair disorders.

Semin. Cutan. Med. Surg.; 33-38.

(49) Hoffman, R.M. (2000):

The hair follicle as a gene therapy target.

Nat. Biotechnol.; 18: 20-21.

(50) Hoffmann, R.M. (2005):

Trichoscan: What is new?

Dermatology.; 211: 54- 62.

- (51) Huang, H.Q.; Lin, Q. M.; Kong, B.; Zeng, R.Y.; Qiao, Y.H.; Chen, H.; Zhang, F. Z. and Xu, L.S. (1999):**
Role of phosphate and kinetic characteristics of complete iron release from native pig spleen ferritin-Fe.
J. Prot. Chem.; 18: 497-504.
- (52) Ito, M.; Kizawa, K.; Hamada, K. and Cotsarelis, G. (2004):**
Hair follicle stem cells in the lower bulge form the secondary germ, a biomedically distinct but functionally equivalent progenitor cell population, at the termination of catagen. Differentiation; 72: 548-557.
- (53) Jahoda, C.A. and Reynolds, A.J. (1996):**
Dermal-epidermal interactions. Adult follicle derived cell populations and hair growth.
Dermatol. Clin.; 14:573-583.
- (54) Jandl, J.H. and Katz, J.H. (1963):**
The plasma-to-cell cycle of transferrin.
J. Clin. Invest.; 42:314-326.
- (55) Kantor, J.; Kessler, L.J.; Brooks, D.G. and Cotsarelis, G. (2003):**
Decreased serum ferritin is associated with alopecia in women.
J. Invest. Dermatol.; 121: 985-988.
- (56) Kasteler, J.S. and Callen, J.P. (1994):**
Scalp involvement in dermatomyositis. Often overlooked or misdiagnosed.
J. A. M. A.; 272:1939-1941.

(57) Kligman, A.M. (1961):

Pathologic dynamics of human hair loss : Telogen effluvium.
Arch. Dermatol.; 83:175-198.

**(58) Knovich, M.A.; Storey, J.A.; Coffman, L.G.; Torti, S.V. and
Torti, F. M. (2009):**

Ferritin for the clinician.
Blood. Reviews.; 23 : 95-104.

(59) Krause, k. and Foitzik, k. (2006):

Biology of the hair follicle : the basics.
Semin. Cutan. Med. Surg.; 25 : 2-10.

(60) Kullavanijaya, P.; Gritiyarangsang, P. and Bisalbutra, P. (1992):

Absence of effects of dimethicone- and nondimethicone-
containing shampoos on daily hair loss rates.
J. Soc. Cosmet. Chem.; 43: 195-206.

**(61) Kumamoto, T.; Shalhevet, D.; Matsue, H.; Mummert,
M.E.; Ward, B.R.; Jester, J.V. and Takashima, A.
(2003):**

Hair follicles serve as local reservoirs of skin mast cell
precursors. Blood.; 102: 1654-1660.

(62) Lacarruba, F.; Dall'Oglio, F. and Nasca, M.R. (2004):

Videodermatoscopy enhances diagnostic capability in some
forms of hair loss.
Am. J. Clin. Dermatol.; 5:205-208.

- (63) Lauer, A.C.; Ramachandran, C.; Lieb L.M.; Niemiec, S. and Weiner, N.D. (1996):**
Targeted delivery to the pilosebaceous unit via liposomes.
Adv. Drug. Deliv. Rev.; 18: 311–324.
- (64) Levi, S.; Santambrogio, P.; Corsi, B.; Cozzi, A. and Arosio, P. (1996):**
Evidence that residues exposed on the three-fold channels have active roles in the mechanism of ferritin iron incorporation.
Biochem. J.; 317: 467- 473.
- (65) Liau, G.; Chan, L.M. and Feng, P. (1991):**
Increased ferritin gene expression is both promoted by cAMP and a marker of growth arrest in rabbit vascular smooth muscle cells.
J. Biol. Chem.; 266 : 18819-18826.
- (66) Lindner, G; Botchkarev, V.A. and Botchkareva, N.V. (1997):**
Analysis of apoptosis during hair follicle regression (catagen).
Am. J. Pathol.; 151: 1601-1617.
- (67) Lynch, S. and Green, R. (2001):**
Assessment of nutritional anemias.
In: Ramakrishnan U, (ed.): Nutritional anemias.
New York: CRC Press.; PP. 23-42.
- (68) Mcelwee, K.L. and Sinclair, R. (2008):**
Hair physiology and its disorders.
In: Finkel, T. and Lowenstien, C., (eds.): Drug discovery today: Diseases mechanisms. Common skin conditions and disorder.
Berlin, Germany: Elsevier.; vol. 5(2) PP.e163-e171.

- (69) Meidan, V.M.; Bonner, M.C. and Michniak, B.B. (2005):**
Transfollicular drug delivery – is it a reality?
Int. J Pharm.; 306:1–14.
- (70) Milman, N. and Kirchhoff, M. (1992):**
Iron stores in 1359, 30-to 60-year old Danish women: evaluation
by serum ferritin and hemoglobin.
Ann. Hematol.; 64: 22-27.
- (71) Milner, Y.; Sudnik, J.; Filippi, M.; Kizoulis, M.; Kashgarian,
M. and Stenn, K. (2002):**
Exogen, shedding phase of the hair growth cycle:
Characterization of a mouse model.
J. Invest. Dermatol.; 119:639-644.
- (72) Mirmirani, P.; Hessol, N.; Maurer, T.; Berger, T.; Greenblatt,
R. and Price, V. (2001):**
Prevalence of hair disorders in the women's interagency HIV
study.
J. Invest. Dermatol. 117:434.
- (73) Moeinvaziri, M.; Mansoori, P.; Holakooee, K.; Safaei, N.Z. and
Abbasi, A. (2009):**
Iron status in diffuse telogen hair loss among women.
Acta. Dermatovenerol. Croat.; 17(4):279-284.
- (74) Nishimura, E.K.; Granter, S.R. and Fisher, D.E. (2005):**
Mechanisms of hair greying: incomplete melanocyte stem cell
maintenance in the niche.
Science.; 307:720-724.

(75) Olsen, E.A. (2003):

Current and novel methods for assessing efficacy of hair growth promoters in pattern hair loss.

J. Am. Acad. Dermatol.; 48(2):253-262.

(76) Oro, A.E. and Scott, M.P. (1998):

Splitting hairs: dissecting roles of signaling systems in epidermal development.

Cell.; 95: 575–578.

(77) Panteleyev, A.A.; Jahoda, C.A. and Christiano, A.M. (2001):

Hair follicle predetermination.

J. Cell. Sci.; 114 : 3419-3431.

(78) Patzelt, A.; Knorr, F.; Blume-Peytavi, U.; Sterry, W. and Lademann, J. (2008):

Hair follicles, their disorders and their opportunities.

In: Finkel, T. and Lowenstien, C., (eds.): Drug discovery today:

Diseases mechanisms. Common skin conditions and disorder.

Berlin, Germany: Elsevier.; vol. 5 (2) PP. 173-181.

(79) Paus, R. and Foitzik, K. (2004):

In search of the “hair cycle clock”: A guided tour.

Differentiation; 72 : 489-511.

(80) Paus, R.; Müller-Röver, S. and Botchkarev, V.A. (1999):

Chronobiology of the hair follicle: Hunting the “hair cycle clock”.

J. Invest. Dermatol. Symp. Proc.; 4 : 338-345.

(81) Peters, E.M.; Botchkarev, V.A.; Botchkareva, N.V.; Tobin, D.J. and Paus, R. (2001):

Hair cycle-associated remodeling of the peptidergic innervation of murine skin and hair growth modulation by neuropeptides.
J. Invest. Dermatol.; 116:236-245.

(82) Pierard, G.E.; Pierard-Franchimont, C.; Marks, R. and Elsner, P. (2004):

EEMCO guidance for the assessment of hair shedding and alopecia.
Skin. Pharmacol. Physiol.; 17(2):98-110.

(83) Punnonen, K.; Irjala, K. and Rajamaki, A. (1997):

Serum transferrin receptor and its ratio to serum ferritin in the diagnosis of iron deficiency.
Blood.; 89 : 1052-1057.

(84) Puolakka, J. (1980):

Serum ferritin in the evaluation of iron status in young healthy women.
Acta. Obstet. Gynecol. Scand.; 95 : 35- 41.

(85) Ramakrishnan, U. and Semba, R.D. (2009):

Iron deficiency and anemia.
In: Semba, R. D. and Bloem, M. W., (eds.): Nutrition and health in developing countries. Second edition .
Totowa, NJ : Humana Press.; Ch. 16. PP. 479-505.

(86) Rebora, A. (1997) :

Telogen effluvium.
Dermatology.; 195 : 209-212.

(87) Rebora, A. and Guarrera, M. (2002):

Kenogen. A new phase of the hair cycle?

Dermatology.; 205: 108–110.

(88) Rebora, A.; Guarrera, M.; Baldari, M. and Vecchio, F. (2005):

Distinguishing androgenetic alopecia from chronic telogen effluvium when associated in the same patient.

Arch. Dermatol.; 143 : 1243-1245.

(89) Ren, B.; Tibbelin, G.; Kajino, T.; Asami, O. and Ladenstein, R. (2003):

The multi-layered structure of Dps with a novel di-nuclear ferroxidase center .

J. Mol. Biol.; 329 : 467-477.

(90) Reynolds, A.J. and Jahoda, C.A.B. (1991):

Hair follicle stem cells? A distinct germinative epidermal cell population is activated in vitro by the presence of hair dermal papilla cells.

J. Cell. Sci.; 99 : 373-385.

(91) Rook, A. and Dawber, R. (1982):

Diffuse alopecia: endocrine, metabolic and chemical influences on the follicular cycle.

In: Rook, A. and Dawber, R., (eds.) : Diseases of the Hair and Scalp.

Oxford, UK : Blackwell Science Publications.; Ch. 5. PP. 115-145.

(92) Ross, E.K.; Vincenzi, C. and Tosti, A. (2006):

Videodermoscopy in the evaluation of hair and scalp disorders.
J. Am. Acad. Dermatol.; 55:799-806.

(93) Roudier-Pujol, C. and Jan, V. (2000):

Alopecies medicamenteuses.
Ann. Dermatol. Venereol. Suppl.; 127: 26-28.

(94) Rushton, D. H. (1993):

Management of hair loss in women.
Dermatol. Clin.; 11: 47-53.

(95) Rushton, D.H. (2003):

Decreased serum ferritin and alopecia in women
J. Invest. Dermatol.; 121 : 985-988.

(96) Rushton, D.H. and Ramsay, I.D. (1992):

The importance of adequate serum ferritin levels during oral
cyproterone acetate and ethinyl oestradiol treatment of diffuse
androgen-dependent alopecia in women.
Clin. Endocrinol (Oxf) ; 36: 421-427.

(97) Rushton, D.H.; Dover, R.; Norris, M.J. and Gilkes, J.J.H.

(2007):

Iron and hair loss in women: what is deficiency? This is the real
question.
J. Am. Acad. Dermatol.; 518-519.

(98) Rushton, D.H.; Dover, R.; Sainsbury, A.W.; Norris, M.J.; Gilkes, J.J.H. and Ramsay, I.D. (2001):

Why should women have lower reference limits for hemoglobin and ferritin concentrations than men?

Br. M. J.; 322 : 1355-1357.

(99) Rushton, D.H.; Dover, R.; Sainsbury, A.W.; Norris, M.J.; Gilkes, J.J. and Ramsay, I.D. (2002):

Iron difficiency is neglected in women's health.

Br. M. J.; 325: 1176.

(100) Rushton, D.H.; James, K.C. and Mortimer, C.H. (1983):

The unit area trichogram in the assessment of androgen-dependent alopecia.

Br. J. Dermatol.; 109 : 429-437.

(101) Rushton, D.H.; Ramsay, I.D.; James, K.C.; Norris, M.J. and Gilkes, J.J. (1990):

Biochemical and trichological characterization of diffuse alopecia in women.

Br. J. Dermatol.; 123 : 187-197.

(102) Santambrogio, P.; Cozzi, A.; Levi, S. and Arosio, P. (1987):

Human serum ferritin G-peptide is recognized by anti-L ferritin subunit antibodies and concanavalin-A.

Br. J. Haematol., 65: 235-237.

(103) Sato, N.; Leopold, P.L. and Crystal, R.G. (1999) :

Induction of the hair growth phase in postnatal mice by localized transient expression of Sonic hedgehog.

J. Clin. Invest.; 104: 855–864.

(104) Shapiro, J.; Wiseman, M. and Lui, H. (2000):

Practical management of hair loss.

Can. Fam. Physician.; 46: 1469 - 1477.

(105) Sinclair, R. (1999):

Diffuse hair loss.

Int. J. Dermatol.; 38: 8-18.

(106) Sinclair, R. (2002):

There is no clear association between low serum ferritin and chronic diffuse telogen hair loss.

Br. J. Dermatol.; 147: 982-984.

(107) Sperling, L.C. (2001):

Hair and systemic disease.

Dermatol. Clin.; 19 : 711-726.

(108) Stefanini, S.; Desideri, A.; Vecchini, P.; Drakenberg, T. and Chiancone, E. (1989):

Identification of the iron entry channels in apoferritin. Chemical modification and spectroscopic studies.

Biochemistry.; 28: 378-382.

(109) Stenn, K.S. (2005):

Exogen is an active, separately controlled phase of the hair growth cycle.

J. Am. Acad. Dermatol.; 52: 374–375.

(110) Stenn, K.S. and Paus, R. (2001):

Controls of hair follicle cycling.

Physiol. Rev.; 81: 449-494.

- (111) Stillman, T.J.; Upadhyay, M.; Norte, V.A.; Sedelnikova, S.E.; Carradus, M.; Tzokov, S.; Bullough, P.A.; Shearman, C.A.; Gasson M.J.; Williams, C.H.; Artymiuk, P. and Green, J. (2005):**

The crystal structures of *Lactococcus lactis* MG1363 Dps proteins reveal the presence of an N-terminal helix that is required for DNA binding.

Mol. Microbiol.; 57: 1101–1112.

- (112) Takagi, H.; Shi, D.; Ha, Y.; Allewell, N.M. and Theil, E. (1998):**

Localized unfolding at the junction of three ferritin subunits. A mechanism for iron release?

J. Biol. Chem.; 273 : 18685-18688.

- (113) Tefferi, A. (2003):**

Anemia in adults: a contemporary approach to diagnosis.

Mayo Clinic Proc.; 78 : 1274-1280.

- (114) Tobin, D.J. and Paus, R. (2001):**

Graying gerontobiology of the hair follicle pigmentary unit.

Exp. Gerontol.; 36:29-54.

- (115) Tobin, D.J.; Gunin, A.; Magerl, M.; Handijski, B. and Paus, R. (2003):**

Plasticity and cytokinetic dynamics of the hair follicle mesenchyme: Implications for hair growth control.

J. Invest. Dermatol.; 120 : 895-904.

(116) Tosti, A. and Pazzaglia, M. (2007):

Drug reactions affecting hair: diagnosis.

Dermatol. Clin.; 25:223–231.

(117) Treffry, A.; Bauminger, E.R.; Hechel, D.; Hodson, N.W.;

Nowik, I.; Yewdall, S.J. and Harrison, P.M. (1993):

Defining the roles of the threefold channels in iron uptake, iron oxidation and iron-core formation in ferritin: a study aided by site-directed mutagenesis.

Biochem. J.; 296 : 721-728.

(118) Treffry, A.; Zhao, Z.; Quail, M.A.; Guest, J.R. and Harrison, P. M. (1998):

How the presence of three iron binding sites affects the iron storage function of the ferritin (EcFtnA) of Escherichia coli

FEBS. Lett.; 432 : 213-218.

(119) Trikha, J.; Waldo, G.S.; Lewandowski, F.A.; Ha, Y.; Theil, E.C.; Weber, P.C. and Allewell, N.M. (1994):

Crystallization and structural analysis of bullfrog red cell L-subunit ferritins.

Proteins.; 18 : 107-118.

(120) Trost, L.B.; Bergfeld, W.F. and Clogeras, E. (2006):

The diagnosis and treatment of iron deficiency and its potential relationship to hair loss.

J. Am. Acad. Dermatol.; 54 : 824-844.

(121) Trowbridge, I.S. and Omary, M.B. (1981):

Human cell surface glycoprotein related to cell proliferation is the receptor for transferrin.

Proc. Natl. Acad. Sci.; 78 : 3039-3043.

(122) Trueb, R.M. (2008):

Diffuse hair loss.

In: Blume-Peytavi, U.; Whiting, D.A.; Tosti, A. and Trub, R.M., (eds.): Hair growth and disorders.

Verlag, Berlin, Heidelberg : Elsevier.; 13 : 259-272.

(123) Trueb, R.M. (2010):

Systematic approach to hair loss in women.

J.D.D.G.; 8 : 1-13.

(124) Umbreit, J. (2005):

Iron deficiency: a concise review.

Am. J. Hematol.; 78:225-231.

(125) Van der Steen, P.; Traupe, H.; Happle, R.; Boezeman, J.;

Strater, R. and Hamm, H. (2003):

The genetic risk for alopecia areata in first degree relatives of severely affected patients. An estimate.

Acta. Dermato.Venereol.; 72(5) : 373-375.

(126) Van Neste, D.J. (2001):

Contrast enhanced phototrichogram (CE-PTG): an improved non-invasive technique for measurement of scalp hair dynamics in androgenetic alopecia - validation study with histology after transverse sectioning of scalp biopsies.

Eur. J. Dermatol.; 11(4) : 326-331.

(127) Van Neste, D.J. (2002):

Assessment of hair loss: clinical relevance of hair growth evaluation methods.

Clin. Exp. Dermatol.; 27(5):358-365.

(128) Van Neste, D.J. and Rushton, D.H. (1997):

Hair problems in women.

Clinics. Dermatol.; 15: 113-125.

(129) Van Neste, D.J. and Tobin, D.J. (2004):

Hair cycle and hair pigmentation: Dynamic interactions and changes associated with aging.

Micron.; 35 : 193-200.

(130) Van Neste, D.J. and Trueb, R.M. (2006):

Critical study of hair growth analysis with computer-assisted methods. J. Eur .

Acad. Dermatol. Venereol.; 20(5):578-383.

(131) Vet, J.A.; Moorselaar, R.J.; Debruyne, F.M. and Schalken, J.A. (1997):

Differential expression of ferritin heavy chain in a rat transitional cell carcinoma progression model.

Biochim. Biophys. Acta.; 1360 : 39-44.

(132) Vogt, A. and Blume-Peytavi, U. (2003):

Biology of the human hair follicle. New knowledge and the clinical significance.

Hautarzt ; 54: 692–698.

- (133) Vogt, A.; Hadam, S.; Heiderhoff, M.; Audring, H.; Lademann, J.; Sterry, W. and Blume-Peytavi, U. (2007):**
Morphometry of human terminal and vellus hair follicles.
Exp. Dermatol.; 16: 946-950.
- (134) Waalen, J.; Feltiti, V. and Beutler, E. (2002):**
Hemoglobin and ferritin concentration in men and women: cross sectional study.
Br. M. J.; 325: 137.
- (135) Waikel, R.L.; Kawachi, Y.; Waikel, P.A.; Wang, X.J. and Roop, D.R. (2001):**
Deregulated expression of c-Myc depletes epidermal stem cells.
Nature. Genet.; 28:165-168.
- (136) Watt, G.D.; Frankel, R.B. and Parafthymiou, G.C. (1985):**
Reduction of mammalian ferritin.
Proc. Natl. Acad. Sci.; 82 : 3640-3643.
- (137) Weismann, K. (1980):**
Zinc metabolism and the skin.
In: Rook, A.S.J. and Savin, J.A., (eds.): Recent advances in dermatology 5.
Edinburgh: Churchill Livingstone.; PP. 109-129.
- (138) Weiss, G. and Goodnough, L.T. (2005):**
Anemia of chronic disease.
N. Engl. J. Med.; 352:1011-1023.
- (139) West, C.E. (1996):**
Strategies to control nutritional anemia.
Am. J. Clin. Nutr.; 64 : 789-790.

(140) Whiting, D.A. (1996):

Chronic telogen effluvium: Increased scalp hair shedding in middle aged women.

J. Am. Acad. Dermatol.; 35: 899-906.

(141) Whiting, D.A. (1996):

Chronic telogen effluvium.

Dermatol. Clin.; 14 : 723-773.

(142) Whiting, D.A. (2004):

The Structure of the Human Hair Follicle. Light Microscopy of Vertical and Horizontal Sections of Scalp Biopsies. Pfizer, Canfield Publishing.

(143) Wu, K.J.; Polack, A. and Dalla-Favera, R. (1999):

Coordinated regulation of iron-controlling genes, H-ferritin and Irp2, by c-Myc.

Science.; 283:676-679.

(144) Yang, X.; Arosio, P. and Chasteen, D. (2000) :

Molecular diffusion into ferritin: pathways, temperature dependence, incubation time, and concentration effects.

Biophys. J.; 78 : 2049-2059.

(145) Younis, I. (1990):

Incidence of skin diseases over a 5 year period in Cairo.

Med. J. Cairo Univ.; 57(3): 215-224.

(146) Younis, I. and El-Azzooni, M. (1998):

Diffuse hair loss in women.

Med. J. Cairo Univ.; 66(2): 455-459.

(147) Yu, M.; Finner, A.; Shapiro, J.; Lo, B.; Barekatain, A. and McElwee, K. (2006):

Hair follicles and their role in skin health.

Exp. Rev. Dermatol.; 1: 855–871.

(148) Zheng, Y.; Eilertsen, K. J.; Ge, L.; Zhang, L.; Sundberg, J.P.; Prouty, S.M.; Stenn, K.S. and Parimoo, S. (1999):

Scd1 is expressed in sebaceous glands and is disrupted in the asebia mouse.

Nature. Genet.; 23:268-270.

(149) Zheng, Y.; Prouty, S.M.; Harmon, A.; Sundberg, J.P.; Stenn, K.S. and Parimoo, S. (2001):

Scd3 a novel gene of the stearyl-CoA desaturase family with restricted expression in skin.

Genomics.; 71:182-191.