



REFERENCES



References

ACOM [American College of Occupational and Environmental Medicine] (2002): “Noise-induced Hearing Loss”. Found online at ACOEM website <http://www.acoem.org/guidelines/article.asp?ID=53>.

Ahmad HO, Dennis JH, Badran O, Ismail M, Ballal SG, Ashoor A, et al. Occupational noise exposure and hearing loss of workers in two plants in eastern Saudi Arabia. *Ann OccupHyg* 2001; 45: 371-80.

Aran JM., Hiel H., Hayashida T., Erre JP., Aurousseau C., Gulhaume A., Dulon D. (1992): “Noise, aminoglycosides, and diuretics”. In: Dancer AL., Henderson D., Salvi RJ. and Hamernik RP. eds. *Noise-Induced Hearing Loss*. St. Louis, MO: Mosby Year Book. Pp. 188–195.

Ashe PC. and Berry MD. (2003): “Apoptotic signaling cascades”. *Prog Neuropsychopharmacol Biol Psychiatry* 27(2):199-214.

Axelsson A. and Barrenas ML. (1992): “ Tinnitus in noise-induced hearing loss”. In: Dancer AL., Henderson D., Salvi RJ. and Hamernik RP. eds. *Noise-Induced Hearing Loss*. St. Louis, MO: Mosby Year Book. Pp. 269–276.

Axelsson A. and Vertes D. (1982): “ Histological findings in cochlear vessels after noise”. In: Hamernik RP., Henderson D. and Salvi R, eds. “New Perspectives on Noise-Induced Hearing Loss”. New York: Raven Press. Pp. 49–68.

Babisch W., Ising H., Gallacher J., Elwood PC., Sweetnam PM., Yarnell JWG., Bainton D. and Baker IA. (1990): “ Traffic noise, work noise and cardiovascular risk factors”. *The caerphilly and Speedwell*

Bedi R. Evaluation of occupational environment in two textile plants in northern India with specific reference to noise. *Ind Health* 2006; 44: 112-6.

Beraneck LL. (1971): “Noise and vibration control”. Mc Graw – Hill Vbook Company, New York.

References

Berger EH., Royster LH. and Thomas WG. (1978): “Presumed noise-induced permanent threshold shift resulting from exposure to an A-weighted Leq of 89 dB”. *Journal of the Acoustical Society of America* 64:192–197.

Berger, EH (2000). Hearing protection devices. In: Berger, EH, Royster, LH, Roster, J.D., Driscoll, DP, and Layne, M, Eds. *The Noise Manual*. Fairfax, VA. American Industrial Hygiene Association, pp. 429-432.

Berglund B. and Lindvall T. (1995): “Community Noise”. *Archives of the Center for Sensory Research*. Stockholm. 2(1), 1-195.

Bruel and Kjaer (1984): “Sound and the human being” 2850, Noerum, Denmark.

Burns A. (1973): “The measurement of sound”. William Clows and Sons, limited (publish), Lodon, Great Britain, P.42 – 122.

Chavalitsakulchai P, Kawakami T, Kongmuang U. Vivatijestsadawut. Noise exposure and permanent hearing loss of textile workers in Thailand. *Industrial health*, 1989; 27:165-173.

Choi DW. (1992): “Excitotoxic cell death”. *J Neurobiol* 23(9):1261-1276.

Davis H. and Silverman S. R. (1970): “Hearing and Deafness”. New York: Holt, Rinehart, & Winston.

Duncan RK. and Saunders JC. (2000): “Stereocilium injury mediates hair bundle stiffness loss and recovery following intense water-jet stimulation”. *J Comp Physiol [A]* 186(11):1095-1106.

Ehrenberger K. and Felix D. (1991): “Glutamate receptors in afferent cochlear neurotransmission in guinea pigs”. *Hear Res* 52(1):73-80.

Feldman A. and Grimes C. (1985): “Employee training programs in occupational hearing conservation”. In A. S. Feldman and C.T. Grimes (Ed), “Hearing conservation in industry”. Baltimore: Williams and Wikins.

References

Franks JR. and Morata TC. (1996): “Ototoxic effects of chemicals alone or in concert with noise”. A review of human studies. In: Axelsson A, Borchgrevink HM, Hamernik RP, Hellstrom PA., Henderson D. and Salvi RJ. eds. “Scientific Basis of Noise-Induced Hearing Loss”. New York: Thieme. Pp. 437–446.

Goldstein M. (1994): “Low- frequency components in complex noise and their perceived loudness and annoyance”. World and health scientific Journal issued by National Institute of Occupational health,5- 17184 Solna, Sweden.

Guingnard J. (1973): “A basis for limiting noise exposure for hearing conservation”. Prepared for Environmental Protection Agency, EPA -550-9-73-001-A, 4-21.

Gunter R, Kowalska S. Lipiwezan A. A permanent noise induced shift in auditory threshold in textile industry workers. Med Pr 1986; 37: 175 - 86.

Gutteridge JM. and Halliwell B. (2000): “Free radicals and antioxidants in the year 2000”. A historical look to the future. Ann N Y Acad Sci 899:136-147.

Hamernik RP., Hsueh KD. (1991): “Impulse noise: Some definitions, physical acoustics and other considerations”. Journal of the Acoustical Society of America 90(1):189–196.

Health and Welfare Canada. (1989): “Federal-Provincial Advisory Committee on Environmental and Occupational Health National Guidelines for Environmental Noise Control Procedures and Concepts for the Drafting of Environmental Noise Regulations”. By laws in Canada. Ottawa, Ontario: Minister of Supply and Services Canada.

Health Council of the Netherlands. (1996): “Effects of noise on health”. Chapter 3 of a report on noise and health prepared by a Committee of the Health Council of The Netherlands In Noise/News International, 4(3), 137-150.

References

Henderson D. and Hamernik RP. (1995): “Biologic bases of noise-induced hearing loss”. *Occup Med* 10(3):513-534.

Henderson D., Hamernik RP. (1982): “Asymptotic threshold shift from impulse noise”. In: Hamernik RP, Henderson D, Salvi R, eds. *New Perspectives on Noise-Induced Hearing Loss*. New York: Raven Press. Pp. 265–281.

Henderson D., Hamernik RP. (1986): “Impulsive noise”. Critical review. *Journal of the Acoustical Society of America* 80(2):569–584.

Henderson D., Subramaniam M. and Boettcher FA. (1993): “Individual susceptibility to noise-induced hearing loss”. An old topic revisited. *Ear and Hearing* 14(3):152–168.

Henselman LW., Henderson D., Shadoan J., Subramaniam M., Saunders S. and Ohlin D. (1995): “Effects of noise exposure, race, and years of service on hearing in U.S. Army soldiers”. *Ear and Hearing* 16(4):382 -391

Horne JA., Pankhurst FL., Reyner LA., Hume K. and Diamond ID. (1994): “A field study of sleep disturbance: Effects of aircraft noise and other factors on 5,742 nights of actimetrically monitored sleep in a large subject sample”. *Sleep*, 17(2), 146-159.

Hu BH., Guo W., Wang PY., Henderson D. and Jiang SC. (2000): “Intense noise-induced apoptosis in hair cells of guinea pig cochleae”. *Acta Otolaryngol* 120(1):19-24.

Juyaratnum, J. Primary health Care approach in occupational health and safety. *African Newsletter on occupational health and safety*, 1991;1 (3): 72-5.

Kill RL. Hearing loss in female jute weavers. *Annal Occupation*. 1983; 18 (10): 97-109.

References

Kryter K. (1985): “Definitions of noise and noise pollution” From Karl D. Kryter’s (Ed) “Effect of noise on man” Academic press, Harcourt, Brace Jovanovich (publish) ch, 1, P1-3.

Larry E., Humes, Lois M., Joellenbeck, and Jane S. (2005): “Noise and Military Service”. Durch, Editors, Committee on Noise-Induced Hearing Loss and Tinnitus Associated with Military Service from World War II to the Present.

Le Page EL. and Johnstone BM. (1980): “Nonlinear mechanical behaviour of the basilar membrane in the basal turn of the guinea pig cochlea. Hear Res 2(3-4):183-189.

Lehnhardt E. (1984): “Klinik der Innenohrschwerhörigkeiten”. Arch Otorhinolaryngol Suppl. 1:58-218.

Leysieffer H., Baumann JW., Muller G. and Zenner HP. (1997): “Ein implantierbarer piezoelektrischer Hörerätewandler für Innenohrschwerhörige, Teil II: Klinisches Implantat”. HNO 45(10):801-815.

Lipscomb M. (1988): “Determination of Noise exposure from David M”. Lipscomb’s (Ed). Hearing conservation in Industry, Schools and Military. Little, Brown (Publish). U.S.A. chapter4. P 35-43.

Majno G. and Joris I. (1995): “Apoptosis, oncosis, and necrosis. An overview of cell death”. Am J Pathol 146(1):3-15.

Mattson MP. (2003): “Excitotoxic and excitoprotective mechanisms: abundant targets for the prevention and treatment of neurodegenerative disorders”. Neuromolecular Med 3(2):65-94.

Melnick W. (1976): “Human asymptotic threshold shift”. In: Henderson D., Hamernik RP., Mills JH. (1973): “Temporary and permanent threshold shifts produced by nine-day exposures to noise”. Speech and Hearing Research 16(3):426–438.

Melnink W. (1985): “Industrial hearing conservation”. Jack Kat’s Ch. 6,P. 61 – 76.

References

Merry CJ and Franks JR (1995). Historical assessment and future directions in the prevention of occupational hearing loss. In: Morata TC and Dunn DE, eds.

Miller JM., Schacht J. and Altschuler R. (2003): “Prevention of noise-induced hearing loss”. In: Henderson D., Prasher D., Kopke R. and Hamernik RP. editors. “Noise induced hearing loss: basic mechanisms, prevention and control”. London: NRN publications, 215-230.

Mills J. and Lilly D. (1971): “Temporary threshold shifts produced by pure tones and by noise in the absence of acoustic reflex”. Journal of the acoustical and the society of America, 50:1556 – 1558.

Mills JH., Gengel RW., Watson CS. and Miller JD. (1970): “Temporary changes of the auditory system due to exposure to noise for one or two days”. Journal of the Acoustical Society of America 48(2, Pt 2):524–530.

Mrena R., Savolainen S., Kuokkanen JT. and Ylikoski J. (2002): “Characteristics of tinnitus induced by acute acoustic trauma”. A long-term follow-up. Audiology and Neuro-Otology 7(2): 122–130.

National Institutes of Health (1990). Noise and Hearing Loss. NIH Consensus Development Conference Consensus Statement 1990, Jan 22-24; 8.

Nicotera TM., Hu BH. and Henderson D. (2003): “The Caspase Pathway in Noise-Induced Apoptosis of the Chinchilla Cochlea”. J Assoc Res Otolaryngol 4(4):466-77.

Nudelmann AA, Costa EA, Feligman J, Ibáñez RN. Perda auditiva induzida pelo ruído. Rio de Janeiro: Revinter; 2001. Occupational Medicine: State of the Art Reviews - Occupational hearing loss. New York: Hanley and Belfus, Inc., pp.669-682.

Ohlemiller KK., McFadden SL., Ding DL., Lear PM. and Ho YS. (2000): “Targeted mutation of the gene for cellular glutathione peroxidase

References

(Gpx1) increases noise-induced hearing loss in mice”. J Assoc Res Otolaryngol 1(3):243-254.

Ohlemiller KK., Wright JS. and Dugan LL. (1999): “Early elevation of cochlear reactive oxygen species following noise exposure”. Audiol Neurotol 4(5):229-236.

Olero UG. Ijaduola GTA. Sowho E. Hearing Threshold in an Auto Assembly Plant: perspects for Hearing Conservation in a Nigerian Factory. Int.Arch. Environ. Health, 1990 62:199-202.

OSHA [Occupational Safety and Health Adminstration] (1974): “Guidelines for noise enforcement”. Washington DC: U.S. Department of Labor, CPL 2-2.35A.

OSHA [Occupational Safety and Health Adminstration] (1981): “Occupational noise exposure”. Hearing conservation amendment, Final rule Fedral Register, 48: 9738 – 9785.

Plontke S., Zenner H.P. and Dietz K. (2004): “Hearing loss from occupational and leisure noise”. GMS Current topics in Otorhiolaryngology-Head and Neck Surgery 3: Doc 06.

Popa R., Anniko M., Takumida M. and Arnold W. (2001): “Localization of nitric oxide synthase isoforms in the human cochlea”. Acta Otolaryngol 121(4):454-459.

Pourbakht A. and Yamasoba T. (2003): “Ebselen attenuates cochlear damage caused by acoustic trauma”. Hear Res 181(1-2):100-108.

Preyer S. and Gummer AW. (1996): “Nonlinearity of mechanoelectrical transduction of outer hair cells as the source of nonlinear basilar-membrane motion and loudness recruitment”. Audiol Neurotol 1(1):3-11.

Puel JL., Ruel J., Gervais dC and Pujol R. (1998): “Excitotoxicity and repair of cochlear synapses after noise-trauma induced hearing loss”. Neuroreport 9(9):2109-2114.

References

Reed JC. (2000): “Mechanisms of apoptosis”. *Am J Pathol* 157(5):1415-1430.

Roberto M., Hamernik RP., Turrentine GA. (1989): “Damage of the auditory system associated with acute blast trauma”. *Ann Otol Rhinol Laryngol Suppl* 140:23-34.

Rodriguez GP. and Gerhardt KJ. (1991): “Influence of outer ear resonant frequency on patterns of temporary threshold shift”. *Ear and Hearing* 12(2):110–114.

Royster LH., Thomas WG., Royster JD. and Lilley D. (1978): “Potential hearing compensation cost by race and sex”. *Journal of Occupational Medicine* 20(12):801–806.

Ruel J., Chen C., Pujol R., Bobbin RP. and Puel JL. (1999): “AMPA-preferring glutamate receptors in cochlear physiology of adult guinea-pig”. *J Physiol* 518 (Pt 3):667-680.

Saunders JC., Schneider ME. and Dear SP. (1985): “The structure and function of actin in hair cells”. *J Acoust Soc Am* 1985 78(1 Pt 2):299-311.

Shakhatreh, F M. Abdul- Baqi K J. Turk M M. Hearing loss in a textile factory. *Saudi M J*, 2000; 21 (1): 58-60.

Shi X., Ren T. and Nuttall AL. (2002): “The electrochemical and fluorescence detection of nitric oxide in the cochlea and its increase following loud sound”. *Hear Res* 164(1-2):49-58.

Sies H., Sharov VS., Klotz LO. and Briviba K. (1997): “A new function for selenoproteins as peroxynitrite reductase”. *J Biol Chem* 272(44):27812-27817.

Spongr VP., Boettcher FA., Saunders SS. and Salvi RJ. (1992): “Effects of noise and salicylate on hair cell loss in the chinchilla cochlea”. *Archives of Otolaryngology–Head and Neck Surgery* 118(2):157–164.

References

Suter AH. (1991): “Noise and Its Effects”. Prepared for the Consideration of the Administrative Conference of the United States.

Taylor W. Noise level of a wide jute loom with and without plastic parts. *Journal of the Textile Institute.* 1967; 9: 377-384.

Thalmann RR. (1976): “Quantitative biochemical techniques for studying normal and noise damaged ears”. In: Henderson D., Hamernik RP., Dosanjh DS., Mills JH. eds. *Effects of Noise on Hearing.* New York: Raven Press. Pp. 129–154.

Thompson S. (1996): “Non-auditory health effects of noise: Updated review”. *Proceedings of Internoise,* 2177-2182.

Tilney LG., Saunders JC., Egelman E. and DeRosier DJ. (1982): “Changes in the organization of actin filaments in the stereocilia of noise-damaged lizard cochleae”. *Hear Res* 7(2):181-197.

Van Campen LE., Dennis JM., Hanlin RC., King SB. And Velderman AM. (1999): “One-year audiologic monitoring of individuals exposed to the 1995 Oklahoma City bombing”. *Journal of the American Academy of Audiology* 10(5):231–247.

Ward W. (1991): “The role of intermittence in P.T.S.”. *Journal of the acoustical Society of America.* 90, 164 – 169.

Ward WD. (1995): “Endogenous factors related to susceptibility to damage from noise”. *Occupational Medicine* 10(3):561–575.

Ward WD., Royster JD. and Royster LH. (2000): “Auditory and nonauditory effects of noise”. In: Berger EH., Royster LH., Royster JD., Driscoll DP. and Layne M. eds. “*The Noise Manual*”l. 5th ed. Fairfax, VA: American Industrial Hygiene Association. Pp. 101–147.

Yamane H., Nakai Y., Takayama M., Iguchi H., Nakagawa T. and Kajima A. (1995): “Appearance of free radicals in the guinea pig inner ear after noise-induced acoustic trauma”. *Eur Arch Otorhinolaryngol* 252(8):504-508.

References

Yamasoba T. and Dolan DF. (1998): “The medial cochlear efferent system does not appear to contribute to the development of acquired resistance to acoustic trauma”. *Hear Res* 120(1-2):143-151.

Zenner HP., Zimmermann U. and Gitter AH. (1987): “Fast motility of isolated mammalian auditory sensory cells”. *Biochem Biophys Res Commun* 149(1):304-308.

Zheng J., Shen W., He DZ., Long KB., Madison LD. and Dallos P. (2000): “Prestin is the motor protein of cochlear outer hair cells”. *Nature* 405(6783):149-155.

Zwerling, C, Whitten PS, Davis CS, and Sprince NL (1997). Occupational Injuries Among Workers With Disabilities, *Journal of the American Medical Association*, 278(24):2163-2166.