

References

Annamaria Morelli, Sandra Filippi, Linda Vignozzi, Rosa Mancina,

Mario Maggi (2006):

Physiology of Erectile Function: An Update on Intracellular Molecular Processes. European Association of Urology (EAU).

European Board of Urology (EBU) update series; 96–108

Antonello Silvestri, Pasquale Galetta, Elena Cerquetani, Giuseppe

Marazzi, Roberto Patrizi, Massimo Fini and Giuseppe

Rosano (2003):

Report of erectile dysfunction after therapy with beta-blockers is related to patient knowledge of side effects and is reversed by placebo.

Eur Heart J; 24: 1928-1932.

Aranda A, Ruilope L, Calvo A (2004):

Erectile dysfunction in essential arterial hypertension and effects of sildenafil: results of a Spanish national study.

Am J Hypertens 17: 139–145.

Bahgat Mittawae, Abdel Rahman El-Nashaar, Ayman Fouda,

Mahmoud Magdy, Rany Shamloul (2006):

Incidence of erectile dysfunction in 800 hypertensive patients: Elsevier Inc.

A Multicenter Egyptian National Study. 67: 575–578.

Bax G, Marin N, Piarulli F, Lamonica M, Bellio F, Fedele D (1998):

Rigiscan evaluation of specific nervous impairment in patients with diabetes and erectile disorders.

Diabetes Care; 21(7):1159-61.

Beevers DG, Gregory YH and O'Brien Eoin, (2007):

Pathophysiology of hypertension.

ABC of hypertension, 5th edition, Blackwell publishing; 3:
12-17.

Bischoff E, Schneider K, Jeremy K (2001):

A conscious-rabbit model to study vardenafil hydrochloride and other agents that influence penile erection.

Int. J. Impot. Res; 13(4): 230-5.

Bivalacqua TJ, Deng W, Champion HC, Hellstrom WJ, Kadowitz PJ (2004):

Gene therapy techniques for the delivery of endothelial nitric oxide synthase to the corpora cavernosa for erectile dysfunction.

Methods Mol. Biol; 279:173–85.

Black HR, Weltin G, Jaffe CC (1991):

The limited echocardiogram: a modification of standard echocardiography for use in the routine evaluation of patients with systemic hypertension.

Am J Cardiol; 67:1027–1030.

Blumentals W, Gomez-Caminero A, Joo S, Vannappagari V (2003):

Is erectile dysfunction predictive of peripheral vascular disease.

The Aging Male, 6, 217-221.

Brenner and Rector's (2004):

Chronic Kidney Disease, Dialysis and Transplantation. In:
The Kidney,

Brenner and Rector's Textbook, 7th ed., Editor: Saunders, pp. 2118-2119.

Brindis RG and Kloner RA (2003):

Sildenafil in patients with cardiovascular disease.
Am. J. Cardiol; 92(9A):26M-36M.

Brock GB, McMahon CG, Chen KK (2002):

Efficacy and safety of tadalafil for the treatment of erectile dysfunction: results of integrated analyses.
J Urol; 168: 1332-6.

Broderick GA (2003):

Oral Pharmacotherapy and the Contemporary Evaluation and Management of Erectile Dysfunction
Rev Urol.; 5(Suppl 7): S9–S20.

Burchardt M, Burchardt T, Anastasiadis A (2001):

Erectile dysfunction is a marker for cardiovascular complications and psychological functioning in men with hypertension.
Int. J. Impot. Res; 13: 276–281.

Buvat J, Maggi M, Gooren L, Guay AT, Kaufman J, Morgentaler A, Schulman C, Tan HM, Torres LO, Yassin A, Zitzmann M (2010):

Endocrine aspects of male sexual dysfunctions.
J Sex Med.;7(4 Pt 2):1627-56.

Cai H and Harrison DG (2000):

Endothelial dysfunction in cardiovascular diseases: the role of oxidant stress.
Circulation Research; 87 (10): 840–4.

Chan SS, Leung DY, Abdullah AS, Lo SS, Yip AW, Kok WM, Ho SY, Lam TH (2010):

Smoking-cessation and adherence intervention among Chinese patients with erectile dysfunction.

Am J Prev Med. Sep; 39(3):251-8.

Chang DW, Wood CG, Kroll SS (2003):

Cavernous nerve reconstruction to preserve erectile function following non-nervesparing radical retropubic prostatectomy: a prospective study.

111(3):1174-1181.

Chew KK, Earle CM, Stuckey BG, Jamrozik K, Keogh EJ (2000):

Erectile dysfunction in general medical practice: Prevalence and clinical correlates.

Int. J. Impot. Res; 12:41-45

Christ G and Hodges S (2006):

Molecular Mechanisms of detrusor and corporal myocyte contraction: identifying targets for pharmacotherapy of bladder and erectile dysfunction.

British Journal of Pharmacology; 147:41-55

Chun J and Carson CC (2001):

Physician-patient dialogue and clinical evaluation of erectile dysfunction.

Urol. Clin. North. Am; 28: 249.

Cordero A, Bertomeu-Martínez V, Mazón P, Fácila L, González-Juanatey JR (2010):

Erectile dysfunction may improve by blood pressure control in patients with high-risk hypertension.

Postgrad Med. Nov;122(6):51-6.

Corvol P, Persu A, Gimenez-Roqueplo AP, Jeunemaitre X (1999):

Seven lessons from two candidate genes in human essential hypertension: angiotensinogen and epithelial sodium channel.

Hypertension; 33(6): 1324–31.

Day D, Ambegaonkar A, Hamot K, McDaniel A (2001):

A new tool for predicting erectile dysfunction.

Adv. Ther; 18:131-9.

Davis JL (2003):

Smoking Can Lead to Erectile Dysfunction.

ebMD Health News Mar 6.

[http://www.webmd.com/erectile-dysfunction/news/](http://www.webmd.com/erectile-dysfunction/news/20030307/smoking-can-lead-to-erectile-dysfunction)

20030307/ smoking-can-lead-to-erectile-dysfunction

Dean A, Dean G, Coulompler B (1994):

EPI-INFO version: a word processing database and statistics program for epidemiology on microcomputer centers and disease control. Atlanta, Georgia. USA.

Dean C, and Lue T (2005):

Physiology of penile erection and pathophysiology of erectile dysfunction.

Urologic Clinics; 32, 379-395.

Della Chiesa A, Pfiffner D, Meier B (2003):

Sexual activity in hypertensive men.

J Hum Hypertens; 17(8):515-521.

Dey J (2002):

Evaluation and treatment of erectile dysfunction in men with diabetes mellitus.

Mayo Clin. Proc; 77:276-82.

Díez J (1999):

Insulin-like growth factor I in essential hypertension.
Kidney Int; 55:744–759.

Djavan B, Milan S, Fong YK (2005):

Dutasteride: a novel dual inhibitor of 5 alpha-reductase for
benign prostatic hyperplasia.
Expert Opin. Pharmacother; 6:311-17.

**Doumas Michel, Alexandros Tsakiris, Stella Douma, Alkiviadis
Grigorakis, Angelos Papadopoulos, Athina Hounta,
Sotirios Tsiodras, Dimitrios Dimitriou, Helen
Giamarellou (2006):**

Factors affecting the increased prevalence of erectile
dysfunction in Greek hypertensive compared to
normotensive subjects.
J Androl 27: 469-477.

Dula E, Bukofzer S, Perdok R, George M (2001):

Double-blind, crossover comparison of 3 mg apomorphine
SL with placebo and with 4 mg apomorphine SL in male
erectile dysfunction.
Eur. Urol; 39:558–63.

Eardley I, Wright P, MacDonagh R, Hole J, Edwards A (2004):

An open-label, randomized, flexible-dose, crossover study
to assess the comparative efficacy and safety of sildenafil
citrate and apomorphine hydrochloride in men with erectile
dysfunction.
BJU. Int; 93: 1271-5

Erectile Dysfunction Alliance UK (1999):

Management guidelines for erectile dysfunction. London:
The Royal Society of Medicine Press.

BMJ 2000; 321:499-503.

**Eric Wespes A, Edouard Amar B, Dimitrios Hatzichristou C, Kosta
Hatzimouratidis C, Francesco Montorsi D, John Pryor
E, Yoram Vardi F (2006):**

EAU Guidelines on erectile dysfunction: an update.

European urology; 49(5):806-15.

Fazio L, Brock G, Kriss D (2004):

Erectile dysfunction: management update. CMAJ;
170:1429-37.

**Feldman HA, Goldstein I, Hatzichristou DG, Krane RJ, McInlay JB
(2000):**

Impotence and its medical and psychosocial correlates:
results of the Massachusetts Male Aging Study.

J. Urol; 151: 54-61.

Ferrario CM and Levy P (2002):

Sexual dysfunction in patients with hypertension:
implications for therapy.

J. Clin. Hypertens; 4(6):424-432.

**Fogari R, Zoppi A, Poletti L, Marasi G, Mugellini A, Corradi L
(2001):**

Sexual activity in hypertensive men treated with valsartan
or carvedilol: a crossover study.

Am. J. Hypertens; 14:27-31.

Franks S, Jacobs HS, Martin N, Nabarro JDN (1978):

Hyperprolactinaemia and Impotence.
Clinical Endocrinology; 8(4): 277-287.

Gambescia N, Weeks G, and Lokhart D (2000):

Erectile Dysfunction: Integrating Couple Therapy, Sex
Therapy and Medical Treatment. New York: W.W. Norton
& Company.
Journal of Family Psychotherapy. Vol 20, Issue 2,3: pp 221
– 240.

**Garban H, Vernet D, Freedman A, Rajfer J, and Gonzalez-Cadavid
N (1995):**

Effect of aging on nitrous oxide-mediated penile erection
in the rat.
American Journal of Physiology; 268: 467-475.

Giuliano FA, Leriche A, Jaudinot EO, deGendre AS (2004):

Prevalence of erectile dysfunction among 7689 patients
with diabetes or hypertension, or both.
Urology; 64:1196-1201.

Goldstein I (2000):

Male sexual circuitry. Working Group for the Study of
Central Mechanisms in Erectile Dysfunction.
Sci. Am; 283: 70.

**Goldstein I, Lue TF, Padma-Nathan H, Rosen RC, Steers WD,
Wicker PA (1998):**

Oral sildenafil in the treatment of erectile dysfunction.
N. Engl. J. Med; 338:1397–1404

Goldstein I, Payton TR, Schechter PJ (2001):

A double-blind, placebo-controlled, efficacy and safety study of topical gel formulation of 1% alprostadil (Topiglan) for the in office treatment of erectile dysfunction.

Urology; 301:5-57.

Gorier FE, McClure RD, Kramer-Levien D (1996):

Endocrine screening for sexual dysfunction using free testosterone determinations.

J. Urol; 156:405-8.

Grimm RH, Grandits GA, Prineas RJ, McDonald RH, Lewis CE, Flack JM (1997):

Long-term effects on sexual function of five antihypertensive drugs and nutritional hygienic treatment in hypertensive men and women. Treatment of Mild Hypertension Study (TOMHS).

Hypertension; 29: 8-14

Haas C, Seftel A, Razmjouei K, Ganz M, Hampei N, and Ferguson K (1998):

Erectile dysfunction in aging: up regulation of endothelial nitric oxide synthase.

Urology; 51: 516-522.

Hellstrom WJ, Gittelman M, Karlin G (2003):

Sustained efficacy and tolerability of vardenafil, a highly potent selective phosphodiesterase type 5 inhibitor, with erectile dysfunction: results of a randomized, double-blind, 26-week placebo-controlled pivotal trial.

Urology; 8:14-61

Hsu GL, Wen HS, Hsieh CH, Liu LJ, Chen YC (2001):

Traumatic glans deformity: reconstruction of distal ligamentous structure.

J Urol; 166:1390.

Irwin MB and Kata EJ (1994):

High attrition rate with intracavernous injection of prostaglandin E1 for impotency.

Urology; 43:84–89.

Jabaloyas JM (2010):

Hormonal etiology in erectile dysfunction.

Arch Esp Urol.; 63(8):621-7.

Jardin A, Wagner G, Khoury S, Guiliano F, Goldstein I, and Nathan H (2000):

Recommendations of the First International Consultation on Erectile Dysfunction, cosponsored by the World Health Organization.

Plymouth UK: Health Publications. Plymouth: Plymbridge Distributors Ltd: 241-303 & 711-726.

Jensen J, Lendorf A, Stimpel H, Frost J, Ibsen H, Rosenkilde P (1999):

The prevalence and etiology of impotence in 101 male hypertensive outpatients.

Am. J. Hypertens; 12:271–275.

Jin Z, Zhu YC, Cui WS, Liu T, Li WR, Yuan YM, Song WD, Xin ZC (2010):

Clinical efficacy and patient satisfaction with penile prosthesis implantation for the treatment of severe erectile dysfunction.

Beijing Da Xue Xue Bao.; 42(4): 413-7.

JNC VI (1997):

Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. The Sixth Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure.

Arch Intern Med; 157:2413–2446.

Jones BE (2000):

Basic mechanisms of sleep–wake states. In: Kryger MH, Roth T, Dement WC, editors.

Principles and Practice of Sleep Medicine. 3rd ed. Philadelphia: W. B. Saunders; 134–54.

Khan MA, Mikhailidis DP, Morgan RJ (2002):

Sildenafil: A urologist's view. Current Medical Research & Opinion, Supplement; 16(1):s48-s58.

Kim Y, de Miguel F, Usiene I (2006):

Injection of skeletal muscle-derived cells into the penis improves erectile function.

Int. J. Impot. Res; 18:329–34.

Kinsey AC, Pomeroy WB and Martin CE (1948):

Sexual Behavior in the Human Male. Philadelphia: W. B. Saunders.

Archives of Sexual Behavior; Volume 31, Number 2, 155-156.

Kirby RS, Culley C, and Goldstein I (1999):

Erectile Dysfunction: A Clinical Guide. Herndon, VA:
Books International.

Textbook of Erectile Dysfunction, vol. 1, Urology 54; pp
431–432.

Kirby RS, O'Leary MP, Carson C (2005):

Efficacy of extended-release doxazosin and doxazosin
standard in patients with concomitant benign prostatic
hyperplasia and sexual dysfunction.

BJU Int; 95:103-9.

Klabunde R, Ryan K, and Paxson C (2007):

Acute hyperglycaemia does not alter coronary vascular
function in isolated perfused rat hearts.

Diabetes, Obesity & Metabolism; 95: 697-705.

Klaus M, Justus P, Bernhard C, Michael S, Joachim W (2005):

Sleep investigations in erectile dysfunction, Department of
Psychiatry, University of Mainz, Untere Zahlbacher Strabe,
Mainz, Germany.

Journal of Psychiatric Research; 39(1): 93-9.

Kloner R (2007):

Erectile dysfunction and hypertension.

Int. J. Impot. Res; 19(3):296-302.

Levme SB (2003):

Erectile dysfunction: why drug therapy isn't always
enough.

Cleve. Clin. J. Med; 70:241-6

Lewis JH, Rosen R, Goldstein I (2004):

Erectile dysfunction in primary care.

Nurse Pract; 29:42-6

Lewis RW, Witherington R, and Gary T (1997):

External vacuum therapy for erectile dysfunction: use and results.

World J. Urol; 15:79-82

Litwin MS, Melmed GY, and Nakazon T (2001):

Life after radical prostatectomy: a longitudinal study.

J. Urol; 166:587

Lohmeier TE (2001):

The sympathetic nervous system and long-term blood pressure regulation.

American Journal of Hypertension 14 (6 Pt 2): 147–154.

Lue TE (2000):

Erectile dysfunction.

N. Engl. J. Med; 342:1802-13.

Luft FC (1998):

Molecular genetics of human hypertension.

J Hypertens; 16:1871–1878

Lundberg L and Biriell JO (1993):

Long-term experience of self-injection therapy with prostaglandin-E1 for erectile dysfunction.

Scand. J. Urol. Nephrol; 30(5):395-397.

Maas R, Schwedhelm E, Albsmeier J (2002):

The pathophysiology of erectile dysfunction related to endothelial dysfunction and mediators of vascular function.

Vasc. Med; 7(3):213-225.

Mak R, De Backer G, Kornitzer M, and De Meyer JM (2002):

Prevalence and correlates of erectile dysfunction in a population-based study in Belgium.

Eur Urol, 41:132

Martin-Morales A, Sanchez-Cruz JJ, Saenz de Tejada I, Rodriguez-Vela L, Jimenez-Cruz JF, Burgos-Rodriguez R (2001):

Prevalence and independent risk factors for erectile dysfunction in Spain: results of the Epidemiologia de la Disfuncion Erectil Masculina Study.

J Urol; 166:569-74.

McCullough AR (2004):

Optimizing patient response to oral erectogenic pharmacotherapy.

Current Sexual Health Reports, Vol. 1(1): 24-28.

McVary KT, Carrier S, and Wessells H (2001):

Smoking and erectile dysfunction: evidence based analysis.

J Urol, 166: 1624

Melander O, Orho M, Fagerudd J (1998):

Mutations and variants of the epithelial sodium channel gene in Liddle's syndrome and primary hypertension. Hypertension;

31(5): 1118-24.

<http://hyper.ahajournals.org/cgi/pmidlookup?view=long&pmid=9576123>.

Melman A and Gingell J (1999):

The epidemiology and pathophysiology of erectile Dysfunction.

J Urol; 161: 5-11.

Mittawae B, El-Nashaar AR, Fouda A, Magdy M, Shamloul R (2006):

Incidence of erectile dysfunction in 800 hypertensive patients: a multicenter Egyptian national study.

Urology; 67(3):575-8.

Moncada I, Jara J, Subira D, Castano I, Hernandez C (2004):

Efficacy of sildenafil citrate at 12 hours after dosing: re-exploring the therapeutic window.

Eur. Urol; 46:357–60

Montague DK, Angermeier KW (2001):

Penile prosthesis implantation.

Urol. Clin. North Am; 28:355–61.

Montorsi F, Briganti A, Salonia A, Rigatti P, and Burnett A (2006):

Can Phosphodiesterase Type 5 Inhibitors Cure Erectile Dysfunction?

European Urology; 49: 979-986.

Morales A, and Heaton JP (2001):

Hormonal erectile dysfunction. Evaluation and management.

Urol Clin North Am, 28: 279

Moreira E, Bestane W, Bartolo E, Fittipaldi J (2002):

Prevalence and determinants of erectile dysfunction in Santos, southeastern Brazil.

Sao Paulo Med. J; 120(2):131-139.

Muammer Kendirci A, Patrick E. Teloken B, Hunter C. Champion C, Wayne JG, Trinity E (2006):

Gene Therapy for Erectile Dysfunction: Fact or Fiction?.

European urology; 50(6): 1208-22.

National Institute of Health (1993):

NIH consensus development panel on impotence.
JAMA 270:83-90.

National Institutes of Health (2005):

Impotence: NIH Publication No. 06-3923, December, 2005, Consensus Statement Online:
<http://consensus.nih.gov/1992/1992Impotence091Html.htm>

Nicolosi A, Moreira ED Jr, Shirai M, Bin Mohd Tambi MI, Glasser DB (2003):

Epidemiology of erectile dysfunction in four countries: cross-national study of the prevalence and correlates of erectile dysfunction.
Urology; 61:201-6.

Nofzinger EA, Buysse DJ, Miewald JM, Meltzer CC, Price JC, Sembrat RC, Ombao H, Reynolds CF, Monk TH, Hall M, Kupfer DJ, Moore RY (2002):

Human regional cerebral glucose metabolism during non-rapid eye movement sleep in relation to waking.
Brain; 125:1105-15.

Oparil S, Zaman MA, Calhoun DA (2003):

Pathogenesis of hypertension.
Ann. Intern. Med.; 139(9): 761-76.

Padma-Nathan H, Hellstrom WJG, Kaiser F (1997):

Treatment of men with erectile dysfunction with transurethral alprostadil.
N. Engl. J. Med; 336:1-7.

Parati G, Di Rienzo M, Ulian L, Santucci C, Girard A, Elghozi J, Mancia G (1998):

Clinical relevance of blood pressure variability.

J Hypertens; 16:25-33.

Paul Arduca (2004):

Erectile Dysfunction: A guide to diagnosis and management. Medical Director, Men's Health Clinic, Freemasons Hospital, Victoria, AUSTRALIA.

Middle East Journal of Family Medicine; Vol. 4 (4): 1-15.

Pegge N, Twomey A, Vaughton K, Gravenort M, Ramsey M, and Price D (2006):

The role of endothelial dysfunction in the pathophysiology of erectile dysfunction in diabetes and in determining response to treatment.

Diabetic Medicine; 23: 873-878.

Pierdomenico SD, Di Nicola M, Esposito AL (2009):

Prognostic Value of Different Indices of Blood Pressure Variability in Hypertensive Patients.

American Journal of Hypertension; 22: 842–7.

Piero Montorsi A, Paolo M, Ravagnani A, Stefano Galli A, Sarah Ghulam Ali A, Alberto Briganti B, Andrea Salonia B, Francesco Montorsi B (2008):

The Triad of Endothelial Dysfunction, Cardiovascular Disease, and Erectile Dysfunction: Clinical Implications.

European Association of Urology; Volume 8, Issue 2, Pages 58-66.

Porst H, Padma-Nathan H, Giuliano F, Anglin G, Varanese L, Rosen R (2003):

Efficacy of tadalafil for the treatment of erectile dysfunction at 24 and 36 hours after dosing: a randomized controlled trial.

Urology; 62:121–5.

Prisant LM, Alpert BS, Robbins CB, Berson AS, Hayes M, Cohen ML, Sheps SG (1995):

American National Standard for nonautomated sphygmomanometers: summary report.

Am. J. Hypertens; 8:210–213.

Putz R and Pabsteds R (2001):

Pelvic diaphragm [floor]: male and female external genitalia. In: Putz R, Pabsteds R, eds.

Sobotta Atlas of Human Anatomy. Philadelphia, PA: Lippincott Williams & Wilkins: pp 222 – 239.

Ralph D, McNicholas T (2000):

UK management guidelines for erectile dysfunction.

BMJ; 321:499-503.

Remigio Vela Navarrete (2008):

Detecting Endothelial Dysfunction in Patients with Erectile Dysfunction: Are There Reliable Laboratory Markers?

European Urology Supplements;

Volume 8, Issue 2, Pages 67-70.

Romanelli F, Sansone A, Lenzi A (2010):

Erectile dysfunction in aging male.

Acta Biomed.;81 Suppl 1:89-94.

Rosen RC, Cappelleri JC, Smith MD (1999):

Development and evaluation of an abridged, 5-item version of the International Index of Erectile Function (IIEF-5) as a diagnostic tool for erectile dysfunction.

International Journal of Impotence. Res; 11:319-326.

Rosen RC, Ralph FR, Lewis AN (1997):

The International Index of Erectile Function (IIEF): a multidimensional scale for assessment of erectile dysfunction.

Urology; 49: 822-830

Saenz de Tejada I, Angulo J, Celtek S, Gonzalez-Cadavid N, Heaton

J, Pickard R (2005):

Pathophysiology of erectile dysfunction.

J Sex Med; 2: 26-39.

Safarinejad MR (2008):

Evaluation of endocrine profile and hypothalamic-pituitary-testis axis in selective serotonin reuptake inhibitor-induced male sexual dysfunction.

J Clin Psychopharmacol.; 28(4):418-23.

Schmidt TD (1993):

The ischio-cavernosus and bulbo-spongiosus muscles in mammalian penile rigidity.

Sleep; 16:171–183.

Schouten BW, Bohnen AM, Groeneveld FP, Dohle GR, Thomas S, Bosch JL (2010):

Erectile dysfunction in the community: trends over time in incidence, prevalence, GP consultation and medication use--the Krimpen study: trends in ED.

J Sex Med. Jul;7(7):2547-53. Epub 2010 May 19.

Seftel AD, Sun P, Swindle R (2004):

The prevalence of hypertension, hyperlipidemia, diabetes mellitus, and depression in men with erectile dysfunction.

J. Urol; 171(6 Part 1): 2341-2345.

Segura J and Ruilope LM (2007):

Obesity, essential hypertension and renin-angiotensin system.

Public Health Nutrition. Volume 10, Issue 10A , pp 1151 - 1155

Seyam R, Albakry A, Ghobish A (2003):

Prevalence of erectile dysfunction and its correlates in Egypt: a community based study.

International Journal of Impotence. Res; 15: 237–245.

Shabbir M, Mikhailidis M, Morgan RJ (2004):

Erectile dysfunction: an underdiagnosed condition associated with multiple risk factors.

Curr Med Res Opm; 20:603-6.

Shabsigh R, Kaufman JM, Steidle C, Padma-Nathan H (2004):

Randomized study of testosterone gel as adjunctive therapy to sildenafil in hypogonadal men with erectile dysfunction who do not respond to sildenafil alone.

J Urol; 63:172:658

Shaeer K, Osegbe D, Siddiqui S (2003):

Prevalence of erectile dysfunction and its correlates among men attending primary care clinics in three countries: Pakistan, Egypt, and Nigeria.

Int. J. Impot. Res. 15: 8-14.

Shafik A (1998):

The mechanism of ejaculation: the glans-vasal and urethro-muscular reflexes.

Arch Androl; 41:71–78

Siroky MB, Azadzo KM (2003):

Vasculogenic erectile dysfunction: newer therapeutic strategies.

Journal of Urology; 170:24 –29.

Sommer F, Schwarzer U, Klotz T, Caspers HP, Haupt G, and Engelmann U (2001):

Erectile dysfunction in cyclists: Is there any difference in penile blood flow during cycling in an upright versus a reclining position?

European Urology, 39: 720

Speckens AEM, Hengeveld MW, Lycklama AN, Van Hemert AM, Hawton KE (1993):

Discrimination between psychogenic and organic erectile dysfunction.

J. Psychosom. Res; 37:135–45.

Strano-Rossi S, Anzillotti L, de la Torre X, Botrè F (2010):

A gas chromatography/mass spectrometry method for the determination of sildenafil, vardenafil and tadalafil and their metabolites in human urine.

Rapid Commun Mass Spectrom. 15; 24(11):1697-706.

Sullivan M, Keoghane S, and Miller M (2001):

Vascular Risk Factors and Erectile Dysfunction.

BJU International; 87: 838-845.

Sun P and Swindle R (2005):

Are men with erectile dysfunction more likely to have hypertension than men without erectile dysfunction? A naturalistic national cohort study.

J. Urol; 174: 244-248.

Takahashi H (2008):

Sympathetic hyperactivity in hypertension (in Japanese).
Nippon Rinsho. Japanese Journal of Clinical Medicine;
66(8): 1495–502.

Taub HC, Lerner SE, Melman A, and Christ GJ (1993):

Relationship between contraction and relaxation in human and rabbit corpus cavernosum.

Urology, 42: 698.

Tsiara AS, Speckens AE, Kattemolle MR, Hengeveld MW (2003):

A prospective long-term follow-up study of patients evaluated for erectile dysfunction: outcome and associated factors.

Int. J. Impot. Res; 7(2):101-110.

Valiquette L, Young JM, Moncada I (2008):

Sustained efficacy and safety of vardenafil for treatment of erectile dysfunction: a randomized, double-blind, placebo-controlled study.

Mayo Foundation for Medical Education and Research,
Mayo Clin. Proc; 80(10):1291-7.

Wagner G, Claes H, Costa S, Cricelli M (2002):

A shared care approach to the management of erectile dysfunction in the community.

International Journal of Impotence. Res; 14: 189–194.

Wespes E, Amar E, Hatzichristou D, Montorsl E Pryor J, Vard Y (2002):

Guidelines on erectile dysfunction. European.
Urology; 41:1-5.

Williams IL, Wheatcroft SB, Shah AM, and Kearney MT (2002):

Obesity, atherosclerosis and the vascular endothelium:
mechanisms of reduced nitric oxide bioavailability in obese
humans.

Int. J. Obes. Relat. Metab. Disord; 26: 754.

**World Health Organization (WHO)/International Society of
Hypertension (ISH) (2003):**

Statement on management of hypertension.
J Hypertens; 21:1983–1992.

Wysowski DK and Swann J (2003):

Use of medications for erectile dysfunction in the United
States, 1996 through 2001.

J. Urol; 169:1040–1042.

Yuan YM, Zhou S, Zhang K (2010):

Methods for evaluation of penile erection hardness.
Zhonghua Nan Ke Xue; 16(7):642-5.

Zedan H, Hareadei AA, Abd-Elsayed AA, Abdel-Maguid EM (2010):

Cigarette smoking, hypertension and diabetes mellitus as
risk factors for erectile dysfunction in upper Egypt.

East Mediterr Health J. Mar; 16(3):281-5.