

References

- Abaza, H.; El-Fadaly, N.; Asal, F. and El-Kalla, F. (1995):** A study of tumour necrosis factor Alpha (TNF- α) and interleukin -6 in spontaneous bacterial peritonitis (SBP). Journal of Hepatol. Gastroenterol. & Inf. Diseases, 3 (5): 1-8.
- Abbitt, P. (1995):** Peritoneal cavity. In: Abbitt, P.L. (ed). New York: McGraw – Hill, 161 – 197.
- Abdel-Kader, S.; Mohran, Z. and Shehata, I. (1995):** A bedside procedure for the diagnosis of spontaneous bacterial peritonitis. Journal of Hepatol. Gastroenterol. & Inf. diseases, 3(5): 23-27.
- Abrink M., Larsson E., Gobl A. et al. (2000):** Expression of lactoferrin in the kidney: implications for innate immunity and iron metabolism Kidney International, 57:2004–2010.
- Aisen, P. and Liebman, A. (1972):** Lactoferrin and transferrin :a comparative study. Biochimica et Biophysica Acta.257:314-323.
- Akriviadis, E.A. and Runyon, B.A. (1990):** The value of an algorithm in differentiating spontaneous from secondary bacterial peritonitis. Gastroenterology, 98: 127 – 133.
- Albillos, A.; Cuervas- Mons, V.; Millan, L. et al. (1990):** Ascitic fluid polymorphonuclear cell count and serum to ascites albumin gradient in the diagnosis of bacterial peritonitis. Gastroenterol. 98: 134-140.
- Almadal, T.P. and Skinhoj, P. (1987):** Spontaneous bacterial peritonitis in cirrhosis, incidence, diagnosis and prognosis. Scand. J. Gastroenterol, 22:295-300.

- Al Sadec, M.M.; Mohammed, M.A.A.; Fathy, E.M.; Metwaly, M.A.A.; Abd Elsama, S.A.U.; Ismail, Y.M. (2004):** Evaluation of urine-screening leucocyte esterase strips in the diagnosis and follow up of treatment response in SBP. From the department of hepatology, gasrtoentriology, infectious diseases and clinical pathology benha faculty of medicine, zagazig university, Egypt.
- Al Sharif, Y.M. (1999):** Role of ultrasonography in the diagnosis of spontaneous bacterial peritonitis. MS.C. Thesis Faculty of Medicine, Cairo University, Egypt.
- Andersen, J.H.; Osbakk, S.A.; Vorland, L.H.; Traavik, T.; Gutteberg, T.J. (2001):** Lactoferrin and cyclic lactoferrin inhibit the entry of human fibroblasts. *Antiviral Research*, 51: 141–149.
- Andreu, M.; Sola, R.; Sitges–Serra, A. et al. (1993):** Risk factors for spontaneous bacterial peritonitis in cirrhotic patients with ascites. *Gastroenterology*, 104: 1133–1138.
- Angeloni, S.; Nicolini, G.; Merli, M. et al. (2003):** validation of automated blood cell counter for the determination of polymorphonuclear cell count in the ascetic fluid of cirrhotic patients with or without spontaneous bacterial peritonitis. *AM J Gastroenterol*, 98:1844-1848.
- Antonsen, S.; Wiggers, P.; Dalhoj, J.; Blaabjerg, O. (1993):** An enzyme linked immunosorbent assay for plasma lactoferrin Concentrations in 362 healthy, adult blood donors. *Scandinavian Journal of Clinical and Laboratory Investigation*, 53: 133–144.
- Ariza, J.; Xial, X.; Esteve, M. et al. (1991):** Aztreonam versus cefotaxime in the treatment of gram-negative spontaneous bacterial peritonitis in cirrhotic patients. *Hepatology*, 14: 91 – 98.

- Arnold, R.R.; Brewer, M.; Gauthier, J.J. (1980):** Bactericidal activity of human lactoferrin: sensitivity of a variety of microorganisms. *Infection and Immunity*, 28: 893-898.
- Arnold, R.R.; Cole, M.F.; McGhee, J.R. (1977):** A bactericidal effect for human lactoferrin. *Science*, 197: 263 –265.
- Arroyo, V.; Gines, P.; Gerbes, A. L. et al. (1996):** Definition and diagnostic criteria of refractory ascites and hepatorenal syndrome in cirrhosis. *Hepatology*, 23:164-176.
- Arroyo, V.; Planas, R.; Gaya, J. et al. (1983):** Sympathetic nervous activity, rennin- angiotensin system and renal excretion of prostaglandin E₂ in cirrhosis: relationship to functional renal failure and sodium and water excretion. *Eur. J. Clin. Invest.* 13:271-278.
- Ashorn, R.G.; Eskola, J.; Tuohimaa, P.J.; Krohn, K.J. (1986):** Effect of the progesterone inducible proteins human lactoferrin and chicken avidin on lymphocyte proliferation. *Human Reproduction*, 1: 149–151.
- Azoulay, E.; Fartoukh, M.; Galliot, R. et al. (2000):** Rapid diagnosis of infectious pleural effusions by use of reagent strips. *CID*. 31: 914 – 919.
- Bac, D.J. (1996):** Spontaneous bacterial peritonitis: an indication for liver transplantation? *Scand. J. Gastroenterol.* 218: 38 – 42.
- Bac, D.V.; De-Marie, S.; Van Blankenstein, M. (1996):** Spontaneous bacterial peritonitis complicating malignancy related ascites. *Dig. Dis. Sci.* 41 (1): 131–132.

- Baggiolini, M.; De Duve, C.; Masson, P.L.; Heremans, J.F. (1970):** Association of lactoferrin with specific granules in rabbit heterophil leukocytes. *The Journal of Experimental Medicine*, 131: 559–570.
- Baker E.N. (1994):** Structure and reactivity of transferrins. *Advances in Inorganic Chemistry*, 41, 363–489.
- Baker E.N., Baker H.M. (2005):** Molecular structure, binding properties and dynamics of lactoferrin. *Cellular and Molecular Life Sciences*, 62: 2531–2539.
- Baker, M.; Blinder, R. and Rice, R. (1986):** Diagnostic imaging of abdominal fluid collections and abscesses. *Crit. Rev. Diagn. Imaging*, 25 (93): 233 – 278.
- Barton, M.H.; Hurley, D.; Norton, N.; Heusner, G.; Costa, L.; Jones, S.; Byars, D.; Watanabe, K. (2006):** Serum lactoferrin and immunoglobulin G concentrations in healthy or ill neonatal foals and healthy adult horses. *Journal of Veterinary Internal Medicine*, 20: 1457–1462.
- Bataller, R.; Gines, P.; Guevara, M. and Arroyo, V. (1997):** Hepatorenal syndrome. *Semin. Liver Dis.* 17:233-47.
- Bates, D.O.; Lodwick, D. and Williams, B. (1999):** Vascular endothelial growth factor and microvascular permeability. *Microcirculation*, 6: 83 – 96.
- Bauer, TM.; Steinbrückner, B.; Brinkmann, FE.; Ditzen, AK.; Schwacha, H.; Aponte, JJ.; Pelz, K.; Kist, M.; Blum, HE. (2001):** Small intestinal bacterial overgrowth in patients with cirrhosis: prevalence and relation with spontaneous bacterial peritonitis. *Am J Gastroenterol*, 96: 2962-2967.

- Bauer, T.M.; Follo, A.; Navasa, M. et al. (2002):** Daily norfloxacin is more effective than weekly rifloxacin in prevention of spontaneous bacterial peritonitis recurrence. *Dig. Dis. Sci*, 47:1356-1361.
- Baynes, R.; Bezwoda, W.; Bothwell, T.; Khan, Q.; Mansoor, N. (1986):** The non-immune inflammatory response, serial changes in plasma iron, iron binding capacity lactoferrin, ferritin and C reactive protein. *Scandinavian Journal of Clinical and Laboratory Investigation*; 46: 695-704.
- Baynes, R.D. and Bezwoda, W.R. (1994):** Lactoferrin and the inflammatory response. *Advances in Experimental Medicine and Biology*, 357: 133–141.
- Beal, A. and Cerra, F. (1994):** Multiple organ failure syndrome in the 1990 systemic inflammatory response and organ dysfunction. *JAMA*. 13:271.
- Bellamy, W.; Takase, M.; Yamauchi, K.; Wakabayashi, H.; Kawase, k.; Tomita, M. (1992):** Identification of the bactericidal domain of lactoferrin. *Biochimica et Biophysica Acta*, 1121: 130–136.
- Bennett, R.M. and Mohla, C. (1976):** A solid phase radioimmunoassay for the measurement of lactoferrin in human plasma: variations with age, sex and diseases. *Journal of Laboratory and Clinical Medicine*, 88: 156–166.
- Berg, RD. and Garlington, AW. (1979):** Translocation of certain indigenous bacteria from the gastrointestinal tract to the mesenteric lymph nodes and other organs in a gnotobiotic mouse model. *Infect Immun*, 23: 403-411.
- Berlov, M.N.; Korableva, E.S.; Andreeva, Y.V.; Ovchinnikova, T.V.,**

- Korkyakov, V.N. (2007):** Lactoferrin from canine neutrophils: isolation and physicochemical and antimicrobial properties. *Biochemistry (Moscow)*, 72: 445-451.
- Bernard, B.; Cadranel, J.F.; Valla, D. et al. (1995):** Prognostic significance of bacterial infection in bleeding cirrhotic patients a prospective study. *Gastroenterology*, 108: 1828.
- Bernard, B.; Grange, J.D.; Khac, E.N. et al. (1999):** Antibiotic prophylaxis for the prevention of bacterial infections in cirrhotic patients with gastrointestinal bleeding: a meta-analysis. *Hepatology*, 29:1655-1661.
- Bezault, J.; Bhimani, R.; Wiprovnick, J.; Furmanski, P. (1994):** Human lactoferrin inhibits growth of solid tumors and development of experimental metastases in mice. *Cancer Research*, 54: 2310–2312.
- Bezwoda, W.R.; Baynes, R.D.; Khan, Q.; Manssor, N. (1985):** Enzyme linked immunosorbent assay for lactoferrin, Plasma and tissue measurements. *Clinica Chimica Acta International Journal of Clinical Chemistry*, 151: 61–69.
- Birgens H.S. (1985):** Lactoferrin in plasma measured by an ELISA technique: Evidence that plasma lactoferrin is an indicator of neutrophil turnover and bone marrow activity in acute leukaemia. *Scandinavian Journal of Haematology*, 34: 326–331.
- Bobadilla, M.; Sifuentes, J. and Garcia-Tsao, G. (1989):** Improved method for bacteriological diagnosis of spontaneous bacterial peritonitis. *J. Clin. Microbiol.* 27: 2145 – 2147.

- Boermester, M.; Vanleeuwen, P. and Coyle, S. (1995):** Interleukin-I blockade attenuates mediator release and dys-regulation of the hemostatic mechanisms during sepsis. *Arch. Surg.* 130:739-748.
- Bone, R.C. (1991):** The pathogenesis of sepsis. *Ann. Intern. Med.* 115: 347 – 469.
- Boxer, L.A.; Coates, T.D.; Haak, R.A.; Wolach, J.B.; Hoffstein, S., Baehner R.L. (1982):** Lactoferrin deficiency associated with altered granulocyte function. *The New England Journal of Medicine*: 307, 404–410.
- Brines, R.D and Brock, J.H. (1983):** The effect of trypsin and chymotrypsin on the in vitro antibacterial and ironbinding proprieties of lactoferrin in human milk and bovine colostrum. Unusual resistance of human apolactoferrin to proteolytic digestion. *Biochimica et Biophysica Acta*, 759: 229–235.
- Brock, J.H. (1980):** Lactoferrin in human milk: its role in iron absorption and protection against enteric infection in the newborn infant. *Archives of Disease in Childhood*, 55: 417–421.
- Brock, J.H. (2002):** The physiology of lactoferrin. *Biochemistry and Cell Biology*, 80: 1–6.
- Brock, J.H.; Arzabe, F.; Lampreave, F.; Pineiro, A. (1976):** The effect of trypsin on bovine transferrin and lactoferrin. *Biochimica et Biophysica Acta*, 446: 214–225.
- Brown, R.D.; Rickard, K.A.; Kronenberg, H. (1983):** Immunoradiometric assay of plasma lactoferrin. *Pathology*, 15: 27–31.
- Broxmeyer, H.E.; Gentile, P.; Bognacki, J.; Ralph, P. (1983):** Lactoferrin, transferrin and acidic isoferritins: regulatory molecules with potential therapeutic value in leukemia. *Blood Cells*, 9: 83–105.

- Butani, R.C.; Shaffer, R.T.; Szykowski, R.D. et al. (2000):** Use of multistix leucocyte estrase dipstick testing for ascitic fluid infection. *Gastroenterology*, 118: 1098 A (abstract).
- Butani, RC.; Shffer, RT.; Szykowski, RD. et al. (2004):** Rapid diagnosis of infected ascitic fluid using leucocyte estrase dipstick testing. *Am J Gastroenterol*, 99:532-537.
- Byl, B.; Roucloux, I.; Crusiaux, A. et al. (1993):** Tumour necrosis factor and interleukin-6 plasma levels in infected cirrhotic patients. *Gastroenterology*, 104: 1492 – 1497.
- Cabrera, J.; Arroyo, V.; Ballesta, A.M. et al. (1982):** Aminoglycoside nephrotoxicity in cirrhosis: value of urinary B2-microglobulin to discriminate functional renal failure from acute tubular damage. *Gastroenterology*, 82: 97 -105.
- Cadranel, J.P.; Denis, J.; Pauwels, A. et al. (1999):** Prevalence and risk factors of bacteruria in cirrhotic patients: A prospective case – control multicenter study in 244 patients. *J. Hepatol.* 31 : 464.
- Caldwell, S.H. and Battle, E.H. (1999):** Ascites and spontaneous bacterial peritonitis. In: Schiff's disease of the liver. Schiff ER, Sorrell MF and Maddrey, WG (Eds.), 8th edition, Lippincott Williams and Wilkins, Wolters Kluwer, Company, Philadelphia, P: 503 – 544.
- Caly, W.R. and Strauss, E. (1993):** A prospective study of bacterial infection in patients with cirrhosis. *J. Hepatol.* 18:353-358.
- Campoli-Richards, D.M.; Monk, J.P.; Price, A. et al. (1988):** Ciprofloxacin: a review of its antibacterial activity, pharmacokinetic properties and therapeutic use. *Drugs*, 3: 373 – 447.
- Cardenas, A.; Uriz, J.; Gines, P. and Arroyo, V. (2000):** Hepatorenal syndrome. *Liver transplantation*, 6 (suppl-1): S 63- S 71.

- Cardenas, A. and Gines, P. (2005):** Management of complications of cirrhosis in patients awaiting liver transplantation. *J. Hepatol.* 42:124-133.
- Casafont, F.; Sanchez, E.; Martin, L. et al. (1997):** Influence of malnutrition on the prevalence of bacterial translocation and spontaneous bacterial peritonitis in experimental cirrhosis in rats. *Hepatology*, 25: 1334 – 1337.
- Castellote, J. (2005):** Can leukocyte esterase reagent strips be used for the diagnosis of spontaneous bacterial peritonitis? *Nat Clin Pract Gastroenterol Hepatol* ,2: 566-567.
- Castellote, J.; Dopez,C.; Gornals, J. et al. (2003) :** Rapid Diagnosis of Spontaneous Bacterial Peritonitis by use of Reagent strips. *Hepatology*, 37; 4; 893 – 896.
- Castellote, J.; Xiol, X.; Verdaguer, R. et al. (1990):** Comparison of two ascitic fluid culture methods in cirrhotic patients with spontaneous bacterial peritonitis. *Am. J. Gastroenterol.* 85 : 1605 – 1608.
- Cejudo-Martin, P.; Ros, J.; Navasa, M. et al. (2001):** Increased production of vascular endothelial growth factor in peritoneal macrophages of cirrhotic patients with spontaneous bacterial peritonitis. *Hepatology*, 34(3): 487 – 493.
- Cereto, F.; Molina, I.; González, A.; Del Valle, O.; Esteban, R.; Guardia, J.; Genescà, J. (2003):** Role of immunosuppression in the development of quinolone-resistant *Escherichia coli* spontaneous bacterial peritonitis and in the mortality of *E. coli* spontaneous bacterial peritonitis. *Aliment Pharmacol*, 17: 695-701.

- Chang, C.S.; Chen, G.H.; Lien, H.C. et al. (1998):** Small intestinal dysmotility and bacterial overgrowth in cirrhotic patients with spontaneous bacterial peritonitis. *Hepatology*, 28 (5): 1187 – 1190.
- Cheng, J.B.; Wang, J.Q.; Bu, D.P.; Liu, G.L.; Zhang, C.G; Wei, H.Y; Zhou, L.Y.; Wang, J.Z. (2008):** Factors affecting the lactoferrin concentration in bovine milk. *Journal of Dairy Science*, 91: 970–976.
- Chinnock, B. and Hendey (2007):** Can clear ascitic fluid appearance rule out spontaneous bacterial peritonitis. *Am. J. Emerg. Med.* 25: 934-937.
- Christou, L.; Pappas, P.; and Falagas, M.S. E. (2007):** Bacterial infection related morbidity and mortality in cirrhosis. *Am. J gastroenterol*, 102:1510-1517.
- Chung, S.; Hayward, C.; Brock, D.J.H.; Van Heyningen, V.(1985):** A monoclonal antibody based immunoassay for human lactoferrin. *Journal of Immunological Methods*, 84: 135–141.
- Cirera, J.; Bauer, T. M.; Navasa, M. et al. (2001):** Bacterial translocation of enteric organisms in patients with cirrhosis. *J. Hepatol.* 34: 32-37.
- Cirioni, O.; Giacometti, A.; Barchiesi, F.; Scalise, G. (2000):** Inhibition of growth of *Pneumocystis carinii* by lactoferrins alone and in combination with pyrimethamine, clarithromycin and minocycline. *The Journal of Antimicrobial Chemotherapy*, 46: 577–582.
- Cornish, J.; Callon, K.E.; Naot, D.; Palmano, K.P.; Banovic, T.; Bava, U.; Watson, M.; Lin, J.M.; Tong, P.C.; Chen, Q.; Chan, V.A.; Reid, H.E.; Fazzalari, N.; Baker, H.M.; Baker, E.N.; Haggarty, N.W.; Grey, A.B.; Reid, I.R. (2004):** Lactoferrin is a potent regulator of bone cell activity and increases bone formation in vivo. *Endocrinology*, 145: 4366-4374.

- Crosa, J.H. (1989):** Genetics and molecular biology of siderophore-mediated iron transport in bacteria. *Microbiological Reviews*, 53: 517–530.
- Damiens, E.; El Yazidi, I.; Mazurier, J.; Duthille, I.; Spik, G.; Boilly-Marer, Y. (1999):** Lactoferrin inhibits G1 cyclin dependent kinases during growth arrest of human breast carcinoma cells. *Journal of Cellular Biochemistry*, 74: 486–498.
- Damiens E., Mazurier J., El Yazidi I., Masson M., Duthille I., Spik G., Boilly-Marer Y. (1998):** Effects of human lactoferrin on NK cell cytotoxicity against haematopoietic and epithelial tumour cells. *Biochimica et Biophysica Acta*, 1402, 277–287.
- Davidsson, L.; Kastenmayer, P.; Yuen, M.; Lonnerdal, B.; Hurrell, R.F. (1994):** Influence of lactoferrin on iron absorption from human milk in infants. *Pediatric Research*, 35: 117–124.
- De Vet, B.J.; Van Gool, J. (1974):** Lactoferrin and iron absorption in the small intestine. *Acta Medica Scandinavica*, 196: 393–402.
- Devi A.S., Das M.R., Pandit M.W. (1994):** Lactoferrin contains structural motifs of ribonuclease. *Biochimica et Biophysica Acta*, 1205: 275–281.
- Edell, S. and Geftter, W. (1979):** Ultrasonic differentiation of types of ascitic fluid. *A.J.R.* 133: 111-114.
- Ekins, A.; Khan, A.G.; Shouldice, S.R.; Schryvers, A.B. (2004):** Lactoferrin receptors in gram-negative bacteria: insights into the iron acquisition process. *Biometals*, 17: 235–243.
- Ersöz, G.; Aydin, A.; Erdem, S.; Yüksel, D.; Akarca, U.; Kumanlioglu, K. (1999):** Intestinal permeability in liver cirrhosis. *Eur J Gastroenterol Hepatol*, 11: 409-412.

- Esaguy, N.; Aguas, A.P.; Vilanova, M.; Silva, M.T. (1991):** Activation of human neutrophils by phorbol ester decreases the cytoplasm compactness and the lactoferrin content of the granulocytes. *Journal of Leukocyte Biology*, 50: 444–452.
- Fairweather Tait, S.J.; Balmer, S.E.; Scott, P.H.; Minski, M.J. (1987):** Lactoferrin and iron absorption in newborn infants. *Pediatric Research*, 22, 651–654.
- Felisart, J.; Rimola, A.; Arroyo, V. et al. (1985):** Cefotaxime is more effective than ampicillin – tobramycin in cirrhotics with severe infections. *Hepatology*, 5: 457 – 462.
- Fernández, J.; Bauer, TM.; Navasa, M.; Rodés, J. (2002):** Diagnosis, treatment and prevention of spontaneous bacterial peritonitis. *Baillieres Best Pract Res Clin Gastroenterol*; 14: 975-990
- Fernandez, J.; Navasa, M.; Planas, R. et al. (2007):** Primary prophylaxis of spontaneous bacterial peritonitis delays hepatorenal syndrome and improves survival in cirrhosis. *Gastroenterology*, 133:818-24.
- Fiuza, C.; Salcedo, M.; Clemente, G.; Tellado, JM. (2002):** Granulocyte colony-stimulating factor improves deficient in vitro neutrophil transendothelial migration in patients with advanced liver disease. *Clin Diagn Lab Immunol*, 9: 433-439.
- Follo, A.; Liovet, J. M.; Navasa, M. et al. (1994):** Renal impairment following spontaneous bacterial peritonitis in cirrhosis. Incidence, clinical course, predictive factors and prognosis. *Hepatology*, 20: 1495-1501.
- Fong, T.L.; Akriviadis, E.A.; Runyon, B.A. and Reynolds, T.B. (1989):** Polymorphonuclear cell count response and duration of antibiotic therapy in spontaneous bacterial peritonitis. *Hepatology*, 9:423– 426.

- Fujihara, T. and Hayashi K. (1995):** Lactoferrin inhibits herpes simplex virus type-1 (HSV-1) infection to mouse cornea. Archives of Virology, 140: 1469–1472.
- Fujita K., Matsuda E., Sekine K., Iigo M., Tsuda H. (2004):** Lactoferrin enhances Fas expression and apoptosis in the colon mucosa of azoxymethane-treated rats. Carcinogenesis, 25, 1961–1966.
- Furmanski, P. Li. Z.P.; Fortuna, M.B.; Swamy, C.V.B.; Das, M.R. (1989):** Multiple molecular forms of human lactoferrin. Identification of a class of lactoferrins that possess ribonuclease activity and lack iron-binding capacity. The Journal of Experimental Medicine, 170: 415- 429.
- Garcia- Tsao, G. (2005):** Bacterial infection in cirrhosis: treatment and prophylaxis. J. Hepatology, 42:S85-S92.
- Garcia- Tsao, G. (2007):** Prevention and treatment of infections in patients with cirrhosis. Clin. Gastroenterology, 21:77-93.
- Garcia- Taso, G. (1999):** Identifying new risk factors for spontaneous bacterial peritonitis: how important is it? Gastroenterology, 117:495-499.
- Garcia-Tsao, G. (1992):** Spontaneous bacterial peritonitis. Gastroenterol. Clin. N.Am. 21:257-275.
- Garcia-Tsao, G. (2004):** Treatment of acute variceal bleeding: general management and prevention of infections. In: Portal hypertension in the 21st century. Groszmann RJ, Bosch J. editors. Dordecht: Kluwer Academic. P: 233-240.
- Garrison, R.N.; Cryer, H.M.; Howard, D.A.; Polk, H.C. (1984):** Clarification of risk factors for abdominal operations in patients with hepatic cirrhosis. Ann. Surg. 199:648-655.

- Gines, P.; Rimola, A.; Plana, S.R. et al. (1990):** Norfloxacin prevents spontaneous bacterial peritonitis recurrence in cirrhosis: results of a double blind placebo-controlled trial. *Hepatology*, 12: 716 – 724.
- Giron – Gonzalez, J.A.; Rodriguez – Ramos, C.; El Vira, J. et al. (2001):** Serial analysis of serum and ascitic fluid levels of soluble adhesion molecules and chemokines in patients with spontaneous bacterial peritonitis. *Clin. Exp. Immunol.* 123 (1): 56 - 61.
- Gitlin, N. (1990):** Ascitic fluid acidosis in bacterial peritonitis: A poor prognostic sign? *Hepatol.* 11: 896 – 897.
- Gomez - Jimenez, J.; Ribera, E.; Gasser, J. et al. (1993):** Randomized trial comparing ceftriaxone with cefonicid for treatment of spontaneous bacterial peritonitis in cirrhotic patients. *Antimicrobial Agents chemother.* 37: 1587 – 1592.
- Goshal, U.C.; Poddar, U.; Yachha, S.K. (1995):** Spontaneous bacterial peritonitis in a child with abdominal lymphoma. *Acta-oncol*,4: 531-534.
- Goulis, J.; Patch, D.; Burroughs, AK. (1999):** Bacterial infection in the pathogenesis of variceal bleeding. *Lancet*, 353: 139-142.
- Goulis, J.; Armonis, A.; Patch, D. et al. (1998):** Bacterial infection is independently associated with failure to control bleeding in cirrhotic patients with gastrointestinal hemorrhage. *Hepatology*, 27:1207-1212.
- Grange, J.D.; Amiot, X.; Grange, V. et al. (1990):** Amoxicillin – Clavulanic Acid therapy of Spontaneous bacterial peritonitis a prospective study of twenty –seven cases in cirrhotic patients. *Hepatol. J.* 11: 360 – 364.

- Green, M.R. and Pastewka, J.V. (1978):** Lactoferrin is a marker for prolactin response in mouse mammary explants .Endocrinology, 103: 1510–1513.
- Groves M.L. (1960):** The isolation of a red protein from milk. Journal of the American Chemical Society, 82, 3345-3350
- Guarner, C.; Runyon, BA.; Young, S.; Heck, M.; Sheikh, MY. (1997):** Intestinal bacterial overgrowth and bacterial translocation in cirrhotic rats with ascites. J Hepatol, 26: 1372-1378.
- Guarner, C. and Runyon, B.A. (1995):** Spontaneous bacterial peritonitis: Pathogenesis, diagnosis and management. Gastroenterology, 3 (4): 311–328.
- Guarner, C. and Runyon, B.A. (1996):** Ascites In: GI/ liver secretes. 1st ed., McNally P.R. (editor): Philadelphia: Hanley and Belfus, P: 202 – 209.
- Guarner, C.; Sola, R.; Soriano, G. et al. (1999):** Risk of the first community –acquired spontaneous bacterial peritonitis in cirrhotics with low ascitic fluid protein. Gastroenterology, 117: 414 - 419.
- Hagiwara S., Kawai K., Anri A., Nagahata H. (2003):** Lactoferrin concentrations in milk from normal and subclinical mastitic cows The Journal of Veterinary Medical Science, 65, 319–323.
- Hagiwara T., Shinoda I., Fukuwatari Y., Shimamura S :(1995)** .Effect of lactoferrin and its peptides on proliferation of rat intestinal epithelial cell line, IEC-18 ‘in the presence of epidermal growth factor. Bioscience ‘Biotechnology, and Biochemistry, 59, 1875–1881.
- Hallak, A. (1989):** Spontaneous bacterial peritonitis: Clinical review. Am. J. Gastroenterol. 84: 345 -350.

- Hallak, A. (1989):** Spontaneous bacterial peritonitis: Clinical review. *Am. J. Gastroenterol.* 84: 345 -350.
- Hansen, N.E.; Malmquist, J.; Thorell, J. (1975):** Plasma myeloperoxidase and lactoferrin measured by radioimmunoassay: relations to neutrophil kinetics. *Acta Medica Scandinavica*, 198: 437–443.
- Harmsen, M.C.; Swart, P.J.; De Bethune, M.P.; Pauwels, R.; De Clercq, E.; The, T.H.; Meijer, D.K.F. (1995):** Antiviral effects of plasma and milk proteins: lactoferrin shows potent activity against both human immunodeficiency virus and human cytomegalovirus replication in vitro. *The Journal of Infectious Diseases*, 172: 380-388.
- Haversen, L.; Ohlsson, B.G.; Hahn-Zoric, M.; Hanson, L.A.; Mattsby-Baltzer, I. (2002):** Lactoferrin down-regulates the LPS-induced cytokine production in monocytic cells via NF-kappa B. *Cellular Immunology*, 220: 83-95.
- He, J. and Furmanski, P. (1995):** Sequence specificity and transcriptional activation in the binding of lactoferrin to DNA. *Nature*, 373, 721–724.
- Hutchens T.W., Henry J.F., Yip T.T., Hachey D.L., Schanler R.J., Motil K.J., Garza C. (1991):** Origin of intact lactoferrin and its DNA-binding fragments found in the urine of human milk-fed preterm infants. Evaluation by stable isotopic enrichment. *Pediatric Research*, 29: 243-250.
- Hellinger, W.C.; Cawley, J.J.; Alvarez, S. et al. (1995):** Clinical comparison of the isolator and Bact/ALERT aerobic blood culture systems. *J. Clin. Microbiol.* 33: 1787 – 1790.

- Hillebrand, D.J. (2002):** Spontaneous bacterial peritonitis. *Curr. Treat. Operations Gastroenterol.* Dec. 5 (6): 479 – 489.
- Hillebrand, D.J. and Runyon, B.A. (1998):** Ascites and spontaneous bacterial peritonitis. In: *Handbook of liver disease*. Friedman LS and Keefe EB (Eds.). Churchill Livingstone, London, Hong Kong, P: 151 – 166.
- Hiscoke, C.; Yoxall, H.; Greig, D. and Lighfoot, N.F. (1990):** Validation of a method for the rapid diagnosis of urinary tract infection suitable for use in general practice. *Br. J. Gen. Pract.* 40: 403 - 405.
- Hoefs, J.C. (1990(a)):** Diagnostic paracentesis: A potential clinical tool. *Gastroenterology*, 98: 230 – 236.
- Hoefs, J.C. (1990(b)):** Spontaneous bacterial peritonitis: prevention and therapy. *Hepatology*, 12: 776 – 781.
- Hoefs, J.C. and Runyon, B.A. (1985):** Spontaneous bacterial peritonitis. *Dis. Mon.* 31: 1-48.
- Hoefs, J.C.; Canawati, H.N.; Sapico, F.I. et al. (1982):** Spontaneous bacterial peritonitis. *Hepatology*, 2: 399-407.
- Horn, S.; Holzer, H. and Hornia, J.H. (1996):** Spontaneous bacterial peritonitis in a patient with nephrogenic ascites during an episode of acute renal transplant rejection. *Am. J. Kidney Dis.* 27: (3): 441–443.
- Huang, Y.S.; Chan, C.Y.; Wu, J.C. et al. (1996):** Serum levels of interleukin-8 in alcoholic liver disease: relationship with disease study, biochemical parameters and survival. *J. Hepatol.* 24: 377 – 384.
- Hutchens, T.W.; Henry, J.F.; Yip, T.T.; Hachey, D.L.; Schanler, R.J.; Motil, K.J.; Garza, C. (1991):** Origin of intact lactoferrin and its

- DNA-binding fragments found in the urine of human milk-fed preterm infants. Evaluation by stable isotopic enrichment. *Pediatric Research*, 29: 243-250.
- Inadomi, J. and Sonnenberg, A. (1997):** Cost-analysis of prophylactic antibiotics in spontaneous bacterial peritonitis. *Gastroenterology*, 113: 1289 – 1294.
- Iyer, S.; Lonnerdal, B. (1993):** Lactoferrin, lactoferrin receptors and iron metabolism. *European Journal of Clinical Nutrition*, 47: 232–241.
- Jameson, G.B.; Anderson, B.F.; Norriss, G.E.; Thomas, D.H.; Baker, E.N.(1998):** Structure of human apolactoferrin at 2.0 Å resolution. Refinement and analysis of ligand-induced conformational change. *Acta Crystallographica. Section D, Biological Crystallography*, 54: 1319-1335.
- Jansen, P. (1997): Spontaneous bacterial peritonitis. Detection, treatment and prophylaxis in patients with liver cirrhosis.** *Neth. J. Med.* 51: 123-128.
- Javid, G.; Khan, B.A.; Shah, A.H. et al. (1998):** Short-course ceftriaxone therapy in spontaneous bacterial peritonitis. *Post graduate Medical Journal*, Oct. 74 (876): 592 -595.
- Johanson B. (1960):** Isolation of an iron containing red protein from human milk. *Acta Chemica Scandinavica*, 14, 510-512.
- Kanyshkova, T.G.; Buneva, V.N.; Nevinsky, G.A. :(2001):** Lactoferrin and its biological functions. *Biochemistry (Moscow)*, 66: 1–7.
- Kelver, M.E.;Kaul, A.; Nowicki, B.; Findley, W.E.; Hutchens, T.W.; Nagamami, M.(1996):** Estrogen regulation of lactoferrin in human endometrium. *American Journal of Reproductive Immunology*, 36: 234–247.

- Kikuchi, M.; Mizoroki, S.; Kubo, T.; Ohiwa, Y.; Kubota, M.; Yamada, N.; Orino, K.; Ohnami, Y.; Watanabe, K. (2003):** Seminal plasma lactoferrin but not transferrin reflects gonadal function in dogs. The Journal of Veterinary Medical Science, 65: 679–684.
- Kirkpatrick, C.H.; Green, I.; Rich, R.R.; Schade, A.L. (1971):** Inhibition of growth of *Candida albicans* by ironunsaturated lactoferrin: relation to host-defense mechanisms in chronic mucocutaneous candidiasis. The Journal of Infectious Diseases, 124: 539–544.
- Klipes, V.E.; Krahauer, H. and Wren, R.E. (1993):** An analysis of liver transplant experience from 37 transplant centers as reported to Medicare. Transplantation, 56: 554 – 561.
- Knapp, R.G., Miller M.C., (1992):** Clinical epidemiology, biostatistics:NMS " National medical series from Williams, wilins. Chapters 1,3,4,13.pages 6-8, 36-39, 55-56, 209-211.
- Konuspayeva, G.; Faye, B.; Loiseau, G.; Levieux, D. (2007):** lactoferrin and immunoglobulin contents in camel's milk (*Camelus bactrianus*, *Camelus dromedarius*, and hybrids) from Kazakhstan. Journal of Dairy Science, 90: 38-46.
- Kutter, D.; Figueiredo, G. and Klemmer, L. (1987):** Chemical detection of leucocytes in urine by means of a new multiple test strip. J. Clin. Chem. Clin. Biochem., 25 : 91 – 94.
- Laffi, G.; Lavilla, G. and Gentilini, P. (1994):** Pathogenesis and management of the hepatorenal syndrome. Sem. Liver Dis. 14: 71-81.

- Lamb, H.M.; Ormrod, D.; Scott, L.J. and Figgitt, D.P. (2002):** Ceftriaxone: an update of its use in the management of community – acquired and nosocomial infections. *Drugs*, 62 (7): 1041 – 1089.
- Le Moine, O.; Deviere, J.; Devaster, J.M. et al. (1994):** Interleukin-6: an early marker of bacterial infection in decompensated cirrhosis. *J. Hepatol.* 20: 819 – 824.
- Leffell, MS.; Spitznagel, JK. (1975):** Fate of human Lactoferrin and myeloperoxidase in phagocytizing human neutrophils: effects of immunoglobulin G subclasses and immune complexes coated on latex beads. *Infect immune*; 12:813-820.
- Legrand, D.; Ellass, E.; Carpentier, M.; Mazurier, J. (2005):** Lactoferrin: a modulator of immune and inflammatory responses. *Cellular and Molecular Life Sciences*, 62: 2549-2559.
- Leitch, E.C. and Willcox, M.D. (1998):** Synergic antistaphylococcal properties of lactoferrin and lysozyme. *Journal of Medical Microbiology*, 47: 837–842.
- Levay, P.F. and Viljoen, M. (1995):** Lactoferrin: a general review. *Haematologica*, 80: 252–267.
- Levy, M.J. and Dipalma, J.A. (1996):** New options for spontaneous bacterial prophylaxis: costly or cost effective? *Am. J. Gastroenterol.* 91: 1050 – 1051.
- Liovet, J.; Planas, R.; Morillas, R. et al. (1993):** Short term prognosis of cirrhotics with spontaneous bacterial peritonitis: multivariate study. *Am. J. Gastroenterology*, 88 (3): 388–392.
- Liovet, J., Moitinho, E.; Sala, M. et al. (2000):** Prevalence and prognostic value of hepatocellular carcinoma in cirrhotic patients presenting with spontaneous bacterial peritonitis. *J. Hepatology*, 33: 423-29.

- Lonnerdal, B. and Iyer, S. (1995):** Lactoferrin: molecular structure and biological function. *Annual Review of Nutrition*, 15:93–110.
- Lorget, F.; Clough, J.; Oliveira, M.; Daury, M.C.; Sabokbar, A.; Offord, E. (2002):** Lactoferrin reduces in vitro osteoclast differentiation and resorbing activity. *Biochemical and Biophysical Research Communications*, 296: 261-266.
- Machnicki, M.; Zimecki, M.; Zagulski, T. (1993):** Lactoferrin regulates the release of tumour necrosis factor alpha and interleukin 6 in vivo. *International Journal of Experimental Pathology*, 74: 433–439.
- Marchetti, M.; Longhi, C.; Conte, M.P.; Pisani, S.; Valenti, P.; Seganti, L. (1996):** Lactoferrin inhibits herpes simplex virus type 1 adsorption to Vero cells. *Antiviral Research*, 29: 221–231.
- Martinez-Bru, C.; Gomez, C.; Cortes, M. et al. (1999):** Ascitic fluid interleukin -8 to distinguish spontaneous bacterial peritonitis and sterile ascites in cirrhotic patients. *Clin. Chem.* 45 (11): 2027 – 2028.
- Martins, CA.; Fonteles, MG.; Barrett, LJ. et al. (1995):** Correlation of Lactoferrin with neutrophilic inflammation in body fluids. *Clin Diagn Lab immunol*, 2:763-765.
- Masson, P.L.; Heremans, J.F.; Dive, C. (1966):** An ironbinding protein common to many external secretions. *Clinica Chimica Acta*, 14: 735–739.
- Masson, P.L.; Heremans, J.F.; Ferin, J. (1968):** Presence of an iron-binding protein (lactoferrin) in the genital tract of the human female. Its immunohistochemical localization in the endometrium. *Fertility and Sterility*, 19: 679-689.

- Masson, P.L.; Heremans, J.F. (1971):** Lactoferrin in milk from different species. *Comparative Biochemistry and Physiology B, Comparative Biochemistry*, 39: 119-129.
- Masson, P.L.; Heremans, J.F.; Schonke, E. (1969):** Lactoferrin, an iron-binding protein in neutrophilic leukocytes. *The Journal of Experimental Medicine*, 130: 643-658.
- Mazurier, J. and Spik, G. (1980):** Comparative study of the iron-binding properties of human transferrins. I. Complete and sequential iron saturation and desaturation of the lactotransferrin. *Biochimica et Biophysica Acta* , 629: 399-408.
- McCormick, P.A.; Greenslade, L.; Kibbler, C.C. et al. (1997):** A prospective randomized trial of ceftazidime versus netilmicin plus mezlocillin in the empirical therapy of presumed sepsis in cirrhotic patients. *Hepatology*, 25: 833 – 836.
- McHutchinson, J.G. and Runyon, B.A. (1994):** Spontaneous bacterial peritonitis. In: *Gastrointestinal and hepatic infections*. Surawicz, CM; Owen, RL (eds). WB Saunders, Philadelphia, P: 455-475.
- McHutchinson, J.G. and Runyon, B.A. (1995):** Spontaneous bacterial peritonitis In: *Gastrointestinal and Hepatic infections*. Surawicz,CM, Owen RL (eds). WB Saunders, Philadelphia, P: 455.
- Mesquita, M.A.; Balbino, E.P.S. ; Albuquerque, R.S. et al. (1997):** Ceftriaxone in the treatment of spontaneous bacterial peritonitis ascitic fluid polymorpho-nuclear count response and short-term prognosis. *Hepato-Gastroenterology*, 44: 1276 – 1280.

- Mesquite, R.; Leite – Mor, M. and Parise, E. (1997):** Fibronectin in the ascitic fluid of cirrhotic patients: correlation with biochemical risk factors for the development of spontaneous bacterial peritonitis. *Braz. J. Med. Biol. Res.* 30: 843 – 847.
- Metz-Boutique, M.H.; Jolles, J.; Mazurier, J.; Schoentgen, F.; Legrand, D.; Spik, G.; Montreuil J.; Jolles, P. (1984):** Human lactotransferrin: amino acid sequence and structural comparisons with other transferrins. *European Journal of Biochemistry*, 145: 659–676.
- Montreuil, J.; Tonnelat, J.; Mullet, S. (1960):** Preparation and properties of lactosiderophilin (lactotransferrin) of human milk. *Biochimica et Biophysica Acta*, 45, 413–421.
- Moore, K.P.; Wendon, J.; Frazer, M. et al. (1992):** Plasma endothelin immunoreactivity in liver disease and the hepatorenal syndrome. *N. Engl. J. Med.* 327: 1774-1778.
- Moosa, A.A.; Quortum, H.A. and Ibrahim, M.D. (1995):** Rapid diagnosis of bacterial meningitis with reagent strips. *Lancet*, 345: 1290 – 1291.
- Navasa, M.; Fernandez, J.; Montoliu, S. et al. (2006):** Randomized, double-blind, placebo-controlled trial evaluating norfloxacin in the primary prophylaxis of spontaneous bacterial peritonitis in cirrhotics with renal impairment, hyponatremia or severe liver failure. *J. Hepatol.* 44 (suppl: 2): 551. [Abstract].
- Navasa, M.; Follo, A.; Filella, X. et al. (1998):** Tumour necrosis factor- α and interleukin -6 in spontaneous bacterial peritonitis in cirrhosis: relationship with the development of renal impairment and mortality. *Hepatology*, 27 (5): 1227 – 1232.

- Navasa, M.; Follo, A.; Liovet, J.M. et al. (1996):** Randomized, comparative study of oral ofloxacin versus intravenous cefotaxime in spontaneous bacterial peritonitis. *Gastroenterology*, 111: 1011-1017.
- Navasa, M.; Rimola, A. and Rodes, J. (1997):** Bacterial infections in liver disease. *Semin liver dis.* 17(4):323-33.
- Navasa, M.; Fernandez, J. and Rodes, J. (1999):** Bacterial infections in liver cirrhosis. *Ital. J. Gastroenterol. Hepatol.* 31:616-25.
- Nelson, K.G.; Takahashi, T.; Bossert, N.L.; Walmer, D.K; McLachlan, J. A. (1991):** Epidermal growth factor replaces estrogen in the stimulation of female genital tract growth and differentiation. *Proceedings of the National Academy of Sciences of the United States of America*, 88: 21–25.
- Niemela, A.; Kulomaa, M.; Vija, P.; Tuohimaa, P.; Saariko-ski, S. (1989):** Lactoferrin in human amniotic fluid .*Human Reproduction*, 4: 99–101.
- Nobre, S. R.; Cabral, J. E.; Gomes, J. J. et al. (2008):** In-hospital mortality in spontaneous bacterial peritonitis: a new predictive model. *Eur. J. Gastroenterol Hepatol.* 20:1176-1181.
- Nousbaum, JB.; Cadranel, JF.; Nahon, P. et al. (2007):** Diagnostic accuracy of the multistix 8 SG reagent strips in diagnosis of spontaneous bacterial peritonitis. *Hepatology* ,45:1275-1281.
- Novella, M.; Sola, R.; Soriano, G. et al. (1997):** Continuous versus inpatient prophylaxis of the first episode of spontaneous bacterial peritonitis with norfloxacin. *Hepatology*, 25: 532–536.

- Oberg, G.; Lindmark, G.; Moberg, L.; Venge, P. (1983):** The peroxidase activity and cellular content of granule proteins in PMN during pregnancy. *British Journal of Haematology*, 55: 701–708.
- Olafson, S. and Blei, A. (1995):** Diagnosis and management of ascites in the age of TIPS. *AJR*. 165: 9 – 15.
- Olafson, T.; Olsson, I.; Venge, P.; Elgefors, B. (1977):** Serum myeloperoxidase and lactoferrin in neutropenia. *Scandinavian Journal of Haematology*, 18: 73–80.
- Omata, Y.; Satake, M.; Maeda, R.; Saito, A.; Shimazaki, K.; Yamauchi, K.; Uzuka, Y.; Tanabe, S.; Sarashina, T.; Mikami, T. (2001):** Reduction of the infectivity of *Toxoplasma gondii* and *Eimeria stiedai* sporozoites by treatment with bovine lactoferricin. *The Journal of Veterinary Medical Science*, 63: 187–190.
- Ortiz, J.; Soriano, G.; Coll, P. et al. (1997):** Early microbiologic diagnosis of spontaneous bacterial peritonitis with Bact/ALERT. *J. Hepatol*. 26 (4): 8.
- Orsi, N. (2004):** The antimicrobial activity of lactoferrin: current status and perspectives. *Biometals*, 17: 189-196.
- Otnaess, A.B.K.; Meberg, A.; Sande, H.A. (1983):** Plasma lactoferrin measured by an enzyme-linked immunosorbent-assay (ELISA). Measurements on adult and infant plasma. *Scandinavian Journal of Haematology*, 31, 235-240.
- Parsi, A.; Saadeh, N.; Zein, N. et al. (2008):** Ascitic fluid lactoferrin for diagnosis of spontaneous bacterial peritonitis. *Gastroenterol*, 135: 803-807.

- Pecquet, S.; Andreumont, A. and Tancrede, C. (1986):** Selective antimicrobial modulation of the intestinal tract by norfloxacin in human volunteer and gonotobiotic mice associated with human focal flora. *Antimicrob. Agent Chemother.* 29 : 1047 – 1052.
- Pelletier, G.; Desur, G.; Ink, O. et al. (1991):** Symptomatic bacterascites : Is it spontaneous bacterial peritonitis ? *Hepatology*, 14: 112 – 115.
- Pelletier, G.; Salmon, D.; Ink, O. et al. (1990):** Culture negative neutrocytic ascites: a less severe variant of spontaneous bacterial peritonitis. *J. Hepatol.* 10: 327 – 331.
- Pentecost, B.T. and Teng, C.T. (1987):** Lactotransferrin is the major estrogen inducible protein in mouse uterine secretions. *The Journal of Biological Chemistry*, 262: 10134-10139.
- Perez-Ruiz, M.; Ros, J.; Morales-Ruiz, M. et al. (1999):** Vascular endothelial growth factor production in peritoneal macrophages of cirrhotic patients: regulation by cytokines and bacterial lipopolysaccharide. *Hepatology*, 29: 1057 – 1063.
- Petschow, B.W.; Talbott, R.D.; Batema, R.P. (1999):** Ability of lactoferrin to promote the growth of *Bifidobacterium* spp. in vitro is independent of receptor binding capacity and iron saturation level. *Journal of Medical Microbiology*, 48: 541–549.
- Pinzello, G.; Simonetti, R.; Camma, C. et al. (1993):** Spontaneous bacterial peritonitis: an update. *Gastroenterology, Int.* 6:54-60.
- Pinzello, G.; Simonetti, R.G.; Craxi, A. et al. (1983):** Spontaneous bacterial peritonitis: a prospective investigation in predominantly non alcoholic cirrhotic patients. *Hepatology*, 3: 545 – 549.

- Pinzello, G.; Virdone, R.; Lojacono, F. et al. (1986):** Is the acidity of ascitic fluid a reliable index making the presumptive diagnosis of spontaneous bacterial peritonitis. *Hepatology*, 3: 545 – 549.
- Propst, T.; Propst, A.; Herold, M. et al. (1993):** Spontaneous bacterial peritonitis is associated with high levels of interleukin -6 and its secondary mediators in ascitic fluid. *Eur. J. Clin. Invest.* 23 (12): 832 – 836.
- Ramachandran, A. and Balasubramanian, KA. (2001):** Intestinal dysfunction in liver cirrhosis: Its role in spontaneous bacterial peritonitis. *J Gastroenterol Hepatol*, 16: 607-612.
- Rasarathnum, B.; Kaye, D.; Jennings, G. et al. (2003):** The effect of selective intestinal decontamination on the hyperdynamic circulatory state in cirrhosis. *Ann. Intern. Med.* 139: 186 – 193.
- Ratledge, C.; Dover, L.G. (2000):** Iron metabolism in pathogenic bacteria. *Annual Review of Microbiology*, 54: 881-941.
- Ricart, E.; Soriano, G.; Novella, M.T. et al. (2000):** Amoxicillin- Clavulanic acid versus cefotaxime in the therapy of bacterial infections in cirrhotic patients. *J. Hepatol.* 32
- Richie E.R., Hilliard J.K., Gilmore R., Gillespie D.J (1987).** Human milk-derived lactoferrin inhibits mitogen and alloantigen induced human lymphocyte proliferation. *Journal of Reproductive Immunology*, 12, 137-148.
- Rimland, D. and Hand, W.L. (1987):** Spontaneous peritonitis: reappraisal. *Am. J. Med. Sci.* 293: 285 – 292.

- Rimola, A. (1991):** Infections in liver disease. In: Oxford textbook of clinical Hepatology, McIntyre, N, Benhamou JP, Bricher J, Rizzetto M, Rodes J. (Eds.), Oxford, Oxford University Press, P : 1272 – 1284.
- Rimola, A.; Garcia-Tsao, G.; Navasa, M. et al. (2000):** Diagnosis treatment and prophylaxis of spontaneous bacterial peritonitis: a consensus document. J. Hepatology, 32: 142–153.
- Rimola, A.; Salmeron, J.M.; Clemente, G. et al. (1995):** Two different dosages of cefotaxime in the treatment of spontaneous bacterial peritonitis in cirrhosis: result of a prospective, randomized, multicenter study. Hepatology, 21: 674 – 679.
- Rimola, A.; Tito, L.; Liach, J. et al. (1992):** Efficacy of ceftizoxime in the treatment of severe bacterial infections in patients with cirrhosis. Drug. Invest. 4: 35– 37.
- Rodriguez-Ramos, C.; Galan, F.; Diaz, F. et al. (2001):** Expression of proinflammatory cytokines and their inhibitors during the course of spontaneous bacterial peritonitis. Digestive Disease and Sciences, Vol. 46: NO. 8: 1668 – 1676.
- Rolachon, A.; Cordier, L.; Bacq, Y. et al. (1995):** Ciprofloxacin and long-term prevention of spontaneous bacterial peritonitis. Results of aprospective controlled trial. Hepatology, 22: 1171-1174.
- Rossi, P.; Giansanti, F.; Boffi, A.; Ajello, M.; Valenti, P.; Chiancone, E.; Antonini, G. (2002):** Ca²⁺ binding to bovine lactoferrin enhances protein stability and influences the release of bacterial lipopolysaccharide. Biochemistry and Cell Biology, 80: 41–48.

- Ruiz- Del-Arbol, L.; Urman, J.; Fernandez, J. et al. (2003):** Systemic, renal and hepatic haemodynamic derangement in cirrhotic patients with spontaneous bacterial peritonitis. *Hepatology*, 38: 1210 – 1218.
- Rumke, P.; Visser, D.; Kwa, H.G.; Hart, A.A. (1971):** Radio-immuno assay of lactoferrin in blood plasma of breast cancer patients, lactating and normal women; prevention of false high levels caused by leakage from neutrophile leucocytes in vitro. *Folia Medica Neerlandica*, 14: 156-168.
- Runyon, B.A. (1986(a)):** Elevated ascitic fluid fibronectin concentration. A non specific finding. *Hepatology*, 3: 219 – 222.
- Runyon, B.A. (1986(b)):** Low-protein concentration ascitic fluid is predisposed to spontaneous bacterial peritonitis. *Gastroenterology*, 91: 1343 -1346.
- Runyon, B.A. (1988):** Patients with deficient ascitic fluid opsonic activity are predisposed to spontaneous bacterial peritonitis. *Hepatology*, 8 (3): 632 – 635.
- Runyon, B.A. (1990):** Monomicrobial non- neutrocytic bacterascites: Avariant of spontaneous bacterial peritonitis. *Hepatology*, 12: 710-715.
- Runyon, B.A. (1998):** Ascites and spontaneous bacterial peritonitis. In *Gastrointestinal and liver disease. Pathophysiology. Diagnosis / Management*, Feldman M, Friedman, LS, Sleisenger MH (Eds), 6th edition, Philadelphia. P: 1310 –1313.
- Runyon, B.A. (1999):** Albumin infusion for spontaneous bacterial peritonitis. *Lancet*, 354: 1838.

- Runyon, B.A. (2002):** Ascites and spontaneous bacterial peritonitis. In: Gastrointestinal and liver disease: pathophysiology, diagnosis, Management, Feldman M. Friedman L. and Sleisenger M. (eds), Philadelphia, London, Sydney, Toronto, P:1517.
- Runyon, B.A. (2003):** Strips and tubes: Refining the diagnosis of spontaneous bacterial peritonitis. *Hepatology*, 37: 745 -747.
- Runyon, B.A. (2004):** Early events in spontaneous bacterial peritonitis. *Gut*, 53: 782–784.
- Runyon, B.A. and Antillon, M.R. (1991):** Ascitic fluid PH and lactate: Insensitive and non specific tests in detecting ascitic fluid infection. *Hepatology*, 13: 929.
- Runyon, B.A. and Hoefs, J.C. (1984):** Culture– negative neutrocytic ascites: A variant of spontaneous bacterial peritonitis. *Hepatology*, 4: 1209 – 1211.
- Runyon, B.A. and Hoefs, T.C. (1985):** Ascitic fluid analysis before, during and after spontaneous bacterial peritonitis. *Hepatology*, 5: 257.
- Runyon, B.A.; Antillon, M.R. and Hutchinson, N. (1992):** Diuresis increases ascitic fluid opsonic activity in patients who survive spontaneous bacterial peritonitis. *J. Hepatol.* 14 (2-3), 249–252.
- Runyon, B.A.; Hoefs, J.C. and Canawati, H.N. (1986):** Polymicrobial bacterascites. *Arch. Intern. Med*, 146: 2173–2175.
- Saez-Royuela, F.; Gomez, F.; Fernandez, M. et al. (1990):** A microbiological and clinical study of 57 cases of spontaneous bacterial peritonitis in liver cirrhosis patients. *Rev. Clin. Exp.* 187 (7): 321-324.

- Saito, N.; Takemori, N.; Hirai, K.; Onodera, R.; Watanabe, S.; Namiki, M. (1993):** Ultrastructural localization of lactoferrin in the granules other than typical secondary granules of human neutrophils. *Human Cell*, 6: 42-48.
- Sam, R.; Sahani, M.; Ulozas, E. et al. (2002):** Utility of a peritoneal dialysis leukocyte strip in the diagnosis of peritonitis. *Artif. Organs*, 26 : 546 – 548.
- Sanchez, L.; Calvo, M.; Brock, J.H. (1992):** Biological role of lactoferrin *Archives of Disease in Childhood*, 67: 657-661.
- Sanders, R. (1998):** Basic physics. In: *Clinical sonography (A practical Guide)*, Sanders RC, (Ed), Philadelphia: Lippincott, P: 2-7.
- Sax, H.C.; Ryan, C.K. and Hardy, D.J. (1996):** Low dose enteral feeding is beneficial during total parenteral nutrition. *Am. J. Surg.* 171: 587 – 590.
- Schryvers, A.B.; Bonnah, R.; Yu, R.H.; Wong, H.; Retzer, M. (1998):** Bacterial lactoferrin receptors. *Advances in Experimental Medicine and Biology*, 443: 123–133.
- Scott, P.H. (1989):** Enzyme immunoassay of lactoferrin in newborn term infants: reference values and influence of diet. *Annals of Clinical Biochemistry*, 26: 407-411.
- Senger, D.R.; Galli, S.J.; Dvorak, A.M. et al. (1983):** Tumour cells secrete a vascular permeability factor that promotes accumulation of ascites fluid. *Science*, 219: 983 – 985.
- Sherlock, S. and Dooley, J. (2002):** Ascites in diseases of the liver and biliary system. 11th edition. Black well scientific publications, P: 127.

- Sherman, M.P.; Bennett, S.H.; Hwang, F.F.; Yu, C. (2004):** Neonatal small bowel epithelia: enhancing anti-bacterial defense with lactoferrin and Lactobacillus GG . *Biometals*, 17: 285–289.
- Sheron, N.; Bird, G.; Koskinas, J. et al. (1993):** Circulating and tissue levels of the neutrophil chemotaxin interleukin -8 are elevated in severe acute alcoholic hepatitis and tissue levels correlate with neutrophil infiltration. *Hepatology*, 18: 41 – 45.
- Silvain, C.; Breux, J.P.; Grollier, G. et al. (1989) :** Les septicemies et les infections du liquide d'ascite du cirrhotique peuvent – elles etre traitees exclusivement par voie orale ? *Gastro enterol. Clin. Biol.*, 13 : 335 -9.
- Singh, N.; Gayowski, T.; Yu, V.L. et al. (1995):** Trimethoprim-sulphamethoxazole for the prevention of spontaneous bacterial peritonitis in cirrhosis: a randomized trial. *Ann. Intern. Med.* 122: 595-598.
- Sinkora, J.; Samankova, P.; Kummer, V.; Leva, L.; Maskova, J.; Rehakova, Z.; Faldyna, M. (2007):** Commercially available rabbit anti-human polyclonal antisera as a useful tool for immune system studies in veterinary species. *Veterinary Immunology and Immunopathology*, 119: 156–162.
- Skolinck, M. (1986):** Ascites search. In: *Guide to the ultrasound Examination of the Abdomen*. Skolinck ML (ed). New York: Springer – Uerlog, P: 11-17.
- Smith, C.R. (1987):** The adverse effects of flouroquinolones. *J. Antimicrob. Chemother.* 20: 709 – 711.

- Smith, J.A.; Bryce, E.A.; Nugi-Yen, J.H. and Robert, F.J. (1995):** Comparison of bactec 9240 and Bact/ALERT blood culture systems in an adult hospital. J. clin. Microbiol. 33: 1905-1908.
- Sorell, W.T.; Quigley, E.M.M.; Jin, G. et al. (1993):** Bacterial translocation in the portal hypertensive rat: studies in basal conditions and on exposure to hemorrhagic shock. Gastroenterology, 104: 1722-1726.
- Sorensen, M.; Sorensen, S.P.L. (1939):** The proteins in whey. Rendus des Travaux du Laboratoire Carlsberg, 23: 55–99.
- Soriano, C.; Coll, P.; Guarner, C. et al. (1995):** Escherichia coli capsular polysaccharide and spontaneous bacterial peritonitis in cirrhosis. Hepatology, 21: 668-673.
- Soriano, G.; Guarner, C.; Tomas, A. et al. (1992):** Norfloxacin prevents bacterial infection in cirrhotics with gastrointestinal hemorrhage. Gastroenterology, 103: 1267-1272.
- Sort, P.; Navasa, M.; Arroyo, V. et al. (1999):** Effect of intravenous albumin renal impairment and mortality in patients with cirrhosis and spontaneous bacterial peritonitis. N. Engl. J. Med. 341: 403-409.
- Such, J. and Runyon, B.A. (1998):** Spontaneous bacterial peritonitis. Clin. Infect. Dis, 27 : 669.
- Such, J.; Hillebrand, D.J.; Guarner, C. et al. (2001):** Tumour necrosis factor – alpha, interleukin -6 and nitric oxide in sterile ascitic fluid and serum from patients who subsequently develop ascitic fluid infection. Dig. Dis. Sci. 46: 2360 -2366.

- Such, J.; Hillebrand, D.J.; Guarner, C. et al. (2004):** Nitric oxide in ascitic fluid is an independent predictor of renal impairment in patients with cirrhosis and spontaneous bacterial peritonitis Eur. J. Gastroenterol. Hepatol. 16(6): 571-577.
- Suzuki, Y.A. and Lonnerdal, B. (2002):** Characterization of mammalian receptors for lactoferrin. Biochemistry and Cell Biology, 80: 75–80.
- Suzuki. Y.A.; Lopez, V.; Lonnerdal, B. (2005):** Mammalian lactoferrin receptors: structure and function. Cellular and Molecular Life Sciences, 62: 2560–2575.
- Sykes, J.A.; Thomas, M.J.; Goldie, D.J.; Turner, G.M. (1982):** Plasma lactoferrin levels in pregnancy and cystic fibrosis. Clinica Chimica Acta, 122: 385–393.
- Tachezy, J.; Kulda, J.; Bahnikova, I.; Suchan, P.; Razga, J.; Schrevel, J. (1996):** Tritrichomonas foetus: iron acquisition from lactoferrin and transferrin. Experimental Parasitology, 83: 216–228.
- Teng, C.T.; Beard, C.; Gladwell, W. (2002):** Differential expression and estrogen response of lactoferrin gene in the female reproductive tract of mouse, rat and hamster. Biology of Reproduction, 67: 1439–1449.
- Terg, R.; Cobas, S.; Fassio, E.; et al. (2000):** Oral ciprofloxacin after a short course of intravenous ciprofloxacin in the treatment of spontaneous bacterial peritonitis: results of a multicenter, randomized study. J. Hepatol. 33: 564 – 569.
- Thevenot, T.; Cadranet, J.F.; Nguyen–Khac, E. et al. (2004):** Diagnosis of spontaneous bacterial peritonitis in cirrhotic patients by use of two reagent strips, Eur. J.Gastro enterol Hepatol. Jun. 16 (16):579–583.

- Thorpe, T.C.; Wilson, M.L.; Turner, J.E. et al. (1990):** Bact / ALERT: an automated colorimetric microbial detection system. J.clin. Microbiol. 28: 1608–1612.
- Thuluvath, P.J.; Morss, S. and Thompson, R. (2001):** Spontaneous bacterial peritonitis in hospital mortality, predictors of survival and health care costs from 1988 to 1998. Am. J. Gastroenterol. 96: 1232-1236.
- Tito, L.; Rimola, A.; Gines, P. et al. (1988):** Recurrence of spontaneous bacterial peritonitis in cirrhosis: frequency and predictive factors. Hepatology, 8: 27–31.
- Toledo, C.; Salmeron, J.M.; Rimola, A. et al. (1993):** Spontaneous bacterial peritonitis in cirrhosis: predictive factors of infection resolution and survival in patients treated with cefotaxime. Hepatology, 17: 251–257.
- Topley, N.; and Williams, T. (1994):** Role of peritoneal membrane in the control of inflammation in peritoneal cavity. Kidney Int.46: 571 – 578.
- Torun, S.; Dolar, E.; Yilmaz, Y. et al. (2007):** Evaluation of leukocyte estrase and nitrite strip tests to detect spontaneous bacterial peritonitis in cirrhotic patients. World J. Gastroenterol.13: 6027-6030.
- Valenti, P. and Antonini, G. (2005):** Lactoferrin: an important host defense against microbial and viral attack. Cellular and Molecular Life Sciences, 62: 2576–2587.

- Vanbiervliet, G.; Rakotoarisoa, C.; Filippit, et al. (2002):** Diagnostic accuracy of a rapid urine–screening test (Multistix 8 SG) in cirrhotic patients with spontaneous bacterial peritonitis. *European J. of Gastroenterology & Hepatology*, Vol 14: No 11.
- Van Snick, J.L.; Masson, P.L.; Heremans, J.F. (1974):** The involvement of lactoferrin in the hyposideremia of acute inflammation. *The Journal of Experimental Medicine*, 140: 1068–1084.
- Van-Thiel, D.H.; Hassanein, T.; Gurakar, A. et al. (1996):** Liver transplantation after an acute episode of spontaneous bacterial peritonitis. *Hepato- gastroenterol.* 43: 1584 – 1588.
- Van Vugt, H.; Van Gool, J.; Ladiges, N.C.; Boers, W. (1975):**
Lactoferrin in rabbit bile: its relation to iron metabolism. *Quarterly Journal of Experimental Physiology and Cognate Medical Sciences*, 60: 79–88.
- Walmer, D.K.; Wrona, M.A.; Hughes, C.L.; Nelson, K.G. (1992):** Lactoferrin expression in the mouse reproductive tract during the natural estrous cycle: Correlation with circulating estradiol and progesterone. *Endocrinology*, 131:1458–1466.
- Wallersted T. S.; Olsson, R.; Simern, M. et al. (2007):** Abdominal tenderness in ascites patients indicates spontaneous bacterial peritonitis. *Eur. J. Internal medicine*, 18: 44-47.
- Wang, W.P.; Iigo, M.; Sato, J.; Sekine, K.; Adachi, I.; Tsuda, H. (2000):** Activation of intestinal mucosal immunity in tumor-bearing mice by lactoferrin. *Japanese Journal of Cancer Research*, 91: 1022–1027.

- Ward, P.P.; Zhou, X.; Conneely, O.M. (1996):** Cooperative interactions between the amino- a carboxyl-terminal lobes contribute to the unique iron-binding stability.
- Webster, S.T.; Brown, K.L.; Lucey, M.R. and Nostrant, T.T. (1996):** Haemorrhagic complications of large volume abdominal paracentesis. *Am.J. Gastroenterol.* 92: 366-368.
- Wells, CL.; Maddaus, MA.; Simmons, RL.; Marston, A.; Bulkley, GB.; Fiddian-Green, RG.; Haglund, UH. (1989):** Bacterial translocation in Splanchnic ischemia and multiple organ failure. Mosby: St. Louis, 195-204.
- Wofson, J. and Hooper, D. (1988):** Norfloxacin, a new targeted flouro-quinolone anti-microbial agent. *Ann. Intern. Med.* 108: 238– 251.
- Zeni, F.; Tardy B.; Vindimian, M. et al. (1993):** High levels of tumour necrosis factor- alpha and interleukin- 6 in the ascitic fluid of cirrhotic patients with spontaneous bacterial peritonitis. *Clin. Infect. Dis.* 17 (2): 218 –223.